A simple genetic architecture and low constraint allows rapid floral evolution in a

diverse and recently radiating plant genus

Jamie L. Kostyun[1,2*], Matthew J.S. Gibson[1], Christian M. King[1,3], and Leonie C. Moyle[1]

[1] Department of Biology, Indiana University, Bloomington, IN 47405

[2] Department of Plant Biology, The University of Vermont, Burlington, VT 05405

[3] Department of Evolution, Ecology and Organismal Biology, The Ohio State University,

Columbus, OH 43210

* Corresponding author: Jamie.Kostyun@uvm.edu, Tel: 802-656-7670

Word count: Summary (200), Introduction (1185), Materials and Methods

(1576), Results (1010), Discussion (1821), Acknowledgements (79), Total Main

Text (5706), Figures (4, all in color), Tables (2), Supplementary Documents (8

Figures, 7 Tables).

Summary

• Genetic correlations among different components of phenotypes, especially

resulting from pleiotropy, can constrain (antagonistic) or facilitate (adaptive) trait

evolution. These factors could especially influence the evolution of traits that are

functionally integrated, such as those comprising the flower. Indeed, pleiotropy is

proposed as a main driver of repeated convergent trait transitions, including the

evolution of phenotypically-similar pollinator syndromes.

• We assessed the role of pleiotropy in the differentiation of floral and other

reproductive traits between two species—*Jaltomata sinuosa* and *J. umbellata*

(Solanaceae)—that have divergent suites of floral traits consistent with bee- and

hummingbird-pollination, respectively. To do so, we generated a hybrid

population and examined the genetic architecture (trait segregation and QTL

distribution) underlying 25 floral and fertility traits.

• We found that most traits had a relatively simple genetic basis (few,

predominantly additive, QTL of moderate to large effect), as well as little

evidence of antagonistic pleiotropy (few trait correlations and QTL co-

localization, particularly between traits of different classes). However, we did

detect a potential case of adaptive pleiotropy among floral size and nectar traits.

• These mechanisms may have facilitated the rapid floral trait evolution observed

within *Jaltomata,* and may be a common component of rapid phenotypic change

more broadly.

Introduction

One feature of phenotypic evolution is the broad variation in observed rates of trait change, both among traits within a single lineage and among lineages. How quickly phenotypes evolve and in what direction, depends not only on their genetic basis--both the number and effect size of causal loci--and the intensity and nature of selection acting upon them, but also on their associations with other traits. Because of genetic covariance, developmental and phylogenetic constraints, and/or correlated selection pressures (Agrawal and Stinchombe 2009), phenotypic traits often do not evolve independently from one another. Among these causal mechanisms, strong genetic covariance arises either because different traits have a shared genetic basis (pleiotropy) or because they are based on genes that are physically adjacent and therefore often co-inherited (via linkage). Strong pleiotropy is often proposed to constrain phenotypic evolution by preventing correlated traits from moving efficiently towards their own (different) fitness optima (antagonistic pleiotropy). However, such shared genetic control may also promote phenotypic change if covariation is aligned in the direction of selection (adaptive pleiotropy) (Agrawal and Stinchombe 2009; Wagner and Zhang 2011; Smith 2016). Nonetheless, despite some detailed studies (e.g. Ettensohn 2013; Shrestha et al. 2014; Manceau et al. 2011; Xu and Schluter 2015), the genetic architecture of many ecologically important traits remains unclear, including the prevalence of strong genetic associations that could shape the course of phenotypic evolution.

How shared architecture influences phenotypic change is especially relevant for suites of traits that are functionally integrated (Armbruster et al. 2014) -- such as the angiosperm flower (Armbruster et al. 2009) -- because the magnitude and direction of

pleiotropy will directly shape if and how these co-varying traits respond to selection.

Further, because pleiotropy can constrain or favor particular developmental trajectories, it

is often proposed to be a main driver of convergent transitions of integrated traits in

different lineages (Preston et al. 2011; Smith 2016). The flower is an especially

promising model for assessing the role of pleiotropy in shaping phenotypic evolution.

Because flowers mediate fitness through their critical reproductive role, their constituent

traits (i.e. reproductive structures, perianth, and other attraction/reward features) are often

highly functionally integrated (Conner 2002; Armbruster et al. 2009). Moreover, repeated

transitions of phenotypically similar or convergent suites of floral traits have been

identified both within and across groups (Fenster et al. 2004; Goodwillie et al. 2010;

Wessinger et al. 2016). For example, multiple parallel shifts from bee-pollination to

hummingbird-pollination are associated with parallel transitions to flowers with red

petals, large amounts of dilute nectar, and narrow corolla tubes within *Penstemon*

(Wessinger et al. 2016); similarly, the evolution of the 'selfing syndrome' (i.e. reduced

overall size, herkogamy, and floral rewards) often accompanies transitions from

outcrossing to predominantly selfing mating systems, and has been documented in

multiple lineages (Stebbins 1974; Goodwillie et al. 2010). Such patterns provide an

opportunity to assess the relative frequency of adaptive vs. antagonistic pleiotropy in

shaping these repeated trait combinations, within a comparative phylogenetic context.

In addition to these ecological and evolutionary features, the known genetic and

molecular bases of floral development (Rijpkema et al. 2006; Smaczniak et al. 2012)

themselves suggest that pleiotropy might be a key component shaping floral phenotypic

change, as well as provide a functionally-informed framework for identifying how

changes in these mechanisms can contribute to the diversification of floral traits. Under

the ABC(DE) model of floral development, the combinatorial action of different gene

products -- primarily different MADS-box transcription factors -- control transitions to

flowering and the specification of floral organ identities and organ maturation, by

promoting or repressing different downstream targets (reviewed in O'Maoileidigh et al.

2014; Bartlett 2017). This combinatorial function, and the ability to regulate shared or

partially shared downstream targets, provides a potential mechanistic explanation for

strong pleiotropy among floral traits. Moreover, several key regulators of floral

development also function during fruit and seed production (Smaczniak et al. 2012;

O'Maoileidigh et al. 2014); such correlated effects on fertility traits are another potential

way that pleiotropy could shape floral phenotypic evolution.

Several empirical approaches have been taken to assess the genetic architecture

underlying floral trait specification and evolution, and to evaluate the strength and

direction of pleiotropy. Classical quantitative genetic analyses have revealed varying

degrees of genetic covariance among floral traits (Gottlieb 1984; Conner et al. 2014),

while numerous QTL (quantitative trait locus) mapping studies have found that loci for at

least some different floral traits appear to co-localize to the same genomic region(s)

(reviewed in Smith 2016). Interestingly, such studies have identified more putative cases

of adaptive pleiotropy (e.g. QTL affect more than one trait in the direction of parental

trait values) than antagonistic, suggesting that adaptive pleiotropy may be a common

mechanism contributing to rapid floral evolution. Because QTL generally span a genomic

region that contains more than one gene, such QTL co-localization is consistent with, but

not definitive evidence of, a role of pleiotropy in shaping floral trait co-variation (e.g. see

Hermann et al. 2013). Nonetheless, identifying strong trait correlations and/or QTL co-

localization represents a critical step in assessing how pervasive pleiotropy could be in

shaping floral trait evolution.

In this study, we examined phenotypic (co)variation among, and the genetic

architecture underlying, floral and other reproductive traits within a segregating hybrid

population derived from two *Jaltomata* (Solanaceae) species with divergent floral traits

(**Figure 1**). Despite only having diversified with the last 5 million years (Sarkinen et al.

2013; Wu et al. 2018), species of *Jaltomata* are highly phenotypically diverse, including

extensive variation in the size, shape, and color of floral traits that is absent among their

close relatives within *Solanum* and *Capsicum*. Indeed, phylogenetic analyses suggest

numerous transitions in floral traits within the genus, including several instances of

convergent evolution (Miller et al. 2011; Wu et al. 2018). Importantly, many of these

transitions appear to involve parallel changes in several different traits within a lineage

(e.g. from flat corollas with small amounts of lightly colored nectar to highly fused

corollas with large amounts of darkly colored nectar), suggesting either a shared genetic

basis and/or correlated selection (perhaps pollinator-mediated selection, Fenster et al.

2004) influences these trait associations.

Here, we identified QTL contributing to floral and other reproductive trait

variation within a recombinant population to: 1) examine the genetic architecture

underlying reproductive trait divergence; 2) assess the role of genetic linkage and/or

pleiotropy (via strong trait correlations and overlapping QTL) in floral trait (co)variation;

and 3) assess evidence for a shared genetic basis between different classes of floral trait

and other reproductive (specifically fertility) traits. We found evidence for a relatively

simple genetic basis underlying most of the examined traits, as well as positive

correlations and significant QTL co-localization among traits within each of three trait

classes (floral morphology, floral color, and fertility). Together, these features might

facilitate rapid changes in these traits. In comparison, we found few associations between

traits from different classes, and therefore little evidence for antagonistic pleiotropy

among these classes. One striking exception was an association between flower size and

nectar traits that acts in the direction exhibited by multiple species in the genus,

suggesting that this could instead be an instance of adaptive pleiotropy.

MATERIALS AND METHODS

Study system: *Jaltomata* (Schlechtendal; Solanaceae) is the sister genus to the

large and economically important *Solanum* (Olmstead et al. 2008; Sarkinen et al. 2013;

Wu et al. 2019), and includes approximately 60-80 species distributed from the

Southwestern United States to the Andean region of South America, in addition to several

species endemic to the Greater Antilles and the Galapagos Islands (Miller et al. 2011;

Mione et al. 2015). Species are highly phenotypically diverse and live in a variety of

habitats (e.g. tropical rainforests, rocky foothills, and *lomas* formations), despite their

recent divergence (<5MYA, Sarkinen et al. 2013).

Here, we focused on a closely related species pair, *J. sinuosa* and *J. umbellata*,

that differ in a suite of floral traits that is representative of major floral suites found in

other species throughout the genus (**Figure 1**; **Table S1**). *Jaltomata sinuosa* has large

rotate flowers with purple petals and a small amount of concentrated nectar (consistent

with bee pollination, Fenster et al. 2004), while *J. umbellata* has small short-tubular

flowers with white petals and a large amount of dilute, but dark red, nectar that is visible

through the corolla tube (consistent with hummingbird pollination, Fenster et al. 2004;

Kostyun and Moyle 2017; Mione et al. 2017). *Jaltomata sinuosa* is distributed along the

Andes in South America from Venezuela to Bolivia, while *J. umbellata* is restricted to

lomas formations along the Peruvian coast. Both species are self-compatible (Kostyun

and Moyle 2017; J.L. Kostyun and T. Mione, unpub.) and shrubby, but differ in leaf traits

such as overall size and shape, and type of trichomes. This species pair also has several

incomplete, intrinsic postzygotic reproductive barriers, including quantitatively reduced

fruit set and hybrid seed viability (Kostyun and Moyle 2017).

Generation of BC$_1$ population and plant cultivation: We developed a

segregating hybrid population by crossing *J. sinuosa* and *J. umbellata*. Viable F1

individuals in both directions of the cross produce flowers that are phenotypically

intermediate between the parental genotypes (**Figure 1**), and retain reduced but sufficient

levels of fertility when back-crossed to parents (Kostyun and Moyle 2017). Because we

were especially interested in the genetic basis of red nectar and a fused corolla tube

(exhibited by *J. umbellata* and F1s but not *J. sinuosa*), we generated the mapping

population by backcrossing a single *J. sinuosa* x *J. umbellata* F1 (as the ovule parent) to

the original parental *J. sinuosa* individual (as the pollen parent). BC1 individuals were

germinated in a growth chamber, and then moved to the Indiana University greenhouse

and grown under the same conditions as the parental and F1 individuals (16 hour light

cycle, watered twice daily, and fertilized weekly).

Trait measurements: We measured 25 floral and other reproductive traits within

our mapping population, F1, and parental genotypes (**Table 1; Table S2; Figure S1**).

Floral morphological traits were measured with hand-held calipers, and nectar volume

per flower was measured to the nearest 1 μL using a pipette. To reduce the potential

effect of daily environmental variation, nectar volume was always measured in the early

afternoon following watering.

Petal and nectar color were quantified using digital photography (Kendal et al.

2013; Garcia et al. 2014): dissected petals and nectar drops were photographed on a

standard background along with white and black color standards. Light conditions were

standardized for all images using RAW Therapee (RAW Therapee Development Team

2012), and color space attributes were measured in ImageJ (Schneider et al. 2012).

Because RGB color attributes are device-dependent (i.e. they can vary depending upon

the specific camera used), color values were also converted into device-independent

L*a*b color attributes, using the ImageJ Color Space Converter plugin (Schwartzwald

2012). For both petal and nectar color, this approach produced eight interrelated color

attributes: Intensity, Red, Green, Blue, Composite RGB, Lightness (L), 'a' color (ranges

from green to magenta), and 'b' color (ranges from cyan to yellow). Broadly, Intensity,

Red, Green, Blue, and Lightness convey information about color brightness, while

Composite RGB, 'a' color, and 'b' color convey information about hue.

We also measured nine fertility traits (**Table 1**) to assess the potential genetic

overlap between floral and other reproductive traits, as well as to examine the genetic

architecture underlying intrinsic postzygotic barriers between this species pair. Fruit and

seed related traits were measured on 2-6 crossed fruit per individual (depending on fruit

set). For F1 and BC1 individuals, crossed fruit were produced using pollen from the *J.*

sinuosa parental individual. To determine seed germination rates (following Farooq et al.

2005), we soaked 10 seeds per individual in 50% bleach for 30 minutes (to soften the

seed coat), rinsed thoroughly, and placed on moist paper within plastic germination

boxes. A week after sowing, seed coats were nicked slightly and seeds were given a drop

of 10 mM giberellic acid (Sigma) to break dormancy. We then scored germination every

2 weeks for 4 months. Pollen viability was estimated from three different flowers per

individual, using established methods (Jewell et al. 2012): Briefly, for each sample all

undehisced anthers from a flower were collected into an eppendorf tube containing

aniline blue histochemical stain, gently ground with a pestle to release pollen, and viable

pollen grains were counted under an EVOS FL Digital Inverted Fluorescence Microscope

(Fisher Scientific). From these data, we also calculated the proportion of viable pollen as

number of viable pollen grains/total pollen grains in the sample.

Statistical analyses on phenotypic data: Following Shapiro-Wilk tests to assess

normality assumptions, we transformed traits that showed a skewed distribution and/or

significantly non-normal residuals. In particular, we arcsine transformed proportional

traits (proportion of corolla fusion, fruit set, and proportion of viable pollen), and log-

transformed inflorescence size, corolla fusion, petal length, style length, herkogamy,

nectar volume, all color attributes, and remaining fertility traits. Significant differences

between parental species for all traits were assessed by t-tests (**Table 1**). Distribution

plots for all traits are provided in Supplementary Materials (**Figures S2-S6**), while

illustrative plots for 15 focal traits are provided in the main text (**Figure 2**). Similarly,

phenotypic correlations within the BC1 population were examined among all traits

(**Table S3**), while relationships among a subset of focal traits are presented in the main

text (**Figure 3**). Given significant correlations among many of the floral morphological

traits and among color attributes (**Table S3**), we also used principle component analyses

(PCAs) to create separate composite metrics (principle components (PCs)) for three

groups of traits (i.e. Morph PC1-PC3; Nectar Color PC1-3; Petal Color PC1-PC3; **Table**

S4) as additional measures of floral variation. Trait correlations, including PCs, are

provided in **Table S5**, and distribution plots are provided in **Figures S3-S5**.

237   ***Genotyping and linkage map construction:*** Genomic DNA was extracted from

young leaf tissue from the 2 parental individuals, 13 F1s (including the F1 parent used to

generate the population), and 269 BC1s, using the Qiagen DNeasy Plant Mini Kit. DNA

quantity and quality were confirmed via Nanodrop (Fisher Scientific) and gel

electrophoresis with λ DNA-HindIII Digest marker (New England BioLabs). Samples

were then sent to Novogene Corporation (Beijing) for genotyping-by-sequencing (GBS).

GBS libraries were prepared using optimized restriction enzymes (MseI and HaeIII), and

following insert size selection, sequenced on an Illumina Hi-Seq to generate 150bp paired

end reads. Raw reads were trimmed and filtered using Trimmomatic (Bolger et al. 2014),

and read quality was checked pre- and post-trimming using fastqc (Andrews 2010). To

identify SNPs, cleaned reads were mapped to the domesticated tomato genome (Tomato

Genome Consortium 2012) using the mem function in BWA (Li 2013). Alignment files

were then input into the STACKS refmap pipeline (Catchen et al. 2013) to determine

genotypes. Reads and genotype data are available in NCBI SRA XXXXXX.

251   To construct the linkage map, we first removed markers that were genotyped in

less than 35% of individuals, or showed significant segregation distortion (i.e. alleles at

>80% or <20% frequency). The linkage map was constructed using the MST and

Kosambi algorithms, implemented in the R package ASMap (Taylor and Butler 2017).

To alleviate map expansion issues, we removed markers which consistently differed from

neighboring markers in terms of genotype assignment, indicating a high likelihood of

genotyping error. The linkage map was then finalized using the ripple function in R

package R/qtl (Broman et al. 2003).

Identifying QTL: We implemented Haley-Knott regression in R/qtl to identify

QTL contributing to each trait. To account for potential environmental contributions to

trait variation, we included date of measurement (Month) and location within the

greenhouse (Bench) as covariates in our QTL scans. Putative QTL were first identified

using the scanone function, followed by permutations for genome-wide LOD significance

thresholds. Two dimensional scans (scantwo function) were used in the stepwise qtl

function to fit multiple QTL models. These models were used to identify significant

QTL, their 1.5 LOD confidence intervals, their effect sizes (i.e. difference in phenotype

mean between homozygotes and heterozygotes), and the total amount of phenotypic

variance explained by each QTL, as well as interactions among QTL, and potential

contributions of covariates. QTL were considered to be co-localized if their 1.5 LOD

intervals overlapped. Significant co-localization was assessed by comparing overlap

among identified QTL to overlap from 10000 randomly generated distributions, for traits

within each category (morphological, color/physiological, or fertility), and between each

trait category. Briefly, a custom Python script was used to generate random distributions

of QTL (by randomly re-distributing the identified QTL among the 12 linkage groups),

and the observed frequency of co-localization in each was recorded for each

randomization to generated count distributions, in R. All code used to generate the

linkage map, identify QTL, and assess QTL co-location, is available on GitHub

(https://github.com/gibsonMatt/jaltomataQTL).

RESULTS

Segregation patterns suggest additive alleles underlie most traits

282   Most traits were significantly different between the two parental species (**Table**

1). F1 means were intermediate for most traits as well, except that petals were generally

brighter (more white) than either parent (**Figures S5+S7**). Other than fruit set and seed

germination rates, all traits were unimodally distributed within the BC1s; phenotypic

values were intermediate between F1s and the recurrent parent (*J. sinuosa*) for many of

these traits, consistent with additive effects (**Figure 2; Figures S2-S6**). Several traits (7

of 25) showed transgressive segregation within the BC1s, including some floral

morphological traits, nectar volume and color, and seed viability and germination rates

(**Table S2; Figures S2-S6**).

Significant correlations observed within – but generally not between – floral

morphology, floral color, and fertility trait categories

294   Within the BC1s, most trait combinations were not strongly associated.

Nonetheless, several correlations remained significant following multiple testing

(Bonferroni) correction (**Table S3**), primarily associations that are expected biologically,

including allometric relationships among floral organs and positive associations among

related fertility traits. For instance, corolla diameter was significantly positively

associated with most other morphological traits, suggesting shared genetic control of

overall floral size (**Figure 3**), while corolla diameter was also significantly negatively

correlated with proportion of corolla fusion (i.e. shorter corolla tubes had wider limbs and

longer tubes had narrower limbs, r = -0.348, $p = 8.68e^{-8}$). These relationships were

recovered with PCA on morphology traits, in which PC1-PC3 explained 76% of the

variance among BC1s (**Table S4**). Based on trait loadings, PC1 corresponds to floral

width vs. depth, PC2 to overall floral size, and PC3 to relative reproductive organ

dimensions. Similarly, related fertility traits also remained strongly correlated, such as

fruit mass with seed set, and number of viable pollen grains with proportion of viable

pollen (**Table S3**). Finally, color attributes within each of nectar color and petal color

were strongly correlated with one another, but these attributes were not associated

between nectar and petals (**Table S3**). From PCAs on nectar color and petal color

attributes (separately), PC1-PC3 for each explained 97% and 94% of the variance among

BC1s, respectively (**Table S4**).

In contrast, there were relatively few significant correlations among different trait

categories. Notable exceptions, however, included a positive relationship between floral

size and nectar volume as well as floral size and certain aspects of nectar color (**Figure 3;**

Table S3). There were also significant positive correlations between pollen viability and

each of several components of flower size (as well as Morph PC2 or "size") (**Tables**

S3+S5). This latter relationship seems to be explained by anther size: across 15 *Jaltomata*

species, mean viable pollen count is significantly associated with anther size prior to

dehiscence ($F = 15.56$, $p = 0.0017$) (J.L. Kostyun, unpub.).

Linkage map construction recovered 12 linkage groups

Mapping high quality reads to the tomato genome identified 25,136 SNPs that

differentiated the two parental species. Following all subsequent filtering (removing

markers genotyped in less than 35% of individuals, with high segregation distortion or

non-Mendelian inheritance, or with high likelihood of genotyping errors), we retained

520 high quality markers. Linkage map construction recovered 12 linkage groups (LGs),

which correspond to the number of chromosomes in the parental species (Mione et al.

1993; Chiarini et al. 2017). Based on orthology with tomato, we were able to confidently

assign 5 of these LGs to chromosomes (**Figure 4**). Total map length was 1593.71 cM

(65.17 cM – 324.48 cM per chromosome/LG), with an average of 2.92 cM between

markers (**Figure 4**).

Few moderate-effect QTL underlie most traits, with little QTL co-localization between

trait categories

We identified a total of 63 QTL for our 25 traits (with 4 additional loci for PC

traits). Most traits had 2-4 QTL, with a range of 0-6 QTL (**Table 2**, **Table S6**). Alleles at

55 of 67 QTL (82%) acted in the direction consistent with parental values (i.e. the allele

from paternal donor *J. umbellata* moved the phenotype of BC1s closer to its species

mean) (**Table 2**), and the amount of phenotypic variation explained per QTL ranged from

2-28%. The latter range suggests that we had reasonable power to identify QTL with

even relatively small effects--explaining as little as 2% of the variance. Consistent with

observed trait segregation patterns, significant interactions among QTL within individual

traits were identified in few cases: for ovary diameter and certain nectar color attributes

(**Table 2**, **Table S6**).

Although every linkage group had at least one QTL, QTL were not distributed

uniformly across the genome, with notable clusters on LG1 and LG9 (**Figure 4, Table 2**).

We also identified several instances of QTL co-localization within trait categories,

especially for morphology and fertility traits, which each had significantly more cases of

co-localization (1.5 LOD overlap) than expected by chance (133 observed vs. upper

bound of 115 expected overlaps, $p = 7.5e^{-4}$; 8 observed vs. upper bound of 4 expected

overlaps, $p = 7.0e^{-6}$, respectively) (**Table S7; Figure S8**). These co-localization instances

included QTL for biologically-related traits (e.g. petal length and corolla diameter, or

fruit mass and seed set; **Table S6**), for which we already observed strong correlations

(**Table S3**). In some cases, co-localized QTL share the same or a very close peak marker

(e.g. petal length, corolla fusion, and ovary diameter on LG12; **Table S6**) which is

suggestive of potential pleiotropy, however we note that the large 1.5 LOD intervals of

some QTL will increase instances of incidental co-localization events.

In contrast, co-localization between different trait categories was never greater

than expected by chance, with co-localization between morphology and color traits

actually significantly less than expected ($p = 0.016$) (**Table S7; Figure S8**). This is

consistent with mostly incidental occurrences of overlap between QTL for traits in

different categories. Nonetheless, we did detect co-localized QTL at the same or very

close peak markers for, for example, nectar color (RGB) and volume on LG3 (**Figure 4;**

Table S6) and for nectar color (a), corolla fusion, and corolla depth on LG7, (**Table S6**),

which provide intriguing cases of potential adaptive pleiotropy (i.e. alleles at QTL that

simultaneously act to increase floral size, nectar darkness, and/or nectar volume).

DISCUSSION

Genetic correlations among different components of the phenotype, especially

resulting from pleiotropy, can constrain or facilitate trait evolution (Agrawal and

Stinchcombe 2009). Pleiotropy could have particularly strong effects on the evolution of

traits that are functionally integrated, such as those comprising the flower (Armbruster et

al. 2009; Smith 2016). To better understand the genetic architecture underlying floral trait

evolution within florally diverse *Jaltomata*, including whether pleiotropy might have

shaped observed variation, we examined patterns of genetic segregation and genetic

architecture for 25 floral and fertility traits in a hybrid (BC1) population between species

with highly divergent floral traits. We found that most of our examined traits have a

relatively simple genetic basis, with few to moderate QTL with largely additive effects.

We also identified strong correlations and significant QTL overlap within trait categories,

but few associations across different types of traits. The exceptions however, between

certain aspects of floral morphology and nectar traits, are consistent with existing trait

associations that are observed across the genus, suggesting that these could be examples

of adaptive pleiotropy. Overall, our data suggest that the rapid floral trait evolution

observed in this group could have been facilitated by a relatively simple genetic basis for

individual floral traits, and a general absence of antagonistic pleiotropy among different

types of reproductive traits, especially morphology and color.

Few genetic changes could underlie floral trait shifts

The relatively simple genetic architecture that we detect for most of our floral

traits might be one mechanism that has permitted rapid floral evolution within the genus.

Indeed, our inference that few QTL controlling corolla traits agrees with comparative

development data from these species (Kostyun et al. 2017) in which we observe that

relatively simple heterochronic changes in corolla trait growth rates distinguish these

rotate vs. tubular corolla forms. Interestingly, our findings are also consistent with

previous studies of floral trait genetics between closely related species (Smith 2016). For

instance, one or few QTL have been found for species differences in nectar volume in

several other systems (e.g. Bradshaw et al. 1998; Stuurman et al. 2004; Wessinger et al.

2014; but see Nakazato et al. 2013), similar to our inference of a single QTL for this trait.

For petal and nectar color, we identified 2 and 5 QTL, respectively, each with moderate

to major effects (**Table 2, Table S6**), similar to other systems that generally identify few

loci of large effect for petal color (e.g. Bradshaw et al. 1998; Wessinger et al. 2014).

Perhaps unlike these cases, however, it is likely that loci controlling color differences in

Jaltomata are regulators of pigment quantity rather than presence/absence biosynthesis,

because both nectar and petal color show gradation in the BC1s rather than discrete color

bins. Moreover, preliminary data from a VIGS (virus-induced gene silencing) pilot study

in *J. sinuosa* indicate that the purple petal pigment is an anthocyanin (J.L. Kostyun and

J.C. Preston, unpub.), whereas for nectar color, preliminary data suggest that an indole-

flavin contributes to red pigment in *J. umbellata* (J.L. Kostyun and D. Haak, unpub.),

consistent with our inference that color variation is unassociated between these different

floral components.

In addition to relatively few contributing loci, many of the examined traits also

appear to be underpinned by additive effects (**Table 1; Figure 3; Figures S2-6**), while

epistatic effects were comparatively rare. Both are factors that might also facilitate more

rapid responses to selection. Other studies have similarly found that floral size traits are

often additive (Gottlieb 1984). Although several floral traits showed transgressive

segregation within our BC1s, which could indicate epistatic interactions, similar patterns

can result from unique combinations of additive alleles that have opposite effects in the

parental species (e.g. deVicente and Tanksley 1993) and we identified individual QTL

with these opposing effects for many of these traits. In comparison, significant interaction

effects among morphological QTL were detected for ovary diameter only (**Table 2**),

consistent with a general lack of epistatic interactions for this class of traits.

The notable exceptions to additivity involved many of the fertility traits (as well

as some components of color, see below). BC1 individuals tended to have lower seed set

and poorer quality seeds (decreased viability and response to germination-inducing

stimuli), and the recombinant BC population contained a subset of highly sterile

individuals. The segregation of recombinant individuals with reduced viability and

fertility often occurs in hybrids (Baack et al. 2015), including in hybrids from additional

Jaltomata species pairs (Kostyun and Moyle 2017). Such patterns are typically due to

deleterious epistatic interactions between loci that have diverged between the two

parental lineages, as has been shown in close relatives including tomatoes (Moyle and

Nakazato 2008; Sherman et al. 2014). These observations in *Jaltomata* are similarly

consistent with a specific role for epistasis among incompatible alleles in the expression

of postzygotic reproductive isolation.

Reduced constraints may also have facilitated rapid floral trait evolution

Because rapid floral evolution may occur either through a lack of antagonistic

pleiotropy or through adaptive pleiotropy, we assessed evidence for these potential

mechanisms within *Jaltomata*. Within trait categories, we detected positive but modest

associations between several floral size traits, and among biologically related fertility

traits (e.g. fruit size and seed set) (**Figure 3; Table S3**), as well as significant co-

localization of QTL for these groups of traits (**Table S7**). Morphological associations in

particular suggest that shared growth regulators (e.g. Sicard and Lenhard 2011; Brock et

al. 2012) contribute to - but do not completely determine - observed variation in floral

organ sizes. In contrast, we detected fewer instances of strong trait correlations and QTL

co-localization between different trait categories (**Table S3**, **Table S7**). This general lack

of antagonistic pleiotropy among different classes of floral and fertility traits may have

facilitated rapid floral evolution in this system by minimizing constraints on the available

combinations of floral traits. Despite this general pattern, we did identify several

instances of QTL co-localization that might represent adaptive pleiotropy. Most notably,

larger flowers generally produced darker (more red) nectar as well as a greater volume of

nectar, as reflected in co-localization of QTL underlying aspects of floral morphology

and both nectar volume and color (**Table 2; Figure 4; Table S6**). Interestingly, this trait

combination (large flowers with copious dark nectar) is actually not exhibited by either

parental species used in this experiment (**Figure 1; Table 1**); however, it is found in

numerous other *Jaltomata* species (see below; Miller et al. 2011; Kostyun and Moyle

2017) and is consistent with recognized pollination syndromes (e.g. Fenster et al. 2004).

***Ecological context for rapid floral change in* Jaltomata**

460   Overall, our findings suggest potential mechanistic explanations for the evolution

of remarkable floral trait diversity among *Jaltomata* species within the last 5 million

years (Sarkinen et al. 2013). Traits with a relatively simple genetic basis that are

uncoupled from other floral and fertility traits have fewer mechanistic constraints, and

therefore can more rapidly respond to selective opportunities as they arise. Although we

have not yet directly assessed the role of selection in shaping floral differences among

species, several features of *Jaltomata* floral biology are consistent with pollinator-

mediated selection on floral traits (van der Niet and Johnson 2012), likely in conjunction

with mating-system related changes (Goodwillie et al. 2010). First, floral trait variation

within *Jaltomata* shows clear hallmarks of selection imposed by pollinator

differentiation. Nearly all species in the earliest diverging *Jaltomata* lineage have

relatively small, ancestrally rotate flowers with small amounts of lightly colored nectar,

and hymenopterans have been observed visiting several of these species (T. Mione, pers.

comm.). In contrast, many species in the South American derived clade—including *J.*

umbellata examined here--have attractive features associated with vertebrate pollination

(Fenster et al. 2004). Several of these are visited by hummingbirds (T. Mione, per.

comm.), notably those with larger flowers with a highly fused corolla (either campanulate

or tubular) and copious amounts of darkly colored nectar; intriguingly, this repeated

natural trait covariation is consistent with the genetic association between floral size and

nectar traits we identified here.

These features indicate that pollinators are a likely source of selection for floral

differentiation among species within *Jaltomata*, but they do not necessarily explain why

Jaltomata as a genus has been uniquely responsive to this pollinator variation, especially

in comparison to its most close relatives. Species from both *Solanum* and *Capsicum* are

found within the same geographical regions as *Jaltomata*, and are therefore exposed to

the same pollinator variation, but are nonetheless almost uniformly rotate and bee-

pollinated (Knapp 2010). Interestingly, one important difference between *Jaltomata* and

these two genera is in their predominant mating system. Self-incompatibility is the

ancestral state in the Solanaceae (Steinbachs and Holsinger 2002) and is broadly

persistent in both *Solanum* and *Capsicum* (Goldberg et al. 2010). In contrast, all

examined *Jaltomata* species are self-compatible (Mione 1992; J.L. Kostyun and T.

Mione, unpub.), indicating that gametophytic self-incompatibility was lost early in the

evolution of this clade. Moreover, the presence of delayed selfing and strong herkogamy

in many species (e.g. Mione et al. 2015; Mione et al. *in review*), in addition to field

observations of pollinators (above; T. Mione, pers. comm.), indicate that species most

likely employ a mixed mating strategy in their native ranges. The absence of genetically-

determined self-incompatibility and the predominance of mixed mating strategies might

have uniquely facilitated the evolution of new floral trait variation in *Jaltomata*,

compared to either *Capsicum* or *Solanum*. Mixed mating strategies are generally

observed to maintain the largest amount of floral trait variation, compared to predominant

selfing or enforced outcrossing (Goodwillie et al. 2005; Rosas-Guerrero et al. 2014). In

addition, they have been predicted to facilitate pollinator shifts--especially to pollinators

that might be more efficient but potentially unreliable (such as hummingbirds)—because

they allow reproductive assurance (via selfing, when pollinators are limited) and increase

the expression of new floral trait variation controlled by recessive alleles (Goodwillie et

al. 2005; Brys et al. 2013; Wessinger and Kelly 2018). Notably, our data indicate that

dark/red colored nectar is at least partially recessive (**Figure 2**), and that red petal

pigmentation is completely recessive (**Figure S7**), consistent with this novel variation

being based on new recessive alleles.

CONCLUSIONS

Genetic correlations among floral traits, especially those due to pleiotropic

effects, likely shape permitted trajectories of floral evolution. To assess how such genetic

associations might have contributed to observed patterns of floral diversity in *Jaltomata*,

we examined segregation patterns and genetic architecture of 25 floral and fertility traits

in a hybrid (BC1) population generated from parents with divergent floral traits. Our data

are consistent with several mechanisms that could have allowed rapid floral trait

evolution in this system: a largely simple genetic basis underlying variation in most of

our floral traits, a general absence of antagonistic pleiotropy constraining floral evolution,

and a potential instance of adaptive pleiotropy governing floral size and nectar traits. This

genetic architecture, in combination with pollinator-mediated selection on a background

of self-compatible mixed mating, might have uniquely positioned this genus for the rapid

floral diversification now evident within *Jaltomata*.

ACKNOWLEDGEMENTS

We thank the IU greenhouse staff for plant care, CJ Jewell and David Haak for

logistical support, undergraduate research assistants (especially Meret Thomas-Huebner,

Devki Shukla, and Shachia Jackson) for assistance with data collection, and Tim Leslie

for assistance with simulations. This work was supported by the IU Biology Department,

National Science Foundation Award (NSF DEB 1136707) to LCM, and National Science

Foundation Graduate Research Fellowship Program (NSF DEB 1342962) and Doctoral

Dissertation Improvement Grant (NSF DEB 1601078) to JLK.

AUTHOR CONTRIBUTIONS

534   JLK and LCM designed the experiment, JLK generated experimental materials,

535   JLK and CMK collected phenotypic data, JLK and MJSG analyzed the data, and JLK and

536   LCM wrote the paper with input from CMK and MJSG.

REFERENCES

Agrawal, A. F., and J. R. Stinchcombe. 2009. How much do genetic covariances alter the

rate of adaptation? *Proc R Soc Lond Biol* **276**:1183-1191.

Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data.

Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Armbruster, W. S., T. F. Hansen, C. Pelabon, R. Perez-Barrales, and J. Maad. 2009. The

adaptive accuracy of flowers: measurement and microevolutionary patterns. *Ann*

Bot **103**:1529-1545.

Armbruster, W. S., C. Pelabon, G. H. Bolstad, and T. F. Hansen. 2014. Integrated

phenotypes: understanding trait covariation in plants and animals. *Proc R Soc*

Lond Biol **369**.

Baack, E., M. C. Melo, L. H. Rieseberg, and D. Ortiz-Barrientos. 2015. The origins of

reproductive isolation in plants. *New Phytol* **207**:968-984.

Bartlett, M. E. 2017. Changing MADS-Box Transcription Factor Protein–Protein

Interactions as a Mechanism for Generating Floral Morphological Diversity.

Integrative and Comparative Biology **57**:1312-1321.

Bradshaw, H. D., K. G. Otto, B. E. Frewen, J. K. McKay, and D. W. Schemske. 1998.

Quantitative trait loci affecting differences in floral morphology between two

species of monkeyflower (*Mimulus*). *Genetics* **149**:367-382.

Brock, M. T., P. X. Kover, and C. Weinig. 2012. Natural variation in GA1 associates

with floral morphology in *Arabidopsis thaliana*. *New Phytol* **195**:58-70.

Broman, K. W., H. Wu, S. Sen, and G. A. Churchill. 2003. R/qtl: QTL mapping in

experimental crosses. *Bioinformatics* 19:889-890.

Brys, R., B. Geens, T. Beeckman, and H. Jacquemyn. 2013. Differences in dichogamy

and herkogamy contribute to higher selfing in contrasting environments in the

annual *Blackstonia perfoliata* (Gentianaceae). *Ann Bot* **111**:651-661.

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: A flexible trimmer for

Illumina Sequence Data. *Bioinformatics*, btu170.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. Stacks:

an analysis tool set for population genomics. *Mol Ecol* 22: 3124-3140.

Chiarini, F. E., D. Lipari, G. E. Barboza, and S. Knapp. 2017. In: IAPT/IOPB

chromosome data 25. K. Marhold and J. Kuccera (eds). *Taxon* 66: 1246–1252.

Conner, J. K. 2002. Genetic mechanisms of floral trait correlations in a natural

population. *Nature* **420**:407-410.

Conner, J. K., I. A. Cooper, R. J. La Rosa, S. G. Perez, and A. M. Royer. 2014. Patterns

of phenotypic correlations among morphological traits across plants and animals.

Phil. Trans. R. Soc. B **369**.

deVicente, M. C., and S. D. Tanksley. 1993. QTL analysis of transgressive segregation in

an interspecific tomato cross. *Genetics* **134**:585-596.

Ettensohn, C. A. 2013. Encoding anatomy: Developmental gene regulatory networks and

morphogenesis. *Genesis* **51**:383-409.

Farooq, M., S. M. A. Basra, N. Ahmad, and K. Hafeez. 2005. Thermal hardening: A new

seed vigor enhancement tool in rice. *J Integr Plant Biol* **47**:187-193.

Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004.

Pollination syndromes and floral specialization. *Annu Rev Ecol Evol Syst* **35**:375-

403.

Garcia, J. E., A. D. Greentree, M. Shrestha, A. Dorin, and A. G. Dyer. 2014. Flower

Colours through the Lens: Quantitative Measurement with Visible and Ultraviolet

Digital Photography. *Plos One* **9**.

Goldberg, E. E., J. R. Kohn, R. Lande, K. A. Robertson, S. A. Smith, and B. Igic. 2010.

Species selection maintains self-Incompatibility. *Science* **330**:493-495.

Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed

mating systems in plants: Occurrence, theoretical explanations, and empirical

evidence. *Annu Rev Ecol Evol Syst* **36**:47-79.

Goodwillie, C., R. D. Sargent, C. G. Eckert, E. Elle, M. A. Geber, M. O. Johnston, S.

Kalisz, D. A. Moeller, R. H. Ree, M. Vallejo-Marin, and A. A. Winn. 2010.

Correlated evolution of mating system and floral display traits in flowering plants

and its implications for the distribution of mating system variation. *New Phytol*

185:311-321.

Gottlieb, L. D. 1984. Genetics and morphological evolution in plants. *Am Nat* **123**:681-

709.

Hermann, K., U. Klahre, M. Moser, H. Sheehan, T. Mandel, and C. Kuhlemeier. 2013.

Tight Genetic Linkage of Prezygotic Barrier Loci Creates a Multifunctional

Speciation Island in *Petunia*. *Current Biology* **23**:873-877.

Jewell, C., A. D. Papineau, R. Freyre, and L. C. Moyle. 2012. Patterns of reproductive

isolation in *Nolana* (chilean bellflower). *Evolution* **66**:2628-2636.

Kendal, D., C. E. Hauser, G. E. Garrard, S. Jellinek, K. M. Giljohann, and J. L. Moore.

2013. Quantifying Plant Colour and Colour Difference as Perceived by Humans

Using Digital Images. *Plos One* **8**.

Knapp. S. 2010. On 'various contrivances': pollination, phylogeny and flower form in the

Solanaceae. *Phil Trans R Soc B* 365:449-460.

Kostyun, J. L. and L. C. Moyle. 2017. Multiple strong postmating and intrinsic

postzygotic reproductive barriers isolate florally diverse species of *Jaltomata*

(Solanaceae)*. Evolution* **71**:1556-1571.

Kostyun, J. L., J. C. Preston, and L. C. Moyle. 2017. Heterochronic developmental shifts

underlie floral diversity within *Jaltomata* (Solanaceae). *EvoDevo* 8(1): 17.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv:1303.3997v1 [q-bio.GN].

Manceau, M., V. S. Domingues, R. Mallarino, and H. E. Hoekstra. 2011. The

Developmental Role of Agouti in Color Pattern Evolution. *Science* **331**:1062-

1065.

Miller, R. J., T. Mione, H. L. Phan, and R. G. Olmstead. 2011. Color by Numbers:

Nuclear Gene Phylogeny of *Jaltomata* (Solanaceae), Sister Genus to *Solanum*,

Supports Three Clades Differing in Fruit Color. *Syst Botany* **36**:153-162.

Mione, T. 1992. *The systematics and evolution of* Jaltomata *(Solanaceae)*. PhD

Dissertation, The University of Connecticut. Available from ProQuest

Dissertations & Theses Global. (303964972).

Mione, T., G. J. Anderson, and M. Nee. 1993. *Jaltomata* I: Circumscription, description

and new combinations for five South American species. *Brittonia* 45: 138-145.

Mione, T., S. Leiva González, and L. Yacher 2015. Two new Peruvian species of

Jaltomata (Solanaceae, Solaneae) with red floral nectar. *Brittonia* **67**:105-112.

Mione, T., K. C. Plourd, S. Leiva González, L. Yacher, C. T. Philbrick, and D. Blume.

2017. Pollination decreases longevity of the protogynous flowers of *Jaltomata*

sinuosa (Solanaceae). *J Biol Nat* **7**:123–9.

Mione, T., J. L. Kostyun, and S. Leiva Gonzalez. In Revision. Self-compatibility,

protogyny and locating the floral nectary of *Jaltomata quipuscoae* (Solanaceae).

Moyle, L. C., and T. Nakazato. 2008. Comparative genetics of hybrid incompatibility:

Sterility in two Solanum species crosses. *Genetics* **179**:1437-1453.

Nakazato, T., L. H. Rieseberg, and T. E. Wood. 2013. The genetic basis of speciation in

the Giliopsis lineage of Ipomopsis (Polemoniaceae). *Heredity* **111**:227-237.

Olmstead, R. G., L. Bohs, H. A. Migid, E. Santiago-Valentin, V. F. Garcia, and S. M.

Collier. 2008. A molecular phylogeny of the Solanaceae. *Taxon* **57**:1159-1181.

Preston, J. C., L. C. Hileman, and P. Cubas. 2011. Reduce, reuse, and recycle:

developmental evolution of trait diversification. *Am. J. Bot.* **98**:397-403.

RAW Therapee Team. 2012. Raw Therapee: cross-platform raw image processing

program. http://rawtherapee.com/.

Rijpkema A, T Gerats, and M. Vandenbussche. 2006. Genetics of floral development in

Petunia. Pages 237-278 *in* L. J. Soltis DE, Soltis PS & Callow JA, editor. Adv Bot

Res: Incorp Adv Plant Pathol: Developmental Genetics of the Flower.

Rosas-Guerrero, V., R. Aguilar, S. Marten-Rodriguez, L. Ashworth, M. Lopezaraiza-

Mikel, J. M. Bastida, and M. Quesada. 2014. A quantitative review of pollination

syndromes: do floral traits predict effective pollinators? *Ecol. Lett* **17**:388-400.

Sarkinen, T., L. Bohs, R. G. Olmstead, and S. Knapp. 2013. A phylogenetic framework

for evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree.

BMC Evolutionary Biology **13**.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25

years of image analysis. *Nature Methods* **9**:671-675.

Schwartzwald, D. 2012. Color Space Converter Plugin for ImageJ.

https://imagej.nih.gov/ij/plugins/color-space-converter.html.

Sherman, N. A., A. Victorine, R. J. Wang, and L. C. Moyle. 2014. Interspecific Tests of

Allelism Reveal the Evolutionary Timing and Pattern of Accumulation of

Reproductive Isolation Mutations. *Plos Genetics* **10**.

Shrestha, R., J. Gomez-Ariza, V. Brambilla, and F. Fornara. 2014. Molecular control of

seasonal flowering in rice, arabidopsis and temperate cereals. *Ann Bot* **114**:1445-

1458.

Sicard, A., and M. Lenhard. 2011. The selfing syndrome: a model for studying the

genetic and evolutionary basis of morphological adaptation in plants. *Ann Bot*

107:1433-1443.

Smaczniak, C., R. G. H. Immink, G. C. Angenent, and K. Kaufmann. 2012.

Developmental and evolutionary diversity of plant MADS-domain factors:

insights from recent studies. *Development* **139**:3081-3098.

Smith, S. D. 2016. Pleiotropy and the evolution of floral integration. *New Phytol* **209**:80-

85.

Stebbins, G.L. 1974. Flowering plants: evolution above the species level. Harvard

University Press, Cambridge, MA.

Steinbachs J.E., and K.E. Holsinger 2002. S-RNase-mediated gametophytic self

incompatibility is ancestral in eudicots. *Mol Biol Evol* **19**: 825–829.

Stuurman, J., M. E. Hoballah, L. Broger, J. Moore, C. Basten, and C. Kuhlemeier. 2004.

Dissection of floral pollination syndromes in *Petunia*. *Genetics* **168**:1585-1599.

Taylor, J., and D. Butler. 2017. R Package ASMap: Efficient Genetic Linkage Map

Construction and Diagnosis. *Journal of Statistical Software* 79:1-29.

Tomato Genome Consortium, The. 2012. The tomato genome sequence provides insights

into fleshy fruit evolution. *Nature* 485:635-641.

van der Niet, T., and S. D. Johnson. 2012. Phylogenetic evidence for pollinator-driven

diversification of angiosperms. *Trends Ecol Evolut* **27**:353-361.

Wagner, G. P., and J. Z. Zhang. 2011. The pleiotropic structure of the genotype-

phenotype map: the evolvability of complex organisms. *Nature Reviews Genetics*

12:204-213.

Wessinger, C. A., L. C. Hileman, and M. D. Rausher. 2014. Identification of major

quantitative trait loci underlying floral pollination syndrome divergence in

Penstemon. *Phil. Trans. R. Soc. B* **369**.

Wessinger, C. A., C. C. Freeman, M. E. Mort, M. D. Rausher, and L. C. Hileman. 2016.

Multiplexed shotgun genotyping resolves species relationships within the North

American genus *Penstemon*. *Am J Bot* **103**:912-922.

Wessinger, C. A., and J. K. Kelly. 2018. Selfing can facilitate transitions between

pollination syndromes. *Am Nat* 191.

Wu, M., J. L. Kostyun, M. W. Hahn, and L. C. Moyle. 2018. Dissecting the basis of

novel trait evolution in a radiation with widespread phylogenetic discordance.

Mol Ecol 27: 3301-3316.

Wu, M., J. L. Kostyun, and L. C. Moyle. 2019. Genome sequence of *Jaltomata* addresses

rapid reproductive trait evolution and enhances comparative genomics in the

hyper-diverse Solanaceae. *Genome Biology and Evolution* in press. Available at:

https://www.biorxiv.org/content/early/2018/05/30/335117.

Xu, S. Q., and P. M. Schluter. 2015. Modeling the two-locus architecture of divergent

pollinator adaptation: how variation in SAD paralogs affects fitness and

evolutionary divergence in sexually deceptive orchids. *Ecology and Evolution*

5:493-502.

714 **Table 1**. Summary of measured floral and fertility traits within parental species, F1s, and

715 BC1s. Phenotypic means and variances provided, while significant differences between

716 parental species were assessed with t-tests on transformed data as appropriate (see

717 Methods). Note that significant differences in seed germination rates could not be tested.

718 * $p<0.05$, ** $p<0.001$; *** $p<0.0001$.

Trait	*J. sinuosa* (n=7)		*J. umbellata* (n=5)	
Floral Morphology/Physiology	**Mean**	**Variance**	**Mean**	**Variance**
Buds per Inflorescence	2.86**	0.03	8.60**	3.24
Calyx Diameter (mm)	15.49**	0.08	8.14**	2.91
Sepal Length (mm)	7.48**	0.02	4.41**	0.56
Corolla Diameter (mm)	29.82***	1.92	15.32***	5.70
Corolla Depth (mm)	1.38**	0.07	10.25**	5.52
Corolla Fusion (mm)	9.45	0.11	10.33	1.55
Corolla Fusion Proportion	0.63*	0.001	0.72*	0.001
Petal Length (mm)	14.85	0.41	14.40	0.16
Stamen Length (mm)	10.80	0.29	10.45	0.22
Anther Length (mm)	2.03*	0.02	1.73*	0.04
Ovary Diameter (mm)	2.28***	0.03	1.37***	0.05
Style Length (mm)	7.90***	0.17	14.64***	1.72
Herkogamy (mm)	-2.20***	0.01	5.37***	4.62
Nectar Volume (uL)	6.86***	1.07	17.80***	13.20
Floral Color				
Nectar Color - Intensity	205.93***	144.59	105.59***	106.17
Nectar Color - Red	207.88	222.00	219.47	274.81
Nectar Color - Green	208.71**	148.02	66.48**	239.95
Nectar Color - Blue	201.35***	105.86	30.79***	114.53
Nectar Color - Composite RGB	-202.17***	115.02	122.20***	48.24
Nectar Color - L	83.49***	20.32	50.42***	31.63
Nectar Color - a	-1.55***	0.95	57.61***	7.10
Nectar Color - b	3.47***	8.20	51.84***	59.31
Petal Color - Intensity	137.05**	127.04	112.33**	136.18
Petal Color - Red	135.85	138.57	128.59	114.35
Petal Color - Green	134.31	119.92	121.87	181.62
Petal Color - Blue	140.97**	123.32	86.55**	134.58
Petal Color - Composite RGB	-139.42***	106.86	-79.82***	208.17
Petal Color - L	83.18	2.59	84.15	4.86
Petal Color - a	0.81*	0.02	-0.93*	0.62
Petal Color - b	-1.26***	0.02	6.71***	1.05
Fertility				
Viable Pollen Grains	25143	36809254	26834	60773556
Proportion Viable Pollen	0.76	0.01	0.71	0.02
Fruit Set	0.93*	0.01	0.55*	0.06
Fruit Mass (g)	0.47**	0.01	0.12**	0.01
Fruit Diameter (cm)	1.01*	0.01	0.61*	0.02

		74.87*	256.92	30.44*	494.43
Seed Set		74.87*	256.92	30.44*	494.43
Viable Seed Set		74.87*	256.92	23.59*	182.31
Viable Seed T50		9.33	--	7.00	--
Viable Seed MGT		22.00	--	14.00	--

719 **Table 1 cont.**

F1s (n=13)		BC1s (n=224)	
Mean	**Variance**	**Mean**	**Variance**
3.95	0.68	3.51	1.63
10.55	0.48	12.10	1.08
5.45	0.13	6.16	0.28
24.58	6.84	28.97	9.92
6.64	0.58	5.22	1.35
9.81	0.40	9.92	1.24
0.67	0.001	0.63	0.002
14.65	1.61	15.78	2.88
10.81	0.59	10.86	1.26
2.01	0.01	2.13	0.04
1.62	0.02	1.82	0.04
12.39	1.04	10.82	1.97
2.95	1.08	1.20	1.02
11.90	12.32	10.76	18.87
172.44	110.66	192.70	272.74
232.93	149.55	213.14	127.46
198.03	248.93	206.89	165.90
74.27	317.30	157.76	1591.28
-39.37	1182.31	-151.51	2429.74
81.31	14.35	83.99	13.42
-0.05	101.74	-4.29	14.91
59.85	75.94	22.91	287.97
147.02	63.03	135.30	119.75
152.60	100.19	137.00	144.35
154.91	246.30	135.92	138.89
137.41	105.01	132.70	113.98
-139.72	391.57	-131.62	142.13
86.93	5.82	83.56	4.60
-0.81	0.19	-0.04	0.36
2.89	3.13	0.75	1.29
22513	16602405	26886	145318716
0.82	0.00	0.74	0.03
0.93	0.01	0.89	0.03
0.24	0.01	0.29	0.01
0.80	0.01	0.85	0.03
36.06	168.37	37.67	361.12
25.46	303.95	26.28	340.08
34.00	1001.78	26.70	607.37
46.04	622.37	39.36	458.11

721 **Table 2**. Summary of identified QTL for key floral and fertility traits. Includes peak

722 location and LOD of QTL, 1.5 LOD intervals, amount of phenotypic variance of trait

723 explained, phenotypic effect size and standard error (back-transformed where applicable),

724 and whether the effect is aligned with parental trait values. Full QTL data, including the

725 full models for all traits, are provided in Table S6.

Trait Category	Trait	LG	Peak Location	Peak LOD
Flower Morph	Inflor. Size (2)	LG 4	55.4	3.04
		LG 1	56	3.02
	Calyx Dia (3)	LG 1	56	9.06
		LG 8	84.3	3.15
		LG 9	66	5.59
	Sepal Length (3)	LG 1	57	8.30
		LG 12	23	3.74
		LG 9	65.9	7.04
	Corolla Dia (3)	LG 4	21.1	3.50
		LG 12	37.4	1.57
		LG 8	73.1	5.87
	Corolla Depth (4)	LG 7	52	17.95
		LG 5	40.8	10.19
		LG 10	55.8	4.48
		LG 9	30	12.38
		LG 5	86	3.57
	Corolla Fusion Prop (4)	LG 4	26	3.77
		LG 1	72	4.60
		LG 9	16	7.19
		LG 7	100	2.46
	Petal Length (3)	LG 12	48.2	2.04
		LG 8	73	3.55
		LG 7	91	4.45
	Stamen Length (4)	LG 4	120.8	3.77
		LG 2	37.5	4.54
		LG 9	0	5.32
	Anther Length (1)	LG 1	18	3.78
	Ovary Dia (4)	LG 7	9.9	3.56
		LG 5	72	2.53

		LG 1	68	5.77
		LG 12	48.2	4.31
	Style Length (2)	LG 2	95.2	2.79
		LG 9	22	9.89
		LG 1	314	4.53
		LG 2	112	7.26
	Herkogamy (5)	LG 11	40	4.79
		LG 8	51.4	3.27
		LG 9	7.3	4.32
		LG 5	116	3.61
		LG 4	59.2	3.55
	PC1 (6)	LG 1	54.2	4.08
		LG 2	31.7	4.73
		LG 8	82	8.08
		LG 9	19	10.34
		LG 1	58	2.62
	PC2 (3)	LG 12	48.2	3.27
		LG 8	58	2.14
	PC3 (2)	LG 1	67.5	4.99
		LG 9	24.7	11.52
Flower Physio	Nectar Volume (1)	LG 3	187.2	2.81
		LG 7	28	9.91
	Nectar PC1 (3)	LG 5	100	13.15
		LG 2	153	5.48
	Nectar PC2 (1)	LG 7	8.4	3.29
Flower Color	Nectar PC3 (1)	LG 3	66.1	3.47
	Petal PC1 (1)	LG 5	65.3	5.83
	Petal PC2 (2)	LG 5	63	18.91
		LG 9	67	10.42
	Petal PC3 (0)	no QTL		
	Fruit Set (0)	no QTL		
	Fruit Dia (2)	LG 1	80	4.67
		LG 11	76.9	3.09
		LG 3	170	2.67
	Fruit Mass (4)	LG 1	96	5.47
Fertility		LG 11	76.5	5.63
		LG 12	0	4.51
	Seed Set (2)	LG 11	76	6.64
		LG 12	0	2.86
	Viable Seed Set (1)	LG 11	76.9	4.42
	Viable Pollen (0)	no QTL		

727   **Table 2 cont.**

1.5 LOD Intervals	%PVE	Effect	Effect SE	Aligned with Parental Value?
15-108	5.50	1.1576	0.0169	YES
34-133	5.46	-1.1628	0.0175	NO
49-81	13.67	-0.8026	0.1202	YES
69-92.55	4.46	-0.4565	0.1196	YES
17-85	8.13	-0.5976	0.1160	YES
49-70.73	12.17	-0.3923	0.0616	YES
10-59	5.22	-0.2539	0.0609	YES
22-73	10.18	-0.3404	0.0584	YES
13-70	4.47	1.3869	0.3440	NO
0-65.17	1.96	-0.9630	0.3606	YES
67-84	7.69	-1.9019	0.3596	YES
48-60.67	21.47	1.1455	0.1164	YES
20-46	11.20	0.8437	0.1187	YES
41-91	4.64	-0.5214	0.1140	NO
11-38	13.94	0.9420	0.1188	YES
80.21-99	5.39	0.0005	0.0000	YES
19-84	5.70	-0.0004	0.0000	NO
57-106	7.02	-0.0005	0.0000	NO
0-67	11.27	0.0008	0.0000	YES
25-100.67	3.29	1.0414	0.0050	NO
0-65.17	2.71	1.0390	0.0053	YES
39-92.55	4.78	-1.0657	0.0053	YES
59-100.67	5.68	0.5510	0.1209	NO
100-133	4.78	-0.4994	0.1194	YES
6-61	5.80	-0.5478	0.1190	YES
0-33	6.86	0.6194	0.1237	NO
8-35.87	6.48	-0.1157	0.0274	YES
0-25	5.56	-0.0936	0.0230	YES
0-84	3.90	0.0915	0.0269	NO
60-107	9.21	-0.1196	0.0229	YES
5-65.17	6.78	-0.1036	0.0231	YES
55-126	3.80	1.0517	0.0061	YES
1-30	14.51	1.1072	0.0063	YES
266-324	6.42	-1.1613	0.0141	NO
30-122	10.60	1.2156	0.0144	YES
5-63	6.81	1.2143	0.0783	YES
26-63	4.57	1.1344	0.0141	YES

3.47-32	6.12	1.1560	0.0140	YES
95-124.77	3.61	0.3880	0.1025	YES
24-76	3.55	-0.3691	0.0983	NO
34-86	4.08	0.3853	0.0958	YES
4-44	4.73	0.4121	0.0952	YES
68.51-92.55	8.08	0.5552	0.0981	YES
12-71	10.34	0.6040	0.0943	YES
43-138	3.72	-0.6666	0.1921	YES
16-62.42	4.69	-0.7556	0.1940	YES
36-92.55	3.03	-0.6066	0.1939	YES
46-86	7.27	-0.6665	0.1371	YES
20-31	18.01	-1.0720	0.1401	YES
172-187.19	5.22	1.2539	0.0366	YES
23-37	11.96	-1.5076	0.2152	YES
83-107	16.45	-1.9822	0.2413	YES
147-171.87	6.31	-1.1060	0.2174	YES
0-22	6.21	0.7061	0.1801	NO
17-121	6.92	-0.5691	0.1412	YES
60-72	11.4	-1.7713	0.3344	YES
59-67	28.06	-1.4839	0.1453	YES
39.15-73	14.09	-0.9231	0.1276	YES
61-103	8.64	-1.0608	0.0054	YES
51-79.93	5.61	-1.0463	0.0052	YES
48-187.19	4.07	-1.0168	0.0023	YES
60-146	8.58	-0.9745	0.0049	YES
54-79.93	8.85	-0.9762	0.0047	YES
0-10.64	7.01	-0.9771	0.0050	YES
63-79.93	12.28	-1.4353	0.0277	YES
0-63	5.07	-1.2857	0.0300	YES
68-79.93	8.85	-1.6267	0.0462	YES

Figure Legends.

Figure 1. Representative flowers of the parental species and their F1 hybrid.

Figure 2. Key trait distributions within the BC1 population, compared to phenotypic

means for *J. sinuosa* (purple line), *J. umbellata* (red line), and their F1s (pink line).

Figure 3. Key floral trait correlations within the BC1 mapping population. Scatterplots

are provided below the diagonal, while Spearman's correlation coefficients and

associated p-values are above the diagonal. Statistically significant correlations following

Bonferroni correction ($p < 3.125 \times 10^{-5}$) are highlighted in red. Correlation values for all

traits are provided in Table S3, and for PC traits provided in Table S5.

Figure 4. Linkage map and distribution of identified QTL (including 1.5 LOD intervals)

for 12 key floral and fertility traits.

Supplementary Information

Table S1. Accession information for material used and generated in this study.

Table S2. Trait measurements for all individuals phenotyped in this study.

Table S3. Correlations among all measured traits within BC1 individuals.

Table S4. Principle component loadings for highly correlated traits.

Table S5. Correlations among PC traits within BC1 individuals

Table S6. Full QTL models for all examined traits.

Table S7. Expected and observed counts of QTL overlap.

Figure S1. Measured morphological traits on mature flowers.

Figures S2-S3: Distributions for floral morphological and physiological traits within the

mapping population.

Figure S4. Distributions for nectar color traits within the mapping population.

Figure S5. Distributions for petal color traits within the mapping population.

Figure S6. Distributions for fertility traits within the mapping population.

Figure S7. Representative examples of petal color variation among F1 individuals,

compared to either parent.

Figure S8. Comparison of observed number of QTL overlaps with counts from 10000

randomly generated simulations, for QTL co-localization overlap within and between

trait categories (morphology, color, and fertility).

J. sinuosa F1 *J. umbellata*

1 cm

	Calyx Dia	Cor Dia	Cor Depth	Cor Fus Prop	Pet Length	Stam Length	Ovary Dia	Style Length	Herk	Nec Vol	Nec Color	Pet Color
Calyx Dia		r=0.656 p<1.02e⁻⁵	r=0.052 p=0.440	r=-0.162 p=0.015	r=0.548 p<1.02e⁻⁵	r=0.105 p=0.118	r=0.571 p<1.02e⁻⁵	r=0.001 p=0.989	r=-0.109 p=0.103	r=0.176 p=0.008	r=-0.040 p=0.551	r=-0.132 p=0.048
		Cor Dia	r=0.216 p=0.001	r=-0.348 p<1.02e⁻⁵	r=0.905 p<1.02e⁻⁵	r=0.380 p<1.02e⁻⁵	r=0.448 p<1.02e⁻⁵	r=0.455 p<1.02e⁻⁵	r=0.113 p=0.092	r=0.286 p<1.33e⁻⁵	r=0.259 p=0.0001	r=-0.067 p=0.321
			Cor Depth	r=0.208 p=0.002	r=0.450 p<1.02e⁻⁵	r=0.122 p=0.067	r=0.198 p=0.003	r=0.277 p=2.68e⁻⁵	r=0.126 p=0.059	r=0.142 p=0.033	r=0.242 p=0.0003	r=0.044 p=0.512
				Cor Fus Prop	r=-0.255 p=0.0001	r=0.088 p=0.191	r=-0.076 p=0.257	r=-0.026 p=0.665	r=0.033 p=0.624	r=-0.070 p=0.295	r=-0.084 p=0.209	r=-0.013 p=0.846
					Pet Length	r=0.447 p<1.02e⁻⁵	r=0.414 p<1.02e⁻⁵	r=0.531 p<1.02e⁻⁵	r=0.163 p=0.015	r=0.239 p=0.0003	r=0.318 p<1.02e⁻⁵	r=0.001 p=0.987
						Stam Length	r=0.015 p=0.827	r=0.511 p<1.02e⁻⁵	r=-0.154 p=0.021	r=0.098 p=0.144	r=0.167 p=0.012	r=0.090 p=0.182
							Ovary Dia	r=-0.056 p=0.406	r=-0.058 p=0.387	r=0.267 p=0.0001	r=0.018 p=0.789	r=0.012 p=0.862
								Style Length	r=0.620 p<1.02e⁻⁵	r=0.159 p=0.018	r=0.326 p<1.02e⁻⁵	r=0.066 p=0.327
									Herk	r=0.015 p=0.820	r=0.221 p=0.0009	r=-0.027 p=0.692
										Nec Vol	r=0.209 p=0.002	r=0.005 p=0.946
											Nec Color	r=0.140 p=0.036
												Pet Color