A simple genetic architecture and low constraint allows rapid floral evolution in a

diverse and recently radiating plant genus

Jamie L. Kostyun[1,2*], Matthew J.S. Gibson[1], Christian M. King[1,3], and Leonie C. Moyle[1]

[1] Department of Biology, Indiana University, Bloomington, IN 47405

[2] Department of Plant Biology, The University of Vermont, Burlington, VT 05405

[3] Department of Evolution, Ecology and Organismal Biology, The Ohio State University,

Columbus, OH 43210

* Corresponding author: Jamie.Kostyun@uvm.edu, Tel: 802-656-7670

Word count: Summary (199), Introduction (1190), Materials and Methods

(1638), Results (1387), Discussion (2014), Acknowledgements (94), Total Main

Text (6229), Figures (4, all in color), Tables (2), Supplementary Documents (8

Figures, 8 Tables).

Summary

• Genetic correlations among different components of phenotypes, especially

resulting from pleiotropy, can constrain or facilitate trait evolution. These factors

could especially influence the evolution of traits that are functionally integrated,

such as those comprising the flower. Indeed, pleiotropy is proposed as a main

driver of repeated convergent trait transitions, including the evolution of

phenotypically-similar pollinator syndromes.

• We assessed the role of pleiotropy in the differentiation of floral and other

reproductive traits between two species—*Jaltomata sinuosa* and *J. umbellata*

(Solanaceae)—that have divergent suites of floral traits consistent with bee- and

hummingbird-pollination, respectively. To do so, we generated a hybrid

population and examined the genetic architecture (trait segregation and QTL

distribution) underlying 25 floral and fertility traits.

• We found that most floral traits had a relatively simple genetic basis (few,

predominantly additive, QTL of moderate to large effect), as well as little

evidence of antagonistic pleiotropy (few trait correlations and QTL co-

localization, particularly between traits of different classes). However, we did

detect a potential case of adaptive pleiotropy among floral size and nectar traits.

• These mechanisms may have facilitated the rapid floral trait evolution observed

within *Jaltomata*, and may be a common component of rapid phenotypic change

more broadly.

Key Words: floral evolution, genetic co-variation, *Jaltomata*, pleiotropy, Solanaceae,

QTL co-localization, QTL mapping

Introduction

One feature of phenotypic evolution is the broad variation in observed rates of

trait change, both among traits within a single lineage and among lineages. How quickly

phenotypes evolve and in what direction, depends not only on their genetic basis--both

the number and effect size of causal loci--and the intensity and nature of selection acting

upon them, but also on their associations with other traits. Because of genetic covariance,

developmental and phylogenetic constraints, and/or correlated selection pressures

(Agrawal & Stinchombe, 2009), phenotypic traits often do not evolve independently from

one another. Among these causal mechanisms, strong genetic covariance arises either

because different traits have a shared genetic basis (pleiotropy) or because they are based

on genes that are physically adjacent and therefore often co-inherited (via linkage).

Strong pleiotropy is often proposed to constrain phenotypic evolution by preventing

correlated traits from moving efficiently towards their own (different) fitness optima

(antagonistic pleiotropy). However, such shared genetic control may also promote

phenotypic change if covariation is aligned in the direction of selection (adaptive

pleiotropy) (Agrawal & Stinchombe, 2009; Wagner & Zhang, 2011; Smith, 2016).

Nonetheless, despite some detailed studies (e.g. Ettensohn, 2013; Shrestha *et al.*, 2014;

Manceau *et al.*, 2011; Xu & Schluter, 2015), the genetic architecture of many

ecologically important traits remains unclear, including the prevalence of strong genetic

associations that could shape the course of phenotypic evolution.

How shared architecture influences phenotypic change is especially relevant for

suites of traits that are functionally integrated (Armbruster *et al.*, 2014) -- such as the

angiosperm flower (Armbruster *et al.*, 2009) -- because the magnitude and direction of

pleiotropy will directly shape if and how these co-varying traits respond to selection.

Further, because pleiotropy can constrain or favor particular developmental trajectories, it

is often proposed to be a main driver of convergent transitions of integrated traits in

different lineages (Preston *et al.*, 2011; Smith, 2016). The flower is an especially

promising model for assessing the role of pleiotropy in shaping phenotypic evolution.

Because flowers mediate fitness through their critical reproductive role, their constituent

traits (i.e. reproductive structures, perianth, and other attraction/reward features) are often

highly functionally integrated (Conner, 2002; Armbruster *et al.*, 2009). Moreover,

repeated transitions of phenotypically similar or convergent suites of floral traits have

been identified both within and across groups (Fenster *et al.*, 2004; Goodwillie *et al.*,

2010; Wessinger *et al.*, 2016). For example, multiple parallel shifts from bee-pollination

to hummingbird-pollination are associated with parallel transitions to flowers with red

petals, large amounts of dilute nectar, and narrow corolla tubes within *Penstemon*

(Wessinger *et al.*, 2016); similarly, the evolution of the 'selfing syndrome' (i.e. reduced

overall size, herkogamy, and floral rewards) often accompanies transitions from

outcrossing to predominantly selfing mating systems, and has been documented in

multiple lineages (Stebbins, 1974; Goodwillie *et al.*, 2010). Such patterns provide an

opportunity to assess the relative frequency of adaptive vs. antagonistic pleiotropy in

shaping these repeated trait combinations, within a comparative phylogenetic context.

In addition to these ecological and evolutionary features, the known genetic and

molecular bases of floral development (Rijpkema *et al.*, 2006; Smaczniak *et al.*, 2012)

themselves suggest that pleiotropy might be a key component shaping floral phenotypic

change, as well as provide a functionally-informed framework for identifying how

changes in these mechanisms can contribute to the diversification of floral traits. Under

the ABC(DE) model of floral development, the combinatorial action of different gene

products -- primarily different MADS-box transcription factors -- control transitions to

flowering and the specification of floral organ identities and organ maturation, by

promoting or repressing different downstream targets (reviewed in O'Maoileidigh *et al.*,

2014; Bartlett, 2017). This combinatorial function, and the ability to regulate shared or

partially shared downstream targets, provides a potential mechanistic explanation for

strong pleiotropy among floral traits. Moreover, several key regulators of floral

development also function during fruit and seed production (Smaczniak *et al.*, 2012;

O'Maoileidigh *et al.*, 2014); such correlated effects on fertility traits are another potential

way that pleiotropy could shape floral phenotypic evolution.

Several empirical approaches have been taken to assess the genetic architecture

underlying floral trait specification and evolution, and to evaluate the strength and

direction of pleiotropy. Classical quantitative genetic analyses have revealed varying

degrees of genetic covariance among floral traits (Gottlieb, 1984; Conner *et al.*, 2014),

while numerous QTL (quantitative trait locus) mapping studies have found that loci for at

least some different floral traits appear to co-localize to the same genomic region(s)

(reviewed in Smith, 2016). Interestingly, such studies have identified more putative cases

of adaptive pleiotropy (e.g. QTL affect more than one trait in the direction of parental

trait values) than antagonistic, suggesting that adaptive pleiotropy may be a common

mechanism contributing to rapid floral evolution. Because QTL generally span a genomic

region that contains more than one gene, such QTL co-localization is consistent with, but

not definitive evidence of, a role of pleiotropy in shaping floral trait co-variation (e.g. see

Hermann *et al.*, 2013). Nonetheless, identifying strong trait correlations and/or QTL co-

localization represents a critical step in assessing how pervasive pleiotropy could be in

shaping floral trait evolution.

In this study, we examined phenotypic (co)variation among, and the genetic

architecture underlying, floral and other reproductive traits within a segregating hybrid

population derived from two *Jaltomata* (Solanaceae) species with divergent floral traits

(**Figure 1**). Despite only having diversified with the last 5 million years (Sarkinen *et al.*,

2013; Wu *et al.*, 2018), species of *Jaltomata* are highly phenotypically diverse, including

extensive variation in the size, shape, and color of floral traits that is absent among their

close relatives within *Solanum* and *Capsicum*. Indeed, phylogenetic analyses suggest

numerous transitions in floral traits within the genus, including several instances of

convergent evolution (Miller *et al.*, 2011; Wu *et al.*, 2018). Importantly, many of these

transitions appear to involve parallel changes in several different traits within a lineage

(e.g. from flat corollas with small amounts of lightly colored nectar to highly fused

corollas with large amounts of darkly colored nectar), suggesting either a shared genetic

basis and/or correlated selection (perhaps pollinator-mediated selection) influences these

trait associations.

Here, we identified QTL contributing to floral and other reproductive trait

variation within a recombinant population to: 1) examine the genetic architecture

underlying reproductive trait divergence; 2) assess the role of genetic linkage and/or

pleiotropy (via strong trait correlations and overlapping QTL) in floral trait (co)variation;

and 3) assess evidence for a shared genetic basis between different classes of floral trait

and other reproductive (specifically fertility) traits. We found evidence for a relatively

simple genetic basis underlying most of the examined traits, as well as positive

correlations and significant QTL co-localization among traits within each of three trait

classes (floral morphology, floral color, and fertility). Together, these features might

facilitate rapid changes in these traits. In comparison, we found few associations between

traits from different classes, and therefore little evidence for antagonistic pleiotropy

among these classes, which otherwise may have constrained trajectories of floral change.

One striking exception was an association between flower size and nectar traits that acts

in the direction exhibited by multiple species in the genus, suggesting that this could

instead be an instance of adaptive pleiotropy.

MATERIALS AND METHODS

Study system: *Jaltomata* (Schlechtendal; Solanaceae) is the sister genus to the

large and economically important *Solanum* (Olmstead *et al*., 2008; Sarkinen *et al*., 2013;

Wu *et al*., 2019), and includes approximately 60-80 species distributed from the

Southwestern United States to the Andean region of South America, in addition to several

species endemic to the Greater Antilles and the Galapagos Islands (Miller *et al*., 2011;

Mione *et al*., 2015). Species are highly phenotypically diverse and live in a variety of

habitats (e.g. tropical rainforests, rocky foothills, and *lomas* formations).

Here, we focused on a closely related species pair, *J. sinuosa* and *J. umbellata*,

that differ in a suite of floral traits that is representative of major floral suites found in

other species throughout the genus (**Figure 1**; **Table S1**). *Jaltomata sinuosa* has large

rotate flowers with purple petals and a small amount of concentrated nectar (consistent

with bee pollination, Fenster *et al*., 2004; Mione *et al*., 2017), while *J. umbellata* has

small short-tubular flowers with yellowish-white petals and a large amount of dilute, but

dark red, nectar that is visible through the corolla tube (Kostyun & Moyle, 2017). The

tubular corollas, large volume of nectar, exerted reproductive organs, and red coloration

observed in this species is consistent with hummingbird pollination (Fenster *et al.*, 2004).

Jaltomata sinuosa is distributed along the Andes in South America from Venezuela to

Bolivia, while *J. umbellata* is restricted to lomas formations along the Peruvian coast.

Both species are self-compatible (Kostyun & Moyle, 2017) and shrubby, but differ in leaf

traits such as overall size and shape, and type of trichomes. This species pair also has

several incomplete, intrinsic postzygotic reproductive barriers, including quantitatively

reduced fruit set and hybrid seed viability (Kostyun & Moyle, 2017).

Generation of BC$_1$ population and plant cultivation: We developed a

segregating hybrid population by crossing *J. sinuosa* and *J. umbellata*. Viable F1

individuals in both directions of the cross produce flowers that are phenotypically

intermediate between the parental genotypes (**Figure 1**), and retain reduced but sufficient

levels of fertility when back-crossed to parents (Kostyun & Moyle, 2017). Because we

were especially interested in the genetic basis of red nectar and a fused corolla tube

(exhibited by *J. umbellata*, present but less pronounced in F1s, but absent in *J. sinuosa*;

Figure 1; Figure S7), we generated the mapping population by backcrossing a single *J.*

sinuosa x *J. umbellata* F1 (as the ovule parent) to the original parental *J. sinuosa*

individual (as the pollen parent). BC1 individuals were germinated in a growth chamber,

and then moved to the Indiana University greenhouse and grown under the same

conditions as the parental and F1 individuals (16 hour light cycle, watered twice daily,

and fertilized weekly).

Trait measurements: We measured 25 floral and other reproductive traits within

our mapping population, F1, and parental genotypes (**Table 1; Table S2; Figure S1**).

Floral morphological traits were measured with hand-held calipers, and nectar volume

per flower was measured to the nearest 1 μL using a pipette. To reduce the potential

effect of daily environmental variation, nectar volume was always measured in the early

afternoon following watering.

Petal and nectar color were quantified using digital photography (Kendal *et al.*,

2013; Garcia *et al.*, 2014): dissected whole petals or nectar drops were photographed on a

standard background along with white and black color standards. Light conditions were

standardized for all images using RAW Therapee (RAW Therapee Development Team,

2012), and color space attributes were measured in ImageJ (Schneider *et al.*, 2012) as the

average value across the entire whole petal or nectar drop sample. Because RGB color

attributes are device-dependent (i.e. they can vary depending upon the specific camera

used), color values were then converted into device-independent L*a*b color attributes,

using the ImageJ Color Space Converter plugin (Schwartzwald, 2012). For petal and

nectar color, this produced three color attributes each: Lightness (L; ranges from 0

[black] to 100 [white]), 'a' color (more negative values correspond to more green, and

more positive values correspond to more magenta), and 'b' color (more negative values

correspond to cyan, and more positive values correspond to yellow).

We also measured nine fertility traits (**Table 1**) to assess the potential genetic

overlap between floral and other reproductive traits, as well as to examine the genetic

architecture underlying intrinsic postzygotic barriers between this species pair. Fruit and

seed related traits were measured on 2-6 crossed fruit per individual (depending on fruit

set). For F1 and BC1 individuals, crossed fruit were produced using pollen from the *J.*

sinuosa parental individual. To determine seed germination rates (following Farooq *et al*.,

2005), we soaked 10 seeds per individual in 50% bleach for 30 minutes (to soften the

seed coat), rinsed thoroughly, and placed on moist paper within plastic germination

boxes. A week after sowing, seed coats were nicked slightly and seeds were given a drop

of 10 mM giberellic acid (Sigma) to break dormancy. We then scored germination every

2 weeks for 4 months. Pollen viability was estimated from three different flowers per

individual, using established methods (Jewell *et al*., 2012): Briefly, for each sample all

undehisced anthers from a flower were collected into an eppendorf tube containing

aniline blue histochemical stain, gently ground with a pestle to release pollen, and viable

pollen grains were counted under an EVOS FL Digital Inverted Fluorescence Microscope

(Fisher Scientific). From these data, we also calculated the proportion of viable pollen as

number of viable pollen grains/total pollen grains in the sample.

Statistical analyses on phenotypic data: Following Shapiro-Wilk tests to assess

normality assumptions, we transformed traits that showed a skewed distribution and/or

significantly non-normal residuals. In particular, we arcsine transformed proportional

traits (proportion of corolla fusion, fruit set, and proportion of viable pollen), and log-

transformed inflorescence size, corolla fusion, petal length, style length, herkogamy,

nectar volume, all color attributes, and remaining fertility traits. Significant differences

between parental species for all traits were assessed by t-tests (**Table 1**). Distribution

plots for all traits are provided in Supplementary Materials (**Figures S3-S6**), while

illustrative plots for 15 focal traits are provided in the main text (**Figure 2**). Similarly,

phenotypic correlations within the BC1 population were examined among all traits

(**Table S3**), while relationships among a subset of focal traits are presented in the main

text (**Figure 3**). Given significant correlations among many of the floral morphological

traits (**Table S3**), we also used principal component analyses (PCAs) to create separate

composite metrics (i.e. Morph PC1-PC3; **Table S4; Figure S2**) as additional measures of

floral variation. Trait correlations, including PCs, are provided in **Table S5**, and

distribution plots are provided in **Figure S4**.

Genotyping and linkage map construction: Genomic DNA was extracted from

young leaf tissue from the 2 parental individuals, 13 F1s (including the F1 parent used to

generate the population), and 269 BC1s, using the Qiagen DNeasy Plant Mini Kit. DNA

quantity and quality were confirmed via Nanodrop (Fisher Scientific) and gel

electrophoresis with λ DNA-HindIII Digest marker (New England BioLabs). Samples

were then sent to Novogene Corporation (Beijing) for genotyping-by-sequencing (GBS).

GBS libraries were prepared using optimized restriction enzymes (MseI and HaeIII), and

following insert size selection, sequenced on an Illumina Hi-Seq to generate 150bp paired

end reads. Raw reads were trimmed and filtered using Trimmomatic (Bolger *et al*., 2014),

and read quality was checked pre- and post-trimming using fastqc (Andrews, 2010). To

identify SNPs, cleaned reads were mapped to the domesticated tomato genome (Tomato

Genome Consortium, 2012) using the mem function in BWA (Li, 2013). Alignment files

were then input into the Stacks refmap pipeline (Catchen *et al*., 2013) to determine

genotypes. A minimum depth of six identical reads was required to create an individual

stack (-m parameter) and automated genotype correction was implemented during export.

Subsequent filtering of genotypes was performed during linkage map construction (see

below). Reads and genotype data are available in NCBI BioProject PRJNA514267.

To construct the linkage map, we first removed markers that were genotyped in

less than 35% of individuals, or showed significant segregation distortion (i.e. alleles at

>80% or <20% frequency). The linkage map was constructed using the MST and

Kosambi algorithms, implemented in the R package ASMap (Taylor & Butler, 2017). To

alleviate map expansion issues, we removed markers which consistently differed from

neighboring markers in terms of genotype assignment, indicating a high likelihood of

genotyping error. The linkage map was then finalized using the ripple function in R

package R/qtl (Broman *et al*., 2003).

Identifying QTL: We implemented Haley-Knott regression in R/qtl to identify

QTL contributing to each trait. To account for potential environmental contributions to

trait variation, we included date of measurement (Month) and location within the

greenhouse (Bench) as covariates in our QTL scans. Putative QTL were first identified

using the scanone function, followed by permutations for genome-wide LOD significance

thresholds. Two dimensional scans (scantwo function) were used in the stepwise qtl

function to fit multiple QTL models. These models were used to identify: significant

QTL, their 1.5 LOD confidence intervals, their effect sizes (i.e. difference in phenotype

mean between homozygotes and heterozygotes), the total amount of phenotypic variance

explained by each QTL, the proportion of parental difference explained (relative

homozygous effect or RHE), interactions among QTL within traits (testing evidence for

epistasis), and potential contributions of covariates. QTL were considered to be co-

localized if their 1.5 LOD intervals overlapped. Significant co-localization was assessed

by comparing overlap among identified QTL to overlap from 10000 randomly generated

distributions, for traits within each category (morphological, color/physiological, or

fertility), and between each trait category. Briefly, a custom Python script was used to

generate random distributions of QTL (by randomly re-distributing the identified QTL

among the 12 linkage groups), and the observed frequency of co-localization in each was

recorded for each randomization to generated count distributions, in R. All code used to

generate the linkage map, identify QTL, and assess QTL co-location, is available on

GitHub (https://github.com/gibsonMatt/jaltomataQTL).

RESULTS

Segregation patterns suggest additive alleles underlie most floral traits

Most traits were significantly different between the two parental species (**Table**

1). F1 means were intermediate for most traits as well, except that petals were generally

brighter (more white) than either parent, perhaps because F1s had lower amounts of the

different pigments found in each parental species (i.e. purple pigment found in *J. sinuosa*

petals, and yellow and red pigments found in *J. umbellata*) (**Figures S5+S7**). Other than

fruit set and seed germination rates, all traits were unimodally distributed within the

BC1s; phenotypic values were intermediate between F1s and the recurrent parent (*J.*

sinuosa) for many of these traits, consistent with additive effects (**Figure 2; Figures S3-**

S6). Several traits (7 of 25) showed transgressive segregation within the BC1s, including

some floral morphological traits, nectar volume and color (all 3 attributes), and seed

viability and germination rates (**Table S2; Figures S3-S6**).

Significant correlations observed within – but generally not between – floral

morphology, floral color, and fertility trait categories

Within the BC1s, most traits were not strongly associated with one another.

Nonetheless, several correlations remained significant following multiple testing

(Bonferroni) correction (**Table S3**), primarily associations that are expected biologically,

including allometric relationships among floral organs and positive associations among

related fertility traits. For instance, corolla diameter was significantly positively

associated with most other morphological traits, suggesting shared genetic control of

overall floral size (**Figure 3**), while corolla diameter was also significantly negatively

correlated with proportion of corolla fusion (i.e. shorter corolla tubes had wider limbs and

longer tubes had narrower limbs, r = -0.348, p = 8.68 x 10^{-8}). These relationships were

recovered with PCA on morphology traits, in which PC1-PC3 explained 76% of the

variance among BC1s (**Table S4**). Based on trait loadings, PC1 corresponds to floral

width vs. depth, PC2 to overall floral size, and PC3 to relative reproductive organ

dimensions. Similarly, related fertility traits also remained strongly correlated, such as

fruit mass with seed set, and number of viable pollen grains with proportion of viable

pollen (**Table S3**).

In contrast, there were relatively few significant correlations among different trait

categories. Notable exceptions, however, included a positive relationship between corolla

diameter and nectar volume, as well as several morphological traits and certain aspects of

color (nectar 'b', petal 'a' and 'b') (**Figure 3; Table S3**). In particular, larger flowers

tended to have more red nectar as well as more purple petals. There were also significant

positive correlations between pollen viability and each of several components of flower

size (as well as Morph PC2 or "size") (**Tables S3+S5**). This latter relationship seems to

be explained by anther size: across 15 *Jaltomata* species, mean viable pollen count is

significantly associated with anther size prior to dehiscence ($F = 15.56$, $p = 0.0017$) (J.L.

Kostyun, unpub.).

Linkage map construction recovered 12 linkage groups

Mapping high quality reads to the tomato genome identified 25,136 SNPs that

differentiated the two parental species. Following all subsequent filtering (removing

markers genotyped in less than 35% of individuals, with high segregation distortion or

non-Mendelian inheritance, or with high likelihood of genotyping errors), we retained

520 high quality markers. Linkage map construction recovered 12 linkage groups (LGs),

which correspond to the number of chromosomes in the parental species (Mione *et al*.,

1993; Chiarini *et al*., 2017). Total map length was 1593.71 cM (65.17 cM – 324.48 cM

per chromosome/LG), with an average of 2.92 cM between markers (**Figure 4**).

Few moderate to large-effect QTL underlie most traits, with little QTL co-localization between trait categories

We identified QTL for 20 of our 25 examined traits, for a total of 63 QTL (with 4

additional loci for Morph PCs). For traits with identified QTL, most had 2-4 QTL

(range=1-7). Alleles at 55 of 67 QTL (82%) acted in a direction consistent with parental

values (i.e. the allele from paternal donor *J. umbellata* moved the phenotype of BC1s

closer to this species mean); at least one QTL with opposite effects was detected for 10 of

the 20 traits (and Morph PC1) (**Table 2**; **Table S6)**. Consistent with trait segregation

patterns observed in the BC1, significant interactions (epistasis) among QTL were

identified for only two traits: ovary diameter and nectar color 'a' (**Table 2**; **Table S6**).

The amount of phenotypic variation explained by each QTL ranged from 2-33%

(**Table 2**; **Table S6**). Of these, 19 traits (and Morph PC1 and PC3) had at least 1

moderate effect QTL (defined as explaining 5-25% of the variance), and 1 trait (petal

color 'b') had a QTL with major effect (defined as >25% variance explained) (**Table 2;**

Table S6). The lower bound of this range suggests that we had reasonable power to

identify QTL with even relatively small effects -- explaining as little as 2% of the

variance. We note however, that like most QTL analyses (Broman, 2001), we likely have

limited power to detect all QTL contributing to our examined traits. Nonetheless, under

strict additivity, QTL account for >50% of the parental difference (relative homozygous

effect, RHE) in 14 of the 20 traits for which we identified QTL (as well as Morph PC1-

3); for 8 of these traits (as well as Morph PC1-3), >75% of the parental difference is

accounted for (**Table 2; Table S6**). That is, for most of our traits, detected loci account

for the majority of parental phenotypic differences. In contrast, for four traits (corolla

diameter, proportion of corolla fusion, herkogamy, and inflorescence size) QTL

accounted for <20% of the parental difference, suggesting additional undetected loci

contribute to these traits. Finally, we estimated RHE >1 for 5 floral traits that also show

the most transgressive segregation among BC1s (nectar color 'b', petal color 'L', petal

length, corolla fusion, and Morph PC2) (**Table S6; Figures S3-5**).

Although every linkage group had at least one QTL, these were not distributed

uniformly across the genome, with notable clusters on LG1 and LG9 (**Figure 4; Table 2;**

Tables S6+S8). We also identified several instances of QTL co-localization within trait

categories, especially for morphology and fertility traits, which each had significantly

more cases of co-localization (1.5 LOD overlap) than expected by chance (133 observed

vs. upper bound of 115 expected overlaps, $p = 6.9$ x 10^{-4}; 8 observed vs. upper bound of 4

expected overlaps, $p = 5.0$ x 10^{-6}, respectively) (**Table S7; Figure S8**). These instances of

co-localization included QTL for biologically-related traits for which we observed strong

correlations, such as those underlying petal length and corolla diameter on LG8 and

LG12, corolla depth and corolla fusion on LG7, LG9 and LG10, and fruit mass and seed

set on LG11 and LG12 (**Tables S3+S8**). In some cases, co-localized QTL share the same

or a very close peak marker (e.g. calyx diameter and inflorescence size on LG1, and petal

length, corolla fusion, and ovary diameter on LG12; **Tables S6+S8**), which suggests

pleiotropic effects, however we note that the large 1.5 LOD intervals of some QTL will

increase instances of incidental co-localization events.

In contrast, co-localization between different trait categories was never greater

than expected by chance (**Table S7; Figure S8**). This is consistent with mostly incidental

overlap between QTL for traits in different categories. Nonetheless, we did detect co-

localization at the same or very close peak markers for three sets of biologically

interesting trait combinations: corolla diameter, proportion of corolla fusion, and petal

color 'L' on LG4; corolla fusion, corolla depth, and nectar color 'a' on LG7; and nectar

volume and nectar color 'b' on LG3 (**Figure 4; Tables S6+S8**). Not all of the traits

within each of these sets is significantly phenotypically correlated in the BC1 (**Table S3**),

especially correlations involving nectar color 'a' and 'b', possibly because these nectar

color traits have additional QTL that are not shared with the corolla traits or nectar

volume. Nonetheless, the colocalized QTL still explain large fractions of the parental

difference for these traits: on LG7 the QTL colocalized with corolla fusion and depth

explains ~20% of the parental difference in nectar color 'a', while the potentially shared

QTL on LG3 explains ~25% of the parental difference in nectar color 'b' and nearly 50%

of the parental difference in nectar volume (**Table 2; Table S6**). Together, these provide

intriguing cases of potential adaptive pleiotropy, such that alleles at QTL may have

simultaneously acted to increase corolla tube depth and nectar darkness, and increase

nectar production and nectar darkness, respectively.

DISCUSSION

Genetic correlations among different components of the phenotype, especially

resulting from pleiotropy, can constrain or facilitate trait evolution (Agrawal &

Stinchcombe, 2009). Pleiotropy could have particularly strong effects on the evolution of

traits that are functionally integrated, such as those comprising the flower (Armbruster *et*

al., 2009; Smith, 2016). Here, we examined the genetic architecture underlying floral trait

evolution within florally diverse *Jaltomata*, including whether pleiotropy might have

shaped observed variation. We found that most of our examined traits have a relatively

simple genetic basis, with few to moderate QTL with largely additive effects. We also

identified strong correlations and significant QTL overlap within trait categories, but few

associations across different types of traits. The exceptions however, between certain

aspects of floral morphology and nectar traits (volume, and both 'a' and 'b' color), are

consistent with existing trait associations that are observed across the genus, suggesting

that these could be examples of adaptive pleiotropy. Although we did not directly assess

the role of selection, our data indicate that the rapid floral trait evolution observed in this

group could have been facilitated by a relatively simple genetic basis for individual floral

traits, and a general absence of antagonistic pleiotropy among different types of

reproductive traits.

Few genetic changes could underlie floral trait shifts

The relatively simple genetic architecture that we detect for most of our floral

traits might be one mechanism that has permitted rapid floral evolution within the genus.

Indeed, our inference that few QTL underlie corolla depth agrees with comparative

development data from these species (Kostyun *et al*., 2017) in which we observe that

relatively simple heterochronic changes in corolla trait growth rates distinguish these

rotate vs. tubular corolla forms. Interestingly, our findings are also consistent with

previous studies of floral trait genetics between closely related species (Smith, 2016), as

well as a quantitative genetics study of floral trait variation between two ecotypes of

Jaltomata procumbens (Mione & Anderson, 2017). In particular, Mione and Anderson

(2017) estimated 1-4 loci contributed to various floral traits, as well as petal spot

patterning. In other systems, one or few QTL have been found for species differences in

nectar volume (e.g. Bradshaw *et al*., 1998; Stuurman *et al*., 2004; Wessinger *et al*., 2014;

but see Nakazato *et al*., 2013), similar to our inference of a single QTL for this trait. For

petal and nectar color (including Lightness, color 'a' and color 'b'), we identified 4 and 7

QTL, respectively, with at least 1 QTL per trait with moderate or large effects (i.e. 10 -

33% PVE) (**Table 2; Table S6**), similar to other systems that generally identify few loci

of large effect for petal color (e.g. Bradshaw *et al*., 1998; Wessinger *et al*., 2014).

Perhaps unlike these cases, however, it is likely that loci controlling color differences in

Jaltomata are regulators of pigment quantity rather than presence/absence biosynthesis,

because both nectar and petal color show gradation in the BC1s rather than discrete color

bins. Based on synteny with domestic tomato, our identified petal color QTL occur on the

same chromosomes as several known anthocyanin biosynthesis genes (i.e. DFR on ch2,

CHS1 and F3H on ch9, and ANS1 on ch10, in tomato), but do not appear to overlap with

the chromosomal locations of these genes. Preliminary data from a VIGS (virus-induced

gene silencing) pilot study in *J. sinuosa* do indicate that the purple petal pigment is an

anthocyanin however (J.L. Kostyun and J.C. Preston, unpub.), whereas for nectar color,

preliminary data suggest that an indole-flavin contributes to red pigment in *J. umbellata*

(J.L. Kostyun and D. Haak, unpub.), consistent with our inference that color variation is

unassociated between these different floral components.

In addition to relatively few contributing loci, many of the examined floral traits

also appear to be underpinned by additive effects (**Table 1; Figure 2; Figures S3-S6**),

while epistatic effects were comparatively rare (see below). Both are factors that might

also facilitate more rapid responses to selection. Other studies have similarly found that

floral size traits are often additive (Gottlieb, 1984), including within *Jaltomata* (Mione &

Anderson, 2017). Although several floral traits showed transgressive segregation within

our BC1s, which could indicate epistatic interactions, similar patterns can result from

unique combinations of additive alleles that have opposite effects in the parental species

(e.g. deVicente & Tanksley, 1993) and we identified individual QTL with these opposing

effects for all traits with transgressive segregation (**Table S6; Figures S3-5**). In addition,

our models only detected significant interactions among QTL for ovary diameter and

nectar color 'a' (**Table S6**), consistent with a general lack of epistatic interactions for

floral traits. Therefore, while we may not have uncovered all epistatic interactions

contributing to our traits--because some QTL might be undetected in our study, and our

models did not assess between trait interactions--both QTL and overall segregation

patterns suggest that floral traits are primarily underpinned by additive rather than

epistatic effects.

In contrast, many of the fertility traits had segregation patterns consistent with

epistatic interactions. BC1 individuals tended to have lower seed set and poorer quality

seeds (decreased viability and response to germination-inducing stimuli), and the

recombinant BC population contained a subset of highly sterile individuals. The

segregation of recombinant individuals with reduced viability and fertility often occurs in

hybrids (Baack *et al*., 2015), including in hybrids from additional *Jaltomata* species pairs

(Kostyun & Moyle, 2017). Such patterns are typically due to deleterious epistatic

interactions between loci that have diverged between the two parental lineages, as has

been shown in close relatives including tomatoes (Moyle & Nakazato, 2008; Sherman *et*

al., 2014). These observations in *Jaltomata* are similarly consistent with a specific role

for epistasis among incompatible alleles in the expression of postzygotic reproductive

isolation.

Reduced constraints may also have facilitated rapid floral trait evolution

Because rapid floral evolution may occur either through a lack of antagonistic

pleiotropy or through adaptive pleiotropy, we assessed evidence for these potential

mechanisms within *Jaltomata*. Within trait categories, we detected positive but modest

associations between several floral size traits, and among biologically related fertility

traits (e.g. fruit size and seed set) (**Figure 3; Table S3**), as well as significant co-

localization of QTL for these groups of traits (**Tables S7-S8**). Morphological associations

in particular suggest that shared growth regulators (e.g. Sicard & Lenhard, 2011; Brock *et*

al., 2012) contribute to observed variation in floral organ sizes, but do not completely

determine it; that is, overall floral size appears to share growth regulators (potentially

represented by co-localized QTL among floral size traits), but different floral organs also

appear to have unique growth regulator(s) (consistent with unique QTL). In contrast, we

detected fewer instances of strong trait correlations and QTL co-localization between

different trait categories (**Tables S3+S7-S8**). This general lack of antagonistic pleiotropy

among different classes of floral and fertility traits may have facilitated rapid floral

evolution in this system by minimizing constraints on the available combinations of floral

traits.

Despite this general pattern, we did identify several instances of QTL co-

localization that might represent adaptive pleiotropy, specifically among different corolla

and nectar traits. Overall, these associations would have produced flowers with larger

corolla limbs, deeper corolla tubes, and a greater volume of darker colored nectar (**Table**

2; Figure 4; Tables S6+S8). Interestingly, this trait combination (large flowers with

copious dark nectar) is actually not exhibited by either parental species used in this

experiment (**Figure 1; Table 1**); however, it is found in numerous other *Jaltomata*

species and is consistent with suspected pollination syndromes within the genus (see

below; Miller *et al*., 2011; Kostyun & Moyle, 2017). Correlated changes in corolla size

and nectar volume have also been observed in several other systems, including those with

documented or hypothesized pollinator shifts (e.g. *Petunia*, *Mimulus*, and *Penstemon*;

Galliot et al. 2006; reviewed in Smith, 2016).

Ecological context for rapid floral change in **Jaltomata**

Overall, our findings suggest potential mechanistic explanations for the evolution

of remarkable floral trait diversity among *Jaltomata* species within the last 5 million

years (Sarkinen *et al*., 2013). Traits with a relatively simple genetic basis that are

uncoupled from other floral and fertility traits have fewer mechanistic constraints, and

therefore can more rapidly respond to selective opportunities as they arise (given

sufficient standing or new genetic variation) (Agrawal & Stinchcombe, 2009; Smith,

2016). Although we have not yet directly assessed the role of selection in shaping floral

differences among species, several features of *Jaltomata* floral biology are consistent

with pollinator-mediated selection on floral traits (van der Niet & Johnson, 2012), likely

in conjunction with mating-system related changes (Goodwillie *et al*., 2010). First, floral

trait variation within *Jaltomata* shows clear hallmarks of selection imposed by pollinator

differentiation. Of the two main lineages within the genus, nearly all species in the

Central and South American clade have relatively small, ancestrally rotate flowers with

small amounts of lightly colored nectar (Wu et al. 2018), and hymenopterans have been

observed visiting several of these species in their native range (T. Mione, pers. comm.).

In contrast, many species in the clade that is restricted to South America—including *J.*

umbellata examined here—have floral features associated with attraction of vertebrate

pollinators (Fenster *et al*., 2004). Several of these species with larger flowers, a highly

fused corolla (either campanulate or tubular), and copious amounts of darkly colored

nectar are visited by hummingbirds (T. Mione, per. comm.); intriguingly, this repeated

natural trait covariation is consistent with the genetic association between floral size and

nectar traits we identified here.

531    These features indicate that shifts among different types of pollinators are a likely

532 source of selection for floral differentiation among species within *Jaltomata*, however,

533 mating system variation in the genus may also have facilitated these transitions. Self-

534 incompatibility is the ancestral state in the Solanaceae (Steinbachs & Holsinger, 2002)

535 and is broadly persistent in close relatives *Solanum* and *Capsicum* (Goldberg *et al*.,

536 2010). In contrast, all examined *Jaltomata* species are self-compatible (Mione, 1992;

537 Kostyun & Moyle, 2017; J.L. Kostyun & T. Mione, unpub.), indicating that gametophytic

538 self-incompatibility was lost early in the evolution of this clade. The presence of delayed

539 selfing and strong herkogamy in many species (e.g. Mione *et al*., 2015; Mione *et al*.,

540 2019), in addition to field observations of pollinators (above; T. Mione, pers. comm.),

541 indicate that species most likely employ a mixed mating strategy in their native ranges.

542 Mixed mating strategies are generally observed to maintain the largest amount of floral

543 trait variation, compared to predominant selfing or enforced outcrossing (Goodwillie *et*

544 *al*., 2005; Rosas-Guerrero *et al.,* 2014). In addition, they have been predicted to facilitate

545 pollinator shifts--especially to pollinators that might be more efficient but potentially

546 unreliable (such as hummingbirds)—because they allow reproductive assurance (via

547 selfing, when pollinators are limited) and increase the expression of new floral trait

548 variation controlled by recessive alleles (Goodwillie *et al*., 2005; Brys *et al*., 2013;

549 Wessinger & Kelly, 2018). Notably, our data indicate that dark/red colored nectar is at

550 least partially recessive (**Figure 2**), and that red petal pigmentation is completely

551 recessive (**Figure S7**), consistent with this novel variation being based on new recessive

552 alleles. Together, these observations are intriguing as they suggest that mixed mating, in

553 conjunction with the specific genetic architecture of floral trait variation, might have

facilitated the evolution of new floral trait variation in *Jaltomata*.

CONCLUSIONS

Genetic correlations among floral traits, especially those due to pleiotropic

effects, likely shape permitted trajectories of floral evolution. To assess how such genetic

associations might have contributed to observed patterns of floral diversity in *Jaltomata*,

we examined segregation patterns and genetic architecture of 25 floral and fertility traits

in a hybrid (BC1) population generated from parents with divergent floral traits. Our data

are consistent with several mechanisms that could have allowed rapid floral trait

evolution in this system: a largely simple genetic basis underlying variation in most of

our floral traits, a general absence of antagonistic pleiotropy constraining floral evolution,

and a potential instance of adaptive pleiotropy governing floral size and nectar traits. This

genetic architecture, in combination with pollinator-mediated selection on a background

of self-compatible mixed mating, might have uniquely positioned this genus for the rapid

floral diversification now evident within *Jaltomata*.

ACKNOWLEDGEMENTS

We thank the IU greenhouse staff for plant care, CJ Jewell and David Haak for

logistical support, undergraduate research assistants (especially Meret Thomas-Huebner,

Devki Shukla, and Shachia Jackson) for assistance with data collection, Tim Leslie for

assistance with simulations, and Mark Rausher and two anonymous reviewers for

feedback on a previous version of the manuscript. This work was supported by the IU

Biology Department, National Science Foundation Award (NSF DEB 1136707) to LCM,

and National Science Foundation Graduate Research Fellowship Program (NSF DEB

1342962) and Doctoral Dissertation Improvement Grant (NSF DEB 1601078) to JLK.

AUTHOR CONTRIBUTIONS

JLK and LCM designed the experiment, JLK generated experimental materials,

JLK and CMK collected phenotypic data, JLK and MJSG analyzed the data, and JLK and

LCM wrote the paper with input from CMK and MJSG.

REFERENCES

Agrawal AF, Stinchcombe JR. 2009. How much do genetic covariances alter the rate of

adaptation? *Proc R Soc Lond Biol* **276**:1183-1191.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Armbruster WS, Hansen TF, Pelabon C, Perez-Barrales R, Maad J. 2009. The

adaptive accuracy of flowers: measurement and microevolutionary patterns. *Ann

Bot* **103**:1529-1545.

Armbruster WS, Pelabon C, Bolstad GH, Hansen TF. 2014. Integrated phenotypes:

understanding trait covariation in plants and animals. *Proc R Soc Lond Biol* **369**.

Baack E, Melo MC, Rieseberg LH, Ortiz-Barrientos D. 2015. The origins of

reproductive isolation in plants. *New Phytol* **207**:968-984.

Bartlett ME. 2017. Changing MADS-Box Transcription Factor Protein–Protein

Interactions as a Mechanism for Generating Floral Morphological Diversity.

Integrative and Comparative Biology **57**:1312-1321.

Bradshaw HD, Otto KG, Frewen BE, McKay JK, Schemske DW. 1998. Quantitative

trait loci affecting differences in floral morphology between two species of

monkeyflower (*Mimulus*). *Genetics* **149**:367-382.

Brock MT, Kover PX, Weinig C. 2012. Natural variation in GA1 associates with floral

morphology in *Arabidopsis thaliana*. *New Phytol* **195**:58-70.

Broman KW. 2001. Review of statistical methods for QTL mapping in experimental

crosses. *Lab Animal (NY)* **30**:44-52.

Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental

crosses. *Bioinformatics* 19:889-890.

Brys R, Geens B, Beeckman T, Jacquemyn H. 2013. Differences in dichogamy and

herkogamy contribute to higher selfing in contrasting environments in the annual

Blackstonia perfoliata (Gentianaceae). *Ann Bot* **111**:651-661.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: A flexible trimmer for Illumina

Sequence Data. *Bioinformatics*, btu170.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013. Stacks: an

analysis tool set for population genomics. *Mol Ecol* 22: 3124-3140.

Chiarini FE, Lipari D, Barboza GE, Knapp S. 2017. IAPT/IOPB chromosome data

25. *Taxon* 66: 1246–1252.

Conner JK. 2002. Genetic mechanisms of floral trait correlations in a natural population.

Nature **420**:407-410.

Conner JK, Cooper IA, La Rosa RJ, Perez SG, Royer AM. 2014. Patterns of

phenotypic correlations among morphological traits across plants and animals.

Phil. Trans. R. Soc. B **369**.

deVicente MC, Tanksley SD. 1993. QTL analysis of transgressive segregation in an

interspecific tomato cross. *Genetics* **134**:585-596.

Ettensohn CA. 2013. Encoding anatomy: Developmental gene regulatory networks and

morphogenesis. *Genesis* **51**:383-409.

Farooq M, Basra SMA, Ahmad N, Hafeez K. 2005. Thermal hardening: A new seed

vigor enhancement tool in rice. *J Integr Plant Biol* **47**:187-193.

Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination

syndromes and floral specialization. *Annu Rev Ecol Evol Syst* **35**:375-403.

Galliot C, Hoballah ME, Kuhlemeier C, Stuurman J. 2006. Genetics of flower size

and nectar volume in *Petunia* pollination syndromes. *Planta* **225**:203-212.

Garcia JE, Greentree AD, Shrestha M, Dorin A, Dyer AG. 2014. Flower Colours

through the Lens: Quantitative Measurement with Visible and Ultraviolet Digital

Photography. *Plos One* **9**.

Goldberg EE, Kohn JR, Lande R, K. Robertson KA, Smith SA, Igic B. 2010. Species

selection maintains self-incompatibility. *Science* **330**:493-495.

Goodwillie C, Kalisz S, Eckert CG. 2005. The evolutionary enigma of mixed mating

systems in plants: Occurrence, theoretical explanations, and empirical evidence.

Annu Rev Ecol Evol Syst **36**:47-79.

Goodwillie C, Sargent RD, Eckert CG, Elle E, Geber MA, Johnston MO, Kalisz S,

Moeller DA, Ree RH, Vallejo-Marin M et al. 2010. Correlated evolution of

mating system and floral display traits in flowering plants and its implications for

the distribution of mating system variation. *New Phytol* **185**:311-321.

Gottlieb LD. 1984. Genetics and morphological evolution in plants. *Am Nat* **123**:681-

709.

Hermann K, Klahre U, Moser M, Sheehan H, Mandel T, Kuhlemeier C. 2013. Tight

Genetic Linkage of Prezygotic Barrier Loci Creates a Multifunctional Speciation

Island in *Petunia*. *Current Biology* **23**:873-877.

Jewell C, Papineau AD, Freyre R, Moyle LC. 2012. Patterns of reproductive isolation

in *Nolana* (chilean bellflower). *Evolution* **66**:2628-2636.

Kendal D, Hauser CE, Garrard GE, Jellinek S, Giljohann KM, Moore JL. 2013.

Quantifying Plant Colour and Colour Difference as Perceived by Humans Using

Digital Images. *Plos One* **8**.

Knapp S. 2010. On 'various contrivances': pollination, phylogeny and flower form in the

Solanaceae. *Phil Trans R Soc B* **365**:449-460.

Kostyun JL, Moyle LC. 2017. Multiple strong postmating and intrinsic postzygotic

reproductive barriers isolate florally diverse species of *Jaltomata* (Solanaceae).

Evolution **71**:1556-1571.

Kostyun JL, Preston JC, Moyle LC. 2017. Heterochronic developmental shifts underlie

floral diversity within *Jaltomata* (Solanaceae). *EvoDevo* **8**:17.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv:1303.3997v1 [q-bio.GN].

Manceau M, Domingues VS, Mallarino R, Hoekstra HE. 2011. The Developmental

Role of Agouti in Color Pattern Evolution. *Science* **331**:1062-1065.

Miller RJ, Mione T, Phan HL, Olmstead RG. 2011. Color by Numbers: Nuclear Gene

Phylogeny of *Jaltomata* (Solanaceae), Sister Genus to *Solanum*, Supports Three

Clades Differing in Fruit Color. *Syst Botany* **36**:153-162.

Mione T. 1992. *The systematics and evolution of* Jaltomata *(Solanaceae)*. PhD

Dissertation, The University of Connecticut. Available from ProQuest

Dissertations & Theses Global. (303964972).

Mione T, Anderson GJ. **2017.** Genetics of floral traits of *Jaltomata procumbens*

(Solanaceae). *Brittonia* **69**:1-10.

Mione T, Anderson GJ, Nee M. 1993. *Jaltomata* I: Circumscription, description and

new combinations for five South American species. *Brittonia* **45**: 138-145.

Mione T, Leiva González S, Yacher L. 2015. Two new Peruvian species of *Jaltomata*

(Solanaceae, Solaneae) with red floral nectar. *Brittonia* **67**:105-112.

Mione T, Plourd KC, Leiva González S, Yacher L, Philbrick CT, Blume D. 2017.

Pollination decreases longevity of the protogynous flowers of *Jaltomata sinuosa*

(Solanaceae). *J Biol Nat* **7**:123-129.

Mione T, Kostyun JL, Leiva González S. 2019. Breeding system features and a novel

method for locating the floral nectar secretion in a South American nightshade

(*Jaltomata quipuscoae*, Solanaceae). *Plant Biosystems* March:1-10.

https://doi.org/10.1080/11263504.2019.1578277.

Moyle LC, Nakazato T. 2008. Comparative genetics of hybrid incompatibility: Sterility

in two *Solanum* species crosses. *Genetics* **179**:1437-1453.

Nakazato T, Rieseberg LH, Wood TE. 2013. The genetic basis of speciation in the

Giliopsis lineage of *Ipomopsis* (Polemoniaceae). *Heredity* **111**:227-237.

Olmstead RG, Bohs L, Migid HA, Santiago-Valentin E, Garcia VF, Collier SM.

2008. A molecular phylogeny of the Solanaceae. *Taxon* **57**:1159-1181.

Preston JC, Hileman LC, Cubas P. 2011. Reduce, reuse, and recycle: developmental

evolution of trait diversification. *Am. J. Bot.* **98**:397-403.

RAW Therapee Team. 2012. Raw Therapee: cross-platform raw image processing

program. http://rawtherapee.com/.

Rijpkema A, Gerats T, Vandenbussche M. 2006. Genetics of floral development in

Petunia. In Soltis DE, Soltis PS, Callow JA, eds. *Adv Bot Res: Developmental*

Genetics of the Flower. Elsevier Ltd, 237-278.

Rosas-Guerrero V, Aguilar T, Marten-Rodriguez S, Ashworth L, Lopezaraiza-

Mikel M, Bastida JM, Quesada M. 2014. A quantitative review of pollination

syndromes: do floral traits predict effective pollinators? *Ecol. Lett* **17**:388-400.

Sarkinen T, Bohs L, Olmstead RG, Knapp S. 2013. A phylogenetic framework for

evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. *BMC*

Evolutionary Biology **13**.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of

image analysis. *Nature Methods* **9**:671-675.

Schwartzwald D. 2012. Color Space Converter Plugin for ImageJ.

https://imagej.nih.gov/ij/plugins/color-space-converter.html.

Sherman NA, Victorine A, Wang RJ, Moyle LC. 2014. Interspecific Tests of Allelism

Reveal the Evolutionary Timing and Pattern of Accumulation of Reproductive

Isolation Mutations. *Plos Genetics* **10**.

Shrestha R, Gomez-Ariza J, Brambilla V, Fornara F. 2014. Molecular control of

seasonal flowering in rice, arabidopsis and temperate cereals. *Ann Bot* **114**:1445-

1458.

Sicard A, Lenhard M. 2011. The selfing syndrome: a model for studying the genetic

and evolutionary basis of morphological adaptation in plants. *Ann Bot* **107**:1433-

1443.

Smaczniak C, Immink RGH, Angenent GC, Kaufmann K. 2012. Developmental and

evolutionary diversity of plant MADS-domain factors: insights from recent

studies. *Development* **139**:3081-3098.

Smith SD. 2016. Pleiotropy and the evolution of floral integration. *New Phytol* **209**:80-

85.

Stebbins GL. 1974. Flowering plants: evolution above the species level. Harvard

University Press, Cambridge, MA.

Steinbachs JE, Holsinger KE. 2002. S-RNase-mediated gametophytic self

incompatibility is ancestral in eudicots. *Mol Biol Evol* **19**: 825–829.

Stuurman J, Hoballah ME, Broger L, Moore J, Basten C, Kuhlemeier C. 2004.

Dissection of floral pollination syndromes in *Petunia*. *Genetics* **168**:1585-1599.

Taylor J, Butler D. 2017. R Package ASMap: Efficient Genetic Linkage Map

Construction and Diagnosis. *Journal of Statistical Software* **79**:1-29.

Tomato Genome Consortium, The. 2012. The tomato genome sequence provides

insights into fleshy fruit evolution. *Nature* **485**:635-641.

van der Niet T, Johnson SD. 2012. Phylogenetic evidence for pollinator-driven

diversification of angiosperms. *Trends Ecol Evolut* **27**:353-361.

Wagner GP, Zhang JZ. 2011. The pleiotropic structure of the genotype-phenotype map:

the evolvability of complex organisms. *Nature Reviews Genetics* **12**:204-213.

Wessinger CA, Hileman LC, Rausher MD. 2014. Identification of major quantitative

trait loci underlying floral pollination syndrome divergence in *Penstemon*. *Phil.*

Trans. R. Soc. B **369**.

Wessinger CA, Freeman CC, Mort ME, Rausher MD, Hileman LC. 2016.

Multiplexed shotgun genotyping resolves species relationships within the North

American genus *Penstemon*. *Am J Bot* **103**:912-922.

Wessinger CA, Kelly JK. 2018. Selfing can facilitate transitions between pollination

syndromes. *Am Nat* **191**.

Wu M, Kostyun JL, Hahn MW, Moyle LC. 2018. Dissecting the basis of novel trait

evolution in a radiation with widespread phylogenetic discordance. *Mol Ecol* **27**:

3301-3316.

Wu M, Kostyun JL, Moyle LC. 2019. Genome sequence of *Jaltomata* addresses rapid

reproductive trait evolution and enhances comparative genomics in the hyper-

diverse Solanaceae. *Genome Biology and Evolution* in press. Available at:

https://www.biorxiv.org/content/early/2018/05/30/335117.

Xu SQ, Schluter PM. 2015. Modeling the two-locus architecture of divergent pollinator

adaptation: how variation in SAD paralogs affects fitness and evolutionary

divergence in sexually deceptive orchids. *Ecology and Evolution* **5**:493-502.

761 **Table 1**. Summary statistics for floral and fertility traits in parental species, F1s, and

762 BC1s. Phenotypic means and variances provided, while significant differences between

763 parental species were assessed with t-tests on transformed data as appropriate (see

764 Methods). Note that differences in seed germination rates were not tested. * p<0.05, **

765 p<0.001; *** p<0.0001.

Trait	*J. sinuosa* (n=7)		*J. umbellata* (n=5)	
Floral Morphology	**Mean**	**Variance**	**Mean**	**Variance**
Buds per Inflorescence	2.86**	0.03	8.60	3.24
Calyx Diameter (mm)	15.49**	0.08	8.14	2.91
Sepal Length (mm)	7.48**	0.02	4.41	0.56
Corolla Diameter (mm)	29.82***	1.92	15.32	5.70
Corolla Depth (mm)	1.38**	0.07	10.25	5.52
Corolla Fusion (mm)	9.45	0.11	10.33	1.55
Corolla Fusion Proportion	0.63*	0.001	0.72	0.001
Petal Length (mm)	14.85	0.41	14.40	0.16
Stamen Length (mm)	10.80	0.29	10.45	0.22
Anther Length (mm)	2.03*	0.02	1.73	0.04
Ovary Diameter (mm)	2.28***	0.03	1.37	0.05
Style Length (mm)	7.90***	0.17	14.64	1.72
Herkogamy (mm)	-2.20***	0.01	5.37	4.62
Floral Color + Physiology				
Nectar Color - L	83.49***	20.32	50.42***	31.63
Nectar Color - a	-1.55***	0.95	57.61***	7.10
Nectar Color - b	3.47***	8.20	71.84***	59.31
Petal Color - L	83.18	2.59	84.15	4.86
Petal Color - a	0.81*	0.02	-0.93*	0.62
Petal Color -b	-1.26***	0.02	6.71***	1.05
Nectar Volume (uL)	6.86***	1.07	17.80	13.20
Fertility				
Viable Pollen Grains	25000	36800000	27000	60770000
Proportion Viable Pollen	0.76	0.01	0.71	0.02
Fruit Set	0.93*	0.01	0.55	0.06
Fruit Mass (g)	0.47**	0.01	0.12	0.01
Fruit Diameter (cm)	1.01*	0.01	0.61	0.02
Seed Set	74.87*	256.92	30.44	494.43
Viable Seed Set	74.87*	256.92	23.59	182.31
Viable Seed T50	9.33	--	7.00	--
Viable Seed MGT	22.00	--	14.00	--

769 **Table 1 cont.**

F1s (n=13)		BC1s (n=224)	
Mean	Variance	Mean	Variance
3.95	0.68	3.51	1.63
10.55	0.48	12.10	1.08
5.45	0.13	6.16	0.28
24.58	6.84	28.97	9.92
6.64	0.58	5.22	1.35
9.81	0.40	9.92	1.24
0.67	0.001	0.63	0.002
14.65	1.61	15.78	2.88
10.81	0.59	10.86	1.26
2.01	0.01	2.13	0.04
1.62	0.02	1.82	0.04
12.39	1.04	10.82	1.97
2.95	1.08	1.20	1.02
81.31	14.35	83.99	13.42
-0.05	101.74	-4.29	14.91
59.85	75.94	22.91	287.97
86.93	5.82	83.56	4.60
-0.81	0.19	-0.04	0.36
2.89	3.13	0.75	1.29
11.90	12.32	10.76	18.87
22513	16602405	26886	145318716
0.82	0.00	0.74	0.03
0.93	0.01	0.89	0.03
0.24	0.01	0.29	0.01
0.80	0.01	0.85	0.03
36.06	168.37	37.67	361.12
25.46	303.95	26.28	340.08
34.00	1001.78	26.70	607.37
46.04	622.37	39.36	458.11

Table 2. QTL for key floral and fertility traits. Includes peak location and LOD of each

QTL, 1.5 LOD intervals, proportion phenotypic variance explained, phenotypic effect

size and standard error, proportion of parental trait difference explained, and whether the

effect is aligned with parental trait values. Comprehensive QTL data, including the full

models for all traits, are provided in Table S6. Note: effect sizes are given as the

difference between homozygote and heterozygote mean phenotypes (i.e. the effect of a

single paternal allele at that QTL), while the proportion of parental trait difference

corresponds to the relative homozygous effect (RHE).

Trait	Linkage Group	Peak Location (1.5 LOD Interval)	Peak LOD
Flower Morphology			
Calyx Diameter (3)	1	56 (49-81)	9.06
	8	84.3 (69-92.55)	3.15
	9	66 (17-85)	5.59
Corolla Diameter (3)	4	21.1 (13-70)	3.50
	8	73.1 (67-84)	5.87
	12	37.4 (0-65.17)	1.57
Corolla Depth (4)	5	40.8 (20-46)	10.19
	7	52 (48-60.67)	17.95
	9	30 (11-38)	12.38
	10	55.8 (41-91)	4.48
Corolla Fusion Prop. (4)	1	72 (57-106)	4.60
	4	26 (19-84)	3.77
	5	86 (80.21-99)	3.57
	9	16 (0-67)	7.19
Petal Length (3)	7	100 (25-100.67)	2.46
	8	73 (39-92.55)	3.55
	12	48.2 (0-65.17)	2.04
Stamen Length (4)	2	37.5 (6-61)	4.54
	4	120.8 (100-133)	3.77
	7	91 (59-100.67)	4.45
	9	0 (0-33)	5.32
Anther Length (1)	1	18 (8-35.87)	3.78

786 **Table 2. cont.**

Ovary Diameter (4)	1	68 (60-107)	5.77
	5	72 (0-84)	2.53
	7	9.9 (0-25)	3.56
	12	48.2 (5-65.17)	4.31
Style Length (2)	2	95.2 (55-126)	2.79
	9	22 (1-30)	9.89
Flower Physiology			
Nectar Volume (1)	3	187.2 (172-187.19)	2.81
Flower Color			
Nectar L (2)	2	154.9 (147-171.87)	2.97
	5	103.9 (74-115.35)	6.44
Nectar a (4)	3	66.1 (60-95)	7.69
	4	14 (0-23.74)	3.99
	7	29.1 (10-42)	2.32
	7	56.2 (43-63)	3.54
Nectar b (4)	2	152 (141-171.87)	4.50
	3	186 (164-187.19)	2.83
	5	100 (94-111)	11.00
	7	26 (20.02-32)	9.02
Petal L (2)	4	23.7 (0-81)	2.40
	5	65.3 (58-71)	3.83
Petal a (4)	5	62 (58-68)	12.11
	8	52 (49-52.48)	8.54
	8	52.5 (52-53)	8.13
	9	64.1 (44-76)	9.89
Petal b (3)	5	64 (60-69)	21.46
	9	52 (52-54)	5.60
	9	55.8 (54.05-57)	4.22
Fertility			
Fruit Diameter (2)	1	80 (61-103)	4.67
	11	76.9 (51-79.93)	3.09
Fruit Mass (4)	1	96 (60-146)	5.95
	3	170 (48-187.19)	2.99
	11	76.5 (54-79.93)	4.19
	12	0 (0-10.64)	4.66
Seed Set (2)	11	76 (63-79.93)	6.64
	12	0 (0-63)	2.86
Viable Seed Set (1)	11	76.9 (68-79.93)	4.42

789   **Table 2. cont.**

%PVE	Effect ± SE	%Parental Trait Difference (RHE)	Aligned with Parental Value?
13.67	-0.803 ± 0.120	0.22	Y
4.46	-0.457 ± 0.120	0.12	Y
8.13	-0.598 ± 0.116	0.16	Y
4.47	1.387 ± 0.344	-0.19	N
7.69	-1.902 ± 0.360	0.26	Y
1.96	-0.963 ± 0.361	0.13	Y
11.20	0.844 ± 0.119	0.19	Y
21.47	1.146 ± 0.116	0.26	Y
13.94	0.942 ± 0.119	0.21	Y
4.64	-0.521 ± 0.114	-0.12	N
7.02	-0.022 ± 0.005	-0.56	N
5.70	-0.021 ± 0.005	-0.53	N
5.39	0.022 ± 0.005	0.56	Y
11.27	0.029 ± 0.005	0.74	Y
3.29	0.018 ± 0.005	-2.66	N
4.78	-0.022 ± 0.005	3.32	Y
2.71	-0.017 ± 0.005	2.50	Y
5.80	-0.548 ± 0.119	3.18	Y
4.78	-0.499 ± 0.119	2.90	Y
5.68	0.551 ± 0.121	-3.20	N
6.86	0.619 ± 0.124	-3.60	N
6.48	-0.116 ± 0.027	0.78	Y

798   **Table 2. cont.**

9.21	-0.120 ± 0.023	0.26	Y
3.90	0.092 ± 0.027	-0.20	N
5.56	-0.094 ± 0.023	0.21	Y
6.78	-0.104 ± 0.023	0.23	Y
3.80	0.022 ± 0.006	0.16	Y
14.51	0.044 ± 0.006	0.33	Y
5.22	0.098 ± 0.027	0.48	Y
5.12	-0.008 ± 0.002	0.08	Y
11.51	-0.014 ± 0.003	0.13	Y
13.04	0.062 ± 0.010	0.23	Y
6.51	-0.043 ± 0.010	-0.16	N
3.72	-0.040 ± 0.012	-0.15	N
5.74	0.050 ± 0.012	0.18	Y
5.25	0.161 ± 0.036	0.27	Y
3.24	0.138 ± 0.038	0.23	Y
13.74	0.288 ± 0.039	0.48	Y
11.03	0.230 ± 0.035	0.39	Y
4.43	0.005 ± 0.002	2.03	Y
7.19	0.007 ± 0.002	2.92	Y
15.28	-0.122 ± 0.016	0.44	Y
10.35	-1.355 ± 0.210	4.81	Y
9.82	1.312 ± 0.209	-4.65	N
12.18	-0.097 ± 0.014	0.35	Y
32.83	0.210 ± 0.019	0.56	Y
7.20	0.486 ± 0.094	1.31	Y
5.35	-0.415 ± 0.093	-1.11	N
8.64	-0.026 ± 0.005	0.52	Y
5.61	-0.020 ± 0.005	0.40	Y
9.35	-0.026 ± 0.005	0.44	Y
4.56	-0.018 ± 0.002	0.31	Y
6.46	-0.021 ± 0.005	0.35	Y
7.22	-0.023 ± 0.005	0.40	Y
12.28	-0.157 ± 0.028	0.65	Y
5.07	-0.109 ± 0.030	0.45	Y
8.85	-0.211 ± 0.046	0.75	Y

Figure Legends.

Figure 1. Representative flowers of the parental species and their F1 hybrid.

Figure 2. Key trait distributions within the BC1 population, compared to phenotypic

means for *J. sinuosa* (purple line), *J. umbellata* (red line), and their F1s (pink line).

Figure 3. Key floral trait correlations within the BC1 mapping population. Scatterplots

are provided below the diagonal, while Spearman's correlation coefficients and

associated p-values are above the diagonal. Statistically significant correlations following

Bonferroni correction ($p < 5.945 \times 10^{-5}$) are highlighted in red. Correlation values for all

traits are provided in Table S3, and for PC traits provided in Table S5.

Figure 4. Linkage map and distribution of identified QTL (including 1.5 LOD intervals)

for 12 key floral and fertility traits.

Supplementary Information

Table S1. Accession information for material used and generated in this study.

Table S2. Trait measurements for all individuals phenotyped.

Table S3. Correlations among all measured traits within BC1 individuals.

Table S4. Principal component loadings for highly correlated traits.

Table S5. Correlations among PC traits within BC1 individuals

Table S6. Full QTL models for all examined traits.

Table S7. Expected and observed counts of QTL overlap.

Table S8. Identified instances of QTL co-localization, based on overlapping 1.5 LOD

intervals.

Figure S1. Measured morphological traits on mature flowers.

Figure S2: Biplots following PCA on floral morphological traits

Figures S3-S4: Distributions for floral morphological traits within the mapping

population.

Figure S5. Distributions for physiological and color traits within the mapping population.

Figure S6. Distributions for fertility traits within the mapping population.

Figure S7. Representative examples of petal and nectar color variation among F1

individuals, compared to either parent.

Figure S8. Comparison of observed number of QTL overlaps with counts from 10000

randomly generated simulations, for QTL co-localization overlap within and between

trait categories (morphology, color, and fertility).

J. sinuosa   F1   *J. umbellata*

1 cm