Title: Chloroplasts navigate towards the pathogen interface to counteract

infection by the Irish potato famine pathogen

Authors: Alexia Toufexi[1†], Cian Duggan[*1†], Pooja Pandey[1], Zachary Savage[1], María

Eugenia Segretin[3], Lok Him Yuen[1], David C. A. Gaboriau[2], Alexandre Y. Leary[1],

Virendrasinh Khandare[1], Andrew D. Ward[4], Stanley W. Botchway[4], Benji C. Bateman[4],

Indranil Pan[5,6], Martin Schattat[7], Imogen Sparkes[8], Tolga O. Bozkurt[1*]

Affiliations:

[1]Department of Life Sciences, Imperial College London, UK

[2]Facility for Imaging by Light Microscopy, NHLI, Faculty of Medicine, Imperial College

London

[3]INGEBI-CONICET, Ciudad Autonoma de Buenos Aires, Argentina

[4]Central Laser Facility, Science and Technology Facilities Council Harwell, UK

[5]Centre for Process systems engineering and Centre for Environmental Policy, Imperial

College London, UK

[6]The Alan Turing Institute, London, UK

[7]Martin-Luther-Universität Halle-Wittenberg, Germany

[8]School of Biological Sciences, University of Bristol, UK

*Correspondence to: o.bozkurt@imperial.ac.uk

[†]These authors contributed equally

Abstract: Chloroplasts are light harvesting organelles that arose from ancient endosymbiotic cyanobacteria. Upon immune activation, chloroplasts switch off photosynthesis, produce anti-microbial compounds, and develop tubular extensions called stromules. We report that chloroplasts navigate to the pathogen interface to counteract infection by the Irish potato famine pathogen *Phytophthora infestans*, physically associating with the specialised membrane that engulfs pathogen haustoria. Outer envelope protein, chloroplast unusual positioning1 (CHUP1), anchors chloroplasts to the host-pathogen interface. Stromules are induced during infection in a CHUP1-dependent manner, embracing haustoria and interconnecting chloroplasts, to form dynamic organelle clusters. Infection-triggered reprogramming of chloroplasts relies on surface immune signalling, whereas pathogen effectors subvert these immune pulses. Chloroplast are deployed focally, and coordinate to restrict pathogen entry into plant cells, a process actively countered by parasite effectors.

Introduction: *Phytophthora infestans* is an oomycete pathogen that causes potato late blight, one of the most historically and economically devastating crop diseases. The pathogen penetrates host cells via haustoria, infection structures that extend from its extracellular invasive hyphae. Haustoria are surrounded by the plant-derived extra-haustorial membrane (EHM), across which effectors secreted by the pathogen translocate inside the host cell (Wang et al., 2017; Whisson et al., 2016, 2007). This interface is key to the success or failure of infection and is therefore targeted by focal immune responses of the plant (Bozkurt et al., 2011; Dagdas et al., 2018; Kwon et al., 2008). Remarkably, despite being continuous with the plasma membrane, there is a stark difference in the biochemical composition of the EHM and the plasma membrane. EHM typically lacks surface localized pattern recognition receptors (PRRs), which activate downstream immune responses through recognition of pathogen associated molecular patterns (PAMPs) (Bozkurt et al., 2015, 2014). Activation of immunity at the cell surface stimulates chloroplasts to shut down photosynthesis, synthesize defence hormone precursors, and generate reactive oxygen species (ROS) (Padmanabhan and Dinesh-Kumar, 2010; Su et al., 2018), suggesting that chloroplasts are committed to the plant defence system. Additionally, chloroplasts produce stroma filled tubules (stromules) that have been implicated in defence (Caplan et al., 2015; Kumar et al., 2018). Pathogens are known to target chloroplasts with effector proteins (de Torres Zabala et al., 2015; Jelenska et al., 2007; Pecrix et al., 2018; Petre et al., 2016). Interestingly, several genes associated with resistance to oomycete pathogens were found to encode chloroplast-localized proteins (Belhaj et al., 2009; van Damme et al., 2009). However, the molecular and physiological mechanism of how chloroplast weaponry is launched against invading pathogens is unclear.

Here we show that to counteract host cell invasion by the Irish potato famine pathogen

P. infestans, chloroplasts navigate to pathogen penetration sites. Photo-relocation

component chloroplast unusual positioning 1 (CHUP1) (Oikawa et al., 2008, 2003)

mediates chloroplast accumulation around the haustorium by facilitating anchoring to the

host-derived perimicrobial membrane interface. Moreover, stromule development

increases upon *P. infestans* infection, in a CHUP1 dependent manner. Stromules

embrace haustoria and mediate physical interactions between chloroplasts, forming

dynamic clusters around the pathogen interface. Notably, infection-triggered stromule

development relies on surface immune signalling, whereas the pathogen subverts this

remotely by switching off surface immune pulses. These results implicate chloroplast

photo-relocation machinery in plant focal immune responses and demonstrate that

chloroplasts play a direct role in antagonizing pathogen invasion.

Results & Discussion:

To gain insights into defence-related chloroplast functions, we investigated chloroplast dynamics during *P. infestans* infection of the solanaceous model plant *Nicotiana benthamiana*. Confocal microscopy of infected leaf epidermal cells stably expressing GFP in chloroplast stroma (CpGFP hereafter) revealed that chloroplasts accumulate at 58% of haustoria (*n* = 299 haustoria) (Fig. 1A-C). Notably, chloroplasts are mainly positioned around infection sites, sieging haustoria in a highly dynamic fashion (Fig. 1C, 1E and Movie S1-3). Furthermore, compared to non-infected controls, we recorded a 5-fold increase in stromule formation in infected CpGFP cells (Fig. 1A-B, 1D). Notably, chloroplasts seized haustoria, tightly embracing the EHM through stromules that extend and coil around the pathogen (Fig. 1A, 1C, Movie S3). Expression of the EHM marker protein RFP:REM1.3 (Bozkurt et al., 2014) in infected CpGFP plants, allowed us to collect further evidence that chloroplasts intimately associate with the EHM (Fig. S1, Movie S4). Remarkably, time lapse-microscopy showed that chloroplasts plunge towards haustoria and embrace the EHM following haustorial penetration (Fig. 1E, Movie S5, Fig. S2), suggesting that chloroplasts may participate in focal immunity.

To determine the extent to which chloroplasts physically associate with the EHM, we next employed optical tweezers in combination with Total Internal Fluorescence Microscopy (TIRF) in infected CpGFP plants. Using optical tweezers, we successfully trapped and moved 17% of chloroplasts (*n* = 29) in non-haustoriated cells, a distance greater than the 10 µm threshold. In comparison, we were unable to trap and move any chloroplasts (0%, *n* = 18) neighbouring haustoria past the threshold, indicating a strong association between the chloroplasts and the EHM. Consistent with this, we recorded a few instances where these chloroplasts were initially pulled away from the EHM, but before they passed the distance threshold they escaped the trap and sprang back

towards their former position (22%, *n* = 18) (Fig. 1G, Movie S6). Taken together, these

results demonstrate that chloroplasts mobilize towards haustoria and tightly embrace the

EHM through induction of stromules, possibly to increase the surface area of interaction,

in a manner similar to chloroplast-nucleus communication upon activation of antiviral

immunity (Kumar et al., 2018).

Remarkably, upon pathogen infection, we noted that chloroplasts frequently formed

dynamic clusters via stromules, occasionally bridging multiple haustoria (Fig. 1A, Fig.

S3, Movie S1, S7, S8). Optical trapping of chloroplasts in pathogen-challenged tissue

illustrated co-migration of chloroplast pairs interconnected by a stromule like extension

(Fig. 1H, and Movie S9), indicating that chloroplasts can be physically linked. Both

observations raise the possibility that chloroplasts might coordinate their defence-related

tasks cooperatively to respond to pathogen attack. However whether plastids can fuse to

form a continuous stromal compartment that enables macromolecule exchange thorough

stromules is debated (Hanson and Hines, 2017; Schattat et al., 2014).

Intriguingly, although rarely observed, we captured two time-lapse image series, which

show collapse of the haustoria during chloroplast steering (Fig. S4, Movie S10, S11).

This prompted the idea that in addition to their biochemical arsenal (de Torres Zabala et

al., 2015; Serrano et al., 2016; Trotta et al., 2014), chloroplasts plausibly harness

mechanical means to oppose pathogen penetration. Supporting this view, we recorded

time-lapse series of chloroplast clusters adjacent to haustoria which moved

simultaneously in the same direction (Movie S12, S13).

Fig. 1. Chloroplasts move towards haustoria and associate with the EHM via stromule induction. (A-B, E-F) Maximum projection confocal micrographs of *N. benthamiana* plants expressing GFP in chloroplast stroma (CpGFP). (A) Stromule

induction and chloroplast accumulation around haustoria with *P. infestans* strain 88069td

compared to (B) Mock inoculated cells. (C) Number of 88069td haustoria associated with

one or more chloroplast. (D) Scatter box-plot shows increased stromule induction (p <

0.01) in infected (*n* = 24 images quantified) vs uninfected (*n* = 24 images quantified)

tissue. (E) Maximum intensity projection time-lapse series showing movement of tracked

chloroplasts (open arrowheads) towards haustorium. Dashed line indicates approximate

path travelled. (F) Single plane image showing stromule (purple arrowhead) wrapped

around haustorium (white arrowhead). (G-H) GFP channel in grayscale from TIRF

microscope. Time-lapse showing laser capture of haustorium associated chloroplast in

CpGFP plant where automated the trapping routine, traps and attempts to move

chloroplast 10 μm. (G) Chloroplast which escapes laser trap, reaching the furthest

possible point from haustorium (3.9 s) before springing back (5.0 s). Dotted line shows

outline of haustorium marked by RFP:REM1.3. (H) Trapped chloroplast (asterisk) linked

by stromule to another chloroplast. When the trapped chloroplast moves, the linked

chloroplast co-migrates. Scale bars: 10 μm.

To elucidate the role of chloroplast positioning in plant cell invasion of *P. infestans*, we

investigated whether a relationship exists between light and pathogen-induced

chloroplast movements. We particularly focused on chloroplast unusual positioning 1

(CHUP1), a protein that regulates chloroplast photo-relocation, movement and anchoring

to the plasma membrane (Oikawa et al., 2008, 2003). Similar to a GFP fused chloroplast

envelope marker protein, translocon at the outer membrane of chloroplasts 64

(TOC64:GFP), CHUP1:GFP labelled the chloroplast outer envelope (Fig 2A & B).

However, CHUP1:GFP displayed an unusually increased fluorescence intensity at foci

across the chloroplast-plasma membrane interface (Fig. 2A). Notably, in haustoriated

cells, CHUP1:GFP accumulated at chloroplast-EHM contact sites (Fig. 2C, Fig. S5),

whereas TOC64:GFP uniformly labelled the chloroplast envelope without producing any

fluorescent foci adjacent to the EHM (Fig. 2D). The discrete punctate localisation of

CHUP1:GFP at the EHM contact sites suggested that CHUP1 could mediate anchoring

of chloroplasts to the EHM which typically lacks plasma membrane proteins (Bozkurt et

al., 2015, 2014; Whisson et al., 2016). These results indicate that CHUP1, a core

component of the chloroplast photo-relocation machinery, might be co-opted for

chloroplast recruitment to the pathogen interface.

Fig. 2. CHUP1 localizes to the chloroplast outer envelope and accumulates at contact points between plastids and the plasma membrane or EHM. Single-plane confocal microscope images of wild-type *N. benthamiana* transiently co-expressing plasma membrane and EHM marker RFP:REM1.3 with CHUP1:GFP (A & C) or TOC64:GFP (B & D). (A) CHUP1:GFP labels chloroplast outer membrane and accumulates at contact points (white arrowheads) with the plasma membrane. (B) TOC64:GFP distributes uniformly across the chloroplast outer membrane. (C) CHUP1:GFP accumulates at *P. infestans* EHM contact points. (D) TOC64:GFP displays a uniform distribution when in contact with the EHM. Scale bars: 10 µm.

We next set out to determine whether CHUP1 is required for pathogen directed

chloroplast movement and EHM docking. Remarkably, downregulation of *CHUP1*

expression through virus-induced gene silencing (VIGS) in infected CpGFP plants,

significantly reduced the number of haustoria that associate with chloroplasts (33%, *n* =

544 haustoria) compared to the silencing control (45%, *n* = 471 haustoria) (Fig. 3A-B,

Fig. S6). We obtained similar results following RNA interference (RNAi) mediated

knockdown of *CHUP1* in both wild-type *N. benthamiana* and CpGFP plants (Fig. S7).

These results indicate that CHUP1 facilitates chloroplast recruitment to the pathogen

interface, where it accumulates, likely through mediating chloroplast anchoring to the

EHM (Fig. 2C). In addition, VIGS mediated knock down of *CHUP1* reduces stromule

induction (8% chloroplasts with stromule(s), *n* = 68 images quantified) compared to

control silencing (2% chloroplasts with stromule(s), *n* = 68 images quantified),

suggesting that CHUP1 is required for pathogen induced stromule development (Fig.

3C). Likewise, in infected CpGFP plants, transient RNAi of CHUP1 led to a 2-fold

decrease in stromule induction (18% chloroplasts with stromule(s), *n* = 11 images

quantified) compared to the GUS-silencing control (39% chloroplasts with stromule(s), *n*

= 10 images quantified) (Fig. S7). Collectively, these results demonstrate that CHUP1 is

essential for both pathogen-directed chloroplast mobility and infection-triggered

stromules formation which is implicated in antimicrobial immunity (Caplan et al., 2015),

pointing to a positive role of CHUP1 in plant immunity.

We next tested the hypothesis that the process of chloroplast positioning and anchoring

to the pathogen interface enhances disease resistance. VIGS of *CHUP1* led to

considerably higher levels of *P. infestans* hyphal growth compared to the silencing

control (Fig. 3D-E). We repeated these infection assays following RNAi-mediated

silencing of *CHUP1*. Consistently, RNAi of *CHUP1* substantially enhanced filamentous

189 growth of *P. infestans* compared to RNAi-GUS control (Fig. 3F-G). Taken together,

190 these results indicate that CHUP1 contributes to plant focal immunity, possibly through

191 coordinating chloroplast navigation to the EHM. However, since CHUP1 is also essential

192 for induction of stromules that intimately associate with the EHM, we infer that processes

193 such as chloroplast steering and docking to the pathogen interface as well as formation

194 of stromules are a series of interconnected events to counteract microbial invasion.

Fig. 3. CHUP1 silencing decreases immunity to *P. infestans*, perihaustorial chloroplast accumulation and stromule induction (A) Maximum-projection confocal micrographs of CpGFP leaf epidermal cells, where *CHUP1* is systemically silenced (VIGS), and infected with 88069td. VIGS-empty vector (EV) included as control. Scale bars: 10 µm. (B) Scatter-boxplot shows reduced percentage of haustoria associated with plastid(s) (p < 0.01) in VIGS-CHUP1 plants (33% of *n* = 544 haustoria), compared to VIGS-EV plants (45% of *n* = 471 haustoria). (C) Scatter-boxplot depicts reduction in stromule induction (p < 0.01) in VIGS-CHUP1 plants (8%, *n* = 68 images quantified), compared to VIGS-EV plants (24%, *n* = 68 images quantified). (D-G) Silencing of *CHUP1* by VIGS or transient RNAi, followed by 88069td inoculation shows CHUP1 contributes to resistance. Hyphal growth measured using fluorescence stereomicroscope at 4 dpi for VIGS (D) and 6 dpi for transient RNAi (F). Scale bars: 10 mm. (E & G) Scatter-boxplots show mean area of hyphal growth for inoculations on VIGS leaves, *n* = 32 leaves per treatment (p < 0.01) (E), and transient RNAi silenced leaves, *n* = 22 leaves for each experiment (p < 0.01).

Finally, to gain further insights into the interplay between chloroplasts and haustoria, we

investigated the means of pathogen induced stromule formation. Stromule development

was previously reported to be induced upon cytoplasmic recognition of bacterial

effectors or viral particles by NLR (nucleotide-binding domain and leucine-rich repeat-

containing) receptors (Caplan et al., 2015; Erickson et al., 2018; Krenz et al., 2012).

However, because *P. infestans* establishes a compatible interaction with *N.*

benthamiana without activating NLR triggered immunity, we reasoned that stromule

development could be stimulated through surface immune recognition (Caplan et al.,

2015). Thus, we monitored stromule formation upon silencing of *BAK1* (Fig. S8), a

surface localized co-receptor that mediates immune signalling through various PRRs

(Chaparro-Garcia et al., 2011; Perraki et al., 2018; Smakowska-Luzan et al., 2018).

Remarkably, in five independent experiments, we noticed a substantial decrease in

infection triggered stromule induction following *BAK1* silencing (4%, $n = 37$ images

quantified) in relation to control silencing (23%, $n = 37$ images quantified) (Fig. 4A-B).

We then tested whether AVR3a, a host-translocated effector of *P. infestans* that

suppresses BAK1-mediated immune signalling (Bos et al., 2010; Chaparro-Garcia et al.,

2011), can perturb pathogen induced stromule development. Notably, following

heterologous expression of AVR3a (Fig. S9), infection-triggered stromule formation

decreased by more than two-fold (Fig. 4C-D). These results demonstrate that stromules

are induced during *P. infestans* host colonization through surface immune signalling,

whereas the pathogen deploys effectors to counteract this process, further supporting

the defence-related role of stromules. Since stromule induction is CHUP1-dependent,

we conclude that CHUP1 contributes to immunity through mediating chloroplast

navigation and chloroplast interaction via diverse signalling stimuli, which are

coordinated to prevent host cell invasion of filamentous pathogens (Fig. 4E, model).

Fig. 4: *P. infestans* induces stromules through surface recognition. (A-B)

Systemically silencing *BAK1* (VIGS-BAK1) reduces stromule induction during infection,

compared with EV control. (A) Maximum projection confocal microscopy images of

CpGFP leaves silenced with VIGS-BAK1 or VIGS-EV, infected with 88069td. (B) Scatter-

boxplot depicts reduction in stromule induction (p < 0.01) in VIGS-BAK1 plants (4%, *n* =

37 images quantified), compared to VIGS-EV plants (23%, *n* = 37 images quantified).

(C) Overexpression of *P. infestans* effector AVR3a reduces stromule induction during

infection with 88069td in CpGFP. (D) Scatter-boxplot shows transient AVR3a expression

reduces stromule induction (9%, *n* = 39 images quantified) compared to EV expression

(22%, *n* = 40 images quantified) (p<0.01). (E) Model: Surface recognition of *P. infestans*

leads to CHUP1-dependent stromule induction. CHUP1 mediates chloroplast movement

to siege haustoria and develop inter-chloroplast clustering at the penetration site.

Our results demonstrate that chloroplasts navigate to the pathogen interface and

establish secure membrane contacts with the EHM to contribute to focal immunity (Fig.

1, Fig. S1-2, Movie S5). These processes are dependent on CHUP1, a key component

of blue-light-induced chloroplast movement and plasma membrane attachment

(Kasahara et al., 2002; Oikawa et al., 2008), indicating that chloroplast photo-relocation

machinery is co-opted for antimicrobial immunity. Remarkably, CHUP1 accumulates at

the chloroplast-EHM interface (Fig. 2C, Fig. S4), possibly to enable stronger membrane

attachment, as has been shown in plasma membrane docking of chloroplasts (Kadota et

al., 2009).

We show that surface immune activation induces stromule development into a large,

intricate web of chloroplasts (Fig. 1A, 1H, Fig. S3, Movie S1, S3, S7-9). These are

reminiscent of mitochondrial networks with poorly understood functions (Hoitzing et al.,

2015). Intricate chloroplast clusters could boost coordination of their defence-related

functions by, for instance, mediating focal deployment of chloroplast weaponry at the

pathogen interface. In addition, a speculative hypothesis is that chloroplast clusters

could also generate mechanical forces to repulse infection structures that invade the

plant cells, as has been suggested by our live cell imaging (Fig. S10-13). However, we

show that adapted pathogens can remotely counteract stromule development by

deploying effectors that shut down surface immune signalling (Fig. 4B). Conceivably, this

would dismantle chloroplast stromules and reduce the surface area of the chloroplasts in

contact with the EHM. These findings further support the notion that stromules are

induced to contribute to pathogen defence (Caplan et al., 2015; Erickson et al., 2018).

Our findings implicate chloroplasts in polarized immune responses of plants against

filamentous pathogens and point to more complex, direct defence-related functions for

chloroplasts.

Materials and Methods

Biological Material

Nicotiana benthamiana plants grown in a growth chamber at 25°C under high light

intensity (16-h-day/8-h-dark photoperiod) were used for all experiments. Transplastomic

GFP-expressing *Nicotiana benthamiana* plants, accumulating GFP in the chloroplast

stroma(Stegemann et al., 2012) were maintained in the same conditions. *Phytophthora*

infestans isolate 88069 (WT)(Van West et al., 1998) and 88069td (TD)(Whisson et al.,

2007), a transgenic strain expressing the red fluorescent marker tandem dimer RFP

(tdTomato), were used. Both isolates were cultured on plates with rye sucrose agar

(RSA) for 12-16 days at 18°C in the dark, as described elsewhere(Song et al., 2009)

prior to use for infection of *N. benthamiana*.

Molecular Cloning and plasmid constructs

The following constructs used in this study were previously published as follows:

RFP:REM1.3(Bozkurt et al., 2014); TOC64:GFP(Breuers et al., 2012); AVR3a cloned in

pICSL86977 was provided by TSLSynBio. CHUP1:GFP construct was cloned using

Gibson Assembly Protocol(Gibson et al., 2009). Briefly, CHUP1:GFP was generated by

amplifying two PCR products (Phusion DNA polymerase, Thermo Scientific) from *N.*

benthamiana cDNA, amplifying *CHUP1-a* (herein *CHUP1*) using two different primer

pairs (CHUP1-A F; GA_CHUP1_R1; GA_CHUP1B_F2; GA_CHUP1B_R2). The PCR

products were assembled and cloned into the destination vector pK7FWG2,0 (GFP) by

Gibson assembly. The purified vector was transformed into *A. tumefaciens* competent

cells. RNAi-silencing constructs for CHUP1 were cloned using the Gateway Cloning

Technology. RNAi-CHUP1 construct was generated to target a region in the middle of

the gene (942-1200bp)(Helliwell and Waterhouse, 2003). The selected sequence was

synthetized and cloned into pK7GWIWG2(II) vector(Karimi et al., 2002). VIGS vector

pTRV2-CHUP1 was assembled by amplifying a 500bp region of *Chup1* (the same region

targeted as previously(Caplan et al., 2015)) from *N. benthamiana* cDNA using primers

VIGS-_CHUP1_F and VIGS_CHUP1_R2. The amplified fragment was then cloned into a

Gateway compatible pTRV2 vector using Gateway Technology (Invitrogen). Silencing

construct TRV2-BAK1 was kindly provided by The Sainsbury Lab(Chaparro-Garcia et

al., 2011).

Transient gene-expression assays in *N. benthamiana*

Agrobacterium tumefaciens GV3101 strain(Hellens et al., 2000) carrying T-DNA

constructs was used to mediate transient gene expression (Agroinfiltration) into 3-4-

week-old *N. benthamiana* leaves, as previously described(Bozkurt et al., 2014, 2011).

Briefly, overnight cultures of transformed *A. tumefaciens* were washed and harvested

with 1500 µl autoclaved dH_2O by centrifugation at 1500*g* twice and resuspended in

agroinfiltration buffer (10 mM 2-(*N*-morpholino-ethanesulfonic acid hydrate (MES

hydrate), 10 mM $MgCl_2$, pH 5.7). For the transient co-expression assays, each *A.*

tumefaciens construct was mixed in agroinfiltration buffer to achieve a final OD_{600} of 0.2

or 0.3 for each *A. tumefaciens*, depending on the experiment. *P. infestans* inoculations

were performed 4 to 24h after infiltrations.

Virus induced gene silencing (VIGS)

Agrobacterium was prepared as above carrying TRV1 and the appropriate TRV2

construct and mixed to a final OD_{600} of 0.4 or 0.2 respectively, in agroinfiltration buffer

supplemented with 100 μM acetosyringone (Sigma) and left in the dark for 2 h prior to

infiltration to stimulate virulence. 14-day old *N. benthamiana* seedlings were infiltrated in

both cotyledons and any true leaves that had emerged. *N. benthamiana* plants were

infiltrated with TRV1 and TRV2-CHUP1 for CHUP1-silencing and TRV1 and TRV2-EV

for the empty vector control. TRV2 containing the *N. benthamiana* sulfur (Su) gene

fragment (TRV2-NbSU) was used as a positive control to indicate viral spread. Plants

were left to grow under standard conditions until experiments could be carried out four

weeks later.

<u>Infection assays</u>

Infection assays to assess the effect of *CHUP1* silencing on *P. infestans* colonization

were performed as follow: for RNAi experiments, *A. tumefaciens* containing RNAi-

CHUP1 or RNAi-GUS (OD_{600} = 0.3) constructs were infiltrated for transient expression

side by side on either halves of independent *N. benthamiana* leaves, 24h prior to

infection. For VIGS experiments, 3 weeks after silencing was initiated leaves were

detached for infection from VIGS-CHUP1 and VIGS-empty vector plants. Zoospores

were harvested from sporangia by addition of cold distilled water and collected after 2h

of incubation at 4°C, adjusting dilution to 50,000 spores/ml. Infection assays were

performed by the addition of 10μl of zoospore droplets to the base, middle and tip of the

abaxial side of each leaf, 3 on each side, as described previously(Saunders et al., 2012;

Song et al., 2009). The infected leaves were maintained in plastic boxes on damp paper

towels at 18°C under 16-h- day/8-h-night conditions, and inoculated with *P. infestans*

88069td. Images of hyphal growth were captured on a Leica DFC300 FX fluorescent

microscope (Leica Microsystems, Germany) using the DSR UV filter at the timepoints

indicated in the Fig. legends. Images were processed in ImageJ (2.0) and quantification

of hyphal growth was performed by measuring the diameter in mm^2 of hyphal growth of

each infected spot.

RT-PCR assay

60 mg of leaf tissue was excised from 4-week-old *N. benthamiana* leaves (RNAi

experiment, 6 days after silencing), and 5-week old leaves (VIGS experiments) and

frozen in liquid N$_2$. RNA was extracted from the leaf tissue using the Plant RNA Isolation

Mini Kit Protocol (Agilent Technologies). RNA quality and concentration was measured

using a NanoDropTM Lite Spectrophotometer (Thermo Scientific). cDNA was

synthesized using as a template 2 μg of RNA following the SuperScript II RT protocol

(Invitrogen). To amplify the cDNA, a standard PCR (RT-PCR) was then performed using

DreamTaq DNA polymerase (5 u/μl) (Thermo Scientific). RNAi-CHUP1 construct effect

on *CHUP1* downregulation was evaluated by RT-PCR using primers CHP1_RT_F1 and

BD-CHUP1-REV. VIGS-CHUP1 silencing was confirmed with CHUP1a_RT_F &

CHUP1a_RT_R, which specifically amplifies *CHUP1a*, and CHUP1b_RT_F &

CHUP1b_RT_R, which specifically amplifies *CHUP1b.* VIGS-BAK1 silencing was

confirmed as previously described(Chaparro-Garcia et al., 2011).

Confocal microscopy

All microscopy analyses were performed on live *N. benthamiana* epidermal cells 2-6

days post agroinfiltrations and infections. Leaf discs were excised and imaged on a

Leica SP5 resonant inverted confocal microscope (Leica Microsystems) using 63X

1.2NA Plan-Apochromat water immersion objective. Specific excitation wavelengths and

filters for emission spectra were set as described previously(Koh et al., 2005). The

Argon laser excitation was set to 488 nm and the Helium-Neon laser to 543 nm and their

fluorescent emissions detected at 495–550 and 570–620 nm to visualize GFP and RFP

fluorescence, respectively. To avoid bleed-through from different fluorophores, images

were acquired using sequential scanning and Maximum Intensity Projections were

created from the Z-stacks. 3D images and movies were generated with confocal files in

12-bit TIFF format imported into NIS-Elements (Version 4.50, Nikon Instruments, UK)

and processed with Advanced Denoising. Movies were made using the Volume View

and Movie Maker modules

Optical trapping setup

Optical trap for chloroplast/stromule capture was setup as described by Sparkes et al

2017, Chapter 13(Hawes, 2018, chap. 13). An optical trap with a two-channel TIRF

microscope (TIRF-M) was combined with a Nikon Ti-U inverted microscope. Optical

trapping was performed using a near infrared trapping laser at 1070 nm using a Nikon

100x, oil immersion, NA 1.49 TIRF objective lens. For GFP and RFP chromophores

fused to the proteins of interest were excited using 488 and 561 nm laser diode,

respectively. Their Fluorescent emissions were detected using two electron multiplying

charge-coupled device (EMCCD, iXon, Andor) cameras. The sample (~5 mm^2 leaf

tissue) was mounted on a computer-controlled variable speed (Märzhäuser) stepper

motor stage. The associated computer-controlled hardware was interfaced using

National Instruments LabVIEW which provides full automation for each trapping routine.

The power of the optical trap laser transmission was set to 40.7 mW. The TIRF image

was recorded from 0 s, the trap was turned on at 1 s, the translation stage movement of

10 µm at 2 µm/s begins at 5 s and ends at 10 s, the trap was deactivated at 11 s, and

the image recording stops at 22 s (relating to 11 s recovery periods).

Bioinformatic and statistical analysis

In order to identify putative orthologs of *Arabidopsis thaliana* for CHUP1 in *N.*

benthamiana genome, a TBLASTN search using the protein sequence of *CHUP1* from

A. thaliana against the *N. benthamiana* cDNA was accomplished in Solgenomics. Two

orthologs (*CHUP1-a* and *CHUP1-b*) of *Arabidopsis CHUP1* gene were identified in *N.*

benthamiana genome. Chloroplast quantification was done automatically using a

MATLAB script. Stromules were manually counted using a semi-automated MATLAB

script. Percentage of chloroplasts with stromules were calculated by dividing the number

of chloroplasts one (or more) stromule(s) by the total number of chloroplasts.

Quantification of hyphal growth was accomplished by measuring the diameter of the

lesion on each inoculated spot using Fiji image-processing software. R package was

used to visualize the values from three to four independent biological replicates by

generating scatter plots. Statistical significance of the differences observed were

assessed by t test when found to be normal by sharpiro test. If data was found to be

non-normally distributed Wilcox statistical test was implemented by R package.

Chloroplast automated counting algorithm through image processing

The image processing algorithms were used to calculate the gradient of the image to

identify the boundaries of the puncta. Enclosed regions formed by the boundaries were

algorithmically identified and counted. This procedure was done for each individual

channel green (in chloroplast stroma) and blue (Chloroplast Auto-fluorescence). The

chloroplast (GFP channel) containing stromule were counted in a semi-automated

fashion.

Acknowledgements: We thank Dr. Alex Jones (Warwick) for initiating collaboration with

IS, Dr. Sebastian Schornack (SLCU) for initiating collaboration with MS, Prof. Peter

Nixon (Imperial) for providing CP-GFP plant seeds. **Funding:** Bozkurt lab funded by

BBSRC (BB/M002462/1). The Facility for Imaging by Light Microscopy (FILM) at Imperial

College London is part-supported by funding from the Wellcome Trust (grant

104931/Z/14/Z) and BBSRC (grant BB/L015129/1). **Author contributions:**

Conceptualization: CD, ZS, MES, IS, MS, TOB; Data curation: AT, CD, PP, ZS, DCAG,

BCB, LHY, IS, TOB; Formal analysis: AT, CD, PP, ZS, BCB, IS, TOB; Funding

acquisition: TOB; Investigation: AT, CD, PP, ZS, MES, LHY, AYL, VK, BCB, TOB;

Methodology: AT, CD, ZS, DCAG, ADW, SWB, BCB, IS, MS, TOB; Project

Administration: CD, TOB; Resources: AT, CD, AYL, VK, IP, IS, TOB; Software: IP;

Supervision: AT, CD, PP, MES, BCB, IS, TOB; Validation: AT, CD, PP, ZS, MES, IS,

MS; Visualisation: CD, ZS, TOB; Writing – original draft: CD, ZS, TOB; Writing – review

& editing: AT, CD, PP, ZS, MES, LHY, DCAG, AYL, ADW, SWB, MS, IS, TOB.

Competing interests: Authors declare no competing interests. **Data and materials

availability:** All data is available in the main text, the supplementary materials or other

raw data is available upon request. We are happy to provide all materials used here

upon request.

References:

Belhaj K, Lin B, Mauch F. 2009. The chloroplast protein RPH1 plays a role in the

immune response of Arabidopsis to Phytophthora brassicae. *Plant J* **58**:287–298.

doi:10.1111/j.1365-313X.2008.03779.x

Bos JIB, Armstrong MR, Gilroy EM, Boevink PC, Hein I, Taylor RM, Zhendong T,

Engelhardt S, Vetukuri RR, Harrower B, Dixelius C, Bryan G, Sadanandom A,

Whisson SC, Kamoun S, Birch PRJ. 2010. Phytophthora infestans effector AVR3a

is essential for virulence and manipulates plant immunity by stabilizing host E3

ligase CMPG1. *Proc Natl Acad Sci* **107**:9909–9914. doi:10.1073/pnas.0914408107

Bozkurt TO, Belhaj K, Dagdas YF, Chaparro-Garcia A, Wu C-H, Cano LM, Kamoun S.

2015. Rerouting of plant late endocytic trafficking toward a pathogen interface.

Traffic **16**:204–26. doi:10.1111/tra.12245

Bozkurt TO, Richardson A, Dagdas YF, Mongrand S, Kamoun S, Raffaele S. 2014. The

Plant Membrane-Associated REMORIN1.3 Accumulates in Discrete Perihaustorial

Domains and Enhances Susceptibility to Phytophthora infestans. *Plant Physiol*

165:1005–1018. doi:10.1104/pp.114.235804

Bozkurt TO, Schornack S, Win J, Shindo T, Ilyas M, Oliva R, Cano LM, Jones AME,

Huitema E, van der Hoorn R a L, Kamoun S. 2011. Phytophthora infestans effector

AVRblb2 prevents secretion of a plant immune protease at the haustorial interface.

Proc Natl Acad Sci U S A **108**:20832–7. doi:10.1073/pnas.1112708109

Breuers FKH, Bräutigam A, Geimer S, Welzel UY, Stefano G, Renna L, Brandizzi F,

Weber APM. 2012. Dynamic Remodeling of the Plastid Envelope Membranes – A

Tool for Chloroplast Envelope in vivo Localizations. *Front Plant Sci* **3**:1–10.

doi:10.3389/fpls.2012.00007

Brodersen P, Achard, Lali Sakvarelidze Rasmussen MB, Dunoyer P, Yamamoto YY,

Sieburth L, Voinnet O. 2008. Widespread Translational Inhibition by Plant miRNAs

and siRNAs. *Science (80-)* **320**:1185–1190.

Caplan JL, Kumar AS, Park E, Padmanabhan MS, Hoban K, Modla S, Czymmek K,

Dinesh-Kumar SP. 2015. Chloroplast Stromules Function during Innate Immunity.

Dev Cell **34**:45–57. doi:10.1016/j.devcel.2015.05.011

Chaparro-Garcia A, Wilkinson RC, Gimenez-Ibanez S, Findlay K, Coffey MD, Zipfel C,

Rathjen JP, Kamoun S, Schornack S. 2011. The receptor-like kinase serk3/bak1 is

required for basal resistance against the late blight pathogen Phytophthora

infestans in Nicotiana benthamiana. *PLoS One* **6**.

doi:10.1371/journal.pone.0016608

Dagdas YF, Pandey P, Tumtas Y, Sanguankiattichai N, Belhaj K, Duggan C, Leary AY,

Segretin ME, Contreras MP, Savage Z, Khandare VS, Kamoun S, Bozkurt TO.

2018. Host autophagy machinery is diverted to the pathogen interface to mediate

focal defense responses against the Irish potato famine pathogen. *Elife* 1–15.

doi:10.7554/eLife.37476

de Torres Zabala M, Littlejohn G, Jayaraman S, Studholme D, Bailey T, Lawson T,

Tillich M, Licht D, Bölter B, Delfino L, Truman W, Mansfield J, Smirnoff N, Grant M.

2015. Chloroplasts play a central role in plant defence and are targeted by

pathogen effectors. *Nat Plants* **1**:15074. doi:10.1038/nplants.2015.74

Erickson JL, Adlung N, Lampe C, Bonas U, Schattat MH. 2018. The Xanthomonas

effector XopL uncovers the role of microtubules in stromule extension and

dynamics in Nicotiana benthamiana. *Plant J* **93**:856–870. doi:10.1111/tpj.13813

Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009.

Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat*

Methods **6**:343–5. doi:10.1038/nmeth.1318

Hanson MR, Hines KM. 2017. Stromules: Probing Formation and Function. *Plant Physiol*

176:pp.01287.2017. doi:10.1104/pp.17.01287

Hawes C. 2018. The Plant Endoplasmic Reticulum, Methods in Molecular Biology. New

York, NY: Springer New York. doi:10.1007/978-1-4939-7389-7

Hellens R, Mullineaux P, Klee H. 2000. Technical Focus:a guide to Agrobacterium binary

Ti vectors. *Trends Plant Sci* **5**:446–51.

Helliwell C, Waterhouse P. 2003. Constructs and methods for high-throughput gene

silencing in plants. *Methods* **30**:289–95.

Hoitzing H, Johnston IG, Jones NS. 2015. What is the function of mitochondrial

networks? A theoretical assessment of hypotheses and proposal for future

research. *BioEssays* **37**:687–700. doi:10.1002/bies.201400188

Jelenska J, Yao N, Vinatzer BA, Wright CM, Brodsky JL, Greenberg JT. 2007. A J

Domain Virulence Effector of Pseudomonas syringae Remodels Host Chloroplasts

and Suppresses Defenses. *Curr Biol* **17**:499–508. doi:10.1016/j.cub.2007.02.028

Kadota A, Yamada N, Suetsugu N, Hirose M, Saito C, Shoda K, Ichikawa S, Kagawa T,

Nakano A, Wada M. 2009. Short actin-based mechanism for light-directed

chloroplast movement in Arabidopsis. *Proc Natl Acad Sci* **106**:13106–13111.

doi:10.1073/pnas.0906250106

Karimi M, Inzé D, Depicker A. 2002. GATEWAY vectors for Agrobacterium-mediated

plant transformation. *Trends Plant Sci* **7**:193–5. doi:10.1016/S1360-1385(02)02251-

3

Kasahara M, Kagawa T, Oikawa K, Suetsugu N, Miyao M, Wada M. 2002. Chloroplast

avoidance movement reduces photodamage in plants. *Nature* **420**:829–832.

doi:10.1038/nature01213

Koh S, André A, Edwards H, Ehrhardt D, Somerville S. 2005. Arabidopsis thaliana

subcellular responses to compatible Erysiphe cichoracearum infections. *Plant J*

44:516–529. doi:10.1111/j.1365-313X.2005.02545.x

Krenz B, Jeske H, Kleinow T. 2012. The induction of stromule formation by a plant DNA-

515 virus in epidermal leaf tissues suggests a novel intra- and intercellular

516 macromolecular trafficking route. *Front Plant Sci* **3**:1–12.

517 doi:10.3389/fpls.2012.00291

518 Kumar AS, Park E, Nedo A, Alqarni A, Ren L, Hoban K, Modla S, McDonald JH,

519 Kambhamettu C, Dinesh-Kumar SP, Caplan JL. 2018. Stromule extension along

520 microtubules coordinated with actin-mediated anchoring guides perinuclear

521 chloroplast movement during innate immunity. *Elife* **7**:1–33.

522 doi:10.7554/eLife.23625

523 Kwon C, Neu C, Pajonk S, Yun HS, Lipka U, Humphry M, Bau S, Straus M, Kwaaitaal M,

524 Rampelt H, El Kasmi F, Jürgens G, Parker J, Panstruga R, Lipka V, Schulze-Lefert

525 P. 2008. Co-option of a default secretory pathway for plant immune responses.

526 *Nature* **451**:835–40. doi:10.1038/nature06545

527 Oikawa K, Kasahara M, Kiyosue T, Kagawa T, Suetsugu N, Takahashi F, Kanegae T,

528 Niwa Y, Kadota A, Wada M. 2003. Chloroplast unusual positioning1 is essential for

529 proper chloroplast positioning. *Plant Cell* **15**:2805–15. doi:10.1105/tpc.016428

530 Oikawa K, Yamasato A, Kong S-G, Kasahara M, Nakai M, Takahashi F, Ogura Y,

531 Kagawa T, Wada M. 2008. Chloroplast Outer Envelope Protein CHUP1 Is Essential

532 for Chloroplast Anchorage to the Plasma Membrane and Chloroplast Movement.

533 *Plant Physiol* **148**:829–842. doi:10.1104/pp.108.123075

534 Padmanabhan MS, Dinesh-Kumar SP. 2010. All Hands on Deck—The Role of

535 Chloroplasts, Endoplasmic Reticulum, and the Nucleus in Driving Plant Innate

536 Immunity. *Mol Plant-Microbe Interact* **23**:1368–1380. doi:10.1094/MPMI-05-10-0113

537 Pecrix Y, Buendia L, Penouilh-Suzette C, Maréchaux M, Legrand L, Bouchez O, Rengel

538 D, Gouzy J, Cottret L, Vear F, Godiard L. 2018. Sunflower resistance to multiple

539 downy mildew pathotypes revealed by recognition of conserved effectors of the

540 oomycete *Plasmopara halstedii*. *Plant J*. doi:10.1111/tpj.14157

Perraki A, DeFalco TA, Derbyshire P, Avila J, Séré D, Sklenar J, Qi X, Stransfeld L,

Schwessinger B, Kadota Y, Macho AP, Jiang S, Couto D, Torii KU, Menke FLH,

Zipfel C. 2018. Phosphocode-dependent functional dichotomy of a common co-

receptor in plant signalling. *Nature* **561**:248–252. doi:10.1038/s41586-018-0471-x

Petre B, Lorrain C, Saunders DGO, Win J, Sklenar J, Duplessis S, Kamoun S. 2016.

Rust fungal effectors mimic host transit peptides to translocate into chloroplasts.

Cell Microbiol **18**:453–465. doi:10.1111/cmi.12530

Saunders DGO, Breen S, Win J, Schornack S, Hein I, Bozkurt TO, Champouret N,

Vleeshouwers VG a a, Birch PRJ, Gilroy EM, Kamoun S. 2012. Host protein BSL1

associates with Phytophthora infestans RXLR effector AVR2 and the Solanum

demissum Immune receptor R2 to mediate disease resistance. *Plant Cell* **24**:3420–

34. doi:10.1105/tpc.112.099861

Schattat MH, Barton KA, Mathur J. 2014. The myth of interconnected plastids and

related phenomena. *Protoplasma* **252**:359–371. doi:10.1007/s00709-014-0666-4

Serrano I, Audran C, Rivas S. 2016. Chloroplasts at work during plant innate immunity. *J

Exp Bot* **67**:3845–3854. doi:10.1093/jxb/erw088

Smakowska-Luzan E, Mott GA, Parys K, Stegmann M, Howton TC, Layeghifard M,

Neuhold J, Lehner A, Kong J, Grünwald K, Weinberger N, Satbhai SB, Mayer D,

Busch W, Madalinski M, Stolt-Bergner P, Provart NJ, Mukhtar MS, Zipfel C,

Desveaux D, Guttman DS, Belkhadir Y. 2018. An extracellular network of

Arabidopsis leucine-rich repeat receptor kinases. *Nature* **553**:342–346.

doi:10.1038/nature25184

Song J, Win J, Tian M, Schornack S, Kaschani F, Ilyas M, Hoorn RAL Van Der, Kamoun

S. 2009. Apoplastic effectors secreted by two unrelated eukaryotic plant pathogens

target the tomato defense protease Rcr3.

Stegemann S, Keuthe M, Greiner S, Bock R. 2012. Horizontal transfer of chloroplast

genomes between plant species. *Proc Natl Acad Sci* **109**:2434–2438.

doi:10.1073/pnas.1114076109

Su J, Yang L, Zhu Q, Wu H, He Y, Liu Y, Xu J, Jiang D, Zhang S. 2018. Active

photosynthetic inhibition mediated by MPK3/MPK6 is critical to effector-triggered

immunity. *PLoS Biol* **16**:1–29. doi:10.1371/journal.pbio.2004122

Trotta A, Rahikainen M, Konert G, Finazzi G, Kangasjärvi S. 2014. Signalling crosstalk in

light stress and immune reactions in plants. *Philos Trans R Soc Lond B Biol Sci*

369:20130235. doi:10.1098/rstb.2013.0235

van Damme M, Zeilmaker T, Elberse J, Andel A, de Sain-van der Velden M, van den

Ackerveken G. 2009. Downy Mildew Resistance in Arabidopsis by Mutation of

HOMOSERINE KINASE. *Plant Cell Online* **21**:2179–2189.

doi:10.1105/tpc.109.066811

Van West P, De Jong AJ, Judelson HS, Emons AMC, Govers F. 1998. The ipiO gene of

phytophthora infestans is highly expressed in invading hyphae during infection.

Fungal Genet Biol **23**:126–138. doi:10.1006/fgbi.1998.1036

Wang S, Boevink PC, Welsh L, Zhang R, Whisson SC, Birch PRJ. 2017. Delivery of

cytoplasmic and apoplastic effectors from *Phytophthora infestans* haustoria by

distinct secretion pathways. *New Phytol*. doi:10.1111/nph.14696

Whisson SC, Boevink PC, Moleleki L, Avrova AO, Morales JG, Gilroy EM, Armstrong

MR, Grouffaud S, Van West P, Chapman S, Hein I, Toth IK, Pritchard L, Birch PRJ.

2007. A translocation signal for delivery of oomycete effector proteins into host

plant cells. *Nature* **450**:115–118. doi:10.1038/nature06203

Whisson SC, Boevink PC, Wang S, Birch PR. 2016. The cell biology of late blight

disease. *Curr Opin Microbiol* **34**:127–135. doi:10.1016/j.mib.2016.09.002

List of Supplementary Materials:

Figures S1-S9

Table S1

Movies S1-S13

Fig. S1: Confocal micrographs showing a chloroplast embracing a haustorium
with its stromules. Leaf epidermal cells from transplastomic CpGFP *N. benthamiana*
plants transiently expressing plasma membrane and EHM marker RFP:REM1.3 were
infected with WT *P. infestans* 8806 and imaged 4dpi. Blue represents chlorophyll
autofluorescence. Scale bar is 10 µm. Magenta arrowheads indicate stromules.

602
603
604 **Fig. S2: Time-lapse series showing chloroplasts accumulation towards a**
605 **haustorium.** Leaf epidermal cells from WT *N. benthamiana* plants expressing PM and
606 EHM marker RFP:REM1.3 (Magenta) and endosomal marker GFP:RAB8 (Yellow)
607 infected with WT *P. infestans* 88069. Cyan is chlorophyll autofluorescence, labelling the
608 chloroplasts. Scale bar is 10 µm.

609
610
Fig. S3: Chloroplasts form long-distance stromule interactions that can bridge
more than one haustorium. Confocal micrographs of leaf epidermal cells from
transplastomic CpGFP *N. benthamiana* plants. Maximum intensity projection (10
images), 5 dpi (days post infection) with red-fluorescent *P. infestans* strain 88069td. 'h'
indicates haustorium, Magenta arrowheads indicate long-distance stromule interactions.
Scale bars are 10 μm. See 3D Movie 7.

617
618
619 **Fig. S4: Time-lapse series showing collapse of haustorium.** *N. benthamiana* leaves
620 transiently co-expressing plasma membrane and EHM marker RFP:REM1.3 with
621 CHUP1:GFP. Arrowhead indicates haustorium that collapses. Scale bars are 10 μm.

622
623
624 **Fig. S5: CHUP1:GFP accumulates at the contact points between the chloroplast**
625 **and the EHM.** *N. benthamiana* leaves transiently co-expressing plasma membrane and
626 EHM marker RFP:REM1.3 with CHUP1:GFP were infected with WT *P. infestans* 88069.
627 Blue represents chlorophyll autofluorescence. Scale bars are 10 µm.

VIGS-CHUP1   VIGS-EV

NbCHUP1-a

NbCHUP1-b

GAPDH

| | VIGS-CHUP1 |
RT-PCR (Full-length)	(2436bp-2935bp)
NbChup1A/B	2976bp

628
629
Fig. S6: VIGS-*CHUP1* silences *CHUP1* compared to the EV control. 12 day old
transplastomic CpGFP *N. benthamiana* plants, were infiltrated with *Agrobacterium*
expressing TRV2:CHUP1 or TRV2:EV. Leaf disks were taken from 5-week old,
uninfected, silenced tissue and RNA was extracted. Semi-quantitative RT-PCR of
CHUP1 shows that it was silenced in VIGS-*CHUP1* tissue compared to EV. RT-PCR of
housekeeping GAPDH was used as an internal control for cDNA loading. For both,
primers that did not amplify the silencing target were use

Fig. S7: RNAi of CHUP1 suppresses stromule formation and haustorial accumulation of chloroplasts (A) Confocal micrographs of leaf epidermal cells from transplastomic CpGFP *N. benthamiana* plants, in which *CHUP1* is locally silenced (RNAi-CHUP1), 5-7 dpi with *P. infestans* (88069td), with RNAi-GUS included as a control. The first scatter-boxplot depicts the effect of RNAi-CHUP1 silencing on the percentage of haustoria in contact with at least one plastid (27% of *n* = 237 haustoria), compared to RNAi-GUS control (50% of *n* = 126 haustoria) (p < 0.01). The second scatter boxplot depicts the effect of RNAi-CHUP1 silencing on the percentage of

chloroplasts with one (or more) stromule(s) (18%, $n = 11$ images quantified) compared to
RNAi-GUS control (39%, $n = 10$ images quantified) ($p < 0.05$). (B) Confocal micrographs
of leaf epidermal cells from WT *N. benthamiana* plants, in which *CHUP1* is locally
silenced (RNAi-CHUP1), 5-7 dpi with *P. infestans* (88069td), with RNAi-GUS included as
a control. The scatter-boxplot depicts the effect of RNAi-CHUP1 silencing on the
percentage of haustoria in contact with at least one plastid (30% of $n = 392$ haustoria),
compared to RNAi-GUS control (55% of $n = 215$ haustoria) ($p < 0.01$). Scale bars are 10
µm. (C) Leaf disks were taken from uninfected tissue infiltrated to express the RNAi
construct at 6 days post infiltration and RNA was extracted. Following semi-quantitative
RT-PCR of *CHUP1* we noted only a slight reduction of *CHUP1* mRNA from RNAi-
CHUP1 tissue compared to RNAi-GUS control. However, clear silencing of CHUP1:GFP
was seen upon western blotting (D), indicating that the RNAi-CHUP1 hairpin silencing
construct mainly acts by blocking CHUP1 translation, as is known to occur for in
RNAi(Brodersen et al., 2008).

A   VIGS-BAK1   VIGS-EV

NbSerk3A/B (BAK1)

GAPDH

B   VIGS-BAK1   VIGS-EV

Top view

Side view

Fig. S8: VIGS-*BAK1* silences *BAK1* compared to the EV control. 12 day old
transplastomic CpGFP *N. benthamiana* plants, were infiltrated with *Agrobacterium*
expressing TRV2:BAK1 or TRV2:EV. (A) Leaf disks were taken from uninfected,
silenced tissue and RNA was extracted. Semi-quantitative RT-PCR of *BAK1* shows that
it was silenced in VIGS-*BAK1* tissue compared to EV. RT-PCR of housekeeping GAPDH
was used as an internal control for cDNA loading. For both, primers that did not amplify
the silencing target were used. (B) Photos of representative *N. benthamiana* plants 5
weeks old, showing symptoms of *BAK1* silencing by VIGS.

A

B

669
670 **Fig. S9: AVR3a expression triggers HR in an R3a-dependent manner, indicating it**
671 **is functional.** AVR3a or R3a were expressed alone, triggering no HR. EV and R3a also
672 triggered no HR. Only expression of AVR3a with R3a triggered HR, as visible by white
673 light or autofluorescence under UV light.

674 **Table S1. Primers used in this study**

Primer name	Sequence (5´to 3´)
CHUP1-A F	AGGCGGCCGCACTAGTATGATAGTCAGGGTAGGTTTAGTGG
GA_CHUP1_R1	GTTCCAAAACTAGTGATGGCTAC
GA_CHUP1B_F2	GCTCAGAAATGCAGGTGATGGT
GA_CHUP1B_R2	TCCTCGCCCTTGCTCACCATTGATCCTGTTTCTTGTGTATTCTCTTCTCC
VIGS-_CHUP1_F	GGTTGATGAACGAGCTGTCCTCAAG
VIGS_CHUP1_R2	TGACACGACTCCTTAATTCTTCAAAAG
CHP1_RT_F1	ATGATAGTCAGGGTAGGTTTAGTG
BD-CHUP1-REV	ACCAGGTCTCACACCTTGTTCTTGTGTACTCTCT
CHUP1a_RT_F	ATGATCGTCAGGGTAGGTTTAGTGGTTGC
CHUP1a_RT_R	TGTTTCTTGTGTATTCTCTTCTCCTGTTTGT
CHUP1b_RT_F	ATGATAGTCAGGGTAGGTTTAGTGGTTGC
CHUP1b_RT_R	TGTTTCTTGTGTATTCTCTTCTCCTGTTTGT

Movie S1: 3D image of Fig. 1A showing chloroplast focal accumulation at
haustoria and stromules interacting with each other and other chloroplasts. 3D
visualisation comprises Z-stack of confocal images of leaf epidermal cells from
transplastomic CpGFP (Yellow) *N. benthamiana* plants infected with red-fluorescent *P.*
infestans strain 88069td (Magenta).

Movie S2: 3D time-lapse series showing chloroplasts sieging haustoria. 3D
visualisation comprises Z-stack of confocal images of leaf epidermal cells from
transplastomic CpGFP (Yellow) *N. benthamiana* plants infected with red-fluorescent *P.*
infestans strain 88069td (Magenta).

Movie S3: 3D time-lapse series showing stromules engulfing a haustorium. 3D
visualisation comprises Z-stack of confocal images of leaf epidermal cells from
transplastomic CpGFP (Yellow) *N. benthamiana* plants infected with red-fluorescent *P.*
infestans strain 88069td (Magenta). Grayscale crop of CpGFP signal highlights the
extent of chloroplast stromules embracing the haustorium.

Movie S4: 3D image of chloroplast and stromules embracing a haustorium. 3D
visualisation comprises Z-stack of confocal images of leaf epidermal cells from
transplastomic CpGFP (Yellow) *N. benthamiana* plants expressing PM and EHM marker
RFP:REM1.3 (Magenta) infected with WT *P. infestans* 88069.

Movie S5: Time-lapse series of Fig. S1 showing chloroplasts accumulation to a
haustorium. Leaf epidermal cells from WT *N. benthamiana* plants expressing PM and
EHM marker RFP:REM1.3 (Magenta) and endosomal marker GFP:RAB8 (Yellow)
infected with WT *P. infestans* 88069. Blue is chlorophyll autofluorescence, labelling the
chloroplasts.

Movie S6: Time-lapse series showing optical trapping of chloroplast in Fig. 1G
which escapes the trap and springs back to the haustorium. TIRF microscopy
combined with laser capture in leaf epidermal cells from transplastomic CpGFP (channel
not shown) *N. benthamiana* plants expressing PM and EHM marker RFP:REM1.3
(Grayscale) infected with WT *P. infestans* 88069. Chloroplast is visible due to chlorophyll
autofluorescence overlapping with RFP emission spectrum. Scale bar is 10 µm.

Movie S7: 3D image of Fig. S3 showing chloroplasts form long-distance stromule
interactions that can bridge more than one haustorium. 3D visualisation comprises
Z-stack of confocal images of leaf epidermal cells from transplastomic CpGFP (Yellow)
N. benthamiana plants infected with red-fluorescent *P. infestans* strain 88069td
(Magenta). 'H' indicates haustorium.

Movie S8: 3D time-lapse series showing dynamic stromule interactions. 3D
visualisation comprises Z-stack of confocal images of leaf epidermal cells from
transplastomic CpGFP (Yellow) *N. benthamiana* plants infected with red-fluorescent *P.*
infestans strain 88069td (Magenta).

Movie S9: Time-lapse series showing optical trapping of chloroplast in Fig. 1H and
comigration of a second chloroplast interacting via a stromule-like extension.
TIRF microscopy combined with laser capture in leaf epidermal cells from transplastomic
CpGFP (Grayscale) *N. benthamiana* plants. Scale bar is 10 µm.

Movie S10: Time-lapse series of Fig. S3 showing collapse of haustorium engulfed
by a chloroplast. Leaf epidermal cells from WT *N. benthamiana* plants expressing
CHUP1:GFP (Yellow) and PM and EHM marker RFP:REM1.3 (Magenta) infected with
WT *P. infestans* 88069. Blue is chlorophyll autofluorescence, labelling the chloroplasts.
"H" indicates haustorium that collapses.

Movie S11: Time-lapse series of Fig. S3 showing collapse of haustorium
associated with a chloroplast. Leaf epidermal cells from WT *N. benthamiana* plants
expressing PM and EHM marker RFP:REM1.3 (Magenta) and endosomal marker
GFP:RAB8 (Yellow) infected with WT *P. infestans* 88069. Blue is chlorophyll
autofluorescence, labelling the chloroplasts. "H" indicates haustorium that collapses.

Movie S12: Time-lapse series showing synchronised steering of chloroplasts
away from haustorium. Leaf epidermal cells from transplastomic CpGFP (Yellow) *N.*
benthamiana plants expressing PM and EHM marker RFP:REM1.3 (Magenta) and GFP-
NbNRC2 (NLR, channel not shown) infected with WT *P. infestans* 88069.

Movie S13: Time-lapse series showing chloroplasts synchronised in pulling away
from haustorium. Leaf epidermal cells from transplastomic CpGFP (Yellow) *N.*
benthamiana plants expressing PM and EHM marker RFP:REM1.3 (Magenta) infected
with WT *P. infestans* 88069. "H" indicates haustorium the chloroplasts pull away from in
a synchronised manner.