Exaggerated in vivo IL-17 responses discriminate recall responses in active TB

Authors

Gabriele Pollara[1][*], Carolin T Turner[1], Gillian S Tomlinson[1], Lucy CK Bell[1], Ayesha Khan[1], Luis Felipe Peralta[2], Anna Folino[3], Ayse Akarca[1], Cristina Venturini[1], Tina Baker[1], Fabio LM Ricciardolo[3], Teresa Marafioti[1], Cesar Ugarte-Gil[2,4], David AJ Moore[4,5], Benjamin M Chain[1], Mahdad Noursadeghi[1]

[1] University College London, London, UK

[2] School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru

[3] Department of Clinical and Biological Sciences, University of Turin, Turin, Italy

[4] TB Centre, London School of Hygiene & Tropical Medicine, London, UK

[5] Laboratorio de Investigación de Enfermedades Infecciosas, Universidad Peruana Cayetano Heredia, Lima, Peru

Correspondence

Dr Gabriele Pollara – Division of Infection & Immunity, University College London, Gower Street, London, WC1E 6BT. g.pollara@ucl.ac.uk

Running title

IL-17 responses in active TB disease

Manuscript word count

7190 words

Online data

This article has an online data supplement, which includes:
- Supplementary figures S1-7
- Supplementary tables S1-7

Conflict of interest statement

The authors have declared that no conflict of interest exists.

Abstract

Background

Host immune responses at the site of *Mycobacterium tuberculosis* (Mtb) infection serve to contain the pathogen, but also mediate the pathogenesis of tuberculosis (TB) and onward transmission of infection. Based on the premise that active TB disease is predominantly a manifestation of immunopathology, we tested the hypothesis that immune responses at the site of host-pathogen interactions would reveal enrichment of immunopathologic responses in patients with active TB that were absent in individuals with equivalent immune memory for Mtb but without disease.

Methods

In cohorts of patients with active TB and cured or latent infection, we undertook molecular profiling at the site of a tuberculin skin test to model in vivo host-pathogen interactions in Mtb infection. Genome-wide transcriptional differences were identified by differential gene expression analyses. Enrichment of immune cells and cytokine activity was derived using specific transcriptional modules. Findings were validated in independent cohorts of patients with active TB, as well as Mtb infected tissues.

Results

Active TB in humans is associated with exaggerated IL-17A/F expression, accumulation of Th17 cells and IL-17A bioactivity, including increased neutrophil recruitment and matrix metalloproteinase-1 expression directly implicated in TB pathogenesis. These features discriminate recall responses in patients with active TB from those with cured or latent infection, and are also evident at the site of TB disease.

Conclusions

Our data are consistent with a model in which elevated Th17 responses within tissues drive immunopathology and transmission in active TB, and support targeting of the IL-17A/F pathway in host-directed therapy for active TB.

Abstract word count

249 words

Introduction

Mycobacterium tuberculosis (Mtb) infection results in a spectrum of clinical outcomes, from asymptomatic latent infection to symptomatic disease. The focus of host-pathogen interactions is characterised histologically by granuloma formation, a chronic inflammatory process that may contain the infection, but can also result in tissue damage that promotes transmission of infection to other individuals (1, 2). The distinctions that tip the balance between protective and pathogenic immune responses remain a fundamental question in tuberculosis research. This knowledge is expected to inform rational vaccine design and development of host-directed therapies (3).

Chronic inflammatory pathology at the site of human tuberculosis has been the subject of extensive descriptive studies, but discriminating between protective and pathological immune responses has been limited to comparing leucocyte phenotype and function in blood from Mtb-exposed patients with and without active disease (1). We have shown that genome-wide transcriptional profiling of biopsies of the tuberculin skin test (TST) can be used to make comprehensive molecular and systems level assessment of in vivo immune responses at the site of standardised host-pathogen interactions (4–7). Importantly, the transcripts enriched within the TST reflect the genome-wide variation in molecular pathology at the site of tuberculosis (TB) disease (5, 7), suggesting the TST represents a valuable surrogate for assessing TB immunopathogenesis in vivo.

On the premise that active TB disease is predominantly a manifestation of immunopathology, in this study we aimed to test the hypothesis that immune responses at the site of host-pathogen interactions, modelled by the TST, would reveal enrichment of immunopathologic responses in patients with active TB that were absent in individuals with equivalent immune memory for Mtb but without disease.

Results

Immune responses at the site of TST in active and cured TB

The TST has been most extensively used to identify patients with T cell memory for mycobacterial antigens, but the clinical response does not differentiate infected individuals with and without active disease (1). We sought to test the hypothesis that molecular profiling of the TST may identify elements of the recall response which are specifically associated with disease. We undertook 48-hour TSTs in patients with microbiologically-confirmed TB disease within the first month of treatment ('active TB') to identify disease associated responses, and compared these to TST responses in patients within one year of curative TB treatment ('cured TB') to identify non-disease

associated recall responses (table S1). Age, gender and site of TB disease were comparable between
the two groups (table S2). As expected, clinical induration in response to the TST was not
significantly different between the two groups (fig 1A), hitherto interpreted to reflect comparable
cell mediated immune memory.

In comparison to skin biopsies from the site of control saline injection, 1910 genes were significantly
enriched in response to the TST in at least one study group. Of these, 1251 were enriched in both
groups (fig 1B). Bioinformatic systems level assessment of the shared response revealed many
prototypic cell mediated immune responses which we had previously described in the TST (fig 1C)
(5). Pairwise assessment of the integrated list of transcripts that were enriched in either group
revealed statistically significant covariance, consistent with the hypothesis that the majority of
responses do not discriminate between the two groups (fig 1D).

Differential gene expression in the TST in active TB

A proportion of genes were differentially enriched between the two groups (fig 1D & tables S3-4). 44
genes were expressed significantly more in patients with active TB (table S3) compared to patients
with cured TB. Amongst these, pathway analysis identified statistically significant enrichment of
transcripts involved in extracellular matrix (ECM) remodelling, such as matrix metalloproteinase 1
(MMP-1), and beta defensins that both exert antimicrobial functions and also provide a chemotactic
gradient for CCR2-expressing cells, including neutrophils (8) (figs 2A & S1A). MMP-1, previously
implicated in pathogenic degradation of the ECM in TB (9), was the most over-expressed gene in
active TB compared to cured TB (fig 2B & table S3). This difference was validated at protein level by
immunohistochemistry, which also revealed that the differences in MMP-1 expression between
patients with active and cured TB was restricted to the inflammatory infiltrates within the TST (fig
2C-E).

Elevated IL-17 responses in active TB

We hypothesised that the genes over-expressed in active TB were regulated by common upstream
signals in the tissue environment. To test this hypothesis, we compared the predicted upstream
regulators of differentially expressed transcripts in the TST of active and cured TB patients using
Ingenuity Pathway Analysis (6). This analysis suggested that IL-17A induced the expression of genes
over-expressed in active TB and not those over-expressed in cured TB (fig 3A). In contrast, IFNγ was
predicted to be an upstream signal for gene expression enriched in both active and cured TB (fig 3A).
IFNγ responses, largely attributed to T helper (Th)-1 polarised CD4+ T cells are necessary for
immunological protection against Mtb (10), but they are insufficient and do not discriminate

between people who do and do not develop disease (1). The role of IL-17A in TB is less clear. This

cytokine belongs to a family of six structurally related cytokines and shares greatest sequence

homology with IL-17F. These bind the same receptor, and consequently exert the same functions,

particularly increased neutrophil recruitment via upregulation of chemokine expression (11).

There is unequivocal evidence that IL-17A/F contribute to host defence against bacterial and fungal

pathogens (12). Importantly however, they are also strongly indicated in the immunopathology of

chronic inflammatory diseases (12, 13). This includes evidence for IL-17A dependent neutrophil

mediated pathology in mouse models of Mtb infection (14–16). Our bioinformatics analysis

suggested increased enrichment of transcripts in the human in vivo recall response of patients with

active TB may be driven by IL-17A/F activity. Therefore, we sought to test the hypothesis that IL-

17A/F activity is exaggerated in active TB. Consistent with this hypothesis the expression of both IL-

17A and IL-17F were enriched in the TST of people with active TB compared to cured TB (fig 3B). In

contrast, IFNγ transcript levels representing the prototypic molecule in cell mediated immune recall

responses was not significantly different (fig 3B). Interestingly, the expression of IL-22, a cytokine

with closely related biological function to IL-17A and IL-17F (17), was also not elevated in active TB

(fig 3B). The differences revealed at the transcriptional level were also reflected by increased

immunofluorescence of IL-17F protein in the TST of people with active TB (fig 3C & 3D).

In order to test the functional significance of the differences in IL-17A/F expression between active

and cured TB, we evaluated differences in the bioactivity of IL-17A between these groups. We

addressed this by generating cellular response modules from the transcriptomes of

cytokine-stimulated keratinocytes (KC) (18, 19). We confirmed that these modules were both

sensitive and specific for their cognate stimuli by assessing their expression in other independent

datasets (fig S2). We then compared the geometric mean expression of these cytokine-specific

transcriptional modules in the TST transcriptomes. The IL-17A-induced gene module was significantly

increased in the TST of people with active TB compared to that of cured TB, but expression of IFNγ,

type I IFN or TNF-inducible gene modules was not significantly different between the two groups (fig

4A). We extended our approach to evaluating the functional bioactivity of other specific cytokines

using transcriptional modules for IL-10- and IL-4/IL-13-inducible gene expression, described in a

previous report (5). Neither of these were significantly different in the TST of people with active and

cured TB (fig S3A). The IL-4/IL-13 bioactivity module was used as a measure of Th2 responses

normally associated with allergy and immune responses to helminths. Another member of the IL-17

family, IL-17E contributes to Th2 responses (11). We found no enrichment of IL-17E expression in the

TST of people with active TB consistent with IL-4/IL-13 bioactivity and distinct from IL-17A/F

bioactivity (fig S3B).

Focusing on mechanisms that may contribute to pathogenesis, enrichment of MMP-1 expression in
the TST of patients with active TB (fig 2) can also be attributed to increased IL-17A/F bioactivity,
through induction of MMP expression by stromal cells (20). Neutrophils can also contribute to the
immunopathology of TB (14–16, 21, 22), and a key function of IL-17A/F is to promote neutrophil
recruitment via induction of neutrophil chemokines (12). Therefore, we tested the hypothesis that
the TST in active TB will also reveal increased neutrophil recruitment, compared to that of patients
with cured TB. We compared the expression of two independently derived transcriptional modules
that have been extensively validated to reflect neutrophil frequency in tissues including the TST (23,
24). We found significantly higher expression of the neutrophil modules in people with active TB
compared to cured TB (fig 4B). These differences were mirrored by gene expression levels of IL-17A-
inducible CXCL1, CXCL8 and S100A9 that drive neutrophil recruitment to sites of inflammation (11,
15) (fig 4C). In contrast, accumulation of monocytes and T cells assessed by their respective gene
expression modules (23), and the expression of the monocyte chemoattractant, CCL2, did not differ
in the TST of people with active and cured TB (fig 4B&C).

Increased frequency of Th17 cells in active TB TST responses

IL-17A/F are predominantly produced by Th17 cells and neutrophils (11). Immunohistochemistry
revealed that in the TST of people with active TB, IL-17F, originated predominantly from
mononuclear cells, rather than from polymorphonuclear cells (fig 5A). Therefore, we tested the
hypothesis that Th17 cells were enriched in the TST of people with active TB compared to that of
cured TB. We derived transcriptional modules specific for differentially polarised CD4+ Th subsets
from a published dataset (25) and demonstrated their specificity in an independent dataset (26) (fig
S4A). We further validated the Th17 module by showing that this correlated closely with the
expression of the IL-17A/F bioactivity module within skin biopsies of patients with psoriasis vulgaris,
representing an alternative Th17-mediated inflammatory condition (27) (fig S4B). The expression of
the Th1- and Th2-associated transcriptional modules in the TST was comparable between patients
with active and cured TB, but expression of the Th17 associated module was significantly increased
in patients with active TB (figs 5B & S3).

In order to investigate the mechanism for increased Th17 responses in the TST of patients with
active TB, we tested the hypothesis that active TB was also associated with increased circulating
Th17 cells. We assessed the expression of the Th subset modules in the transcriptome of PPD-
stimulated PBMC from an independent cohort of individuals with active TB disease and latent TB
infection (28). In contrast to that observed in the TST, patients with active TB revealed enrichment
for the Th1-associated gene expression module in PBMC but showed no difference in the

Th17-associated module (fig 5C). As a result, we explored the alternative hypothesis that the
inflammatory environment generated at the site of TST promotes the Th17 differentiation observed
in patients with active TB. We investigated the expression of cytokines implicated in Th17
differentiation, IL-1β, IL-6, IL-23 and TGFβ (11). The expression of each of these was significantly
correlated with enrichment for Th17 cells in the TST of active TB. In contrast there was no
correlation with IL-12 and IL-4 expression that drive Th1 and Th2 cell differentiation respectively (fig
5D). These data support a model in which the local tissue microenvironment may promote Th17
differentiation within tissues in active TB.

IL-17 activity is not a feature of latent TB and is not confounded by demographic
background, extrapulmonary disease or time on treatment

In order to validate our findings, we compared the TST transcriptome of a second independent
cohort of people with active TB with that of individuals with latent TB infection. Consistent with our
previous results, this comparison revealed significant enrichment for IL-17A/F bioactivity and
neutrophil infiltration in people with active TB (fig 6A). Furthermore, Th17 cells were enriched in
those with active TB despite there being no overall difference in T cell numbers compared to the TST
in individuals with latent TB (fig 6A). Of note, the comparison to latent TB in this analysis, also
excluded the possibility that lower levels of IL-17A/F bioactivity in the TST of people with cured TB
compared to active TB was an off-target effect of the antimicrobial treatment for active TB.

Taken together our data show that active TB is associated with exaggerated Th17 recall responses
and IL-17 bioactivity within the tissue microenvironment of host-pathogen interactions. Our findings
were replicated using two different methods for transcriptional profiling (microarray figs 4-5 and
RNA-Seq fig 6), and were consistent in geographically distinct cohorts (fig S5A). Significantly
exaggerated Th17 recall and IL-17A/F bioactivity in people with active TB compared to either cured
or latent TB were preserved in analyses including only UK cohorts (figs S5B & S5C), confirming that
the differences were not due to confounding by other variables in the different cohorts. In addition,
we found no difference in these responses between people with pulmonary and extrapulmonary TB
disease (fig S6). Of note, people with active TB were assessed at different time points within the first
month of treatment (median 11 days, IQR 5-28 days). Rapid changes in the peripheral blood
transcriptome associated with active TB have been reported (29, 30), but we found no diminution of
the exaggerated Th17 recall responses and IL-17A bioactivity invoked by the TST challenge in this
time frame (fig S7), suggesting that the mechanisms that drive these responses in active are not
swiftly reversed by treatment.

 IL-17 activity is present at the site of TB disease

Having established that IL-17A/F production by T cells was a prominent feature of the in vivo tissue
recall response to Mtb stimulation, we sought to determine whether this was also evident at the site
of TB disease. The transcriptome of human TB granuloma (31) showed an enrichment of T cells, Th17
cells and IL-17A bioactivity compared to normal lung tissue (fig 6B). In addition, we examined the
transcriptome of human Mtb-infected lymphadenitis compared to other 'reactive' causes of
lymphadenopathy devoid of granulomatous inflammation or malignancy (32). Interestingly, despite
the fact that reactive lymph nodes were enriched for the T cell transcriptional module, the Th17 and
IL-17A/F bioactivity modules were enriched within Mtb infected lymph nodes (fig 6C), further
confirming enrichment for IL-17A/F bioactivity to be a feature at the site of TB disease.

Discussion

Active TB disease is characterised by chronic inflammation that can result in significant tissue
destruction, necessary for the onward transmission of Mtb (2). Identifying the processes that govern
this immunopathology offers the opportunity to intervene therapeutically, limiting tissue damage
and transmission of infection. Pathogenic immune pathways have been difficult to identify because
most components of the immune response to Mtb do not discriminate between different clinical
outcomes of infection (1). The present study provides compelling new evidence that IL-17A/F
responses may mediate immunopathology in active TB. The inclusion of multiple cohorts and diverse
demographic backgrounds increased the generalisability of our findings and circumvented the
limitations of our cross-sectional study design. The findings are consistent with the well-established
role of IL-17A/F in the immunopathology of chronic inflammatory human disease exemplified by
psoriasis, for which blockade of the IL-17A/F axis provides an effective treatment (13).

Importantly, the primary physiological role of IL-17A/F is to promote host defence against bacterial
and fungal infection, cogently demonstrated by mice deficient for IL-17A/F or IL-17 receptors, and by
humans with inborn errors of IL-17 immunity (12). Mouse models also suggest a protective role for
IL-17A/F following BCG vaccination (33) and in the early stages of Mtb infection, particularly in the
context of more virulent Mtb strains (34, 35), through localising T cells near Mtb-infected
macrophages (34) and by preventing formation of necrotic granuloma (36). However, in mice
rendered susceptible to TB disease as a result of IFNγ deficiency or through receiving repeated BCG
vaccinations, IL-17A/F responses drive neutrophil-mediated pathology (14–16, 21). We infer from
these data that IL-17A/F responses may play a dichotomous role in TB by contributing to protection
in early infection, but to the pathology of disease if early infection is not controlled and
multibacillary bacterial replication and chronic immune activation ensues. Additional support for this

model is evident in IL-27R deficient mice, which exhibit exaggerated Th17 responses because they

lack IL-27 inhibition of RORγT (37). These mice show enhanced clearance of Mtb, but also increased

immunopathology dependent on IL-17A. Interestingly, a human candidate gene study identified host

genetic variation associated with increased secretion of IL-17A to be correlated with both protection

from incident TB but also more severe TB disease (38). Taken together, we hypothesise that

exaggerated IL-17A/F recall responses are the consequence of chronic multibacillary infection, but in

turn mediate increased pathology in established disease. Of the other IL-17 family members that

signal via alternative receptors, much less is known about the functional role of IL-17B, IL-17C and IL-

17D (11). These cytokine responses were not specifically tested in the present study. IL-17E, known

to promote Th2 responses (11), was not differentially enriched in active TB. Equally, we found no

evidence of elevated IL-10 responses in active TB, indicating an uncoupling of IL-17A/F activity from

regulatory responses, driving immunopathology rather than the control of Mtb replication (39).

Another cytokine, IL-22, does have functional overlap with IL-17A/F (17), but did not discriminate

between people with and without disease in our study.

Our data support the hypothesis that exaggerated IL-17A/F responses arise from Th17 cells, but they

do not unequivocally exclude other cellular sources. Our immunohistochemical analysis did not show

any clear evidence for IL-17F production by polymorphonuclear cells in the TST, but we were not

able to test whether other lymphocyte populations, such as γδ T cells, made a significant

contribution (35, 40, 41). Future studies will require single cell resolution to definitively confirm the

source of exaggerated IL-17A/F responses in this model, as well as to determine whether active TB

shows enrichment for 'pathogenic' Th17 cells that express both IFNg and IL-17A/F (42). Consistent

with previous studies (40, 43), we found no evidence of increased circulating Th17 cells in active TB,

although this was measured in a separate active TB cohort to the ones that underwent TST

assessment. Nevertheless, this observation underscores the unique value of molecular level

assessment of immune responses in the tissue microenvironment at the site of host-pathogen

interactions, modelled by the TST. Importantly, it also suggests the model that differential Th17

responses within the TST are governed by immune signalling networks within the tissue

microenvironment after T cell recruitment. In active TB patients, the tissue expression of cytokines

that promote Th17 differentiation correlated with the transcriptional module for Th17 cells. The

source of these cytokines is likely to be myeloid cells (11, 13, 44). Although we cannot prove a causal

relationship at this stage, our data are consistent with a model in which infiltrating monocytes from

patients with active TB secrete elevated levels of cytokines that promote the differentiation of

recruited T cells to a Th17 phenotype. Of note, active TB patients have higher frequency of

circulating CD14+ CD16+ non-classical monocytes (45), which can potentiate the differentiation of
CD4 T cells to a Th17 phenotype in patients with chronic inflammatory conditions (46).

Our findings challenge the long-established view that curative treatment of TB does not lead to
contemporaneous changes to immunological recall responses (47). Our data are consistent with the
hypothesis that IL-17A-inducible neutrophil chemotaxis and expression of MMP-1 represent key
mediators of immunopathology in TB by promoting granuloma formation, bacterial replication and
matrix degradation that can lead to cavitation and onward transmission (9, 48). Our results support
future studies to evaluate the impact of modulating IL-17A/F activity to ameliorate the pathology
associated with chronic TB disease. The availability of therapies that block the IL-17A/F cytokine axis,
developed and licensed for chronic inflammatory diseases (13), offers invaluable opportunities to
transition from proof of concept pre-clinical studies, for example in non-human primate models, to
first in man experiments.

Methods

Study populations

The study comprised recruitment of several different populations. The discovery 'Active TB' group
that formed the basis of most of the analyses in figs 1-5 was the HIV seronegative patients from the
same cohort described in our previous publication (5), who were recruited from TB clinics in London,
UK and Cape Town, South Africa, and were all within one month of commencing antibiotic therapy.
The comparator 'Cured TB' group was an independent cohort recruited from TB clinics in London, UK
and Lima, Peru who fulfilled the inclusion/exclusion criteria (table S1). All 'Cured TB' patients were
less than 2 years after completion of curative anti-TB antibiotic therapy for drug sensitive disease. In
addition, a separate validation cohort of patients with active TB was recruited from TB clinics in
London (fig 6). These patients were also within one month of commencing antibiotic therapy. This
population was compared to individuals with latent TB recruited from TB clinics in London and Lima,
who fulfilled inclusion/exclusion criteria (table S1). All study participants were HIV seronegative and
for those with active or cured TB, the presence of Mtb infection was confirmed by routine culture or
molecular based methods according to local clinic protocols. The demographic, clinical and
laboratory data for each study group is summarised in table S2.
312

Study approval

314   Recruitment of patients with cured and latent TB was approved by UK National Research Ethics
315   Committee (reference number: 14/LO/0505) and Universidad Peruana Cayetano Heredia
316   Institutional Ethics Committee (reference number: 62349). Recruitment of patients for the validation
317   active TB cohort was approved UK National Research Ethics Committee (reference number:
318   16/LO/0776).
319

Study schedule and sampling

On recruitment to the study, all participants received 0.1 mL intradermal injection of two units
tuberculin (Serum Statens Institute) or saline in the volar aspect of one forearm, and this site was
marked with indelible ink. At 48 hours, the clinical response at the injection site was evaluated by
measurement of the maximum diameter of inflammatory induration and two 3 mm adjacent punch
biopsies were obtained from marked TST or saline injection site as previously described (4). One
biopsy was placed in RNAlater (Thermo Fisher) and stored at -70°C, and the other biopsy was placed
in 10% formalin neutral buffered solution (Sigma-Aldrich) and stored at room temperature for at
least 1 week prior to paraffin embedding.
329

Sample processing

The TST transcriptome from active TB patients in the discovery cohort was derived directly from the
data repository E-MTAB-3254 (ArrayExpress - https://www.ebi.ac.uk/arrayexpress/). Skin samples
from all other participants was stored in RNAlater at -70oC after collection. For processing, TST
samples were equilibrated to room temperature for 30 minutes before being transferred to CK14
lysing kit tubes (Bertin Instruments) containing 350µl of Buffer RLT (Qiagen) supplemented with 1%
2-Mercaptoethanol (Sigma). Tubes were pulsed for 6 cycles on a Precellys Evolution homogeniser
(Bertin Instruments), each cycle consisting of 23 seconds of homogenisation at speeds of 6300 rpm.
Samples were rested on ice for 2 minutes between cycles. After homogenisation, cellular debris and
lysing beads were precipitated by centrifugation and RNA isolated from the supernatant using
RNeasy Mini Kit (Qiagen). Total RNA from TST samples of patients with cured TB was purified,
labelled and hybridised on Agilent 8x60k microarrays as previously described (5). For the validation
cohort of active TB and individuals with latent TB, the KAPA mRNA HyperPrep Kit (Roche Diagnostics)
was used to construct stranded mRNA-Seq libraries from up to 500 ng intact total RNA after which
paired-end sequencing was carried out using the 75 cycle high-output kit on the NextSeq 500
desktop sequencer (Illumina). Each run contained 24 samples and was demultiplexed using bcl2fastq

by Illumina ([https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-](https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-software.html)
[software.html](https://support.illumina.com/sequencing/sequencing_software/bcl2fastq-conversion-software.html)). Paired end reads were mapped to the Ensembl human transcriptome reference
sequence (homo sapiens GRCh38, latest version available). Mapping and generation of read counts
per transcript were done using Kallisto (49). R/Bioconductor package tximport was used to import
the mapped counts data and summarise the transcripts-level data into gene level (50).

Whole genome transcriptional profiling and analysis software

Raw microarray data was processed and normalised as previously described (51). Raw RNASeq
counts were normalised within-sample into TPM (transcripts per million) to remove feature-length
and library-size effects (52). Log2 transformed TPM were used for further analysis. Data matrices
from non-TST datasets were obtained from processed data series downloaded at the ArrayExpress
repository. Probe identifiers were converted to gene symbols using platform annotations provided
with each dataset. In circumstances where downloaded datasets were not log2 transformed, this
was performed on the entire processed data matrix. Significant gene expression differences
between datasets were calculated from normalised expression matrices using MultiExperiment
Viewer v4.9 (http://www.tm4.org/mev.html). Pathway analysis was performed in InnateDB (53) and
visualized as network diagrams in Gephi v0.8.2 beta. Upstream regulator analysis was performed
using Ingenuity Pathway Analysis (Qiagen), focusing on cytokines with predicted activation z-score
>2. Genes predicted to be regulated in either active or cured TB formed the basis of the network
diagram in fig 3. The expression of transcriptional modules was determined by calculating the
geometric mean expression of all the module constituent genes found in the dataset being analysed,
using R scripts generated in our previous publication (23), and which are available to download and
use from the Github repository (https://github.com/MJMurray1/MDIScoring). Venn diagrams were
constructed using the BioVenn tool (http://www.biovenn.nl/).

Transcriptomic data repositories

372 All transcriptional datasets used in this study are described in table S5. Accession numbers refer to
373 datasets in the ArrayExpress repository (https://www.ebi.ac.uk/arrayexpress/). The TST
374 transcriptome of the discovery cohort of patients with active TB was derived from dataset E-MTAB-
375 3254, the cured TB TST transcriptome was derived from dataset E-MTAB-6815, and the validation
376 active TB and latent TB TST transcriptomes were derived from dataset E-MTAB-6816.

Module derivation and expression

Immune cell modules used were ones rated with the highest Module Discriminatory Index (MDI)
score for module sensitivity and specificity as determined in our previous publication (23). These
were "M19" (T cells), "M37.1" (neutrophils) and "Monocyte 2-fold" (monocytes). We also made use
of another neutrophil module "ImSig" that was derived and validated elsewhere (24). We have
previously generated and validated the specificity of macrophage response modules to IL-10 or
IL4/IL-13 stimulation (5), and these are also utilised in this manuscript.

To derive keratinocyte (KC) cytokine-response modules, we made use of previously published
transcriptomic data (GSE12109 & GSE36287) from primary human keratinocytes (KCs) stimulated
with a selection of cytokines (18, 19). Significant transcriptional responses (paired t-test with α of
p<0.05 and no multiple testing correction) of genes over-expressed >4-fold in the cognate cytokine
condition relative to unstimulated KC were initially identified. Modules were then derived from
genes that were not also upregulated > 2-fold by non-cognate cytokine conditions compared to
unstimulated KC. The KC IL-17 response module utilised a cut-off of 2-fold between IL-17 stimulated
KC and unstimulated KC as too few genes were upregulated >4-fold compared to unstimulated KC.
The constituent genes of the KC modules are shown in table S6. The KC TNFa response module
generated in this way has already been published (6), but the other cytokine modules are newly
described. The gene components of the KC cytokine-response modules are available in table S6, and
their specificity was evaluated in both datasets in which they were derived and in independent
datasets of in vitro cytokine-stimulated KC (fig S1A).

The transcriptome of CD4+ T cells polarised towards different T helper phenotypes was derived from
dataset GSE54627 (25). To derive specific modules for each differentiation state, we used gene
expression from cells stimulated with anti-CD3 and anti-CD28, identifying genes over-expressed in
the phenotype of interest compared to all other conditions by paired t-test with α of p<0.05 and no
multiple testing correction. Each module was derived from the unique genes over expressed by
more than 1.5-fold in the cognate condition compared to all other stimulation conditions (except for
Th1 module where a 2 -fold cut-off was used). The gene components of T helper modules are
available in table S7, and their specificity was evaluated in the dataset from which they were
derived, in an independent dataset of polarised CD4 T cells, and in skin biopsies of patients with
psoriasis vulgaris (fig S1B) (25–27).

Immunostaining

Immunostaining of IL-17F was performed on 10 μm sections as previously described (54). Briefly,
monoclonal mouse-anti-IL-17F, 1:50, code MA5-16229 [Thermo Fisher Scientific, Waltham, MA, USA]
and DAPI (sc-24941, Santa Cruz Biotechnology, Texas, Dallas, USA) for detecting nuclei were used.
Irrelevant primary antibodies from Sigma (Milan, Italy; irrelevant mouse, 1:50, code I8765) were
applied at the same concentration of the related specific primary antibodies for immunostaining of
negative control slides, and as a further negative control, secondary antibody (AF568, goat anti-
mouse; Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) alone was used (data not shown).
Light-microscopic analysis was performed at a magnification of 40x with a Leica DM4000B
microscope equipped with DFC-320 Leica digital camera (Leica Microsystems, Wetzlar, Germany).
Zeiss confocal microscope (LSM800, Carl Zeiss, Germany) was used to acquire confocal images.
Quantification of IL-17F was performed on 10x confocal images of samples through the
measurement of mean grey value (MGV) using software ImageJ (https://imagej.net/Fiji).

Immunostaining of MMP-1 was performed as previously described (55, 56). Scanned slide images
were obtained with use of NanoZoomer Digital Pathology System (Hamamatsu, Japan.
Quantification of MMP-1 staining was performed blindly by extracting MMP-1 associated 3, 3 -
diaminobenzidine (DAB) stain using standard deconvolution protocols in ImageJ. Cellular infiltrates
were manually selected as depicted in Fig 2 and DAB stain quantified by staining intensity as
proportion of the area selected using ImageJ. The selected region was moved without resizing to
adjacent tissue that did not contain cellular infiltration to calculate background MMP-1 intensity.
Three cellular infiltrates and background tissue regions were analysed for each tissue samples. Six
TST samples were quantified in each group.

Author contributions

Substantial contributions to the conception or design of the work;	GP, CT, GST, LCB LFP, TM, CU, DAM, BMC, MN
Substantial contributions to the acquisition, analysis, or interpretation of data for the work;	GP, CT, GST, LCB, AK, LFP, AF, AA, CV, TB, FLR, TM, BMC, MN
Drafting the work or revising it critically for important intellectual content;	GP, CT, GST, LCB, BMC, MN
Final approval of the version submitted for publication	All authors
Accountability for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved	GP, MN

Acknowledgements

The work presented in this manuscript received funding from the Wellcome Trust (WT101766/Z/13/Z to GP and 207511/Z/17/Z to MN), Medical Research Council (MR_N007727_1 to GST) and National Institute for Health Research Biomedical Research Centre at University College London Hospitals.

We thank our team of research nurses, in particular Cristina F Turienzo and Victoria Dean, for their assistance in recruiting patients and collecting samples, and the UCL-UCLH Pathogen Genomics Unit for RNASeq sample processing (https://www.ucl.ac.uk/infection-immunity/pathogen-genomics-unit).

References

1. O'Garra A et al. The Immune Response in Tuberculosis. *Annu. Rev. Immunol.* 2013;31(1):475–527.

2. Elkington PT, Ugarte-Gil CA, Friedland JS. Matrix metalloproteinases in tuberculosis. *Eur. Respir. J.* 2011;38(2):456–464.

3. Tiberi S et al. Tuberculosis: progress and advances in development of new drugs, treatment regimens, and host-directed therapies. *Lancet Infect. Dis.* 2018;18(7):e183–e198.

4. Tomlinson GS et al. Transcriptional profiling of innate and adaptive human immune responses to mycobacteria in the tuberculin skin test. *Eur. J. Immunol.* 2011;41(11):3253–3260.

5. Bell LCK et al. In Vivo Molecular Dissection of the Effects of HIV-1 in Active Tuberculosis. *PLOS Pathog* 2016;12(3):e1005469.

6. Byng-Maddick R et al. Tumor Necrosis Factor (TNF) Bioactivity at the Site of an Acute Cell-Mediated Immune Response Is Preserved in Rheumatoid Arthritis Patients Responding to Anti-TNF Therapy. *Front. Immunol.* 2017;8:932.

7. Lachmandas E et al. Tissue Metabolic Changes Drive Cytokine Responses to Mycobacterium tuberculosis [Internet]. *J. Infect. Dis.* [published online ahead of print: 2018]; doi:10.1093/infdis/jiy173

8. Röhrl J, Yang D, Oppenheim JJ, Hehlgans T. Human β-Defensin 2 and 3 and Their Mouse Orthologs Induce Chemotaxis through Interaction with CCR2. *J. Immunol.* 2010;184(12):6688–6694.

9. Elkington P et al. MMP-1 drives immunopathology in human tuberculosis and transgenic mice. *J. Clin. Invest.* 2011;121(5):1827–1833.

10. Jouanguy E et al. Partial interferon-gamma receptor 1 deficiency in a child with tuberculoid

bacillus Calmette-Guérin infection and a sibling with clinical tuberculosis.. *J. Clin. Invest.*

1997;100(11):2658–2664.

11. Iwakura Y, Ishigame H, Saijo S, Nakae S. Functional Specialization of Interleukin-17 Family

Members. *Immunity* 2011;34(2):149–162.

12. Li J, Casanova J-L, Puel A. Mucocutaneous IL-17 immunity in mice and humans: host defense vs.

excessive inflammation. *Mucosal Immunol.* 2018;11(3):581–589.

13. Hawkes JE, Chan TC, Krueger JG. Psoriasis pathogenesis and the development of novel targeted

immune therapies. *J. Allergy Clin. Immunol.* 2017;140(3):645–653.

14. Cruz A et al. Pathological role of interleukin 17 in mice subjected to repeated BCG vaccination

after infection with Mycobacterium tuberculosis. *J. Exp. Med.* 2010;207(8):1609–1616.

15. Gopal R et al. S100A8/A9 Proteins Mediate Neutrophilic Inflammation and Lung Pathology during

Tuberculosis. *Am. J. Respir. Crit. Care Med.* 2013;188(9):1137–1146.

16. Desvignes L, Ernst JD. Interferon-γ-Responsive Nonhematopoietic Cells Regulate the Immune

Response to Mycobacterium tuberculosis. *Immunity* 2009;31(6):974–985.

17. Ronacher K, Sinha R, Cestari M. IL-22: An Underestimated Player in Natural Resistance to

Tuberculosis? [Internet]. *Front. Immunol.* 2018;9. doi:10.3389/fimmu.2018.02209

18. Nograles KE et al. Th17 cytokines interleukin (IL)-17 and IL-22 modulate distinct inflammatory

and keratinocyte-response pathways. *Br. J. Dermatol.* 2008;159(5):1092–1102.

19. Swindell WR et al. Heterogeneity of Inflammatory and Cytokine Networks in Chronic Plaque

Psoriasis. *PLOS ONE* 2012;7(3):e34594.

20. Singh S et al. Interleukin-17 regulates matrix metalloproteinase activity in human pulmonary

tuberculosis. *J. Pathol.* 2018;244(3):311–322.

21. Nandi B, Behar SM. Regulation of neutrophils by interferon-γ limits lung inflammation during

tuberculosis infection., Regulation of neutrophils by interferon-γ limits lung inflammation during

tuberculosis infection. *J. Exp. Med. J. Exp. Med.* 2011;208, 208(11, 11):2251, 2251–2262.

22. Marais S et al. Neutrophil-Associated Central Nervous System Inflammation in Tuberculous

Meningitis Immune Reconstitution Inflammatory Syndrome. *Clin. Infect. Dis.* 2014;59(11):1638–

1647.

23. Pollara G et al. Validation of Immune Cell Modules in Multicellular Transcriptomic Data. *PLOS*

ONE 2017;12(1):e0169271.

24. Nirmal AJ et al. Immune Cell Gene Signatures for Profiling the Microenvironment of Solid

Tumors. *Cancer Immunol. Res.* 2018;6(11):1388–1400.

25. Touzot M et al. Combinatorial flexibility of cytokine function during human T helper cell

differentiation [Internet]. *Nat. Commun.* 2014;5. doi:10.1038/ncomms4987

26. Ramesh R et al. Pro-inflammatory human Th17 cells selectively express P-glycoprotein and are

refractory to glucocorticoids. *J. Exp. Med.* 2014;211(1):89–104.

27. Russell CB et al. Gene expression profiles normalized in psoriatic skin by treatment with

brodalumab, a human anti-IL-17 receptor monoclonal antibody.. *J. Immunol. Baltim. Md 1950*

2014;192(8):3828–3836.

28. Lu C et al. Novel Biomarkers Distinguishing Active Tuberculosis from Latent Infection Identified

by Gene Expression Profile of Peripheral Blood Mononuclear Cells. *PLoS ONE* 2011;6(8):e24290.

29. Bloom CI et al. Detectable changes in the blood transcriptome are present after two weeks of

antituberculosis therapy. *PloS One* 2012;7(10):e46191.

30. Cliff JM et al. Distinct Phases of Blood Gene Expression Pattern Through Tuberculosis Treatment

Reflect Modulation of the Humoral Immune Response. *J. Infect. Dis.* 2013;207(1):18–29.

31. Kim M-J et al. Caseation of human tuberculosis granulomas correlates with elevated host lipid

metabolism. *EMBO Mol. Med.* 2010;2(7):258–274.

32. Tomlinson GS et al. Transcriptional Profiling of Endobronchial Ultrasound-Guided Lymph Node

Samples Aids Diagnosis of Mediastinal Lymphadenopathy. *CHEST* 2016;149(2):535–544.

33. Khader SA et al. IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell

responses after vaccination and during Mycobacterium tuberculosis challenge. *Nat. Immunol.*

2007;8(4):369–377.

34. Gopal R et al. Unexpected Role for IL-17 in Protective Immunity against Hypervirulent

Mycobacterium tuberculosis HN878 Infection. *PLoS Pathog.* 2014;10(5):e1004099.

35. Yoshida YO et al. Essential Role of IL-17A in the Formation of a Mycobacterial Infection-Induced

Granuloma in the Lung. *J. Immunol.* 2010;184(8):4414–4422.

36. Domingo-Gonzalez R et al. Interleukin-17 limits hypoxia-inducible factor 1α and development of

hypoxic   granulomas   during   tuberculosis   [Internet].   *JCI   Insight*   2017;2(19).

doi:10.1172/jci.insight.92973

37. Erdmann H et al. The increased protection and pathology in Mycobacterium tuberculosis -

infected IL-27R-alpha-deficient mice is supported by IL-17A and is associated with the IL-17A-

induced expansion of multifunctional T cells. *Mucosal Immunol.* 2018;11(4):1168–1180.

38. Rolandelli A et al. The IL-17A rs2275913 single nucleotide polymorphism is associated with

protection to tuberculosis but related to higher disease severity in Argentina. *Sci. Rep.* 2017;7:40666.

39. Gideon HP et al. Variability in Tuberculosis Granuloma T Cell Responses Exists, but a Balance of

Pro- and Anti-inflammatory Cytokines Is Associated with Sterilization. *PLOS Pathog*

2015;11(1):e1004603.

40. Coulter F et al. IL-17 Production from T Helper 17, Mucosal-Associated Invariant T, and γδ Cells in

Tuberculosis Infection and Disease. *Front. Immunol.* 2017;8:1252.

41. Lockhart E, Green AM, Flynn JL. IL-17 production is dominated by gammadelta T cells rather than

CD4 T cells during Mycobacterium tuberculosis infection. *J. Immunol. Baltim. Md 1950*

2006;177(7):4662–4669.

42. Hu D et al. Transcriptional signature of human pro-inflammatory T H 17 cells identifies reduced

IL10 gene expression in multiple sclerosis. *Nat. Commun.* 2017;8(1):1600.

43. Chen X et al. Reduced Th17 Response in Patients with Tuberculosis Correlates with IL-6R

Expression on CD4+ T Cells. *Am. J. Respir. Crit. Care Med.* 2010;181(7):734–742.

44. Krausgruber T et al. IRF5 promotes inflammatory macrophage polarization and TH1-TH17

responses. *Nat. Immunol.* 2011;12(3):231–238.

45. Lastrucci C et al. Tuberculosis is associated with expansion of a motile, permissive and

immunomodulatory CD16[+] monocyte population via the IL-10/STAT3 axis. *Cell Res.*

2015;25(12):1333.

46. Rossol M, Kraus S, Pierer M, Baerwald C, Wagner U. The CD14brightCD16+ monocyte subset is

expanded in rheumatoid arthritis and promotes expansion of the Th17 cell population. *Arthritis*

Rheum. 2012;64(3):671–677.

47. Clifford V, He Y, Zufferey C, Connell T, Curtis N. Interferon gamma release assays for monitoring

the response to treatment for tuberculosis: A systematic review. *Tuberc. Edinb. Scotl.*

2015;95(6):639–650.

48. Parasa VR et al. Inhibition of Tissue Matrix Metalloproteinases Interferes with Mycobacterium

tuberculosis-Induced Granuloma Formation and Reduces Bacterial Load in a Human Lung Tissue

Model [Internet]. *Front. Microbiol.* 2017;8. doi:10.3389/fmicb.2017.02370

49. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification.

Nat. Biotechnol. 2016;34(5):525–527.

50. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates

improve gene-level inferences. *F1000Research* 2015;4:1521.

51. Chain B et al. Error, reproducibility and sensitivity: a pipeline for data processing of Agilent

oligonucleotide expression arrays. *BMC Bioinformatics* 2010;11:344.

52. Conesa A et al. A survey of best practices for RNA-seq data analysis. *Genome Biol.* 2016;17:13.

53. Breuer K et al. InnateDB: systems biology of innate immunity and beyond--recent updates and

continuing curation. *Nucleic Acids Res.* 2013;41(Database issue):D1228-1233.

54. Ricciardolo FLM et al. Identification of IL-17F/frequent exacerbator endotype in asthma. *J.*

Allergy Clin. Immunol. 2017;140(2):395–406.

55. Marafioti T et al. Novel markers of normal and neoplastic human plasmacytoid dendritic cells.

Blood 2008;111(7):3778–3792.

56. Akarca AU et al. BRAF V600E mutation-specific antibody, a sensitive diagnostic marker revealing

minimal residual disease in hairy cell leukaemia. *Br. J. Haematol.* 2013;162(6):848–851.

Figure 1. TST transcriptome in patients with active and cured TB disease. A) Induration at the site of TST was recorded by routine clinical assessments in both populations (mm) and was not different between the two groups (Mann-Whitney test). B) Venn diagram depicting genes significantly upregulated 2-fold in the TST of patients with active or cured TB relative to control saline injection. (p<0.01 by t-test with no multiple testing correction). C) TST transcriptome common to both active and cured TB summarised as a network diagram. Purple nodes represent Reactome database functional pathways, yellow nodes represent genes and edges reflect relationship between pathways and genes. Pathway node diameters are proportional to the respective pathway –log10 p value enrichment statistic. D) Pairwise dot-plot of 1910 gene integrated TST signature in patients with either active or cured TB. Dotted line reflects line of perfect covariance. p value derived from linear regression model between the two variables. Red and blue dots represent genes >2 fold

581   differentially expressed (p<0.01 by t-test with no multiple testing correction) between active and
582   cured TB.

Figure 2. MMP-1 over-expression in active TB. A) Network diagram of genes and Reactome pathways over-expressed in the TST of patients with active TB compared to cured TB. Purple nodes represent Reactome database functional pathways, yellow nodes represent genes and edges reflect relationship between pathways and genes. Pathway node diameters are proportional to the respective pathway −log10 p value enrichment statistic. B) mRNA expression in TST of patients with active and cured TB of MMP1 gene. C) Representative MMP-1 immunohistochemistry staining in inflammatory infiltrates with TST samples from patients with active and cured TB. D) Differential MMP-1 staining intensity for 18 cellular infiltrates in each group of patients relative to adjacent zones of skin with no cellular infiltration. E) MMP-1 staining intensity in TST zones outside inflammatory infiltrates. * = p<0.01 by Mann-Whitney test.

Figure 3. IL-17 is over expressed in active TB. A) Network diagram depicting upstream cytokine analysis of genes differentially expressed in TST of patients with active and cured TB. Red and blue nodes represent significant genes overexpressed in active and cured TB respectively as identified in fig 1D. Green nodes represent predicted cytokines regulating the gene expression of red and blue nodes. Edges depict relationship between upstream regulators and differentially expressed genes. B) Expression of selected cytokine transcript within the TSTs of active and cured TB patients. C) Expression of IL-17F by immunofluorescence in TST of patients with active TB. Left panel = phase contrast image, right panel = IL-17F positivity (white). D) Quantification of IL-17F staining throughout TST sections from patients with active and cured TB, determined by pixel intensity as a proportion of area sampled. Each dot represents IL-17F expression in the cross-section of an entire TST biopsy from one patient. * p<0.01 by Mann-Whitney test.

Figure 4. TST challenge in active TB is characterised by enrichment of IL-17 responses. Enrichment in TST relative to saline injection of A) keratinocyte response modules of cytokine bioactivity, B) immune cell modules and C) CXCL1, CXCL8, S100A9 and CCL2 genes. Neutrophil (Neut.) modules M37.1 and ImSig validated for sensitivity and specificity in references (23, 24) respectively. * = $p < 0.01$ by Mann-Whitney test.

Figure 5. Active TB is characterised by elevated Th17 cells. A) IL-17F immunohistochemistry in TST of patients with active TB. Black arrows point to mononuclear cells that express IL-17F and white arrows point to polymorphonuclear cells that do not express IL-17F. B) Th17 and Th1 cell module enrichment in TST relative to saline injection. C) Expression of Th17 and Th1 modules in PPD-stimulated PBMC from patients with active or latent TB (data originated from dataset GSE27984). * = p<0.01 by Mann-Whitney test. D) Relationship between expression of Th17 transcriptional module and individual cytokine gene expression in TST of cytokines implicated in polarising CD4+ T cells to a Th17 phenotype. Data points represent patients with active TB, Spearman rank correlation coefficients (r^2) and p values.

Figure 6. Th17 cells and IL-17 activity are a feature of both TST reactions and sites of human TB disease. A) Enrichment in TST relative to saline injection of transcriptional modules for IL-17 bioactivity, neutrophils, Th17 cells and T cells in TST of independent cohort of patients with active TB and a separate cohort of individuals with latent TB infection. B) Expression of T cells, Th17 cells and IL-17 bioactivity modules from the site of human TB granuloma relative to healthy lung tissue (dataset GSE20050) and C) in human TB infected lymph nodes (LN) relative to reactive lymph nodes that do not display evidence of granulomatous inflammation or cancer (dataset E-MTAB-2547). * = p<0.01 by Mann-Whitney test.