Exaggerated in vivo IL-17 responses discriminate recall responses in active TB

Authors

<u>Gabriele Pollara</u>[1*], Carolin T Turner[1], Gillian S Tomlinson[1], Lucy CK Bell[1], Ayesha Khan[1], Luis Felipe

Peralta[2], Anna Folino[3], Ayse Akarca[1], Cristina Venturini[1], Tina Baker[1], Fabio LM Ricciardolo[3], Teresa

Marafioti[1], Cesar Ugarte-Gil[2,4], David AJ Moore[4,5], Benjamin M Chain[1], Mahdad Noursadeghi[1]

[1] University College London, London, UK

[2] School of Medicine, Universidad Peruana Cayetano Heredia, Lima, Peru

[3] Department of Clinical and Biological Sciences, University of Turin, Turin, Italy

[4] TB Centre, London School of Hygiene & Tropical Medicine, London, UK

[5] Laboratorio de Investigación de Enfermedades Infecciosas, Universidad Peruana Cayetano Heredia,

Lima, Peru

Correspondence

Dr Gabriele Pollara – Division of Infection & Immunity, University College London, Gower Street,

London, WC1E 6BT. g.pollara@ucl.ac.uk

Running title

IL-17 responses in active TB disease

Manuscript word count

3405 words

Online data

This article has an online data supplement, which includes:

• Supplementary figures 1-8
• Supplementary tables E1-8
• Supplementary methods

Support

The work presented in this manuscript received funding from the Wellcome Trust
(WT101766/Z/13/Z to GP and 207511/Z/17/Z to MN), Medical Research Council (MR_N007727_1 to
GST) and National Institute for Health Research Biomedical Research Centre at University College
London Hospitals.

Author contributions

Substantial contributions to the conception or design of the work;	GP, CT, GST, LCB LFP, TM, CU, DAM, BMC, MN
Substantial contributions to the acquisition, analysis, or interpretation of data for the work;	GP, CT, GST, LCB, AK, LFP, AF, AA, CV, TB, FLR, TM, BMC, MN
Drafting the work or revising it critically for important intellectual content;	GP, CT, GST, LCB, BMC, MN
Final approval of the version submitted for publication	All authors
Accountability for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved	GP, MN

Abstract

Host immune responses at the site of *Mycobacterium tuberculosis* (Mtb) infection serve to contain the

pathogen, but also mediate the pathogenesis of tuberculosis (TB) and onward transmission of

infection. Interferon gamma (IFNγ) responses do not discriminate between protection and

pathogenicity, but IL-17A/F responses, known to drive pathology in diverse chronic inflammatory

diseases, have also been associated with TB pathogenesis in animal models. At the site of in vivo

immune recall responses to Mtb modelled by the tuberculin skin test, we show for the first time that

active TB in humans is also associated with exaggerated IL-17A/F expression, accumulation of Th17

cells and IL-17A/F bioactivity, including increased neutrophil recruitment and matrix

metalloproteinase-1 expression directly implicated in TB pathogenesis. These features discriminate

recall responses in patients with active TB from those with cured or latent infection and are also

evident at the site of TB disease. Our data support targeting of this pathway in host-directed therapy

for TB.

Abstract word count

156 words

Introduction

Mycobacterium tuberculosis (Mtb) infection results in a spectrum of clinical outcomes, from asymptomatic latent infection to symptomatic disease. The focus of host-pathogen interactions is characterised histologically by granuloma formation, a chronic inflammatory process that may contain the infection, but can also result in tissue damage that promotes transmission of infection to other individuals (1, 2). The distinctions that tip the balance between protective and pathogenic immune responses remain a fundamental question in tuberculosis research. This knowledge is expected to inform rational vaccine design and development of host-directed therapies (3).

Chronic inflammatory pathology at the site of human tuberculosis has been the subject of extensive descriptive studies, but discriminating between protective and pathological immune responses has been limited to comparing leucocyte phenotype and function in blood from Mtb-exposed patients with and without active disease (1). We have shown that genome-wide transcriptional profiling of biopsies of the tuberculin skin test (TST) can be used to make comprehensive molecular and systems level assessment of in vivo immune responses at the site of standardised host-pathogen interactions (4–7). Importantly, the transcripts enriched within the TST reflect the genome-wide variation in molecular pathology at the site of tuberculosis (TB) disease (5, 7), suggesting the TST represents a valuable surrogate for assessing TB immunopathogenesis in vivo.

On the premise that active TB disease is predominantly a manifestation of immunopathology, in this study we aimed to test the hypothesis that immune responses at the site of host-pathogen interactions, modelled by the TST, would reveal enrichment of immunopathologic responses in patients with active TB that were absent in individuals with equivalent immune memory for Mtb but without disease.

Results

Immune responses at the site of TST in active and cured TB

The TST has been most extensively used to identify patients with T cell memory for mycobacterial antigens, but the clinical response does not differentiate infected individuals with and without active disease (1). We sought to test the hypothesis that molecular profiling of the TST may identify elements of the recall response which are specifically associated with disease. We undertook 48-hour TSTs in patients with microbiologically-confirmed TB disease within the first month of treatment ('active TB') to identify disease associated responses, and compared these to TST responses in patients within one year of curative TB treatment ('cured TB') to identify non-disease associated recall responses (table

E1). Age, gender and site of TB disease were comparable between the two groups (table E2). As

expected, clinical induration in response to the TST was not significantly different between the two

groups (fig 1A), hitherto interpreted to reflect comparable cell mediated immune memory.

In comparison to skin biopsies from the site of control saline injection, 1910 genes were significantly

enriched in response to the TST in at least one study group. Of these, 1251 were enriched in both

groups (fig 1B). Bioinformatic systems level assessment of the shared response revealed many

prototypic cell mediated immune responses which we had previously described in the TST (fig 1C) (5).

Pairwise assessment of the integrated list of transcripts that were enriched in either group revealed

statistically significant covariance, consistent with the hypothesis that the majority of responses do

not discriminate between the two groups (fig 1D).

Differential gene expression in the TST in active TB

A proportion of genes were differentially enriched between the two groups (fig 1D & tables E3-4). 44

genes were expressed significantly more in patients with active TB (table E3) compared to patients

with cured TB. Amongst these, pathway analysis identified statistically significant enrichment of

transcripts involved in extracellular matrix (ECM) remodelling, such as matrix metalloproteinase 1

(MMP-1), and beta defensins that both exert antimicrobial functions and also provide a chemotactic

gradient for CCR2-expressing cells, including neutrophils (8) (figs 2A & E1A). MMP-1, previously

implicated in pathogenic degradation of the ECM in TB (9), was the most over-expressed gene in

active TB compared to cured TB (fig 2B & table E3). This difference was validated at protein level by

immunohistochemistry, which also revealed that the differences in MMP-1 expression between

patients with active and cured TB was restricted to the inflammatory infiltrates within the TST (fig 2C-

E).

Elevated IL-17 responses in active TB

We hypothesised that the genes over-expressed in active TB were regulated by common upstream

signals in the tissue environment. To test this hypothesis, we compared the predicted upstream

regulators of differentially expressed transcripts in the TST of active and cured TB patients using

Ingenuity Pathway Analysis (6). This analysis suggested that IL-17A induced the expression of genes

over-expressed in active TB and not those over-expressed in cured TB (fig 3A). In contrast, IFNγ was

predicted to be an upstream signal for gene expression enriched in both active and cured TB (fig 3A).

IFNγ responses, largely attributed to T helper (Th)-1 polarised CD4+ T cells are necessary for

immunological protection against Mtb (10), but they are insufficient and do not discriminate between

people who do and do not develop disease (1). The role of IL-17A in TB is less clear. This cytokine

belongs to a family of six structurally related cytokines and shares greatest sequence homology with

IL-17F. These bind the same receptor, and consequently exert the same functions, particularly

increased neutrophil recruitment via upregulation of chemokine expression (11).

There is unequivocal evidence that IL-17A/F contribute to host defence against bacterial and fungal

pathogens (12). Importantly however, they are also strongly indicated in the immunopathology of

chronic inflammatory diseases (12, 13). This includes evidence for IL-17A dependent neutrophil

mediated pathology in mouse models of Mtb infection (14–16). Our bioinformatics analysis suggested

increased enrichment of transcripts in the human in vivo recall response of patients with active TB

may be driven by IL-17A/F activity. Therefore, we sought to test the hypothesis that IL-17A/F activity

is exaggerated in active TB. Consistent with this hypothesis the expression of both IL-17A and IL-17F

were enriched in the TST of people with active TB compared to cured TB (fig 3B). In contrast, IFNγ

transcript levels representing the prototypic molecule in cell mediated immune recall responses was

not significantly different (fig 3B). Interestingly, the expression of IL-22, a cytokine with closely related

biological function to IL-17A and IL-17F (17), was also not elevated in active TB (fig 3B). The differences

revealed at the transcriptional level were also reflected by increased immunofluorescence of IL-17F

protein in the TST of people with active TB (fig 3C & 3D).

In order to test the functional significance of the differences in IL-17A/F expression between active

and cured TB, we evaluated differences in the bioactivity of IL-17A between these groups. We

addressed this by generating cellular response modules from the transcriptomes of

cytokine-stimulated keratinocytes (KC) (18, 19). We confirmed that these modules were both sensitive

and specific for their cognate stimuli by assessing their expression in other independent datasets (fig

E2 & table E6). We then compared the geometric mean expression of these cytokine-specific

transcriptional modules in the TST transcriptomes. The IL-17A-induced gene module was significantly

increased in the TST of people with active TB compared to that of cured TB, but expression of IFNγ,

type I IFN or TNF-inducible gene modules was not significantly different between the two groups (fig

4A). We extended our approach to evaluating the functional bioactivity of other specific cytokines

using transcriptional modules for IL-10- and IL-4/IL-13-inducible gene expression, described in a

previous report (5). Neither of these were significantly different in the TST of people with active and

cured TB (fig E3A). The IL-4/IL-13 bioactivity module was used as a measure of Th2 responses normally

associated with allergy and immune responses to helminths. Another member of the IL-17 family, IL-

17E contributes to Th2 responses (11). We found no enrichment of IL-17E expression in the TST of

people with active TB consistent with IL-4/IL-13 bioactivity and distinct from IL-17A/F bioactivity (fig

E3B).

Focusing on mechanisms that may contribute to pathogenesis, enrichment of MMP-1 expression in

the TST of patients with active TB (fig 2) can also be attributed to increased IL-17A/F bioactivity,

through induction of MMP expression by stromal cells (20). Neutrophils can also contribute to the

immunopathology of TB (14–16, 21, 22), and a key function of IL-17A/F is to promote neutrophil

recruitment via induction of neutrophil chemokines (12). Therefore, we tested the hypothesis that the

TST in active TB will also reveal increased neutrophil recruitment, compared to that of patients with

cured TB. We compared the expression of two independently derived transcriptional modules that

have been extensively validated to reflect neutrophil frequency in tissues including the TST (23, 24).

We found significantly higher expression of the neutrophil modules in people with active TB compared

to cured TB (fig 4B). These differences were mirrored by gene expression levels of IL-17A-inducible

CXCL1, CXCL8 and S100A9 that drive neutrophil recruitment to sites of inflammation (11, 15) (fig 4C).

In contrast, accumulation of monocytes and T cells assessed by their respective gene expression

modules (23), and the expression of the monocyte chemoattractant, CCL2, did not differ in the TST of

people with active and cured TB (fig 4B&C).

Increased frequency of Th17 cells in active TB TST responses

IL-17A/F are predominantly produced by Th17 cells and neutrophils (11). Immunohistochemistry

revealed that in the TST of people with active TB, IL-17F, originated predominantly from mononuclear

cells, rather than from polymorphonuclear cells (fig 5A). Therefore, we tested the hypothesis that

Th17 cells were enriched in the TST of people with active TB compared to that of cured TB. We derived

transcriptional modules specific for differentially polarised CD4+ Th subsets from a published dataset

(25) and demonstrated their specificity in an independent dataset (26) (fig E4A & table E7). We further

validated the Th17 module by showing that this correlated closely with the expression of the IL-17A/F

bioactivity module within skin biopsies of patients with psoriasis vulgaris, representing an alternative

Th17-mediated inflammatory condition (27) (fig E4B). The expression of the Th1- and Th2-associated

transcriptional modules in the TST was comparable between patients with active and cured TB, but

expression of the Th17 associated module was significantly increased in patients with active TB (figs

5B & E3).

In order to investigate the mechanism for increased Th17 responses in the TST of patients with active

TB, we tested the hypothesis that active TB was also associated with increased circulating Th17 cells.

We assessed the expression of the Th subset modules in the transcriptome of PPD-stimulated PBMC

from an independent cohort of individuals with active TB disease and latent TB infection (28). In

contrast to that observed in the TST, patients with active TB revealed enrichment for the

Th1-associated gene expression module in PBMC but showed no difference in the Th17-associated

module (fig 5C). As a result, we explored the alternative hypothesis that the inflammatory

environment generated at the site of TST promotes the Th17 differentiation observed in patients with

active TB. We investigated the expression of cytokines implicated in Th17 differentiation, IL-1β, IL-6,

IL-23 and TGFβ (11). The expression of each of these was significantly correlated with enrichment for

Th17 cells in the TST of active TB. In contrast there was no correlation with IL-12A and IL-4 expression

that drive Th1 and Th2 cell differentiation respectively (fig 5D). To test the hypothesis that conditions

favouring Th17 differentiation were present in vivo, we generated and validated a transcriptional

response module for IL-1 activity in human fibroblasts (fig E5 and table E8) in order to quantify

functional IL-1 activity in the TST. This analysis confirmed increased expression of IL-1 inducible genes

in the TST of patients with active TB (fig 5E). Of note, this finding was consistent with the bioinformatic

prediction of IL1 signalling upstream of some transcripts that were differentially enriched

predominantly in active TB (fig 3A). Overall, these data support a model in which the local tissue

microenvironment, in part driven by elevated IL-1 activity, may promote Th17 differentiation within

tissues in active TB.

IL-17 activity is not a feature of latent TB and is not confounded by demographic background,

extrapulmonary disease or time on treatment

In order to validate our findings, we compared the TST transcriptome of a second independent cohort

of people with active TB with that of individuals with latent TB infection. Consistent with our previous

results, this comparison revealed significant enrichment for IL-17A/F bioactivity and neutrophil

infiltration in people with active TB (fig 6A). Furthermore, Th17 cells were enriched in those with active

TB despite there being no overall difference in T cell numbers compared to the TST in individuals with

latent TB (fig 6A). Of note, the comparison to latent TB in this analysis, also excluded the possibility

that lower levels of IL-17A/F bioactivity in the TST of people with cured TB compared to active TB was

an off-target effect of the antimicrobial treatment for active TB.

Taken together our data show that active TB is associated with exaggerated Th17 recall responses and

IL-17 bioactivity within the tissue microenvironment of host-pathogen interactions. Our findings were

replicated using two different methods for transcriptional profiling (microarray figs 4-5 and RNA-Seq

fig 6), and were consistent in geographically distinct cohorts (fig E6A). Significantly exaggerated Th17

recall and IL-17A/F bioactivity in people with active TB compared to either cured or latent TB were

preserved in analyses including only UK cohorts (figs E6B & E6C), confirming that the differences were

not due to confounding by other variables in the different cohorts. In addition, we found no difference

in these responses between people with pulmonary and extrapulmonary TB disease (fig E7). Of note,

people with active TB were assessed at different time points within the first month of treatment

(median 11 days, IQR 5-28 days). Rapid changes in the peripheral blood transcriptome associated with

active TB have been reported (29, 30), but we found no diminution of the exaggerated Th17 recall

responses and IL-17 bioactivity invoked by the TST challenge in this time frame (fig E8), suggesting that

the mechanisms that drive these responses in active are not swiftly reversed by treatment.

IL-17 activity is present at the site of TB disease

Having established that IL-17A/F production by T cells was a prominent feature of the in vivo tissue

recall response to Mtb stimulation, we sought to determine whether this was also evident at the site

of TB disease. The transcriptome of human TB granuloma (31) showed an enrichment of T cells, Th17

cells and IL-17A bioactivity compared to normal lung tissue (fig 6B). In addition, we examined the

transcriptome of human Mtb-infected lymphadenitis compared to other 'reactive' causes of

lymphadenopathy devoid of granulomatous inflammation or malignancy (32). Interestingly, despite

the fact that reactive lymph nodes were enriched for the T cell transcriptional module, the Th17 and

IL-17A/F bioactivity modules were enriched within Mtb infected lymph nodes (fig 6C), further

confirming enrichment for IL-17A/F bioactivity to be a feature at the site of TB disease.

Discussion

Active TB disease is characterised by chronic inflammation that can result in significant tissue

destruction, necessary for the onward transmission of Mtb (2). Identifying the processes that govern

this immunopathology offers the opportunity to intervene therapeutically, limiting tissue damage and

transmission of infection. Pathogenic immune pathways have been difficult to identify because most

components of the immune response to Mtb do not discriminate between different clinical outcomes

of infection (1). The present study provides compelling new evidence that IL-17A/F responses may

mediate immunopathology in active TB. The inclusion of multiple cohorts and diverse demographic

backgrounds increased the generalisability of our findings and circumvented the limitations of our

cross-sectional study design. The findings are consistent with the well-established role of IL-17A/F in

the immunopathology of chronic inflammatory human disease exemplified by psoriasis, for which

blockade of the IL-17A/F axis provides an effective treatment (13).

Importantly, the primary physiological role of IL-17A/F is to promote host defence against bacterial

and fungal infection, cogently demonstrated by mice deficient for IL-17A/F or IL-17 receptors, and by

humans with inborn errors of IL-17 immunity (12). Mouse models also suggest a protective role for IL-

17A/F following BCG vaccination (33) and in the early stages of Mtb infection, particularly in the

context of more virulent Mtb strains (34, 35), through localising T cells near Mtb-infected

macrophages (34) and by preventing formation of necrotic granuloma (36). However, in mice

rendered susceptible to TB disease as a result of IFNγ deficiency or through receiving repeated BCG

vaccinations, IL-17A/F responses drive neutrophil-mediated pathology (14–16, 21). We infer from

these data that IL-17A/F responses may play a dichotomous role in TB by contributing to protection in

early infection, but to the pathology of disease if early infection is not controlled and multibacillary

bacterial replication and chronic immune activation ensues. Additional support for this model is

evident in IL27R deficient mice, which exhibit exaggerated Th17 responses because they lack IL-27

inhibition of RORγT (37). These mice show enhanced clearance of Mtb, but also increased

immunopathology dependent on IL-17A. Interestingly, a human candidate gene study identified host

genetic variation associated with increased secretion of IL-17A to be correlated with both protection

from incident TB but more severe TB disease (38). Taken together, we hypothesise that exaggerated

IL-17A/F recall responses are the consequence of chronic multibacillary infection, but in turn mediate

increased pathology in established disease. Of the other IL-17 family members that signal via

alternative receptors, much less is known about the functional role of IL-17B, IL-17C and IL-17D (11).

These cytokine responses were not specifically tested in the present study. IL-17E, known to promote

Th2 responses (11), was not differentially enriched in active TB. Equally, we found no evidence of

elevated IL-10 responses in active TB, indicating an uncoupling of IL-17A/F activity from regulatory

responses, driving immunopathology rather than the control of Mtb replication (39). Another

cytokine, IL-22, does have functional overlap with IL-17A/F (17), but did not discriminate between

people with and without disease in our study.

Our data support the hypothesis that exaggerated IL-17A/F responses arise from Th17 cells, but they

do not unequivocally exclude other cellular sources. Our immunohistochemical analysis did not show

any clear evidence for IL-17F production by polymorphonuclear cells in the TST, but we were not able

to test whether other lymphocyte populations, such as γδ T cells, made a significant contribution (35,

40, 41). Future studies will require single cell resolution to definitively confirm the source of

exaggerated IL-17A/F responses in this model, as well as to determine whether active TB shows

enrichment for 'pathogenic' Th17 cells that express both IFNg and IL-17A/F (42). Consistent with

previous studies (40, 43), we found no evidence of increased circulating Th17 cells in active TB,

although this was measured in a separate active TB cohort to the ones that underwent TST

assessment. Nevertheless, this observation underscores the unique value of molecular level

assessment of immune responses in the tissue microenvironment at the site of host-pathogen

interactions, modelled by the TST. Importantly, it also suggests the model that differential Th17

responses within the TST are governed by immune signalling networks within the tissue

microenvironment after T cell recruitment. In active TB patients, the tissue expression of cytokines

that promote Th17 differentiation correlated with the transcriptional module for Th17 cells, and

elevated IL-1 activity was present in the TST. The source of these cytokines is likely to be myeloid cells

(11, 13, 44). Although we cannot prove a causal relationship at this stage, our data are consistent with

a model in which infiltrating monocytes cells from patients with active TB secrete elevated levels of

cytokines that promote the differentiation of recruited T cells to a Th17 phenotype. Of note, active TB

patients have higher frequency of circulating CD14+ CD16+ non-classical monocytes (45), which can

potentiate the differentiation of CD4 T cells to a Th17 phenotype in patients with chronic inflammatory

conditions (46).

Our findings challenge the long-established view that curative treatment of TB does not lead to

contemporaneous changes to immunological recall responses (47). Our data are consistent with the

hypothesis that IL-17A-inducible neutrophil chemotaxis and expression of MMP-1 represents a key

mediator of immunopathology in TB by promoting granuloma formation, bacterial replication and

matrix degradation that can lead to cavitation and onward transmission (9, 48). Our results support

future studies to evaluate the impact of modulating IL-17A/F activity to ameliorate the pathology

associated with chronic TB disease. The availability of therapies that block the IL-17A/F cytokine axis,

developed and licensed for chronic inflammatory diseases (13), offers invaluable opportunities to

transition from proof of concept pre-clinical studies, for example in non-human primate models, to

first in man experiments.

Methods

Study populations

298 The cohorts of patients recruited to the study are described in detail in Supplementary Methods.

299 Inclusion and exclusion criteria are in table E1 and the demographic, clinical and laboratory data for

300 each study group is summarised in table E2.

Study approval

302 Recruitment of patients with cured and latent TB was approved by UK National Research Ethics

303 Committee (reference number: 14/LO/0505) and Universidad Peruana Cayetano Heredia Institutional

304 Ethics Committee (reference number: 62349). Recruitment of patients for the validation active TB

305 cohort was approved UK National Research Ethics Committee (reference number: 16/LO/0776).

Study schedule and sampling

307   On recruitment to the study, all participants received 0.1 mL intradermal injection of two units
308   tuberculin (Serum Statens Institute) or saline in the volar aspect of one forearm, and this site was
309   marked with indelible ink. Measurement of the TST site, biopsy collection and sample storage was
310   performed as previously described (4) and is expanded in more detail in Supplementary Methods.

Sample processing

312   The TST transcriptome from active TB patients in the discovery cohort was derived directly from the
313   data repository E-MTAB-3254 (ArrayExpress - https://www.ebi.ac.uk/arrayexpress/). Skin samples
314   from all other participants were processed for RNA extraction and transcriptional profiling by either
315   microarrays or RNA-Seq as described in Supplementary Methods.

Whole genome transcriptional profiling and analysis software

317   Raw microarray data was processed and normalised as previously described (49). Raw RNASeq counts
318   were normalised within-sample into TPM (transcripts per million) to remove feature-length and
319   library-size effects (50). Log2 transformed TPM were used for further analysis, including
320   transcriptional module derivation and expression that is described in further detail in Supplementary
321   Methods.

Transcriptomic data repositories

323   All transcriptional datasets used in this study are described in table E5. Accession numbers refer to
324   datasets in the ArrayExpress repository (https://www.ebi.ac.uk/arrayexpress/). The TST
325   transcriptome of the discovery cohort of patients with active TB was derived from dataset E-MTAB-
326   3254, the cured TB TST transcriptome was derived from dataset E-MTAB-6815, and the validation
327   active TB and latent TB TST transcriptomes were derived from dataset E-MTAB-6816.

Immunostaining

329   Immunostaining of IL-17F was performed on 10 μm sections as previously described (51).
330   Immunostaining of MMP-1 was performed as previously described (52). More detailed methods for
331   both these targets, as well as approaches to quantify staining intensity, is available in Supplementary
332   Methods.

Acknowledgements

We thank our team of research nurses, in particular Cristina F Turienzo and Victoria Dean, for their

assistance in recruiting patients and collecting samples, and the UCL-UCLH Pathogen Genomics Unit

for RNASeq sample processing (https://www.ucl.ac.uk/infection-immunity/pathogen-genomics-unit).

None of the authors have commercial affiliations or consultancies, stock or equity interests, or patent-

licensing arrangements that could be considered a conflict of interest regarding the submitted

manuscript.

References

1.   O'Garra A, Redford PS, McNab FW, Bloom CI, Wilkinson RJ, Berry MPR. The Immune Response in

Tuberculosis. *Annu Rev Immunol* 2013;31:475–527.

2.   Elkington PT, Ugarte-Gil CA, Friedland JS. Matrix metalloproteinases in tuberculosis. *Eur Respir J*

2011;38:456–464.

3.   Tiberi S, du Plessis N, Walzl G, Vjecha MJ, Rao M, Ntoumi F, Mfinanga S, Kapata N, Mwaba P,

McHugh TD, Ippolito G, Migliori GB, Maeurer MJ, Zumla A. Tuberculosis: progress and advances

in development of new drugs, treatment regimens, and host-directed therapies. *Lancet Infect Dis*

2018;18:e183–e198.

4.   Tomlinson GS, Cashmore TJ, Elkington PTG, Yates J, Lehloenya RJ, Tsang J, Brown M, Miller RF,

Dheda K, Katz DR, Chain BM, Noursadeghi M. Transcriptional profiling of innate and adaptive

human immune responses to mycobacteria in the tuberculin skin test. *Eur J Immunol*

2011;41:3253–3260.

5.   Bell LCK, Pollara G, Pascoe M, Tomlinson GS, Lehloenya RJ, Roe J, Meldau R, Miller RF, Ramsay A,

Chain BM, Dheda K, Noursadeghi M. In Vivo Molecular Dissection of the Effects of HIV-1 in Active

Tuberculosis. *PLOS Pathog* 2016;12:e1005469.

6.   Byng-Maddick R, Turner CT, Pollara G, Ellis M, Guppy NJ, Bell LCK, Ehrenstein MR, Noursadeghi M.

Tumor Necrosis Factor (TNF) Bioactivity at the Site of an Acute Cell-Mediated Immune Response

Is Preserved in Rheumatoid Arthritis Patients Responding to Anti-TNF Therapy. *Front Immunol*

2017;8:932.

7.   Lachmandas E, Rios-Miguel AB, Koeken VACM, van der Pasch E, Kumar V, Matzaraki V, Li Y, Oosting

M, Joosten LAB, Notebaart RA, Noursadeghi M, Netea MG, van Crevel R, Pollara G. Tissue

Metabolic Changes Drive Cytokine Responses to Mycobacterium tuberculosis. *J Infect Dis*

2018;doi:10.1093/infdis/jiy173.

8.   Röhrl J, Yang D, Oppenheim JJ, Hehlgans T. Human β-Defensin 2 and 3 and Their Mouse Orthologs

Induce Chemotaxis through Interaction with CCR2. *J Immunol* 2010;184:6688–6694.

9.   Elkington P, Shiomi T, Breen R, Nuttall RK, Ugarte-Gil CA, Walker NF, Saraiva L, Pedersen B, Mauri

F, Lipman M, Edwards DR, Robertson BD, D'Armiento J, Friedland JS. MMP-1 drives

immunopathology in human tuberculosis and transgenic mice. *J Clin Invest* 2011;121:1827–1833.

10. Jouanguy E, Lamhamedi-Cherradi S, Altare F, Fondanèche MC, Tuerlinckx D, Blanche S, Emile JF,

Gaillard JL, Schreiber R, Levin M, Fischer A, Hivroz C, Casanova J-L. Partial interferon-gamma

receptor 1 deficiency in a child with tuberculoid bacillus Calmette-Guérin infection and a sibling

with clinical tuberculosis. *J Clin Invest* 1997;100:2658–2664.

11. Iwakura Y, Ishigame H, Saijo S, Nakae S. Functional Specialization of Interleukin-17 Family

Members. *Immunity* 2011;34:149–162.

12. Li J, Casanova J-L, Puel A. Mucocutaneous IL-17 immunity in mice and humans: host defense vs.

excessive inflammation. *Mucosal Immunol* 2018;11:581–589.

13. Hawkes JE, Chan TC, Krueger JG. Psoriasis pathogenesis and the development of novel targeted

immune therapies. *J Allergy Clin Immunol* 2017;140:645–653.

14. Cruz A, Fraga AG, Fountain JJ, Rangel-Moreno J, Torrado E, Saraiva M, Pereira DR, Randall TD,

Pedrosa J, Cooper AM, Castro AG. Pathological role of interleukin 17 in mice subjected to repeated

BCG vaccination after infection with Mycobacterium tuberculosis. *J Exp Med* 2010;207:1609–

1616.

15. Gopal R, Monin L, Torres D, Slight S, Mehra S, McKenna KC, Fallert Junecko BA, Reinhart TA, Kolls

J, Báez-Saldaña R, Cruz-Lagunas A, Rodríguez-Reyna TS, Kumar NP, Tessier P, Roth J, Selman M,

Becerril-Villanueva E, Baquera-Heredia J, Cumming B, Kasprowicz VO, Steyn AJC, Babu S, Kaushal

D, Zúñiga J, Vogl T, Rangel-Moreno J, Khader SA. S100A8/A9 Proteins Mediate Neutrophilic

Inflammation and Lung Pathology during Tuberculosis. *Am J Respir Crit Care Med* 2013;188:1137–

1146.

16. Desvignes L, Ernst JD. Interferon-γ-Responsive Nonhematopoietic Cells Regulate the Immune

Response to Mycobacterium tuberculosis. *Immunity* 2009;31:974–985.

17. Ronacher K, Sinha R, Cestari M. IL-22: An Underestimated Player in Natural Resistance to

Tuberculosis? *Front Immunol* 2018;9:.

18. Nograles KE, Zaba LC, Guttman E, Fuentes-Duculan J, Suarez-Farinas M, Cardinale I, Khatcherian

A, Gonzalez J, Pierson KC, White TR, Pensabene C, Coats I, Novitskaya I, Lowes MA, Krueger JG.

Th17 cytokines interleukin (IL)-17 and IL-22 modulate distinct inflammatory and keratinocyte-

response pathways. *Br J Dermatol* 2008;159:1092–1102.

19. Swindell WR, Xing X, Stuart PE, Chen CS, Aphale A, Nair RP, Voorhees JJ, Elder JT, Johnston A,

Gudjonsson JE. Heterogeneity of Inflammatory and Cytokine Networks in Chronic Plaque

Psoriasis. *PLOS ONE* 2012;7:e34594.

20. Singh S, Maniakis-Grivas G, Singh UK, Asher RM, Mauri F, Elkington PT, Friedland JS. Interleukin-

17 regulates matrix metalloproteinase activity in human pulmonary tuberculosis. *J Pathol*

2018;244:311–322.

21. Nandi B, Behar SM. Regulation of neutrophils by interferon-γ limits lung inflammation during

tuberculosis infection., Regulation of neutrophils by interferon-γ limits lung inflammation during

tuberculosis infection. *J Exp Med J Exp Med* 2011;208, 208:2251, 2251–2262.

22. Marais S, Wilkinson KA, Lesosky M, Coussens AK, Deffur A, Pepper DJ, Schutz C, Ismail Z, Meintjes

G, Wilkinson RJ. Neutrophil-Associated Central Nervous System Inflammation in Tuberculous

Meningitis Immune Reconstitution Inflammatory Syndrome. *Clin Infect Dis* 2014;59:1638–1647.

23. Pollara G, Murray MJ, Heather JM, Byng-Maddick R, Guppy N, Ellis M, Turner CT, Chain BM,

Noursadeghi M. Validation of Immune Cell Modules in Multicellular Transcriptomic Data. In: Mills

K, editor. *PLOS ONE* 2017;12:e0169271.

24. Nirmal AJ, Regan T, Shih BB, Hume DA, Sims AH, Freeman TC. Immune Cell Gene Signatures for

Profiling the Microenvironment of Solid Tumors. *Cancer Immunol Res* 2018;6:1388–1400.

25. Touzot M, Grandclaudon M, Cappuccio A, Satoh T, Martinez-Cingolani C, Servant N, Manel N,

Soumelis V. Combinatorial flexibility of cytokine function during human T helper cell

differentiation. *Nat Commun* 2014;5:.

26. Ramesh R, Kozhaya L, McKevitt K, Djuretic IM, Carlson TJ, Quintero MA, McCauley JL, Abreu MT,

Unutmaz D, Sundrud MS. Pro-inflammatory human Th17 cells selectively express P-glycoprotein

and are refractory to glucocorticoids. *J Exp Med* 2014;211:89–104.

27. Russell CB, Rand H, Bigler J, Kerkof K, Timour M, Bautista E, Krueger JG, Salinger DH, Welcher AA,

Martin DA. Gene expression profiles normalized in psoriatic skin by treatment with brodalumab,

a human anti-IL-17 receptor monoclonal antibody. *J Immunol Baltim Md 1950* 2014;192:3828–

3836.

28. Lu C, Wu J, Wang H, Wang S, Diao N, Wang F, Gao Y, Chen J, Shao L, Weng X, Zhang Y, Zhang W.

Novel Biomarkers Distinguishing Active Tuberculosis from Latent Infection Identified by Gene

Expression Profile of Peripheral Blood Mononuclear Cells. *PLoS ONE* 2011;6:e24290.

29. Bloom CI, Graham CM, Berry MPR, Wilkinson KA, Oni T, Rozakeas F, Xu Z, Rossello-Urgell J,

Chaussabel D, Banchereau J, Pascual V, Lipman M, Wilkinson RJ, O'Garra A. Detectable changes in

the blood transcriptome are present after two weeks of antituberculosis therapy. *PloS One*

2012;7:e46191.

30. Cliff JM, Lee J-S, Constantinou N, Cho J-E, Clark TG, Ronacher K, King EC, Lukey PT, Duncan K,

Helden PDV, Walzl G, Dockrell HM. Distinct Phases of Blood Gene Expression Pattern Through

Tuberculosis Treatment Reflect Modulation of the Humoral Immune Response. *J Infect Dis*

2013;207:18–29.

31. Kim M-J, Wainwright HC, Locketz M, Bekker L-G, Walther GB, Dittrich C, Visser A, Wang W, Hsu F-

F, Wiehart U, Tsenova L, Kaplan G, Russell DG. Caseation of human tuberculosis granulomas

correlates with elevated host lipid metabolism. *EMBO Mol Med* 2010;2:258–274.

32. Tomlinson GS, Thomas N, Chain BM, Best K, Simpson N, Hardavella G, Brown J, Bhowmik A, Navani

N, Janes SM, Miller RF, Noursadeghi M. Transcriptional Profiling of Endobronchial Ultrasound-

Guided Lymph Node Samples Aids Diagnosis of Mediastinal Lymphadenopathy. *CHEST*

2016;149:535–544.

33. Khader SA, Bell GK, Pearl JE, Fountain JJ, Rangel-Moreno J, Cilley GE, Shen F, Eaton SM, Gaffen SL,

Swain SL, Locksley RM, Haynes L, Randall TD, Cooper AM. IL-23 and IL-17 in the establishment of

protective pulmonary CD4+ T cell responses after vaccination and during Mycobacterium

tuberculosis challenge. *Nat Immunol* 2007;8:369–377.

34. Gopal R, Monin L, Slight S, Uche U, Blanchard E, A. Fallert Junecko B, Ramos-Payan R, Stallings CL,

Reinhart TA, Kolls JK, Kaushal D, Nagarajan U, Rangel-Moreno J, Khader SA. Unexpected Role for

IL-17 in Protective Immunity against Hypervirulent Mycobacterium tuberculosis HN878 Infection.

In: Lewinsohn DM, editor. *PLoS Pathog* 2014;10:e1004099.

35. Yoshida YO, Umemura M, Yahagi A, O'Brien RL, Ikuta K, Kishihara K, Hara H, Nakae S, Iwakura Y,

Matsuzaki G. Essential Role of IL-17A in the Formation of a Mycobacterial Infection-Induced

Granuloma in the Lung. *J Immunol* 2010;184:4414–4422.

36. Domingo-Gonzalez R, Das S, Griffiths KL, Ahmed M, Bambouskova M, Gopal R, Gondi S, Muñoz-

Torrico M, Salazar-Lezama MA, Cruz-Lagunas A, Jiménez-Álvarez L, Ramirez-Martinez G, Espinosa-

Soto R, Sultana T, Lyons-Weiler J, Reinhart TA, Arcos J, Garcia-Hernandez M de la L, Mastrangelo

MA, Al-Hammadi N, Townsend R, Balada-Llasat J-M, Torrelles JB, Kaplan G, Horne W, Kolls JK,

Artyomov MN, Rangel-Moreno J, Zúñiga J, *et al.* Interleukin-17 limits hypoxia-inducible factor 1α

and development of hypoxic granulomas during tuberculosis. *JCI Insight* 2017;2:.

37. Erdmann H, Behrends J, Ritter K, Hölscher A, Volz J, Rosenkrands I, Hölscher C. The increased

protection and pathology in Mycobacterium tuberculosis -infected IL-27R-alpha-deficient mice is

supported by IL-17A and is associated with the IL-17A-induced expansion of multifunctional T

cells. *Mucosal Immunol* 2018;11:1168–1180.

38. Rolandelli A, Pino REHD, Pellegrini JM, Tateosian NL, Amiano NO, Barrera S de la, Casco N,

Gutiérrez M, Palmero DJ, García VE. The IL-17A rs2275913 single nucleotide polymorphism is

associated with protection to tuberculosis but related to higher disease severity in Argentina. *Sci*

Rep 2017;7:40666.

39. Gideon HP, Phuah J, Myers AJ, Bryson BD, Rodgers MA, Coleman MT, Maiello P, Rutledge T,

Marino S, Fortune SM, Kirschner DE, Lin PL, Flynn JL. Variability in Tuberculosis Granuloma T Cell

Responses Exists, but a Balance of Pro- and Anti-inflammatory Cytokines Is Associated with

Sterilization. *PLOS Pathog* 2015;11:e1004603.

40. Coulter F, Parrish A, Manning D, Kampmann B, Mendy J, Garand M, Lewinsohn DM, Riley EM,

Sutherland JS. IL-17 Production from T Helper 17, Mucosal-Associated Invariant T, and γδ Cells in

Tuberculosis Infection and Disease. *Front Immunol* 2017;8:1252.

41. Lockhart E, Green AM, Flynn JL. IL-17 production is dominated by gammadelta T cells rather than

CD4 T cells during Mycobacterium tuberculosis infection. *J Immunol Baltim Md 1950*

2006;177:4662–4669.

42. Hu D, Notarbartolo S, Croonenborghs T, Patel B, Cialic R, Yang T-H, Aschenbrenner D, Andersson

KM, Gattorno M, Pham M, Kivisakk P, Pierre IV, Lee Y, Kiani K, Bokarewa M, Tjon E, Pochet N,

Sallusto F, Kuchroo VK, Weiner HL. Transcriptional signature of human pro-inflammatory T H 17

cells identifies reduced IL10 gene expression in multiple sclerosis. *Nat Commun* 2017;8:1600.

43. Chen X, Zhang M, Liao M, Graner MW, Wu C, Yang Q, Liu H, Zhou B. Reduced Th17 Response in

Patients with Tuberculosis Correlates with IL-6R Expression on CD4+ T Cells. *Am J Respir Crit Care*

Med 2010;181:734–742.

44. Krausgruber T, Blazek K, Smallie T, Alzabin S, Lockstone H, Sahgal N, Hussell T, Feldmann M,

Udalova IA. IRF5 promotes inflammatory macrophage polarization and TH1-TH17 responses. *Nat*

Immunol 2011;12:231–238.

45. Lastrucci C, Bénard A, Balboa L, Pingris K, Souriant S, Poincloux R, Saati TA, Rasolofo V, González-

Montaner P, Inwentarz S, Moraña EJ, Kondova I, Verreck FA, Sasiain M del C, Neyrolles O,

Maridonneau-Parini I, Lugo-Villarino G, Cougoule C. Tuberculosis is associated with expansion of

a motile, permissive and immunomodulatory CD16+ monocyte population via the IL-10/STAT3

axis. *Cell Res* 2015;25:1333.

46. Rossol M, Kraus S, Pierer M, Baerwald C, Wagner U. The CD14brightCD16+ monocyte subset is

expanded in rheumatoid arthritis and promotes expansion of the Th17 cell population. *Arthritis*

Rheum 2012;64:671–677.

47. Clifford V, He Y, Zufferey C, Connell T, Curtis N. Interferon gamma release assays for monitoring

the response to treatment for tuberculosis: A systematic review. *Tuberc Edinb Scotl* 2015;95:639–

650.

48. Parasa VR, Muvva JR, Rose JF, Braian C, Brighenti S, Lerm M. Inhibition of Tissue Matrix

Metalloproteinases Interferes with Mycobacterium tuberculosis-Induced Granuloma Formation

and Reduces Bacterial Load in a Human Lung Tissue Model. *Front Microbiol* 2017;8:.

49. Chain B, Bowen H, Hammond J, Posch W, Rasaiyaah J, Tsang J, Noursadeghi M. Error,

reproducibility and sensitivity: a pipeline for data processing of Agilent oligonucleotide expression

arrays. *BMC Bioinformatics* 2010;11:344.

50. Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, Szcześniak MW,

Gaffney DJ, Elo LL, Zhang X, Mortazavi A. A survey of best practices for RNA-seq data analysis.

Genome Biol 2016;17:13.

51. Ricciardolo FLM, Sorbello V, Folino A, Gallo F, Massaglia GM, Favatà G, Conticello S, Vallese D,

Gani F, Malerba M, Folkerts G, Rolla G, Profita M, Mauad T, Di Stefano A, Ciprandi G. Identification

of IL-17F/frequent exacerbator endotype in asthma. *J Allergy Clin Immunol* 2017;140:395–406.

52. Akarca AU, Shende VH, Ramsay AD, Diss T, Pane-Foix M, Rizvi H, Calaminici MR, Grogan TM, Linch

D, Marafioti T. BRAF V600E mutation-specific antibody, a sensitive diagnostic marker revealing

minimal residual disease in hairy cell leukaemia. *Br J Haematol* 2013;162:848–851.

Figure legends

Figure 1. TST transcriptome in patients with active and cured TB disease. A) Induration at the site of TST was recorded by routine clinical assessments in both populations (mm) and was not different between the two groups (Mann-Whitney test). B) Venn diagram depicting genes significantly upregulated 2-fold in the TST of patients with active or cured TB relative to control saline injection. (p<0.01 by t-test with no multiple testing correction). C) TST transcriptome common to both active and cured TB summarised as a network diagram. Purple nodes represent Reactome database functional pathways, yellow nodes represent genes and edges reflect relationship between pathways and genes. Pathway node diameters are proportional to the respective pathway –log10 p value enrichment statistic. D) Pairwise dot-plot of 1910 gene integrated TST signature in patients with either active or cured TB. Dotted line reflects line of perfect covariance. p value derived from linear regression model between the two variables. Red and blue dots represent genes >2 fold differentially expressed (p<0.01 by t-test with no multiple testing correction) between active and cured TB.

Figure 2. MMP-1 over-expression in active TB. A) Network diagram of genes and Reactome pathways over-expressed in the TST of patients with active TB compared to cured TB. Purple nodes represent Reactome database functional pathways, yellow nodes represent genes and edges reflect relationship between pathways and genes. Pathway node diameters are proportional to the respective pathway –log10 p value enrichment statistic. B) mRNA expression in TST of patients with active and cured TB of MMP1 gene. C) Representative MMP-1 immunohistochemistry staining in inflammatory infiltrates with TST samples from patients with active and cured TB. D) Differential MMP-1 staining intensity for 18 cellular infiltrates in each group of patients relative to adjacent zones of skin with no cellular infiltration. E) MMP-1 staining intensity in TST zones outside inflammatory infiltrates. * = p<0.01 by Mann-Whitney test.

Figure 3. IL-17 is over expressed in active TB. A) Network diagram depicting upstream cytokine analysis of genes differentially expressed in TST of patients with active and cured TB. Red and blue nodes represent significant genes overexpressed in active and cured TB respectively as identified in fig 1D. Green nodes represent predicted cytokines regulating the gene expression of red and blue nodes. Edges depict relationship between upstream regulators and differentially expressed genes. B) Expression of selected cytokine transcript within the TSTs of active and cured TB patients. C) Expression of IL-17F by immunofluorescence in TST of patients with active TB. Left panel = phase contrast image, right panel = IL-17F positivity (white). D) Quantification of IL-17F staining throughout

TST sections from patients with active and cured TB, determined by pixel intensity as a proportion of

area sampled. Each dot represents IL-17F expression in the cross-section of an entire TST biopsy from

one patient. * p<0.01 by Mann-Whitney test.

Figure 4. TST challenge in active TB is characterised by enrichment of IL-17 responses. Enrichment in

TST relative to saline injection of A) keratinocyte response modules of cytokine bioactivity, B) immune

cell modules and C) CXCL1, CXCL8, S100A9 and CCL2 genes. Neutrophil (Neut.) modules M37.1 and

ImSig validated for sensitivity and specificity in references (23, 24) respectively. * = p<0.01 by Mann-

Whitney test.

Figure 5. Active TB is characterised by elevated Th17 cells. A) IL-17F immunohistochemistry in TST of

patients with active TB. Black arrows point to mononuclear cells that express IL-17F and white arrows

point to polymorphonuclear cells that do not express IL-17F. B) Th17 and Th1 cell module enrichment

in TST relative to saline injection. C) Expression of Th17 and Th1 modules in PPD-stimulated PBMC

from patients with active or latent TB (data originated from dataset GSE27984). * = p<0.01 by Mann-

Whitney test. D) Relationship between expression of Th17 transcriptional module and individual

cytokine gene expression in TST of cytokines implicated in polarising CD4+ T cells to a Th17 phenotype.

Data points represent patients with active TB. Spearman rank correlation coefficients (r^2) and p values.

D) Enrichment in TST relative to saline injection of IL-1β fibroblast response modules indicative of IL-

1 cytokine bioactivity. * = p<0.01 by Mann-Whitney test.

Figure 6. Th17 cells and IL-17 activity are a feature of both TST reactions and sites of human TB

disease. A) Enrichment in TST relative to saline injection of transcriptional modules for IL-17

bioactivity, neutrophils, Th17 cells and T cells in TST of independent cohort of patients with active TB

and a separate cohort of individuals with latent TB infection. B) Expression of T cells, Th17 cells and IL-

17 bioactivity modules from the site of human TB granuloma relative to healthy lung tissue (dataset

GSE20050) and C) in human TB infected lymph nodes (LN) relative to reactive lymph nodes that do not

display evidence of granulomatous inflammation or cancer (dataset E-MTAB-2547). * = p<0.01 by

Mann-Whitney test.

Figure 1

A

TST induration (mm)

Active TB   Cured TB

B

Active TB

1251

Cured TB

D

Cured TB gene expr. (log$_2$)

slope= 1.004
p<0.0001

Active TB gene expr. (log$_2$)

C

Interferon alpha/beta signaling

Interferon Signaling

Interferon gamma signaling   Cytokine Signaling   Signaling by Interleukins

Hemostasis

Generation of second messenger molecules

TCR signaling

Immune System

Innate Immune System

Adaptive Immune System

Interactions between Lymphoid and non-Lymphoid cells

Figure 2

Figure 3

Figure 4

Figure 5

A

B

TST

Th17

Module enrichment (log$_2$)

*

Active TB   Cured TB

Th1

Active TB   Cured TB

C

Blood

Th17

Module expression (log$_2$)

Active TB   Latent TB

Th1

*

Active TB   Latent TB

D

Gene expr. (log$_2$)

IL1B	IL6	IL23A	TGFB	IL12A	IL4

Th17 module expr. (log$_2$)

$r^2 = 0.631$ $r^2 = 0.683$ $r^2 = 0.693$ $r^2 = 0.396$ $r^2 = 0.035$ $r^2 = 0.011$
$p = 0.0002$ $p < 0.0001$ $p < 0.0001$ $p = 0.0090$ $p = 0.4851$ $p = 0.6963$

E

IL-1 response module

Module enrichment (log$_2$)

*

Active TB   Cured TB

Figure 6