Targeting of RBM10 to S1-1 Nuclear Bodies:

Targeting Sequences and its Biological Significance

Running Title: RBM10 and S1-1 nuclear bodies

Ling-Yu Wang[1,2,#], Sheng-Jun Xiao[1,3,#], Hiroyuki Kunimoto[1], Kazuaki Tokunaga[4], Hirotada Kojima[1], Masatsugu Kimura[5], Takahiro Yamamoto[6], Naoki Yamamoto[7,8], Zhao Hong[1], Koji Nishio[9], Hideo Yamane[10], Tokio Tani[4], Koichi Nakajima[1], Hiroyoshi Iguchi[10], and Akira Inoue[1,10*]

Departments of [1]Immunology, [5]Radioisotope Center, and [10]Otolaryngology, Osaka City University Graduate School of Medicine, Osaka 545-8585, Japan; Departments of [2]Human Genetics and [3]Pathology, Guilin Medical University, Guilin 541004, P. R. China; [4]Department of Biological Sciences, Graduate School of Science and Technology, Kumamoto University, 2-39-1 Kurokami, Kumamoto 860-8555, Japan; [6]Graduate School of Life Science, Hokkaido University, Sapporo 060-0808, Japan; [7]Psychiatry Department, Tokyo Metropolitan Tama Medical Center, Tokyo 183-8524, Japan; [8]Department of Biological Sciences, Tokyo Metropolitan University, Tokyo 192-0397, Japan; [9]Department of Anatomy and Neurosciences, Graduate School of Medicine, Nagoya University, Nagoya 466-8550, Japan.

*Corresponding author: Akira Inoue, Department of Otolaryngology, Osaka City University

Graduate School of Medicine, 1-4-3 Asahi-machi, Abeno-ku, Osaka 545-8585, Japan ;

E-mail: ainoue@med.osaka-cu.ac.jp

#These authors contributed equally to this work.

Abstract

RBM10 is an RNA-binding protein that regulates alternative splicing (AS). This protein

localizes to the extra-nucleolar nucleoplasm and S1-1 nuclear bodies (NBs). We investigated

the biological significance of RBM10 localization to S1-1 NBs, which is poorly understood.

Our analyses revealed that RBM10 possesses two S1-1 NB-targeting sequences (NBTSs), one

in the KEKE motif region and another in the C_2H_2 Zn finger (ZnF). These NBTSs acted

synergistically and were sufficient for localization of RBM10 to S1-1 NBs. Furthermore, the

C_2H_2 ZnF not only acted as an NBTS, but was also essential for regulation of AS by RBM10.

RBM10 did not participate in S1-1 NB formation. We confirmed the previous finding that

localization of RBM10 to S1-1 NBs increases as cellular transcriptional activity decreases and

vice versa. These results indicate that RBM10 is a transient component of S1-1 NBs and is

sequestered in these structures via its NBTSs when cellular transcription decreases. We

propose that the NB-targeting activity of the C_2H_2 ZnF is induced when it is not bound to pre-

mRNA or the splicing machinery complex under conditions of reduced transcription.

Introduction

RBM10 (previously called S1-1) is an RNA-binding protein (Inoue et al., 1996) that regulates

alternative splicing (AS) of pre-mRNAs (Wang et al., 2013) via exon skipping, as shown for

the Drosophila protein Discs large homolog 4 (Zheng et al., 2013), the Notch pathway

regulator NUMB (Bechara et al., 2013), and the apoptosis-related protein Fas (Inoue et al.,

2014), or via selection of an internal 5′-splice site in the target exon, as shown for the

apoptosis-related protein Bcl-x (Inoue et al., 2014). Loss of RBM10 function causes TARP

syndrome (talipes equinovarus, atrial septal defect, Robin sequence, and persistent left

superior vena cava), which is characterized by various developmental anomalies, such as cleft

palate and malformation of the heart (Johnston et al., 2010; Gripp et al., 2011), as well as

other diseases, including cancers such as lung adenocarcinoma (Imielinski et al., 2012;

Cancer Genome Atlas Research Network, 2014) (for a review, see Loiselle and Sutherland,

2018).

Nuclear bodies (NBs) are membrane-less compartments of the nucleus. There are

various types of NBs, and their formation, compositions, and roles in relation to the biological

activities of the nucleus have been studied (for a review see Mao et al., 2011 and Staněk and

Fox 2017). While some types of NBs, such as Cajal bodies, nuclear speckles, and

paraspeckles, have been extensively investigated, the biological significance of S1-1 NBs and

their relationship with RBM10 remain largely unknown. RBM10 localizes to S1-1 NBs and S1-1 granules in the nucleus. S1-1 NBs are about 0.5–2.0 μm in size, with usually 10–20 per nucleus. S1-1 granules, which localize to the perichromatin fibrils that comprise transcription and splicing sites, are much smaller and more abundant than S1-1 NBs. Similar to paraspeckles, S1-1 NBs are often closely associated with nuclear speckles. The structure of S1-1 NBs changes according to transcriptional activity; when cellular transcription is globally reduced, RBM10 accumulates in S1-1 NBs and these structures become enlarged (Inoue et al., 2008).

In the present study, we identified the S1-1 NB-targeting sequences (NBTSs) within RBM10 that participate in the targeting and localization of this protein to S1-1 NBs. In addition, we elucidated the biological significance of this localization in relation to the AS regulatory function of RBM10. We discuss that RBM10 is reversibly regulated by sequestration in S1-1 NBs in response to changes in cellular transcriptional activity.

Results

Specific sequences are responsible for targeting of RBM10 to S1-1 NBs

Wild type and deletion mutants of the 852-amino acid (aa) isoform of RBM10 linked in-frame to the N-terminus of EGFP (S1-1/EGFP-N1) were expressed in ARL (adult rat liver epithelial) cells. The subcellular localizations of these proteins were assessed by examining EGFP fluorescence and by staining with anti-S1-1 antiserum, which recognized an epitope within aa 8–63 of RBM10 (Supporting inform. Fig. S1). In contrast with the aa 4–414 deletion, the aa 4–742 deletion completely abolished localization to S1-1 NBs (Fig. 1A and B), suggesting that an NBTS is present within aa 415–742.

The location of the tentative NBTS was further defined as being within aa 411–680 using deletion constructs 1–4 [Fig. 2A and 2C (a)]. The product of construct 3 localized weakly to S1-1 NBs, suggesting that an NBTS is present within aa 543–586 [Fig. 2C (a)] and that another NBTS is present in the unexamined aa 681–742 region. The NBTS within aa 543–586 was further analyzed using constructs 5–8 [Fig. 2B and 2C (b)]. The products of constructs 5 and 6 localized weakly to S1-1 NBs, indicating that the NBTS is present within aa 532–577 [Fig. 2C (b)]. Based on these results, an upstream NBTS (NBTS1) was concluded to be present within aa 543–577 [Fig. 2C (c)]. A KEKE motif, which is characterized by

alternating positive and negative aa residues (Realini et al., 1994), is located at aa 549–568

[Fig. 3 (1)].

RBM10 contains another NBTS

The other NBTS located within aa 681–742 was examined using the constructs pNB1/EGFP-

C3 and pNB2/EGFP-C3, where *NB1* and *NB2* encoded aa 476–645 and aa 476–760 of

RBM10, respectively (Fig. 4A). NB1 and NB2 lacked the epitope recognized by the anti-S1-1

antiserum; thus, the antiserum was used to detect endogenous intact RBM10 localized to S1-1

NBs as a control.

NB1 contained the KEKE region/NBTS1 and a weak nuclear localization sequence

(NLS3) (Xiao et al., 2013). Consequently, NB1 was not only detected in S1-1 NBs, but also

in the nucleoplasm and cytoplasm (Fig. 4A). By contrast, NB2 harbored a strong NLS (NLS1)

in its C-terminus and consequently localized predominantly to the nucleus. Moreover, similar

to endogenous RBM10, NB2 localized to distinct S1-1 NBs owing to an additional NBTS

within aa 681–742.

This region (aa 681–742) harboring the second putative NBTS contains a C_2H_2-type

Zn finger (ZnF) at aa 681–706 [Fig. 3 (2)] and a downstream sequence (designated after-ZnF)

at aa 707–742. These regions and a control region (aa 644–670) were examined by similar

deletion analysis. Only the ZnF deletion mutant localized weakly to S1-1 NBs (Fig. 4B). To

confirm that the ZnF possesses NB-targeting activity, the cysteine residues critical for ZnF

function at aa 683 and 686 were substituted with alanine or the histidine residue at aa 699 was

substituted with serine [Figs 3 (2) and 4C]. The localizations of these two ZnF mutants to S1-

1 NBs were weaker than those of parental NB2 and the after-ZnF deletion mutant (Δ after-

ZnF). However, the ZnF mutants still localized weakly to S1-1 NBs due to the presence of

NBTS1. The second NBTS in the ZnF region was called NBTS2 [Fig. 3 (2)].

Verification of NBTS1 and NBTS2

As shown above, the NB-targeting activity of NB2 was retained upon single deletion of

NBTS1 (KEKE region) or NBTS2 (C_2H_2 ZnF). Importantly, NB2 did not clearly localize to

S1-1 NBs upon simultaneous deletion of both NBTSs [Fig. 5A (1)], demonstrating their

synergistic action in NB localization.

Using ImageJ Plot Profile, NB localization was analyzed in a more quantitative

manner. RBM10 does not localize to the nucleolus (Inoue et al., 2008); thus, analysis was

performed using lines that traversed the extra-nucleolar region [Fig. 5A (2)]. In contrast with

NB2 and Δ after-ZnF, the NB2 mutants lacking the KEKE region (NBTS1) or the C_2H_2 ZnF

region (NBTS2) showed blunt peaks, and the double deletion mutant showed no distinct

peaks. Consistently, the number of S1-1 NBs larger than 0.5 μm decreased upon deletion of

NBTSs [Fig. 5A (3)]. The results indicate that NBTS1 and NBTS2 synergistically enhance

NB localization.

The NBTSs were further verified by employing clones that expressed the KEKE or

C_2H_2 ZnF region linked to the FLAG sequence. In contrast with the control octamer repeat

sequence (OCRE), KEKE and C_2H_2 ZnF localized to S1-1 NBs (Fig. 5B). Based on these

results, we conclude that both the KEKE region (NBTS1) and the C_2H_2 ZnF region (NBTS2)

possess NB-targeting activity.

The C_2H_2 ZnF is essential for the AS function of RBM10

RBM10 regulates AS. We assessed whether the identified NBTSs are associated with this

function of RBM10. Wild-type RBM10 and RBM10 harboring mutations in NBTS1 (KEKE

region) or NBTS2 (C_2H_2 ZnF) were expressed in RBM10-knockout HepG2 cells and their AS

activities in pre-mRNA splicing of *fas* were examined. In this system, wild-type RBM10

causes skipping of exon 6 in *fas* pre-mRNA (Inoue et al., 2014). pre-mRNA splicing of *fas*

was not affected by mutation of the critical KEKKE sequence in the KEKE region/NBTS1 to

AAKAA, but was almost completely abolished by mutation of the CLLC sequence in the C_2H_2

ZnF/NBTS2 to ALLA (Fig. 6, see also Fig. 3), demonstrating that the C_2H_2 ZnF is essential

for regulation of AS by RBM10. Importantly, these results also indicate that the C_2H_2 ZnF has

two functions: regulation of AS and localization of RBM10 to S1-1 NBs.

RBM10 does not participate in the structural organization of S1-1 NBs

The possibility that RBM10 localizes to S1-1 NBs and participates in the formation of these

structures was examined. We previously showed that RNA is stored in nuclear domains called

TIDRs (transcription-inactivation-dependent RNA domains) when HeLa cells are injected

with Cy3-labeled *ftz* model pre-mRNA and successively incubated with actinomycin D (Act

D) (Tokunaga et al., 2006) and that TIDRs and S1-1 NBs are the same structural entities

(Inoue et al., 2008). Using this experimental system, we examined whether S1-1 NBs/TIDRs

formed in the absence of RBM10. For this purpose, RBM10 was knocked down using

specific siRNAs in HeLa cells (Fig. 7a), Cy3-labeled *ftz* pre-mRNA was introduced into cells,

and cells were incubated with Act D. S1-1 NBs/TIDRs formed even in the absence of

RBM10, as visualized by Cy3 fluorescence of *ftz* RNA (Fig. 7b). Thus, we conclude that

RBM10 does not participate in the structural organization of S1-1 NBs.

RBM10 is transiently sequestered in S1-1 NBs when transcription is reduced

We previously showed that localization of RBM10 to S1-1 NBs is closely associated with

reduced transcription (Inoue et al., 2008). When cells were incubated with Act D, the

localization of RBM10 to S1-1 NBs increased in a time-dependent manner. These NBs

became larger and fewer in number [Fig. 8A (1)]; however, the cellular level of RBM10 does

not change upon incubation with Act D (Inoue et al., 2008). Enlarged S1-1 NBs also form

upon heat shock, serum starvation, and confluent cell culture, where cellular transcriptional

activity is commonly decreased, and upon treatment with the RNA polymerase II (pol II)

inhibitor α-amanitin (Inoue et al., 208). We verified these previous results using other pol II

inhibitors, namely, flavopiridol and 5,6-dichloro-1-β-D-ribo-furanosylbenzimidazole (DRB)

[Fig. 8A (2)]. These reagents inhibit CDK9, the kinase subunit of the positive transcription

elongation factor P-TEFb, which is essential for productive transcription (Chao and Price,

2001). These two inhibitors were examined because flavopiridol also inhibits the cell cycle

kinases CDK1 and CDK4, while DRB also inhibits casein kinase II (CK2) (Meggio et al.,

1990, Bensaude, 2011). RBM10 accumulated in S1-1 NBs upon treatment with either

inhibitor [Fig. 8A (2)], suggesting that this accumulation was due to inhibition of P-TEFb. In

support of this, the effects of DRB [Fig. 8A (2)] were not replicated by treatment with the

potent CK2 inhibitor LY294002 (Gharbi et al., 2007) or knockdown of CK2 with specific

siRNAs (data not shown). Thus, the formation of enlarged S1-1 NBs upon treatment with

flavopiridol and DRB is due to their inhibition of P-TEFb, and RBM10 accumulates in S1-1

NBs when transcription is arrested.

The changes in S1-1 NBs were reversible. When cells were washed with fresh

medium, the enlarged S1-1 NBs formed in the presence of α-amanitin or DRB returned to

their initial state within 4 and 1 h, respectively [Fig. 8A (2)]. The difference in the amount of

time required to reverse the effect is likely due to differences in the affinities of these

inhibitors for their binding sites or because α-amanitin also triggers pol II degradation

(Nguyen et al, 1996).

Figure 8B (1) shows the effect of DRB on cellular transcriptional activity.

Transcriptional activity decreased almost to the basal level upon treatment with DRB for 45

min. By contrast, the protein level of RBM10 remained constant upon treatment with DRB

for 3 h [Fig. 8B (2)], as previously shown upon treatment with Act D for 7 h (Inoue et al.,

2008). In addition, following a chase with fresh medium, transcriptional activity returned

close to the initial level after 1 h, and accordingly enlarged S1-1 NBs returned to their original

state [Fig. 8B (1) and A (2)]. The RBM10 protein level remained constant during the chase

period [Fig. 8B (2)]. Thus, the localization of RBM10 to S1-1 NBs changes in a dynamic and

reversible manner according to the level of cellular transcriptional activity, without a change

in the protein level of RBM10. The dynamic and reversible localization of RBM10 to S1-1

NBs, as well as the finding that RBM10 does not participate in the formation of S1-1 NBs

(Fig. 7), indicate that RBM10 is a transient component of S1-1 NBs and is sequestered in

these structures when cellular transcription decreases. These results also suggest that S1-1 NB

sequesters proteins, in addition to other possible functions.

Discussion

The present study demonstrates that RBM10 has two NBTSs, called NBTS1 and NBTS2,

which are located in the KEKE motif and C_2H_2 ZnF regions, respectively. Each NBTS alone

can target and localize to S1-1 NBs; however, both are required for increased localization of

RBM10 to S1-1 NBs. Thus, NBTS1 and NBTS2 act in a synergistic manner. NBTS1 and

NBTS2 are separated by approximately 110 aa residues and have no sequence homology

(Figs 1 and 3). Thus, these NBTSs likely interact with spatially close but distinct structures

within S1-1 NBs, the geometry of which seems to enable their synergistic action.

Under reduced transcription conditions, RBM10 localizes to S1-1 NBs and these

structures become enlarged. When transcriptional activity is restored, enlarged S1-1 NBs

return to their initial state and RBM10 leaves from these structures (Inoue et al, 2008, and

Fig. 8). The present study showed that RBM10 does not participate in the formation of S1-1

NBs and that the cellular level of RBM10 is maintained upon inhibition of transcription and

during the subsequent chase period. These findings indicate that RBM10 is a transient

component of S1-1 NBs and is stored or sequestered in these structures when transcription

decreases.

The identified NBTSs were sufficient for localization of RBM10 to S1-1 NBs. All

known NBs, including S1-1 NBs (Inoue et al., 2008), are membrane-less compartments (for a

review, Mao et al., 2011); therefore, we suggest that targeting and localization of transient NB

proteins to NBs are based on the association of specific sequences of the proteins, here

NBTSs, with acceptor sites in NBs. This may be obvious, but it is worth noting.

The C_2H_2 ZnF/NBTS2 was bi-functional: it was not only important for NB-

targeting/localization, but was also essential for regulation of AS by RBM10. Hernández et al.

(2016) reported that deletions from the C-terminus, including the G-patch and C_2H_2 ZnF,

result in progressive and partial loss of the AS regulatory activity of RBM10, indicating that

the integrity of RBM10 is required for its full function. Our results support this previous study

and further demonstrate that substitution of the two critical cysteine residues of the C_2H_2 ZnF

with alanine was sufficient to completely abolish the AS regulatory activity of RBM10 (Fig.

6).

How the ZnF performs the dual functions of NB-targeting and AS regulation is an

intriguing question. When transcription is suppressed, the level of nucleoplasmic pre-mRNAs

decreases and the C_2H_2 ZnF no longer binds to its substrates or the splicing machinery

complex, which functions together with RBM10. This unoccupied state seems to induce the

NB targeting activity of the C_2H_2 ZnF, thereby increasing the localization of RBM10 to S1-1

NBs. Thus, the C_2H_2 ZnF likely acts as a sensor to enhance sequestration of RBM10 in S1-1

NBs.

The *RBM10* gene resides on the X chromosome, and one of its copies is inactivated

by X-inactivation in females (Coleman, et al., 1996, Thiselton, et al., 2002, Goto and Kimura,

2009). This implies that a double dose of RBM10 is unfavorable to cells and that the cellular

level of active RBM10 must be controlled. In this regard, sequestration of RBM10 in S1-1

NBs may be a mechanism to rapidly and efficiently regulate the active nucleoplasmic level of

RBM10. Specifically, RBM10 is promptly released from S1-1 NBs and localizes to

transcription/splicing sites in the nucleoplasm when transcription increases, while it is stored

in S1-1 NBs when transcription decreases. This sequestration may prevent excessive RBM10

from engaging in unfavorable AS of irrelevant pre-mRNAs, which would produce products

that elicit unfavorable effects on cells.

Dysfunction of RBM10 leads to various diseases. Specific defects in NBTS activity

may also cause RBM10-related diseases. Further studies are required to elucidate the

molecular mechanisms that regulate sequestration of RBM10 in S1-1 NBs, including the

identification and characterization of the acceptor sites for RBM10 in S1-1 NBs and the

resident core proteins that comprise S1-1 NBs.

Experimental procedures

Cells and cell culture. HeLa cells derived from human cervical carcinoma, HepG2 cells

derived from human hepatocellular carcinoma, and HEK293T cells derived from human

embryonic kidney and transformed with the SV40 large T-antigen were cultured in

Dulbecco's Modified Eagle Medium (Nissui). ARL cells (rat liver epithelial cells; Japan

Health Sci. Found., JCRB0248) were cultured in Williams' Medium E (Gibco BRL). The

media were supplemented with 5% fetal calf serum (Gibco BRL), 100 µg/mL streptomycin,

and 100 IU/mL penicillin. Fetal calf serum was used at a concentration of 10% for culture of

HepG2 cells. In inhibition experiments, ARL cells were incubated with 5 µg/mL Act D, 50

µg/mL α-amanitin, 0.5 mM DRB, or 0.2 µM flavopiridol.

Immunostaining. Immunofluorescence analysis was performed as described previously

(Inoue et al., 2008). RBM10 and S1-1 NBs were stained with anti-S1-1 antiserum (Inoue et

al., 2008; 1:250 working dilution) and rhodamine-conjugated goat anti-rabbit IgG (1:200

dilution; ICN, 55662). Stained cells were examined using a fluorescence microscope

(Olympus, type IX70; numerical aperture, 0.4; and objective lens, ×20). Data were captured

using image acquisition software (DP Controller). Images were processed for contrast and/or

brightness.

Immunoblotting. Western blotting was performed as previously described (Inoue et al.,

2008). Total proteins (equivalent to 0.2 μg of cellular DNA) were electrophoresed on 10%

sodium dodecyl sulfate-polyacrylamide gels (e-PARGEL; Atto).

Primers, RBM10 mutant plasmids, and transfection. The PCR primers used to prepare the

various constructs are listed in Table S1. Each primer contained a restriction sequence at one

end. The PCR conditions were as follows: elongation (30 cycles, 5 s/kb) at 72°C, denaturation

(10 s) at 95°C or 98°C, and annealing (15 s) at 3.5–5.0°C below the lower T_m of each primer

set. PCR products were digested with the corresponding restriction enzymes, purified, and

ligated into pEGFP-N1, pEGFP-C3 (Clontech), or p3xFLAG-Myc-CMV-24 (Sigma-Aldrich).

Competent JM109 *Escherichia coli* cells (Takara) were transformed with the constructs, the

resulting colonies were cultured, and recombinant plasmids were isolated using the Pure

Yield Plasmid Miniprep System (Promega). The sizes and DNA sequences of the purified

plasmids were verified by agarose gel electrophoresis and sequencing with a 3130x1 DNA

Sequencer (ABI), respectively. Transfection was performed using Lipofectamine LTX and

PLUS Reagent (Invitrogen) according to the manufacturer's recommendations.

Constructs and mutagenesis of RBM10. The constructs ND1, ND11, and ND15, and that

encoding aa 1–469 were produced by sequential deletion from pS1-1/EGFP-N1 (Inoue et al.,

2008) as described previously (Xiao et al., 2013). To generate pNB1/EGFP-C3 and

pNB2/EGFP-C3, sequences encoding aa 476–645 and aa 476–760 of RBM10 were PCR-

amplified and ligated into the *Sac*I and *Pst*I sites of pEGFP-C3. Deletion or substitution

mutagenesis of these plasmids was performed using a PrimeSTAR Mutagenesis Basal Kit

(Takara) and the primers listed in Table S1. The sequences encoding aa 475–533, the KEKE

region (aa 529–590 or aa 529–571), and the C_2H_2 ZnF region (aa 676–711) were similarly

amplified (Table S1) and ligated into the *Hin*dIII and *Kpn*I sites of the p3xFLAG-Myc-CMV-

24 vector. The two KEKE constructs yielded similar results.

RBM10 knockdown. HeLa cells were transfected with 40 nM specific human RBM10-

siRNA5 or control AllStar siRNA (Qiagen) using Lipofectamine 2000 (Invitrogen) and

incubated for 3 days, in accordance with the manufacturer's protocol. A second specific

siRNA, which was designed to target a different region of RBM10 mRNA, yielded similar

results, indicating that the results obtained were not due to an off-target effect.

Nuclear injection of *ftz* pre-mRNA. Cy3-labeled *ftz* pre-mRNA was microinjected into the

nuclei of HeLa cells at approximately 80% confluency using a Microinjector 5170

(Eppendorf) as previously described (Tokunaga et al., 2006). Cells were incubated for 2 h in

the presence of Act D (5.0 μg/mL). Enlarged S1-1 NBs/TIDRs were observed by fluorescence

microscopy (Tokunaga et al., 2006).

RBM10 knockout. RBM10-knockout HepG2 cells were generated using the CRISPR/Cas9-

mediated genome editing system. Two guide RNAs (cgttcatatcctcgcgagta and

cagccgagaccacgactacc) designed to target exon 3 (14-bp apart) were cloned into pX330-U6-

Chimeric_BB-CBh-hSpCas9 (Addgene) according to the manufacturer's instructions. HepG2

cells were transfected with the plasmids together with pEF-puromycin using Lipofectamine

3000 (Thermo Fisher Scientific Inc.) and incubated with 1 μg/mL puromycin for 2–3 days to

obtain cell colonies. Knockout of RBM10 in the cloned cells was confirmed by Western

blotting using anti-S1-1 antiserum and by DNA sequencing.

Generation of lentiviruses expressing wild-type and mutant RBM10. RBM10 mutants

harboring mutations of KEKKE (to AAKAA) or CLLC (to ALLA), as well as a double

mutant (to AAKAA/ALLA), were prepared by double-strand plasmid mutagenesis using the

primer pairs listed in Table S1 and a Quikchange Site-Directed Mutagenesis Kit (Agilent

Tech.). After verification by DNA sequencing, each RBM10 cDNA was ligated into the *Xho*I

and *Not*I sites of the lentiviral vector pCSII-EF-MCS-IRES-Venus (a gift from H. Miyoshi,

RIKEN, Tsukuba, Japan). Lentivirus expressing wild-type or mutant RBM10 was produced

by co-transfection of 293T cells with pCSII-EF-3xFLAG-wild type or -mutated RBM10,

pCMV-VSV-G-RSV-Rev, and pCAG-HIVgp (Miyoshi et al., 1997).

Assessment of transcriptional activity. ARL cells at ~80% confluency in 3-cm plates were

incubated with or without DRB (0.5 mM). [^{3}H]-uridine was added to the cultures 30 min

before reaction termination. Cells were lysed in 0.2 mL of 0.2 M NaCl, 0.1% SDS, and Tris-

HCl, pH 8.0. Plates were rinsed with 0.3 mL of 0.2 M NaCl and Tris-HCl, pH 8.0. The

combined cell lysates were mixed with three volumes of ethanol and incubated at −30°C for

30 min. Precipitates were collected by centrifugation and digested in 0.2 mL of 2 mM MgCl$_2$,

10 mM Tris-HCl, pH 8.0, first with DNase I (10 μg) and RNase A (5 μg), and then with

proteinase K (10 μg), each for 12 h. Aliquots of the digests were mixed with scintillation

cocktail to measure radioactivity.

Acknowledgments

S.X. and L.W. are recipients of Human Resource Development Scholarships from the Japan

International Cooperation Center (JICE), Japan. This work was supported by grants-in-aid for

scientific research from the Japan Society for the Promotion of Science (JSPS) (KAKENHI

grant numbers 21550162, J152640005, and 15K05569 to A.I., and 21590317 to K.N.).

References

Aravind, L. & Koonin, E. V. (1999). G-patch: a new conserved domain in eukaryotic RNA-processing proteins and type D retroviral polyproteins. *Trends Biochem. Sci. 24*, 342-344. https://doi.org/10.1016/S0968-0004(99)01437-1

Bechara, E. G., Sebestyén, E., Bernardis, I., Eyras, E. & Valcárcel, J. (2013). RBM5, 6, and 10 differentially regulate NUMB alternative splicing to control cancer cell proliferation. *Mol. Cell 52*, 720-733. doi: 10.1016/j.molcel.2013.11.010

Bensaude, O. (2011). Inhibiting eukaryotic transcription. *Transcription 2*, 103-108. DOI: 10.4161/trns.2.3.16172

Cancer Genome Atlas Research Network. (2014). Comprehensive molecular profiling of lung adenocarcinoma. *Nature 511*, 543-50. DOI: 10.1038/nature13385.

Chao, S. H. & Price, D. H. (2001). Flavopiridol inactivates P-TEFb and blocks most RNA polymerase II transcription in vivo. *J. Biol. Chem. 276*, 31793-31799. DOI 10.1074/jbc.M102306200

Coleman, M. P., Ambrose, H. J., Carrel, L., Németh, A. H., Willard, H. F. & Davies, K. E. (1996). A novel gene, DXS8237E, lies within 20 kb upstream of UBE1 in Xp11.23 and has a different X inactivation status. *Genomics 31*, 135-138. https://doi.org/10.1006/geno.1996.0022

Gharbi, S. I., Zvelebil, M. J., Shuttleworth, S. J., Hancox, T., Saghir, N., Timms, J. F. &

Waterfield, M. D. (2007). Exploring the specificity of the PI3K family inhibitor LY294002.

Biochem. J. 404, 15-21. DOI: 10.1042/BJ20061489

Goto, Y. & Kimura, H. (2009). Inactive X chromosome-specific histone H3 modifications

and CpG hypomethylation flank a chromatin boundary between an X-inactivated and an

escape gene. *Nucl. Acids Res. 37*, 7416-7428. DOI:10.1093/nar/gkp860

Gripp, K. W., Hopkins, E., Johnston, J. J., Krause, C., Dobyns, W. B. & Biesecker, L. G.

(2011). Long-term survival in TARP syndrome and confirmation of RBM10 as the disease-

causing gene. *Am. J. Med. Genet. A. 155A*, 2516-2520. DOI: 10.1002/ajmg.a.34190

Hernández, J., Bechara, E., Schlesinger, D., Delgado, J., Serrano, L. & Valcárcel, J. (2016)

Tumor suppressor properties of the splicing regulatory factor RBM10. *RNA Biol. 13*, 466-72.

doi: 10.1080/15476286.2016.1144004

Imielinski, M., Berger, A. H., Hammerman, P. S., Hernandez, B., Pugh, T. J., Hodis, E., Cho,

J., Suh, J., Capelletti, M., Sivachenko, A., et al. (2012). Mapping the hallmarks of lung

adenocarcinoma with massively parallel sequencing. *Cell 150*, 1107-1120. DOI:

10.1016/j.cell.2012.08.029

Inoue, A., Takahashi, K. P., Kimura, M., Watanabe, T. & Morisawa, S. (1996). Molecular

cloning of an RNA binding protein. S1-1. *Nucl. Acids Res. 24*, 2990-2997.

https://doi.org/10.1093/nar/24.15.2990

Inoue, A., Tsugawa, K., Tokunaga, K., Takahashi, K. P., Uni, S., Kimura, M., Nishio, K.,

Yamamoto, N., Honda, K., Watanabe, T., Yamane, H. & Tani, T. (2008). S1-1 nuclear

domains: characterization and dynamics as a function of transcriptional activity. *Biol. Cell*

100, 523-535. DOI: 10.1042/BC20070142.

Inoue, A., Yamamoto, N., Kimura, M., Nishio, K., Yamane, H. & Nakajima K. (2014).

RBM10 regulates alternative splicing. *FEBS Lett. 588*, 942-947.

https://doi.org/10.1016/j.febslet.2014.01.052

Johnston, J. J., Teer, J. K., Cherukuri, P. F., Hansen, N. F., Loftus, S. K., NIH Intramural

Sequencing Center (NISC), Chong, K., Mullikin, J. C. & Biesecker, L. G. (2010). Massively

parallel sequencing of exons on the X chromosome identifies RBM10 as the gene that causes

a syndromic form of cleft palate. *Am. J. Hum. Genet. 14*, 743-748.

doi: 10.1016/j.ajhg.2010.04.007

Loiselle, J. J. & Sutherland, L. C. (2018) RBM10: Harmful or helpful-many factors to

consider. *J. Cell. Biochem. 119*, 3809-3818. doi: 10.1002/jcb.26644

Mao, Y. S., Zhang, B. & Spector, D. L. (2011) Biogenesis and function of nuclear bodies.

Trends in Genetics 27, 295-306. doi:10.1016/j.tig.2011.05.006

Martin, B. T., Serrano, P., Geralt, M. & Wüthrich, K. (2016). Nuclear magnetic resonance

structure of a novel globular domain in RBM10 containing OCRE, the Octamer repeat

sequence motif. *Structure 24*, 158-164. https://doi.org/10.1016/j.str.2015.10.029

Meggio, F., Shugar, D. & Pinna, L. A. (1990). Ribofuranosyl-benzimidazole derivatives as

inhibitors of casein kinase-2 and casein kinase-1. *Eur. J. Biochem. 187*, 89-94.

Miyoshi, H., Takahashi, M., Gage, F. H. & Verma, I. M. (1997). Stable and efficient gene

transfer into the retina using an HIV-based lentiviral vector. *Proc. Natl. Acad. Sci. USA 94*,

10319-10323.

Nguyen, V. T., Giannoni, F., Dubois, M. F., Seo, S. J., Vigneron, M., Kédinger, C., et al.

(1996). In vivo degradation of RNA polymerase II largest subunit triggered by α-amanitin.

Nucl. Acids Res. 24, 2924-2929.

Realini, C., Rogers, S. W. & Rechsteiner, M. (1994). KEKE motifs: Proposed roles in

protein- protein association and presentation of peptides by MHC Class I receptors. *FEBS*

Letters 348, 109-113. https://doi.org/10.1016/0014-5793(94)00569-9

Sloan, K. E. & Bohnsack, M. T. (2018) Unravelling the Mechanisms of RNA Helicase

Regulation. *Trends in Biochemical Sciences 43*, 237-250. doi.org/10.1016/j.tibs.2018.02.001.

Staněk, D. & Fox, A. H. (2017). Nuclear bodies: news insights into structure and function.

Curr Opin Cell Biol. 46, 94-101. doi: 10.1016/j.ceb.2017.05.001.

Thiselton, D. L., McDowall, J., Brandau, O., Ramser, J., d'Esposito, F., Bhattacharya, S. S.,

Ross, M. T., Hardcastle, A. J. & Meindl, A. (2002). An integrated, functionally annotated

gene map of the DXS8026–ELK1 interval on human Xp11. 3–Xp11. 23: Potential hotspot for

neurogenetic disorders. *Genomics 79*, 560-572. https://doi.org/10.1006/geno.2002.6733

Tokunaga, K., Shibuya, T., Ishihama, Y., Tadakuma, H., Ide, M., Yoshida, M., Funatsu, T.,

Ohshima, Y. & Tani, T. (2006). Nucleocytoplasmic transport of fluorescent mRNA in living

mammalian cells: nuclear mRNA export is coupled to ongoing gene transcription. *Genes*

Cells 11, 305-317. https://doi.org/10.1111/j.1365-2443.2006.00936.x

Wang, Y., Gogol-Döring, A., Hu, H, Fröhler, S., Ma, Y., Jens, M., Maaskola, J., Murakawa,

Y., Quedenau, C., Landthaler, M., et al. (2013). Integrative analysis revealed the molecular

mechanism underlying RBM10-mediated splicing regulation. *EMBO Mol. Med. 5*, 1431-

1442. doi: 10.1002/emmm.201302663

Xiao, S., Wang, L., Kimura, M. Kojima, H., Kunimoto, H., Nishiumi, F., Yamamoto, N.,

Nishio, K., Fujimoto, S., Kato, T., et al. (2013). S1-1/RBM10: Multiplicity and cooperativity

of nuclear localization domains. *Biol. Cell 105*, 162-174.

https://doi.org/10.1111/boc.201200068

Zheng, S., Damoiseaux, R., Chen, L. & Black, D. L. (2013) A broadly applicable high-

throughput screening strategy identifies new regulators of Dlg4 (Psd-95) alternative splicing.

Genome Res. 23, 998–1007. doi: 10.1101/gr.147546.112

Figure legends

Fig. 1 RBM10 is targeted to NBs.

A. Structure of the 852-aa isoform of RBM10. RRM I and II: RNA recognition motif I and II, respectively; E4: a 78-aa region encoded by exon 4 and expressed in the 930-aa isoform; C_4 ZnF and C_2H_2 ZnF: Cys$_4$-type and Cys$_2$His$_2$-type ZnFs, respectively; NLS1: the principal nuclear localization sequence (Xiao et al., 2013); G-patch: a motif sometimes found in eukaryotic RNA-processing proteins (Aravind and Koonin, 1999; Sloan and Bohnsack, 2018). The OCRE, KEKE motif, and amphipathic α-helix (AHH) are described in Fig. 3. **B.** Presence of an NBTS within aa 415–742. Plasmids ND1, ND11, and ND15 were derived from pS1-1(RBM10)/EGFP-N1 by deleting the regions corresponding to aa 2–4, 4–414, and 4–742, respectively, and expressed in ARL cells. Green, EGFP; red, RBM10 stained with anti-S1-1 antiserum. Some S1-1 NBs are indicated by arrows. Scale bar: 10 μm.

Fig. 2 Identification of NBTS1 within aa 543–577.

A. and **B.** The indicated regions of pS1-1(RBM10)/EGFP-N1 were deleted to yield constructs 1–4 [**C** (a)] and constructs 5–8 [**C** (b)], and expressed in ARL cells. Green, EGFP; red, RBM10 stained with anti-S1-1 antiserum. Some S1-1 NBs are indicated by arrows. Scale bar: 10 μm. **C.** Deletion diagrams. Lines in (a) and (b) indicate the deleted regions in A and B,

with the corresponding residue numbers provided at each end. Green lines: constructs whose

products localized weakly to S1-1 NBs. (c) Summary of A and B. NBTS1 was located within

aa 543–577 (red line). The box shows the location of the KEKE motif.

Fig. 3 Primary sequences of the KEKE, C_2H_2 ZnF, and OCRE regions.

(1) KEKE region. Alternating positive and negative aa residues in the KEKE region are

indicated by boldface. "Amphi α-helix" is a region predicted to form a perfect AAH with five

helical turns (Inoue et al., 1996). (2) C_2H_2 ZnF. The cysteine and histidine residues critical for

ZnF function are highlighted by boldface. (3) OCRE (octamer repeat, also called NLS3).

Brackets 1–5 represent the five repeating octamers (8 aa) originally assigned based on

hydrophilicity/hydrophobicity profiles (Inoue et al., 1996). The aa 486–540 region (aa 564–

618 of the 930-aa isoform) forms a globular domain with an anti-parallel β-sheet fold (Martin

et al., 2016).

Fig. 4 Identification of NBTS2.

A. NB1 and NB2. The RBM10 sequences encoding aa 476–645 and 476–760 were ligated

into pEGFP-C3 and expressed in ARL cells. Green bars in the upper panel represent NBTS1

at aa 543–577 (the KEKE region). The blue bar represents the region expected to contain

another NBTS within aa 681–742. Purple bars indicate NLS3 (aa 482–540) and NLS1 (principal NLS, aa 743–759). Lower panels show expression of pNB1/EGFP-C3 and pNB2/EGFP-C3. Green, EGFP; red, endogenous RBM10 stained with anti-S1-1 antiserum. Representative S1-1 NBs are indicated by arrows. **B.** Determination of NBTS2. The region corresponding to aa 644–670 (control) (1), aa 685–703 (C_2H_2ZnF) (2), or aa 711–742 (after-ZnF) (3) was deleted from pNB2/EGFP-C3 and these constructs were expressed in ARL cells. Green, EGFP; red, RBM10 stained with anti-S1-1 antiserum. The lower diagram indicates the location of NBTS1 (green bar), the deleted regions (1–3), and the location of the C_2H_2 ZnF (box, aa 681–706). **C.** Confirmation of NBTS2. Two cysteine residues and one histidine residue in the C_2H_2 ZnF of NB2/EGFP-C3 were substituted with alanine and serine, respectively. Green, EGFP. Scale bar: 10 μm.

Fig. 5 Verification of NBTSs.

A. NB localization of NB2 and its mutants. (1) The KEKE region (aa 524–598), the C_2H_2 ZnF region (aa 685–703), both the KEKE and C_2H_2 ZnF regions, or the after-ZnF region (aa 711–742) were deleted from pNB2/EGFP-C3 and these constructs were expressed in ARL cells. Green: EGFP. Some S1-1 NBs and nucleoli are indicated by arrows and arrowheads, respectively. (2) Analysis by ImageJ Plot Profile. The nuclear distributions of the mutants

were analyzed along the indicated lines, which traversed the extra-nucleolar nucleoplasm and

S1-1 NBs. (3) Number of S1-1 NBs. NBs larger than 0.5 μm were counted by two individuals

independently using enlarged images of cells (Fig. S2), and the numbers were averaged. **B.**

Verification of the NB-targeting activity of the KEKE region and C_2H_2 ZnF. Sequences

encoding the KEKE region (aa 529–590), the C_2H_2 ZnF (aa 676–711), and the control OCRE

region (aa 475–533) were PCR-amplified, ligated to the FLAG-tag in p3xFLAG-Myc-CMV-

24, and expressed in ARL cells. Cells were stained with an antibody against FLAG (green),

and nuclei were counterstained with 4´,6-diamidino-2-phenylindole (blue). Scale bar: 10 μm.

Fig. 6 NBTS2/C_2H_2 ZnF is essential for regulation of AS by RBM10.

RBM10-knockout HepG2 cells were transduced with lentivirus carrying wild-type RBM10 or

RBM10 in which the KEKKE sequence of NBTS1 was mutated to AAKAA or the CLLC

sequence of NBTS2 was mutated to ALLA. After 3 days, AS of *fas* transcripts was examined

for skipping versus inclusion of exon 6 (skipping/inclusion ratio). **A.** mRNA expression levels

of RBM10 and its mutants. **B.** AS of *fas* pre-mRNA. Skipping/inclusion ratios of exon 6 were

obtained as described by Inoue et al (2014). The results of two separate experiments were

averaged, with deviations indicated by vertical lines at the top of the bars. (–) indicates

RBM10-knockout HepG2 cells. WT, AAKAA, and ALLA indicate RBM10-knockout HepG2

cells transduced with wild-type RBM10 or RBM10 with mutation of AAKAA or ALLA, respectively. Two independent experiments in B yielded similar results, as indicated by the small deviations of the averaged exon 6-skipping/inclusion values. However, RBM10 mRNA expression levels in one experiment were roughly 50% lower than those in A (data not shown), suggesting that the expressed construct proteins in both experiments were at saturation levels in performing AS of *fas* pre-mRNA.

Fig. 7 RBM10 is not essential for the structural organization of S1-1 NBs.

A. HeLa cells were treated with RBM10-specific siRNA (1) or control AllStar siRNA (2). Immunoblotting of RBM10 and GAPDH is shown. The molecular sizes of the 930- and 852-aa isoforms in kDa are presented on the left. **B.** Nuclei of siRNA-treated HeLa cells were microinjected with Cy3-labeled *ftz* pre-mRNA and cells were incubated with 5 µg/mL Act D for 2 h. Fluorescence images show RBM10 labeled with anti-S1-1 antiserum (left panels) and Cy3-labeled *ftz* pre-mRNA (right panels). Arrows indicate representative enlarged S1-1 NBs.

Fig. 8 RBM10 is transiently sequestered in S1-1 NBs when transcription is suppressed.

A. (1) Effects of Act D on S1-1 NBs. ARL cells were incubated with Act D (5.0 µg/mL) for the indicated durations and stained with anti-S1-1 antiserum. Arrows indicate some S1-1

NBs. (2) Formation of enlarged S1-1 NBs upon suppression of transcription with various inhibitors. ARL cells were incubated with Act D, α-amanitin, flavopiridol, DRB, or LY294002 for 3 h. Enlarged S1-1 NBs in cells treated with α-amanitin and DRB reverted back to their initial state upon further incubation (chasing) in fresh media for 4 and 1 h, respectively. Cells were stained with anti-S1-1 antiserum. Bar: 10 μm. **B.** (1) Transcriptional activity in DRB-treated ARL cells under the conditions described in A (2). Western blotting of RBM10 in DRB-treated cells. (a) and (b) indicate the 930- and 852-aa isoforms of RBM10, respectively.

Fig. 1

A

B

	EGFP	anti-RBM10	merged
ND1 Δ2–4			
ND11 Δ4–414			
ND15 Δ4–742			

Fig. 2

A

1
Δ⁴¹¹⁻⁴⁸⁴

2
Δ⁴⁷⁶⁻⁵⁴²

3
Δ⁵²⁴⁻⁵⁹⁸

4
Δ⁵⁸⁷⁻⁶⁸⁰

B

5
Δ⁵³²⁻⁵⁴⁹

6
Δ⁵⁵⁸⁻⁵⁶⁴

7
Δ⁵⁷⁸⁻⁵⁸¹

8
Δ⁵⁸⁷⁻⁶¹¹

C

Deleted regions and the location of NBTS1

NB

(a)
(1) 411—484
(2) 476—542
(3) 524 — 586 / 543 — 587—598
680 (4)

1. **++**
2. **++**
3. **–/+**
4. **++**

(b)
(5) 532—549
(6) 558—564
578—581 (7)
587—611 (8)

5. **–/+**
6. **–/+**
7. **++**
8. **++**

(c)
From (a) and (b) : 543 — 577

aa 400   500   **KEKE**   600   700
549 ———— 568

Fig. 3

KEKE motif   Amphi. α-helix

        529   533   543   549   568   571   577

(1)   KEKE region: YWDGERRTYIPALEQSA**D**GH**KD**TGASS**KE**G**KEKKEK**H**K**T**K**TAQQIAKDMERWARSLNK

                                KEKE region (**NBTS1)**

         676   681   699   706   711

(2) C_2H_2 ZnF: TDWQKLA**C**LL**C**RRQFPSKEALIR**H**QQLSGL**H**KQNLEIHRRA

                    C2H2 ZnF (**NBTS2**)

         486   489   529   540

(3)   OCRE: APESYSQYPVPDVSTYQYDETSGYYYDPQTGLYYDPNSQYYYNAQSQQYLYWDGERRTYIPALEQS

                       1   2   3   4   5

                               OCRE (Octamer Repeat)

Fig. 4

A

NB1 476 — 645   NLS3   NBTS1
NB2 476 — 760   NLS1

400   500   543   577   600   681   742   800

NB1   endogenous RBM10   NB2   endogenous RBM10

B

(1) Δ aa 644–670   endogenous RBM10   (2) Δ aa 685–703   endogenous RBM10   (3) Δ aa 711–742   endogenous RBM10

NB2 476 — 760   NBTS1   (1) (2) (3)

500   600   644   681   706   742   C₂H₂ ZnF

C

NB2   NB2   C683A, C686A   H699S   Δ after ZnF

Fig. 5

A

NB2	ΔKEKE-region	ΔC$_2$H$_2$-ZnF	Δ (KEKE-region + C$_2$H$_2$-ZnF)	Δ after-ZnF

(1)

(2)

(3) 34 ± 8   11 ± 5   12 ± 5   5 ± 2   30 ± 7

B

OCRE (NLS3)   KEKE region   C$_2$H$_2$ ZnF

Fig. 6

A

B

Fig. 7

Fig. 8

A

(1)

0 h	0.5 h	3 h	7 h

(2)

Control   ActD   α-Amanitin   chase 4 h

flavopiridol   DRB   chase 1 h   LY294002

B

(1)

[³H]uridine (dpm ×10⁻³)

+ chase 1 h

Time : 3.0 0.75 1.75 3.0 3.0 (h)

DRB : −   +

(2)

Cont.   DRB (3h)   0   1   chase (h)

— (a)
— (b)

1.00   1.01   1.02   density