Diversity, biogeography, and evidence for endemism of SAR11 bacteria from the Arctic Ocean

Susanne Kraemer[1], Arthi Ramachandran[1], David Colatriano[1], Connie Lovejoy[2], and David A. Walsh[1*]

[1]Department of Biology, Concordia University, 7141 Sherbrooke St. West, Montreal, Quebec H4B 1R6, Canada

[2]Département de biologie, Institut de Biologie Intégrative et des Systèmes (IBIS) and Québec-Océan, Université Laval, Québec, G1K 7P4, Canada

*Corresponding author: Phone: (514) 848-2424 (ext. 3477)

E-mail: david.walsh@concordia.ca

Key words: ITS phylogeny, metagenome assembled genomes, metatranscriptomics, ecotype, evolution

Abstract

The Arctic Ocean is relatively isolated from other oceans and consists of strongly stratified

water masses with distinct histories, nutrient, temperature and salinity characteristics, therefore

providing an optimal environment to investigate local adaptation. The globally distributed SAR11

bacterial group consists of multiple ecotypes that are associated with particular marine

environments, yet relatively little is known about Arctic SAR11 diversity. Here, we examined

SAR11 diversity using ITS analysis, metagenome-assembled genomes (MAGs) and

metatranscriptomic datasets. Arctic SAR11 assemblages were comprised of the S1a, S1b, S2,

and S3 clades, and structured by water mass and depth. The fresher surface layer was dominated

by an ecotype (S3-derived P3.2) previously associated with Arctic and brackish water. In contrast,

deeper waters of Pacific origin were dominated by the P2.3 ecotype of the S2 clade, within which

we identified a novel subdivision (P2.3s1) that was rare outside the Arctic Ocean. Arctic S2-

derived SAR11 MAGs were restricted to high latitudes and included MAGs related to the recently

defined S2b subclade, a finding consistent with bi-polar ecotypes and Arctic endemism. These

results place the stratified Arctic Ocean into the SAR11 global biogeography and have identified

SAR11 lineages for future investigation of adaptive evolution in the Arctic Ocean.

Introduction

The SAR11 (Pelagibacterales) group accounts for roughly 30% of bacteria in the ocean surface and 25% of mesopelagic bacteria [1–3]. High phylogenetic diversity and divergence into ecological lineages (i.e. ecotypes) of SAR11 tends to mirror distinct conditions in the oceanic environment [2, 4–6]. SAR11 are classified into clades and subclades based on 16S rRNA genes and further classified into phylotypes based on rRNA internal transcribed spacers (ITS) (Table 1). Three major SAR11 clades (S1, S2 and S3) are currently recognized, with several subclades defined within S1 (S1a, S1b, S1c). Within this diversity approximately 12 phylotypes have been described. To date, many of the phylotypes are thought to have restricted distributions. For example, there are three phylotypes in subclade S1a: P1a.1 is associated with cold environments [7], while P1a.2 and P1a.3 are associated with temperate and tropical environments, respectively [5, 7]. Subclade S1b is generally associated with tropical environments [5, 7, 8], while subclade S1c is generally associated with deep marine samples [5]. Recently, the clade S2 was divided into subclades S2a and S2b. The S2a subclade is further divided into an oxygen minimum zone subclade (S2a.a) and a tropical subclade (S2a.b) [8]. The S2 subclade phylotypes have been further divided by environment. P2.1, is associated with tropical environments [7], P2.2 is reported from cold, especially Antarctic, waters [7, 9], and the P2.3 phylotype is more ambiguous; associated with cold waters and tropical deep-sea environments [5, 7]. To date, three phylotypes have been distinguished for clade S3; P3.1 is associated with coastal, mesohaline surface [10] and tropical samples [7], P3.2 is found in brackish [10], temperate [7], as well as Arctic samples [11] and the third S3 phylotype, LD12, is found in freshwater [12] (Table 1).

The Arctic Ocean is small, geographically isolated, and more influenced by freshwater and ice compared to other oceans [13–15]. These distinct characteristics would favour the evolution of locally adapted microbial assemblages, however evidence for this is difficult to document. Biogeographic patterns of taxa reported from the Arctic vary, with reports of cosmopolitan, bi-polar and endemic species, suggesting a potential for different levels of specialization to local

conditions [6, 16–18]. Although SAR11 are reported widely and account for 25 to 30% of Arctic

Ocean bacterial assemblages [19, 20], comparatively little information exists on Arctic SAR11

diversity [7]. The extensive knowledge of SAR11 diversity and ecology in other oceans makes it

an attractive clade to investigate ecotypes adapted to specific Arctic Ocean conditions. The

objective of this study was to place SAR11 from the Arctic Ocean within a global context using

ITS phylogenetic analysis and comparative genomics using metagenome-assembled genomes

(MAGs). We then examined the distribution of SAR11 along a latitudinal transect of the stratified

waters of the Canada Basin in the western Arctic Ocean. We targeted three water masses in the

upper 200 m, the surface layer, the deep chlorophyll maximum (DCM), which corresponds to a

halocline formed by Pacific Summer Water, and the Pacific Winter Water (PWW) layer [21], These

three water masses were previously found to have distinct microbial communities [22] and we

hypothesised that 1) different SAR11 ecotypes would be favored within the Arctic water masses,

and 2) the distinct characteristics of the Arctic Ocean would favor the existence of endemic SAR11

ecotypes.

Methods

Sampling and metagenomic data generation

Samples were collected aboard the Canadian Coast Guard Icebreaker CCGS Louis S. St-

Laurent from the Western Arctic Ocean from latitudes 73˚ to 79˚ N in September 2015, during the

Joint Ocean Ice Study cruise in the Canada Basin (Table 2). Sample collection and preservation,

DNA extraction, and metagenomic data generation are described previously [23], details are

provided in the Supplementary Information.

96
ITS phylogenetic analysis

SAR11 ITS sequences were retrieved from twelve assembled metagenomes, clustered

and filtered (Supplementary information). We combined the Arctic ITS sequences with reference

sequences from SAR11 genomes and ITS sequences from two previous biogeographic studies

[2, 5] and assigned Arctic ITS sequences to phylotypes. We determined phylotype distribution

across the samples with PCoA ordination of the Bray-Curtis dissimilarity as described in the

Supplementary Information.

Metagenome-assembled genomes

We conducted metagenomic binning using the Metabat2 pipeline [24] on a 12 sample

coassembly [23, 25]. Further SAR11 bin filtering and cleaning was conducted as described in the

Supplementary Information. Seven SAR11 MAGs with low contamination values (<4%) and

relatively high completeness values estimated using CheckM [26] (36 % to 47%) were selected

for further analysis (Table 3). The distribution of orthologs across SAR11 MAGs and reference

genomes was analyzed with ProteinOrtho [27]. A set of 39 single copy ortholog genes were

concatenated and used for phylogenetic analysis using MEGA-cc with a JTT substitution model

[28], as further described in the Supplementary Information.

Arctic SAR11 transcriptome

RNA samples from surface water, DCM, and PWW at station CB4 (75.24˚ N), were

preserved in RNAlater and stored at -80 C. Detailed sample processing is described in the

Supplementary Information. Metatranscriptomic data was generated by paired-end Illumina

sequencing at the Department of Energy Joint Genome Institute as in [23]. Unassembled

metatranscriptomic data was recruited against the protein-coding gene sequences from SAR11

MAGs and the S3 reference genome IMCC9063 using bbmap with a 98% cutoff [25]. Recruited

reads per kilobase metagenome (RPKB) were calculated as: ((#Number of mapped reads of gene

× Total number of mapped reads) / #Total reads in metatranscriptome) × 1000. To obtain MAG-

level expression we utilized the average number of mapped reads across all genes.

Comparative genome content, average nucleotide identity and signatures of selection of Arctic MAGs

To identify Arctic-specific SAR11 genes, we compared all genes within the Arctic MAGs to those found within 41 SAR11 reference genomes using ProteinOrtho [27]. The protein functions of genes only found in Arctic MAGs were retrieved following the IMG annotation [29] or the SwissProt database [30]. We calculated average nucleotide identity (ANI) between Arctic MAGs and reference genomes representing the different subgroups following the method implemented within IMG [29].

We compared the genome content of MAG SAR11-484 to two closely related S2a reference genomes, HIMB058 and AAA795E07. Orthologous genes were identified across the three genomes using ProteinOrtho [27]. We calculated dN/dS between these shared genes using the Young and Nielson method in yn00 as implemented in paml [31] as described in the Supplemental Information, where values greater than one are indicative of diversifying selection.

Fragment recruitment and SAR11 biogeography.

To determine the prevalence of Arctic MAGs across global marine biomes, we performed reciprocal best hit analysis [23, 32] for 168 metagenomic samples from a range of marine biomes. Metagenomic datasets included 134 TARA ocean datasets [33] which were randomly sub-sampled to a size of approximately 1 GB of reads each to facilitate analysis. These were added to the Arctic metagenomes and 22 Antarctic datasets that included 20 marine and 2 ACE lake samples [2]. We examined the distribution of the seven Arctic MAGs, 48 representative SAR11 reference genomes, and 38 TARA ocean MAGs previously described as SAR11 [34] as described in the Supplemental Information. Best hits from the reciprocal blast were filtered to a minimum length of 100 bp and a minimum identity of 98% and the number of recorded hits per reference genome and metagenomic sample were used to calculate the number of reads recruited per

megabase genome per gigabase metagenome (RPMG). See Supplemental Table 1 for all

metagenomic datasets and reference genomes.

We utilized the RPMG matrix for PCoA ordination of the Bray-Curtis dissimilarity of the

RPMG matrix. The envdist function as implemented in vegan with 999 permutations was used for

Post-Hoc tests of environmental variables. [35].

Results

Environmental setting

The stratified upper waters of the Canada Basin in the Western Arctic Ocean sampled

during the late summer–autumn of 2015 were typical for the region, with warmer and fresher

summer waters above colder slightly saltier winter Pacific-origin water (Table 2). Salinity at the

surface ranged from 25.6-27.3 and nitrate concentrations were below the detection limit (0.5 μM).

Below the surface water, relative chlorophyll *a* fluorescence at the DCM was calculated to range

from 0.23 to 0.33 mg m^{-3}. The DCM depth varied from 25-79 m, with the shallowest value at CB11,

the most northerly station along the western edge of the Canada Basin (Figure 1A-B). Nitrate

concentrations increased with depth to up to 16 mg m^{-3} in the PWW, where chlorophyll *a* was low

(0.05 μmol m^{-3}).

SAR11 ITS sequence diversity

No SAR11 16S rRNA sequences were present in the metagenomes. Within the

metagenomic assemblies, we identified 140 high quality SAR11 ITS sequences belonging to 111

unique ITS sequence variants and combined these with previously published sequences for

phylogenetic analysis. Sixty Arctic ITS sequences did not cover the full ITS region and were

assigned to phylotypes using their best BLAST hit against full-length SAR11 ITS sequences. In

total, 6 distinct phylotypes were evident from ITS sequences (Figure 1C), from all major clades

S1, S2, and S3 (Figure 1C, Figure 2A-D).

Clade S1. Within S1, we identified the S1a and S1b subclades. The Arctic ITS S1a

sequences mostly clustered apart from previously published P1a sequences (Figure 2A). These

P1a-related sequences were common in samples from the PWW and DCM layer, but nearly

absent from the surface layer (Figure 2A, Figure 3A). Within subclade S1b (Figure 2B), Arctic

Ocean ITS sequences were found in two clusters. One cluster corresponded to a previously

described P1b.a group [7], while the other grouped within a previously designated non-

monophyletic tropical P1b cluster [2], hereafter termed P1b.b (Figure 2B), but we were unable to

connect these groupings to the three previously described P1b.1-3 phylotypes [5]. P1b.a

contained three full-length Arctic sequences and was found nearly exclusively in the PWW (Figure

2B, Figure 3A).

Clade S2. The majority of ITS sequences were assigned to phylotypes within subclade

S2 (Figure 2C). In our phylogenetic analysis, we recovered a monophyletic group within the P2.3

phylotype, with Arctic sequences clustering with two deep water Red Sea sequences, forming a

novel cluster P2.3s1 (Figure 2C). Apart from a single DCM sample, which contained a high

number of P1a sequences (coverage of ~670, compared to an average coverage of ~40, Fig. 3A),

sequence types within the P2.3s1 phylotype were the most frequently detected in Arctic waters,

specifically in DCM and PWW samples (Figure 2C, Figure 3A). The published ITS sequences

previously assigned to P2.1 and P2.2 were interspersed with one another, and we therefore refer

to this subgroup as P2.1–2.2. P2.1–2.2 was distributed relatively evenly across sampling depths

and locations (Figure 2C, Figure 3A).

Clade S3. We recovered two ITS sequence type variants belonging to the brackish Arctic

phylotype P3.2 (Figure 2D). P3.2 was common in the less saline surface water samples and

absent below the DCM (Figure 2D**,** Figure 3A).

Phylotype abundance and biogeography

PCoA ordination of samples based on the relative abundance of phylotypes showed that SAR11 assemblages were structured along the first axis in relation to the water layer sampled (Figure 3B). In contrast, the second axis had little explanatory power and structure was driven by the highly abundant P1a sequence type in the DCM sample of station CB8. PWW samples contained a high contribution of diverse phylotypes including P1a, P1b.a, P1b, P2.1–P2.2 and P2.3 and P2.3s1, while P3.2 was most frequent in surface layer samples. P2.1–P2.2, was not associated with contributions to a specific depth class of samples likely because multiple poorly resolved ecotypes are contained within this group (Figure 3A, B). Secondary fitting of environmental variables onto the ordination did not yield significant results.

Characteristics of Arctic Ocean SAR11 MAGs

We binned SAR11 scaffolds from the Arctic Ocean metagenome co-assembly based on tetranucleotide frequency and coverage across samples. After automated binning and manual curation, we selected seven high quality SAR11 MAGs for further analysis (Table 3). Quality was based on high N50 values (19-100Kb), a low number of scaffolds (5-26), relatively high completeness (35-47%), and low contamination (0-4%) values.

SAR11 MAGs did not contain rRNA genes or ITS sequences, and thus their placement into 16S rRNA or ITS-based phylogenies was not possible. Instead we investigated their phylogeny using a concatenated gene tree of Arctic MAGs, reference genomes, and SAGs from the Red Sea and the Eastern Tropical North Pacific oxygen minimum zone (ETNP OMZ) [8] (Figure 4A). The Arctic MAG SAR11–312 belonged to subclade S1a and was prevalent at and below the DCM layer, but not at the surface (Figure 4B). This MAG was more closely related to phylotype P1a.1 than to P1a.3; a placement supported by its maximum ANI value in comparison to a P1a.1 reference genome (80%). Its phylotype could not be determined, as the ITS regions for most of the reference genomes were missing.

The remaining six MAGs were members of the S2 clade of SAR11. Four of the MAGs

(SAR11–112, SAR11–272, SAR11–410, and SAR11-484) were members of the S2a subclade,

while two (SAR11–144 and SAR11–196) were most closely related to two SAGs from the S2b

subclade that originated from the ETNP OMZ (Figure 4A). The S2a MAGs exhibited differential

distributions in the stratified waters of the Arctic Ocean. Three closely related MAGs (SAR11–

112, SAR11–272, SAR11–410) were detected at all depths, while a more distantly related S2a

MAG (SAR11–484) was only detected in the surface and DCM samples (Figure 4B). The

phylogenetic placement of these MAGs was supported by their maximum ANI values in

comparison to the S2a reference genome HIMB058 (>75%). In contrast, MAGs SAR11–144 and

SAR11–196 were detected in DCM and PWW samples (Figure 4B), and are the most likely MAGs

to represent the novel P2.3s1 phylotype defined in this study. Due to the absence of any S2b

reference genomes with high completeness, we were unable to calculate ANI of SAR11–144 and

SAR11–196 with S2b.

239
Arctic SAR11 transcriptomes

To assess transcriptional activity of the Arctic SAR11 assemblages, we recruited

metatranscriptome data from surface, DCM, and PWW depths at one station (CB4, 75.24° N)

against the MAGs. RPKB values for all genes within the MAGs roughly followed their coverage in

the metagenomes (Figure 4C). S1a MAG SAR11–312 was transcriptionally active in DCM and

PWW waters. Transcripts from S2a MAGs SAR11–112, SAR11–272 and SAR11–410 were found

at all depths, however they were mostly associated with the DCM layer, and, in the case of MAG

SAR11–272, the PWW layer. In contrast, the more distantly related S2a MAG SAR11–484 was

found to be active in both DCM and surface waters. S2b-related MAGs SAR11–144 and SAR11–

196 were only active in DCM and PWW samples. As we were unable to assemble a S3 genome

from the Arctic metagenomes, we utilized the S3-P3.2 genome IMCC9063 to test its

transcriptional activity in the Arctic. IMCC9063 transcript abundance was high in the surface layer.

Transcripts from most SAR11 genes were not detected or were very rare in the

metatranscriptomes. We detected only 13 genes with RPKB values higher than one, belonging to

MAGs SAR11–144, SAR11–196 and SAR11–312. All but one of these highly expressed genes

were found in the DCM, reflecting the relatively high activities of these MAGs there. Of the 13

genes, six are of unknown function. The remaining seven highly expressed genes are involved in

nucleotide transport and metabolism (two genes), amino acid transport and metabolism (two

genes), co-enzyme transport and metabolism (one gene), cell membrane biogenesis (one gene)

and cell cycle control (one gene) (Supplemental Table 2).

Comparison of gene content

To investigate the potential of local adaptation through variation in gene content we

identified SAR11 genes specific to the Arctic Ocean by comparing the distribution of orthologs

between Arctic MAGs and 60 reference genomes representing the known phylogenetic diversity

of SAR11. Of the 2 648 orthologs we identified across SAR11 genomes, 233 were only found in

Arctic MAGs and four of these were present in more than one MAG. The majority of these

orthologs were poorly characterized proteins (58%, 136 orthologs: COG categories S and R). A

further 24% (55 orthologs) were involved in metabolism, 3% (7 orthologs) in cellular processes

and signaling and 7% (16 orthologs) in information storage and processing (Supp. Table 3).

To determine patterns of adaptation, we investigated the relationship between Arctic S2a

MAG SAR11–484 and its two most closely related reference genomes: Pacific HIMB058 and Red

Sea SAG AAA795E07 (Figure 4D). SAR11–484 shared more orthologs with HIMB058 than with

the Red Sea SAG (though this could reflect the higher completeness of the HIMB058 genome).

Of the 1 170 orthologs identified in these three genomes, we identified 221 complete genes which

were common to all three genomes (Figure 4D) and tested these genes for diversifying selection.

None of the genes showed dN/dS values > 1 in any comparison between the three genomes.

Thus, none of the orthologs investigated here showed signatures that may indicate diversifying

selection and adaptation to local conditions.

Global biogeography of SAR11 genomes

We investigated the biogeographic distribution of Arctic SAR11 populations using Arctic

MAGs and available SAR11 genomes. In addition to being present in Arctic DCM and PWW

samples, MAG SAR11–312, belonging to subclade S1a, was also detected in the mesopelagic

zone of colder oceans (Figure 5A). Arctic S2a MAGs were present in metagenomic datasets that

were mostly from polar regions, with the exception of SAR11–112 and SAR11–272, which were

present in Mediterranean samples. The Arctic MAGs most closely related to the S2b subclade

were even more narrowly distributed. Outside of Arctic samples, SAR11–144 was only detected

in South Atlantic DCM layer and North Atlantic mesopelagic samples. In contrast, MAG SAR11–

196 was only found in Arctic DCM and PWW samples (Figure 5A).

Other than the Arctic MAGs, most SAR11 genomes recruited poorly across both Arctic

and Antarctic metagenomic samples, suggesting they are not significant members of Arctic

SAR11 assemblages. However, there were some exceptions: S1a genome HTCC1062, which

was the most widely distributed genome, was prevalent in the Antarctic and to a lesser degree

the Arctic. The generalist S2a genome HIMB058, which is prevalent in most marine biomes, was

present in Arctic and Antarctic samples (Figure 5A). Lastly, S3 genome IMCC9063, characterized

as a fresher, arctic phylotype [10], was found across surface samples of several marine biomes,

but was most prevalent in Arctic surface waters, followed by Antarctic samples.

To further explore SAR11 biogeography in relation to environmental conditions of ocean

biomes, we performed PCoA ordination of the Bray-Curtis distance matrix of the RPMG-values

(Figure 5B). Overall, metagenomic samples clustered according to their biome and sampling

depth. Along the first axis, samples separated according to a temperature and salinity gradient,

with colder, fresher samples such as the Arctic and Antarctic environments investigated here

distinguished from warmer, saltier environments. Along the second axis, metagenomic samples

were separated according to their depth. Deep samples were characterized by high nitrate

concentrations while more shallow samples had higher oxygen and Chl *a* concentrations. Notably,

Arctic and Antarctic surface and DCM samples clustered together (Figure 5B, as polar biome

samples in lower left quadrant) while Arctic PWW samples were clearly distinct in their

compositions (Figure 5B, blue circles). S1a MAG SAR11–312 had a vastly different distribution

compared to most other S1a genomes, which were found largely in warmer coastal and trades

biomes (e.g. RS39, HIMB5, HIMB083). Arctic S2a MAGs were associated with polar DCM and

surface samples. MAGs SAR11–144 and 196 were differentially distributed from all other

genomes investigated here, including the most closely related North Pacific A6S6 and B3S13.

These results were robust to expanding the analysis to MAGs assembled from the initial TARA

Oceans project, which did not include Arctic samples (Supp. Figure 1).

315
Discussion

Ribosomal RNA amplicon surveys of the marine environment have documented the

occurrence of SAR11 in Arctic marine systems [36–38], but the diversity and biogeography of the

different evolutionary lineages has not been investigated in detail. Here we found evidence for

the presence of at least six SAR11 groups in the Arctic. The global success of the SAR11 group

across marine environments points at a capacity to adapt to a wide array of environmental

conditions. While Arctic and Antarctic Oceans are superficially similar environments with respect

to solar radiation, the presence of sea ice and algae blooms following ice melt in spring [18], the

Arctic Ocean is surrounded by land and freshwater input from large rivers makes the Arctic Ocean

a more estuary-like marine biome compared to the Antarctic. The rivers also bring in terrestrial-

derived organic matter which represents a potential substrate and selective force on the bacterial

communities [23]. Increasing evidence suggests microbial Arctic endemism in other bacterial

groups [23], but this is the first genome-level study to our knowledge within the SAR11 group.

The biogeography of ITS phylotypes offers a hint that local selective forces are at work

but the relatively coarse nature of the marker masks potential differentiation at the genome level,

such that ITS-based phylotypes may not necessarily discern discrete bacterial populations that

are locally adapted. Recent advances in sequencing techniques and genome assembly

algorithms have resulted in the utilization of MAGs and SAGs to extend the biogeographies of

bacteria beyond the phylogenetic marker level and infer patterns of local adaptation at the

genome level [8, 24, 39]. Utilizing this approach, gene content of individual SAR11 genomes has

been linked to niche partitioning based on nutrients [40], metabolic adaptation to environmental

productivity [41] and oxygen minimum zones [8]. However, this approach may have limitations

since assembling SAR11 genomes from metagenomes is notoriously difficult due to their high

levels of polymorphism [42]. Moreover, recent work suggests that gene content differences alone

may not be sufficient to explain SAR11 biogeographic patterns [43]. Nonetheless, our combined

MAG-based and marker gene-based results converged, providing evidence of novel arctic

ecotypes.

Structure of Arctic Ocean SAR11 assemblages

Arctic phylotypes belonging to S1a, S2 and S3 showed preferences linked to different

water masses [5]. The fresher Arctic surface waters were dominated by a P3.2 phylotype that had

been previously associated with Arctic and brackish waters [10, 11]. In contrast, PWW depths

were dominated by the cold [2] and deep [5] phylotype P2.3. The majority of all recovered ITS

phylotypes from the Arctic Ocean belonged to the P2.3 phylotype, with one highly abundant

phylotype (P2.3s1, Figure 2C) that was previously only represented by a few sequences from

deep in the Red Sea [5]. The Arctic Ocean may constitute the centre of the P2.3s1 range

distribution [44] and its potential for local adaptation as well as its niche requirements needs

further exploration. In contrast to P3.2 and P2.3, the P2.1–P2.2 phylotype showed a broader

distribution across depths and samples. However, our phylogenetic analyses failed to recover the

two distinct previously published phylotypes P2.1 (tropical) and P2.2 (cold) [2].

To further investigate the link between SAR11 diversity and the environment we utilized

metagenomic binning to assemble seven Arctic Ocean SAR11 MAGs that belonged to or were

closely related to subclades S1a and S2a and S2b (Table 1), and investigated their biogeographic

distribution and transcriptional activity in the Arctic Ocean. Three of the four S2a MAGs recovered

here were widely distributed across depth classes and most stringently corresponded to the

distribution of the P2.1–P2.2 phylotype. The two MAGs most closely related to S2b reference

genomes were mostly present in DCM and PWW waters, mirroring the distribution of the P2.3

and P2.3s1 phylotypes. However, in the absence of direct ITS evidence we were unable to test

directly that S2b-related Arctic MAGs correspond to the extremely common P2.3s1 ITS phylotype.

Surprisingly, we failed to assemble a S3 MAG, which based on the abundance of corresponding

ITS sequences, the high fragment recruitment of the S3 reference genome IMCC9063, and the

high transcriptome signal of this genome in surface waters, should have been recoverable.

Moreover, we were unable to detect any long (>10 kb) SAR11 scaffolds mapping to S3. The

absence of any other strong S3 signal, apart from the presence of ITS sequences, is intriguing

and warrants further investigation.

Placing Arctic SAR11 into the global biogeography

To determine the biogeography of Arctic MAGs, in comparison to reference genomes

representing SAR11 diversity, we performed fragment recruitment analysis across 168

metagenomic datasets from different marine biomes, including the Arctic and two Antarctic

habitats: the Southern Ocean and Ace Lake, a saline Antarctic lake. While Arctic MAGs belonging

to clade S1 and subclade S2a were also detected in other metagenomic samples, one MAG,

which was most closely related to subclade S2b, SAR11–196, was found only in samples from

the Arctic Ocean, providing evidence in support of endemism. However, due to the low genome

completeness of this MAG and the lack of closely related reference genomes it was not feasible

to link this apparent endemism to gene content [8, 41] or selection acting on genes responsible

for the apparent local adaptation to the Arctic, as was recently done for a different and wide-

spread SAR11 population [43]. Arctic S2a MAGs were sparsely distributed across global

metagenomes but were consistently found in Arctic, Antarctic and South Atlantic biomes,

indicating a bi-polar distribution of this subclade. However, two of the three MAGs showed

recruitment to the warm and salty Mediterranean Sea in both the surface and the DCM layer.

Other reference genomes within the subclade were isolated from warm and oxygen-poor

environments (e.g. A6S6 and A10S10 [8]), indicating that cross-binning across closely related

genomes during the fragment recruitment is theoretically possible. Lastly, in contrast to the

majority of other S1a reference genomes investigated, the Arctic S1a MAG SAR11–312 was

found across the mesopelagic layer of most Oceans.

More broadly speaking, SAR11 assemblages were structured by their biomes and

environmental features. The majority of SAR11 reference genomes (including most S1a and S1b

genomes), as well as MAGs assembled from the global TARA ocean metagenomes [33], are

associated with warmer, more shallow environments from the Trades, Coastal and Westerlies

Biomes. A second group of reference genomes (mostly S1c) were most common in deeper water

with lower oxygen and higher nitrate concentrations, as previously reported for S1c genotypes

found in nitrate replete oceanic OMZs [8]. Polar biome samples formed a distinct third

environmental group. Within the polar samples, PWW samples clustered away from all other

environments, providing support for the hypothesis that distinct SAR11 assemblages in the

samples are selected by environmental conditions, specifically with respect to the Arctic S2b

MAGs, which were most frequent here. In contrast to PWW samples, other Arctic samples

clustered closely with samples from the Southern Ocean, pointing towards the bi-polar distribution

for, among others, the Arctic S2a MAGs that were the most frequent in these samples.

Different degrees of endemism within bacterial communities in the Arctic Ocean have

been reported. Patterns vary for different groups of bacteria, but in general Arctic bacterial

communities are complex assemblages of bacteria with cosmopolitan, bi-polar, and Arctic-

specific distributions, indicating varying degrees of adaptation to local conditions. [6, 17, 18, 45–

47]. In the present study, we set out to describe and place Arctic Ocean SAR11 into the global

SAR11 biogeography using both marker gene and MAG-based approaches. Assembly of the

highly genome-streamlined SAR11 clade, which shows high rates of recombination, from

metagenomes is difficult, complicating comparisons with 16S rRNA-based approaches.

Nevertheless, we detected a previously nearly undescribed ITS phylotype, P2.3s1, which was the

most common phylotype in most Arctic PWW and DCM samples. Moving from phylotypes to

MAGs, we detected Arctic SAR11 genomes with both apparently endemic and bi-polar

biogeographic distributions. The selective forces shaping these biogeographic distributions as

well as the resulting adaptive responses on the genome level merit future investigation.

Acknowldgements

Data were collected aboard the CCGS Louis S. St-Laurent in collaboration with

researchers from Fisheries and Oceans Canada at the Institute of Ocean Sciences and Woods

Hole Oceanographic Institution's Beaufort Gyre Exploration Program and are available at

http://www.whoi.edu/beaufortgyre. We would like to thank both the captain and crew of the CCGS

Louis S. St-Laurent, the chief scientist, William J. Williams, and the scientific team aboard. The

work was conducted in collaboration with the U.S. Department of Energy Joint Genome Institute,

a DOE Office of Science User facility, and was supported under Contract No. DE-AC02-

05CH11231. Funding from the Canadian Natural Science and Engineering Research Council

(NSERC) Discovery grants (D.W. and C.L.) and the Canada Research Chair Program (D.W.) are

acknowledged.

References

1.   Morris RM, Rappé MS, Connon SA, Vergin KL, Siebold WA, Carlson CA, et al. SAR11

clade dominates ocean surface bacterioplankton communities. *Nature* 2002.

2.   Brown M V, Lauro FM, DeMaere MZ, Muir L, Wilkins D, Thomas T, et al. Global

biogeography of SAR11 marine bacteria. *Mol Syst Biol* 2012; **8**: 1–13.

3.   Giovannoni SJ. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Ann Rev*

Mar Sci 2017.

4.   Carlson C a, Morris R, Parsons R, Treusch AH, Giovannoni SJ, Vergin K. Seasonal

dynamics of SAR11 populations in the euphotic and mesopelagic zones of the

northwestern Sargasso Sea. *ISME J* 2009; **3**: 283–95.

5.   Ngugi DK, Stingl U. Combined Analyses of the ITS Loci and the Corresponding 16S

rRNA Genes Reveal High Micro- and Macrodiversity of SAR11 Populations in the Red

Sea. *PLoS One* 2012.

6.   Pommier T, Pinhassi J, Hagström Å. Biogeographic analysis of ribosomal RNA clusters

from marine bacterioplankton. *Aquat Microb Ecol* 2005.

7.   Brown M V., Lauro FM, Demaere MZ, Muir L, Wilkins D, Thomas T, et al. Global

biogeography of SAR11 marine bacteria. *Mol Syst Biol* 2012.

8.   Tsementzi D, Wu J, Deutsch S, Nath S, Rodriguez-R LM, Burns AS, et al. SAR11

bacteria linked to ocean anoxia and nitrogen loss. *Nature* 2016.

9.   García-Martínez J, Rodríguez-Valera F. Microdiversity of uncultured marine prokaryotes:

The SAR11 cluster and the marine Archaea of Group I. *Mol Ecol* 2000.

10.   Herlemann DPR, Woelk J, Labrenz M, Jürgens K. Diversity and abundance of

'Pelagibacterales' (SAR11) in the Baltic Sea salinity gradient. *Syst Appl Microbiol* 2014;

37: 601–604.

11.   Oh HM, Kang I, Lee K, Jang Y, Lim S Il, Cho JC. Complete genome sequence of strain

IMCC9063, belonging to SAR11 subgroup 3, isolated from the Arctic Ocean. *J Bacteriol* .

2011.

12. Bahr M, Hobbie JE, Sogin ML. Bacterial diversity in an arctic lake: A freshwater SAR11

cluster. *Aquat Microb Ecol* 1996.

13. McLaughlin FA, Carmack EC, Macdonald RW, Melling H, Swift JH, Wheeler PA, et al.

The joint roles of Pacific and Atlantic-origin waters in the Canada Basin, 1997-1998.

Deep Res Part I Oceanogr Res Pap 2004.

14. Macdonald RW, McLaughlin FA, Carmack EC. Fresh water and its sources during the

SHEBA drift in the Canada Basin of the Arctic Ocean. *Deep Res Part I Oceanogr Res*

Pap 2002.

15. Guéguen C, McLaughlin FA, Carmack EC, Itoh M, Narita H, Nishino S. The nature of

colored dissolved organic matter in the southern Canada Basin and East Siberian Sea.

Deep Res Part II Top Stud Oceanogr 2012.

16. Pedrós-Alió C, Potvin M, Lovejoy C. Diversity of planktonic microorganisms in the Arctic

Ocean. *Prog Oceanogr* 2015.

17. Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, Gonzalez JM, et al.

Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in the

surface ocean. *Proc Natl Acad Sci* 2013.

18. Ghiglione J-F, Galand PE, Pommier T, Pedros-Alio C, Maas EW, Bakker K, et al. Pole-to-

pole biogeography of surface and deep marine bacterial communities. *Proc Natl Acad Sci*

2012.

19. Comeau AM, Li WKW, Tremblay JÉ, Carmack EC, Lovejoy C. Arctic ocean microbial

community structure before and after the 2007 record sea ice minimum. *PLoS One* 2011.

20. Alonso-Sáez L, Sánchez O, Gasol JM, Balagué V, Pedrós-Alio C. Winter-to-summer

changes in the composition and single-cell activity of near-surface Arctic prokaryotes.

Environ Microbiol 2008.

21. Shimada K, Itoh M, Nishino S, McLaughlin F, Carmack E, Proshutinsky A. Halocline

483   structure in the Canada Basin of the Arctic Ocean. *Geophys Res Lett* 2005.

484   22.   Monier A, Comte J, Babin M, Forest A, Matsuoka A, Lovejoy C. Oceanographic structure

485   drives the assembly processes of microbial eukaryotic communities. *ISME J* 2015.

486   23.   Colatriano D, Tran PQ, Guéguen C, Williams WJ, Lovejoy C, Walsh DA. Genomic

487   evidence for the degradation of terrestrial organic matter by pelagic Arctic Ocean

488   Chloroflexi bacteria. *Commun Biol* 2018.

489   24.   Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately

490   reconstructing single genomes from complex microbial communities. *PeerJ* 2015.

491   25.   Bushnell B. BBMap: a fast, accurate, splice-aware aligner. *Joint Genome Instritute,*

492   *department of energy* . 2014.

493   26.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: Assessing the

494   quality of microbial genomes recovered from isolates, single cells, and metagenomes.

495   *Genome Res* 2015.

496   27.   Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho:

497   Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics* 2011.

498   28.   Kumar S, Stecher G, Peterson D, Tamura K. MEGA-CC: Computing core of molecular

499   evolutionary genetics analysis program for automated and iterative data analysis.

500   *Bioinformatics* 2012.

501   29.   Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Tennessen K, et al.

502   The standard operating procedure of the DOE-JGI Metagenome Annotation Pipeline

503   (MAP v.4). *Stand Genomic Sci* 2016.

504   30.   Bairoch A, Apweiler R. The SWISS-PROT protein sequence data bank and its

505   supplement TrEMBL in 1999. *Nucleic Acids Res* . 1999.

506   31.   Yang Z. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol* 2007.

507   32.   Landry Z, Swa BK, Herndl GJ, Stepanauskas R, Giovannoni SJ. SAR202 genomes from

508   the dark ocean predict pathways for the oxidation of recalcitrant dissolved organic matter.

MBio 2017.

33.   Karsenti E, Acinas SG, Bork P, Bowler C, de Vargas C, Raes J, et al. A holistic approach

to marine Eco-systems biology. *PLoS Biol* 2011.

34.   Delmont TO, Quince C, Shaiber A, Esen OC, Lee STM, Lucker S, et al. Nitrogen-Fixing

Populations Of Planctomycetes And Proteobacteria Are Abundant In The Surface Ocean.

doi.org 2017.

35.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, et al. Vegan:

Community Ecology Package. *URL https//cran.r-project.org,*

https//github.com/vegandevs/vegan 2016.

36.   Bano N, Hollibaugh JT. Phylogenetic composition of bacterioplankton assemblages from

the Arctic Ocean. *Appl Environ Microbiol* 2002.

37.   Kirchman DL, Cottrell MT, Lovejoy C. The structure of bacterial communities in the

western Arctic Ocean as revealed by pyrosequencing of 16S rRNA genes. *Environ*

Microbiol 2010.

38.   Galand PE, Potvin M, Casamayor EO, Lovejoy C. Hydrography shapes bacterial

biogeography of the deep Arctic Ocean. *ISME J* 2010; **4**: 564–576.

39.   Cameron TJ, Temperton B, Swan BK, Landry ZC, Woyke T, Delong EF, et al. Single-cell

enabled comparative genomics of a deep ocean SAR11 bathytype. *ISME J* 2014.

40.   Grote J, Thrash JC, Huggett MJ. Streamlining and Core Genome Conservation among

Highly Divergent Members of the SAR11 Clade. 2012; **3**: 1–13.

41.   Schwalbach MS, Tripp HJ, Steindler L, Smith DP, Giovannoni SJ. The presence of the

glycolysis operon in SAR11 genomes is positively correlated with ocean productivity.

Environ Microbiol 2010; **12**: 490–500.

42.   Viklund J, Ettema TJG, Andersson SGE. Independent genome reduction and

phylogenetic reclassification of the oceanic SAR11 clade. *Mol Biol Evol* 2012.

43.   Delmont TO, Kiefl E, Kilinc O, Esen OC, Uysal I, Rappe MS, et al. The global

535  biogeography of amino acid variants within a single SAR11 population is governed by

536  natural selection. *bioRxiv* 2017.

537 44. Kirkpatrick M, Barton NH. Evolution of a Species' Range. *Am Nat* 1997.

538 45. Barton AD, Dutkiewicz S, Flierl G, Bragg J, Follows MJ. Patterns of diversity in marine

539  phytoplankton. *Science (80-)* 2010.

540 46. Thomas MK, Kremer CT, Klausmeier CA, Litchman E. A global pattern of thermal

541  adaptation in marine phytoplankton. *Science (80-)* 2012.

542 47. Ladau J, Sharpton TJ, Finucane MM, Jospin G, Kembel SW, O'Dwyer J, et al. Global

543  marine bacterial diversity peaks at high latitudes in winter. *ISME J* 2013.

544 48. Cabello-Yeves PJ, Zemskay TI, Rosselli R, Coutinho FH, Zakharenko AS, Blinov V V., et

545  al. Genomes of novel microbial lineages assembled from the sub-ice waters of Lake

546  Baikal. *Appl Environ Microbiol* 2018.

547 49. Jimenez-Infante F, Ngugi DK, Vinu M, Blom J, Alam I, Bajic VB, et al. Genomic

548  characterization of two novel SAR11 isolates from the Red Sea, including the first strain

549  of the SAR11 Ib clade. *FEMS Microbiol Ecol* 2017.

Tables

Table 1: SAR11 clades, subclades and phylotypes, as well as their environmental associations

as described in the literature.

Table 2: Western Arctic Ocean metagenome sampling sites, their environmental feature and

environmental parameters including sampling depth (Depth), water temperature (Temp.), Salinity,

Chl a Fluorescence, CDOM, Oxygen and Nitrate (NO_3).

Table 3: Summary statistics of MAGs from the Western Arctic Ocean.

Figure legends

Figure 1: Study metadata and overview of SAR11 ITS phylogeny. A) Sampling locations of the

Western Arctic Ocean metagenomes. B) Environmental profiles of sampling locations showing

temperature (˚C), salinity, chlorophyll a fluorescence (mg m^{-3}) and nitrate (mmol m^{-3}). C) Maximum

likelihood ITS phylogeny including Arctic and reference sequences. Subclades which contain

Arctic ITS sequence types are underlined in orange.

Figure 2: ITS phylogeny of SAR11 subclades. The longest sequence type per Arctic cluster was

used in the phylogeny and only sequence types containing complete ITS regions are included.

Arctic samples are indicated by coloured squares; Surface (green), DCM (orange) and PWW

(blue), Subclades from Table 1: A) S1a, B) S1b , C) S2 and D) S3.

Figure 3: Distribution of SAR11 phylotypes in the Arctic Ocean. A) ITS region coverage of Arctic

SAR11 phylotypes across samples. B) Principal coordinate analysis ordination of Bray-Curtis

dissimilarities of Arctic samples based on the coverage of different ITS phylotypes. Scaling 2 is

shown. Dark blue diamonds: PWW samples, blue diamonds: SCM samples, light blue samples:

surface samples. Asterisks show phylotypes' weighted average frequency.

Figure 4: Phylogenetic context, distribution, transcriptional activity and gene orthology of Arctic

SAR11 MAGs. A) Maximum likelihood phylogeny based on 39 concatenated orthologous loci.

Only bootstrap values higher than 0.6 are shown on the tree. Colored squares and circles at the

tips indicate the environment and temperature of origin (if known), matching phylotypes if known

are indicated. B) Coverage of Arctic MAG scaffolds across Western Arctic ocean metagenomes.

C) RPKB values of the transcriptomes of station CB4 of Arctic MAGs. D) Venn diagram of gene

orthology between the Arctic S2a MAG SAR11-484 and the reference genomes AAA795E07 and

HIMB058.

Figure 5: Global biogeography of SAR11 genomes. A) RPMG table of Arctic MAGs and reference

genomes by ocean layer and ocean region. Abbreviations for oceans are Arctic (AO), Southern

(SO), South Pacific (SP), North Pacific (NP), South Atlantic (SA), North Atlantic (NA), Indian

Ocean (IO), Mediterranean (MS) and Red (RS) Seas. For clarity only RPMG values > 3 are shown

in the figure. B) Principal coordinate analysis ordination of Bray-Curtis dissimilarities of

metagenome samples based on RPMG values of arctic MAGs and reference genomes. Scaling

2 is shown. Arrows indicate significant environmental variables after *Post-Hoc* testing. Asterisks

show the weighted average of each genomes' frequency.

Supplemental information

Supplemental Table 1: List of all genomes and metagenomic samples used in this study.

Supplemental Table 2: List of transcribed genes of Arctic SAR11 MAGs, as well as their COGs

Supplemental Table 3: List of genes unique to arctic SAR11 MAGs, as well as their COGs.

Supplemental Figure 1: Principal coordinate analysis ordination of Bray-Curtis dissimilarities of

metagenome samples based on RPMG values of arctic MAGs and reference genomes, including

TARA ocean-derived SAR11 MAGs. Scaling 2 is shown. Arrows indicate significant environmental

variables after Post-Hoc testing. Asterisks show the weighted average of each genomes'

frequency.

Table 1: SAR11 clades, subclades and phylotypes, as well as their environmental associations as described in the literature.

Clade (16S rRNA)	Subclade (16S rRNA)	Phylotype (ITS)	Reference genome available?	Environmental description
S1	S1a	P1a.1	Y	-Cold [2] -S1: Freshwater Lake Baikal [48]
		P1a.2	N	-Temperate [2]
		P1a.3	Y	-Tropical [2] -Tropical, depth generalist [5]
	S1b	P1b.1	Y	-P1b: Tropical [2] -P1b: Tropical, epi- and bathypelagic [49] -S1b: Tropical, oxygen generalist [8] -P1b.3: Tropical, upper euphotic layer [5]
		P1b.2	Y	
		P1b.3	Y	
	S1c	NA2	Y	-Deep, tropical [5, 8]
S2	S2a.a	-	Y	-OMZ specialist, tropical [8]
	S2a.b	P2.1	Y	-P2.1: Tropical [2]
		P2.2	N	-P2.2: Cold [2] -P2.2: Antarctic [9]
	S2b	P2.3	Y	-P2.3: Cold [2] -P2.3: Deep, tropical [5] -S2b: Tropical, OMZ [8] -P2.3s1: Arctic SCM and PWW (this study)
S3		P3.1	Y	-Coastal, mesohaline, surface [10] -Tropical [2]
		P3.2	Y	-Brakish [10] -Temperate [2] -Arctic [11]
		LD12	Y	-Freshwater [12]

Table 2: Western Arctic Ocean metagenome sampling sites, their environmental feature and environmental parameters including sampling depth (Depth), water temperature (Temp.), Salinity, Chl a Fluorescence, CDOM, Oxygen and Nitrate (NO_3).

Station	Lat.	Long.	Feature	Depth [m]	Temp. [˚C]	Salinity	Fluorescence [mg m^{-3}]	CDOM [mg m^{-3}]	Oxygen [mmol m^{-3}]	NO_3 [mmol m^{-3}]
CB2_154	73.22	-150.22	SUR	7	-1.26	25.70	0.15	2.54	391.85	0
CB2_152	73.22	-150.22	DCM	67	-0.86	31.48	0.33	3.9	351.56	4.45
CB2_150	73.22	-150.22	PWW	177	-1.45	33.18	0.05	4.21	281.85	15.98
CB4_138	75.26	-150.07	SUR	5	-1.39	26.14	0.12	2.58	394.08	0
CB4_136	75.26	-150.07	DCM	79	-0.021	31.16	0.23	3.9	329.14	4.65
CB4_134	75.26	-150.07	PWW	208	-1.47	33.14	0.05	4.36	282.39	16.13
CB8_130	77.1	-150.23	SUR	5	-1.46	27.2	0.19	2.78	395.598	0
CB8_128	77.1	-150.23	DCM	58	-0.15	30.97	0.3	3.89	383.72	0.35
CB8_126	77.1	-150.23	PWW	213	-1.45	33.14	0.05	4.42	281.09	16.24
CB11_90	79.25	-150.06	SUR	5	-1.48	27.31	0.24	2.67	394.88	0
CB11_88	79.25	-150.06	DCM	25	-1.04	29.69	0.25	3.22	401.18	0
CB11_86	79.25	-150.06	PWW	190	-1.46	33.15	0.05	4.35	281.89	15.82

Table 3: Summary statistics of MAGs from the Western Arctic Ocean.

Subgroup	MAG	Size (mb)	Cov (x)	GC (%)	Completeness (%)	Contamination (%)	N50 (kb)	# of scaffolds	# of genes	# of unique genes
S1a	312	0.42	27.83	27.3	47.17	1.89	21.23	20	462	33
S2a	112	0.42	8.66	28.31	38.68	1.89	18.76	20	462	49
S2a	272	0.3	17.96	29.52	35.85	0	98.98	5	320	10
S2a	410	0.42	16.77	30.68	24.53	1.89	18.96	26	462	36
S2a	484	0.44	38.18	28.54	35.09	0	38.17	14	481	35
S2	144	0.63	27.78	29.21	41.51	3.77	30.34	23	672	47
S2	196	0.39	23.43	28.5	46.99	0	93.24	8	430	29

A

B

A

		AO			ACE	SO			SP				NP			SA			NA				IO			MS		RS	
		SURF	DCM	PWW	SURF	SURF	DCM	MES	SURF	MIX	DCM	MES	SURF	DCM	MES	SURF	DCM	MES	SURF	MIX	DCM	MES	SURF	DCM	MES	SURF	DCM	SURF	DCM
S1a	HTCC1062																												
	HTCC1013																												
	SAR11-312																												
	AAA795A20																												
	HIMB083																												
	AAA795O19																												
	AAA795I19																												
	HTCC8051																												
	HTCC7211																												
	HTCC7214																												
	HTCC7217																												
	HTCC9022																												
	RS39																												
	AAA795D22																												
	AAA795O20																												
	HIMB1321																												
	HIMB5																												
S1b	RS40																												
	AAA795E22																												
	AAA795M22																												
S1c	AAA240E13																												
	AAA288E13																												
	AAA288G21																												
	AAA288N07																												
S2a	A10S10																												
	SAR11-272																												
	SAR11-410																												
	SAR11-112																												
	AAA795E07																												
	HIMB058																												
	SAR11-484																												
S2b	A6S6																												
	B3S13																												
	SAR11-144																												
	SAR11-196																												
LD12	AAA023L09																												
	AAA028D10																												
	AAA027C06																												
	AAA027J10																												
	AAA280B11																												
P3.1	HIMB114																												
P3.2	IMCC9063																												

RPMG

B

14.03%

Depth

Nitrate

AAA240E13
AAA288E13
AAA288N07
AAA023L09
AAA288G21

A10S10
SAR11-312
A6S6
B3S13

SAR11-144

SAR11-196

SAR11-272
SAR11-410
SAR11-112

IMCC9063
HTCC1013
HTCC1062
SAR11-484

Salinity

HTCC8051
AAA280B11 HTCC7217
HIMB1321
HTCC7214 RS40
HTCC9022
HTCC7211 HIMB058 HIMB083
AAA795O19 AAA795O20
HIMB5 AAA795A20 AAA795M22
RS39 AAA795I19 AAA795E22
HIMB114 AAA795E07 AAA795D22

Temperature

Chl a

Oxygen

27.78%

Marine Biome
● Coastal Biome
● Polar Biome
● Trades Biome
● Westerlies Biome

Feature
◇ SRF
+ MIX
■ DCM
● PWW
▲ MES

Supplemental information

Supplemental methods

Sampling and DNA extractions

Samples from 4 stations (CB2, CB4, CB8, and CB11) were analysed, with a total of 12 water samples collected from 10-LNiskin bottles mounted on a rosette system. Samples were from the surface layer (Surface: 5 to 7 m), the DCM (25 to 79 m) and the PWW layers (177 to 213 m). The rosette was equipped with profilers for conductivity, temperature and depth (Sea-Bird Electronics Inc., California, USA) coupled with sensors for dissolved oxygen (DO), coloured dissolved organic matter (CDOM) and chlorophyll *a* (Chl *a*) fluorescence. Sample depth was determined during the downward casts, to ensure that target water masses were sampled, with the DCM identified from the Chl *a* fluorescence and PWW centered on a salinity of 31. Nutrient samples and Chl *a* were collected directly from the Niskin bottles and analysed on board as previously described [1]. Consistency of the targeted and actual sampling depths were verified by bottle salinities analyzed onboard using a Guideline Portasal and IAPSO standard seawater [2], while dissolved oxygen (DO) was determined by Winkler-based titration (Kit B, Scripps Institute of Oceanography, USA). Water for planktonic nucleic acid (DNA) analysis was sequentially filtered through a 3-µm polycarbonate filter and 0.22 µm pore size Sterivex® filters (Millipore, Billerica, MA, USA). Filters were preserved in RNAlater for 20 to 40 min and then kept at -80˚C until laboratory processing. DNA was extracted from the Sterivex® filters (3 to 0.22 µm size fraction) following the protocol described previously and stored at -80˚C [3]. Physical oceanographic profiles were visualised using Ocean Data View [4].

Metagenomic sequencing and assembly

The 12 Arctic Ocean samples were sequenced at the Department of Energy Joint Genome Institute (Walnut Creek, CA, USA) using an Illumina HiSeq 2500-1TB platform as described in

Colatriano et al. [3]. Raw reads were assembled using MEGAHIT [5] and gene annotation was performed with the Integrated Microbial Genomes (IMG) database tool (version 4.11.0) [6].

Internal transcribed spacer (ITS) phylogenetic analysis

We used the complete ITS region from a randomly chosen SAR11 reference genome (HIMB114) to extract ITS regions from each of the twelve single sample metagenome assemblies using BLASTn [7]. All hits were filtered to a maximum e-value of 10e-35 and a minimum scaffold size of 350 bp. The retrieved sequences were aligned using the MUSCLE algorithm as implemented in MEGA v.7 [8] and poorly aligned sequences removed after visual inspection. Remaining sequences were clustered using CD-hit [9] at an identity of 100%. Arctic ITS sequences not covering both of the ITS-region tRNAs were removed from the alignment. To assign Arctic sequences to specific phylotypes, we analysed the Arctic ITS sequences with reference sequences from published genomes and ITS sequences used in two previous biogeographic studies [10, 11]. All sequences were realigned using MUSCLE as implemented in MEGA v.7 [8], and the resulting alignment curated in Mesquite [12]. Maximum likelihood trees using GTR+Gamma models of nucleotide substitution were calculated in MEGA v.7 with 100 bootstraps [8]. Subsequently, we assigned phylotype membership of the Arctic ITS sequences (and their CD-hit clusters) based on tree topology. ITS CD-hit clusters representing short ITS fragments were blasted against a custom ITS database of full-length Arctic ITS samples using BLASTn and assigned by their best hit [7].

We performed PCoA ordination of the Bray-Curtis dissimilarity of the coverage of each Arctic ITS phylotype within each sample, correcting for negative eigenvalues using the Caillez method [13]. Ordination was performed using the R packages phyloseq, ade4 and vegan [14–16] and visualized using the package ggplot2 [17]. *Post-hoc* tests were performed using the envdist function as implemented in vegan [16] with 999 permutations. All computational parallelization was performed with parallel [18].

Metagenomic binning

We conducted metagenomic binning using the Metabat2 pipeline [19] on a coassembly of all twelve samples [3]. Fastq reads of all Arctic samples were mapped to scaffolds larger than 10 kb using bbmap [20] with a minimum id of 100 to achieve a high degree of between-bin differentiation. 44 bins belonging to the SAR11 cluster were identified using the checkM lineage_wf function [21]. These bins were cleaned with the refineM filter_bins and taxon_filter options using a taxonomic reference genome database with added SAR11 reference genomes. Scaffolds with aberrant coverage across individual samples were removed after visual inspection.

Concatenated gene phylogeny

Single-copy orthologous genes were identified in seven Arctic SAR11 MAGs, in 41 SAR11 reference genomes representing the clade's phylogenetic diversity, as well as in eight outgroup genomes using ProteinOrtho [22]. We removed all loci present in less than five of the seven MAGs, resulting in 39 orthologs. We aligned their amino acid sequences using the MUSCLE algorithm as implemented in MEGA v.7 [8]. Aligned orthologs were concatenated in Mesquite [12] and positions were masked if they had ZORRO-calculated confidence scores <0.5 [23], resulting in a concatenated alignment of 12 516 amino acids. A maximum likelihood phylogeny was constructed using MEGA-cc with a JTT substitution model, a gamma-distribution with an invariant sites model of rate variation and nearest-neighbour interchange heuristic search method, and bootstrapped 100 times [24].

Arctic SAR11 transcriptome

RNA samples were collected during a cruise in September 2017 and stored as described previously for DNA samples. To extract the RNA, RNAlater was expelled and the filters inundated with 1.5mL Lysis buffer from the mirVana RNA isolation kit (Invitrogen, California, USA), followed

by two minutes of vortexing. 150 µl of miRNA homogenate were added to the filter, followed by vortexing and ten minutes of incubation on ice. The lysate was then removed from the filter and RNA extracted using the phenol-chloroform extraction protocol as described in the mirVana RNA isolation standard protocol (Invitrogen, CA, USA).

Comparative genome content, average nucleotide identity and signatures of selection of Arctic MAGs

We compared the genome content of the Arctic S2a MAG SAR11-484 to two closely related S2a reference genomes, HIMB058 and AAA795E07. Orthologous genes were identified across the three genomes using ProteinOrtho [22]. We investigated the selective pressures acting on these shared genes by performing codon-based nucleotide alignments with MUSCLE as implemented in MEGA v.7 [8]. Pairwise dN/dS ratios of each gene were calculated using the Young and Nielson method in yn00 as implemented in paml [25].

Fragment recruitment and SAR11 biogeography.

Seven Arctic MAGs, 48 representative SAR11 reference genomes, and 38 TARA ocean MAGs previously described as SAR11 [26] were cleaned of rRNA sequences, blasted against the metagenomic datasets using blastall (e-value = 0.00001), and all hits recorded. Subsequently, all recruited reads were extracted from the metagenomic datasets and used in a reciprocal best-hit blast against a database of SAR11 MAGs and reference genomes. Best hits from the reciprocal blast were filtered to a minimum length of 100 bp and a minimum identity of 98% and the number of recorded hits per reference genome and metagenomic sample used to calculate the number of reads recruited per megabase genome per gigabase metagenome (RPMG).

PCoA ordination of the Bray-Curtis dissimilarity of the RPMG matrix was conducted using the R packages; vegan, ade4 and phyloseq [14–16]. Negative eigenvalues were corrected using

the Caillez method [13]. The depth, temperature, salinity, nitrate and Chl *a* concentrations for each sample were projected onto the ordination.

1.   Onda DFL, Medrinal E, Comeau AM, Thaler M, Babin M, Lovejoy C. Seasonal and Interannual Changes in Ciliate and Dinoflagellate Species Assemblages in the Arctic Ocean (Amundsen Gulf, Beaufort Sea, Canada). *Front Mar Sci* 2017.

2.   Shimada K, Itoh M, Nishino S, McLaughlin F, Carmack E, Proshutinsky A. Halocline structure in the Canada Basin of the Arctic Ocean. *Geophys Res Lett* 2005.

3.   Colatriano D, Tran PQ, Guéguen C, Williams WJ, Lovejoy C, Walsh DA. Genomic evidence for the degradation of terrestrial organic matter by pelagic Arctic Ocean Chloroflexi bacteria. *Commun Biol* 2018.

4.   Schlitzer R. Ocean Data View.

5.   Li D, Liu CM, Luo R, Sadakane K, Lam TW. MEGAHIT: An ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 2015.

6.   Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Tennessen K, et al. The standard operating procedure of the DOE-JGI Metagenome Annotation Pipeline (MAP v.4). *Stand Genomic Sci* 2016.

7.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990.

8.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol* 2016.

9. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: Accelerated for clustering the next-generation sequencing data. *Bioinformatics* 2012.

10. Brown M V, Lauro FM, DeMaere MZ, Muir L, Wilkins D, Thomas T, et al. Global biogeography of SAR11 marine bacteria. *Mol Syst Biol* 2012; **8**: 1–13.

11. Ngugi DK, Stingl U. Combined Analyses of the ITS Loci and the Corresponding 16S rRNA Genes Reveal High Micro- and Macrodiversity of SAR11 Populations in the Red Sea. *PLoS One* 2012.

12. Maddison W, Maddison D. Mequite: a modular system for evolutionary analysis. Version 3.04. *http://mesquiteproject.org* 2015.

13. Legendre P, Andersson MJ. Distance-based redundancy analysis: Testing multispecies responses in multifactorial ecological experiments. *Ecol Monogr* 1999.

14. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. *PLoS One* 2013.

15. Dray S, Dufour A-B. The **ade4** Package: Implementing the Duality Diagram for Ecologists. *J Stat Softw* 2007.

16. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, et al. Vegan: Community Ecology Package. *URL https//cran.r-project.org, https//github.com/vegandevs/vegan* 2016.

17. Wickham H. Elegant Graphics for Data Analysis. *Media* 2015.

18. Tang D, Gupta G. An efficient parallel dynamic programming algorithm. *Comput Math with Appl* 1995.

19. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. *PeerJ* 2015.

20.   Bushnell B. BBMap: a fast, accurate, splice-aware aligner. *Joint Genome Instritute, department of energy* . 2014.

21.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: Assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015.

22.   Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics* 2011.

23.   Wu M, Chatterji S, Eisen JA. Accounting for alignment uncertainty in phylogenomics. *PLoS One* 2012.

24.   Kumar S, Stecher G, Peterson D, Tamura K. MEGA-CC: Computing core of molecular evolutionary genetics analysis program for automated and iterative data analysis. *Bioinformatics* 2012.

25.   Yang Z. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol Biol Evol* 2007.

26.   Delmont TO, Quince C, Shaiber A, Esen OC, Lee STM, Lucker S, et al. Nitrogen-Fixing Populations Of Planctomycetes And Proteobacteria Are Abundant In The Surface Ocean. *doi.org* 2017.