The client-binding domain of the cochaperone Sgt2 has a helical-hand structure that binds a short hydrophobic helix

(120 of 120)

Ku-Feng Lin, Michelle Y. Fry*, Shyam M. Saladi*, and William M. Clemons, Jr.

* contributed equally to this work

Correspondence: W.M.C, clemons@caltech.edu
Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125

Keywords: Tail-anchored proteins, Protein targeting, co-chaperones, Sti1, Hop

Abstract (150 words – currently 149)

The targeting and insertion of tail-anchored (TA) integral membrane proteins (IMP) into the correct membrane is critical for cellular homeostasis. The fungal protein Sgt2, and its human homolog SGTA, binds hydrophobic clients and is the entry point for targeting of ER-bound TA IMPs. Here we reveal molecular details that underlie the mechanism of Sgt2 binding to TA IMP clients. We establish that the Sgt2 C-terminal region is flexible but conserved and sufficient for client binding. A molecular model for this domain reveals a helical hand forming a hydrophobic groove, consistent with a higher affinity for TA IMP clients with hydrophobic faces and a minimal length of 11 residues. This work places Sgt2 into a broader family of TPR-containing co-chaperone proteins.

Introduction

An inherently complicated problem of cellular homeostasis is the biogenesis of hydrophobic IMPs which are synthesized in the cytoplasm and must be targeted and inserted into a lipid bilayer. Accounting for ~25% of transcribed genes [1], IMPs are primarily targeted by cellular signal binding factors that recognize a diverse set of hydrophobic α-helical signals as they emerge from the ribosome [2-4]. One important class of IMPs are tail-anchored (TA) proteins whose hydrophobic signals are their single helical transmembrane domain (TMD) located near the C-terminus and are targeted post-translationally to either the ER or mitochondria [5-9]. In the case of the canonical pathway for ER-destined TA IMPs, each is first recognized by homologs of mammalian SGTA (small glutamine tetratricopeptide repeat protein) [4,6,10,11]. Common to all signal binding factors is the need to recognize, bind, and then hand off a hydrophobic helix. How such factors can maintain specificity to a diverse set of hydrophobic clients that must subsequently be released remains an important question.

Homologs of *Saccharomyces cerevisiae* Sgt2 (ySgt2) and *Homo sapiens* SGTA (referred to here as hSgt2 for simplicity), collectively Sgt2, are involved in a variety of cellular processes regarding the homeostasis of membrane proteins including the targeting of TA IMPs [9,12-14], retrograde transport of membrane proteins for ubiquitination and subsequent proteasomal degradation [15], and regulation of mislocalized membrane proteins (MLPs) [16,17]. Among these, the role of Sgt2 in the primary pathways responsible for targeting TA clients to the endoplasmic reticulum (ER) is best characterized, *i.e.* the fungal Guided Entry of Tail-anchored proteins (GET) or the mammalian Transmembrane Recognition Complex (TRC) pathway. In the GET pathway, Sgt2 functions by binding a cytosolic TA client then transferring the TA client to the ATPase chaperone Get3 (human homolog is also Get3) with the aid of the heteromeric Get4/Get5 complex (human Get4/Get5/Bag6 complex) [13,18-20]. In this process, TA client binding to Sgt2, after hand-off from Hsp70, is proposed as the first committed step to ensure that ER TA clients are delivered to the ER membrane while mitochondrial TA clients are excluded [3,13,21]. Subsequent transfer of the TA client from Sgt2 to the ATP bound Get3 induces conformational changes in Get3 that trigger ATP hydrolysis, releasing Get3 from Get4 and favoring binding of the Get3-TA client complex to the Get1/2 (human Get1/Get2) receptor at the ER leading to release of the TA client into the membrane [22-26]. Deletions of yeast GET genes (*i.e. get1Δ*, *get2Δ*, or *get3Δ*) cause cytosolic aggregation of TA clients dependent on Sgt2 [26,27].

In addition to targeting TA IMPs, there is evidence hSgt2 promotes degradation of IMPs

through the proteasome by cooperating with the Bag6 complex, a heterotrimer containing Bag6,

hGet4, and hGet5, which acts as a central hub for a diverse physiological network related to protein

targeting and quality control [19,28-30]. The Bag6 complex can associate with ER membrane-

embedded ubiquitin regulatory protein UbxD8, transmembrane protein gp78, proteasomal

component Rpn10c, and an E3 ubiquitin protein ligase RNF126 thereby connecting hSgt2 to ER

associated degradation (ERAD) and proteasomal activity. Depletion of hSgt2 significantly inhibits

turnover of ERAD IMP clients and elicits the unfolded protein response[16]. Furthermore, the

cellular level of MLPs in the cytoplasm could be maintained by co-expression with hSgt2, which

possibly antagonize ubiquitination of MLPs to prevent proteasomal degradation [15,17]. These

studies demonstrate an active role of hSgt2 in triaging IMPs in the cytoplasm and the breadth of

hSgt2 clients including TA IMPs, ERAD, and MLPs all harboring one or more TMD. Roles for

hSgt2 in disease include polyomavirus infection [31], neurodegenerative disease [27,32],

hormone-regulated carcinogenesis [33,34], and myogenesis [35], although the underlying

molecular mechanisms are still unclear.

The architecture of Sgt2 includes three structurally independent domains that define the three

different interactions of Sgt2 (Fig. 1*A*) [12,36-39]. The N-terminal domain forms a homo-dimer

composed of a four-helix bundle with 2-fold symmetry that primarily binds to the ubiquitin-like

domain (UBL) of Get5/Ubl4A for TA IMP targeting [36,40] or interacts with the UBL on the N-

terminal region of BAG6 [41] where it is thought to initiate downstream degradation processes

[15,28,29]. The central region comprises a co-chaperone domain with three repeated TPR motifs

arranged in a right handed-superhelix forming a 'carboxylate clamp' for binding the C-terminus

of heat-shock proteins (HSP) [12,42]. The highly conserved TPR domain was demonstrated to be

critical in modulating propagation of yeast prions by recruiting HSP70 [27] and may associate

with the proteasomal factor Rpn13 to regulate MLPs [43]. More recently, it was demonstrated that

mutations to residues in the TPR domain which prevent Hsp70 binding impair the loading of TA

clients onto ySgt2 [21], consistent with a direct role of Hsp70 in TA IMP targeting via the TPR

domain. The C-terminal methionine-rich domain of Sgt2 is responsible for binding to hydrophobic

clients such as TA IMPs [11,37,44]. Other hydrophobic segments have been demonstrated to

interact with this domain such as the membrane protein Vpu (viral protein U) from human

immunodeficiency virus type-1 (HIV1), the TMD of tetherin [44], the signal peptide of myostatin

[35], and the N-domain of the yeast prion forming protein Sup35 [27]. All of these studies suggest

that the C-terminus of Sgt2 binds broadly to hydrophobic stretches, yet structural and mechanistic

information for client recognition is lacking.

In this study, we provide the first structural characterization of the C-domains from Sgt2

(Sgt2-C) and show that, in the absence of substrate, it is relatively unstructured. We demonstrate

that a conserved region of the C-domain, defined here as C_{cons}, is sufficient for client binding.

Analysis of the C_{cons} sequence identifies six amphipathic helices whose hydrophobic residues are

required for client binding. Based on this, we computationally generate an *ab initio* structural

model that is validated by point mutants and disulfide crosslinking. Artificial TA clients are then

used to define the properties within clients critical for binding to Sgt2-C. The results show that

Sgt2-C falls into a larger STI1 family of TPR-containing co-chaperones and allow us to propose a

mechanism for TA client binding.

Results

The flexible Sgt2-C domain

Based on sequence alignment (Fig. 1*A*), the Sgt2-C contains a conserved core of six predicted

helices flanked by unstructured loops that vary in length and sequence. Previous experimental

work suggested that this region is particularly flexible, as this domain in the *Aspergillus fumigatus*

is sensitive to proteolysis [12]. Similarly, for ySgt2-TPR-C, the sites sensitive to limited

proteolysis primarily occur within the loops flanking the conserved helices (Fig. 1*A, red arrows*

and S1*B*). This flexible nature of the C-domain likely contributes to its anomalous passage through

a gel-filtration column where Sgt2-C elutes much earlier than the similarly-sized, but well-folded

Sgt2 TPR-domain (Fig. 1*B*). The larger hydrodynamic radius matches previous small-angle X-ray

scattering measurement of the ySgt2 TPR-C domain that indicated a partial unfolded characteristic

in a Kratky plot analysis. The circular dichroism (CD) spectra for both homologs suggests that the

C-domain largely assumes a random-coil conformation, with 40-45% not assignable to a defined

secondary structure category (Fig. 1*C*) [45]. The well-resolved, sharp, but narrowly dispersed

chemical shifts of the backbone amide protons in ^1H-^{15}N HSQC spectra of Sgt2-C (Fig. 1*D,E*),

indicate a significant degree of backbone mobility, similar to natively unfolded proteins [46] and

consistent with results seen by others [47], further highlighting the lack of stable tertiary structure.

[12]. Taken all together, Sgt2-C appears to be a flexible domain.

The conserved region of the C-domain is sufficient for substrate binding

We then asked if the flexible Sgt2-C is the site of client binding in the co-chaperone and if so,

where within this domain is the binding region. During purification Sgt2-C is susceptible to

proteolytic activity being cut at several specific sites (Fig. 1*A*). Proteolysis occurred primarily at

Leu$_{327}$ and in the poorly conserved N-terminal region (between Asp$_{235}$-Gly$_{258}$). Given the

intervening region, between Gly$_{258}$ and Leu$_{327}$ on ySgt2, is conserved (Fig 1*A*), it and the

corresponding region on hSgt2, may mediate TA client binding (Fig. 2*A, grey*). To test this, we

established a set of his-tagged Sgt2 constructs of various lengths (Fig. 2*C*). These Sgt2-C mutants

were co-expressed with an MBP-tagged TA client, Sbh1, and binding was detected by the presence

of captured TA clients in nickel elution fractions (Fig. 2*B*). As previously seen [13], we confirm

that Sgt2-TPR-C alone is sufficient for capturing a TA client (Fig. 2*C*). As one might expect, the

C-domain was also sufficient for binding the TA client. A predicted six α-helical methionine-rich

region of Sgt2-C (Fig. 1*A*), hereafter referred to as Sgt2-C$_{cons}$, is sufficient for binding to Sbh1.

For ySgt2, a minimal region H1-H5 (ΔH0) poorly captures Sbh1, while for hSgt2 the equivalent

minimal region is sufficient for capturing the client at a similar level as the longer C$_{cons}$ domain

(Fig. 2*C*). The predicted helices in Sgt2-C$_{cons}$ are amphipathic and their hydrophobic patches could

be used for client binding (Fig. 2*D*).

To test this, each of the six helices in Sgt2-C$_{cons}$ was mutated to replace the larger hydrophobic

residues with alanines, dramatically reducing the overall hydrophobicity. For all of the helices,

alanine replacement of the hydrophobic residues significantly reduces binding of Sbh1 to Sgt2-C

(Fig. 2*E & F*). While these mutants expressed at similar levels to the wild-type sequence, one

cannot rule out that some of these changes may affect the tertiary structure of this domain. In

general, these results imply that these amphipathic helices are necessary for client binding since

removal of the hydrophobic faces disrupts binding. The overall effect on binding by each helix is

different, with mutations in helices 1-3 having the most dramatic reduction in binding suggesting

that these are more crucial for Sgt2-TA client (Sgt2-TA) complex formation. It is also worth noting,

as this is a general trend, that hSgt2 is more resistant to mutations that affect binding (Fig. 2*F*) than

ySgt2, which likely reflect different thresholds for binding.

Molecular modeling of Sgt2-C domain

Despite the need for a molecular model, the C-domain has resisted structural studies, likely due to the demonstrated inherent flexibility. Based on the six conserved α-helical amphipathic segments (Fig. 1*A*) that contain hydrophobic residues critical for TA client binding (Fig. 2*D-E),* we expect some folded structure to exist. Therefore, we performed *ab initio* molecular modeling of Sgt2-C using a variety of prediction methods resulting in a diversity of putative structures [48-52]. As expected, all models showed buried hydrophobic residues as this is a major criterion for *in silico* protein folding. Residues outside the ySgt2-C$_{cons}$ region adopted varied conformations consistent with their expected higher flexibility. Pruning these N- and C-terminal regions to focus on the ySgt2-C$_{cons}$ region (Fig. S2*A*) revealed a potential binding interface for a hydrophobic substrate, examples are seen in Quark models (1, 4, & 6 shown), Robetta 1 & 2, and I-TASSER 2 & 3, whereas others models had no clearly distinguishable groove. Given the intrinsic flexibility of the Sgt2-C domain, it is possible that models without a groove are found in the non-TMD bound structural ensemble.

For a working model of TMD-bound ySgt2-C, we chose the highest scored Quark structures where a general consistent architecture was seen (Fig. 3*A*) [48]. The overall model contained a potential TA client binding site, a hydrophobic groove formed by the amphipathic helices. The groove is approximately 15 Å long, 12 Å wide, and 10 Å deep, which is sufficient to accommodate three helical turns of an α-helix, ~11 amino acids (Fig. 3*B*).

To validate the model, we interrogated the accuracy of the predicted structural arrangement by determining distance constraints from crosslinking experiments. We selected four pairs of residues in close spatial proximity and one pair far apart based on the Quark models (Fig. 4*A*). Calculating a C$_\beta$-C$_\beta$ distance between residue pairs for each model (Fig. 4*E*), the Quark models 2 and 3 were the most consistent with an expected distance of 9Å or less for the close pairs. In all alternative models, the overall distances are much larger and should not be expected to form disulfide bonds *in vitro* if they represent a TMD-bound state. For Robetta, a number of the models have pairs of residues within 9Å and Robetta's per-residue error estimate suggests relatively high confidence in the C$_{cons}$ region (Fig. S2*B*).

As a control, we first confirmed that the cysteine-mutant pairs do not affect the function of ySgt2. We utilized an *in vitro* transfer assay where a yeast Hsp70 homolog Ssa1 loaded with a TA client delivers the client to ySgt2 [21,49,50] (Fig. 4*C*). Purified Ssa1 is mixed with detergent solubilized strep-tagged Bos1-TMD (a model ER TA client) that contained a p-benzoyl-l-

phenylalanine (BPA) labeled residue, Bos1_BPA, are diluted to below the critical micelle

concentration resulting in soluble complexes of Bos1_BPA/Ssa1. Full-length ySgt2 variants were

each tested for the ability to capture Bos1_BPA from Ssa1. After the transfer reaction, each was UV-

treated to generate Bos1 crosslinks. Successful capture of the TA clients by ySgt2 was detected

using an anti-strep Western blot and the appearance of a Bos1_BPA/ySgt2 crosslink band (Fig. 4*D*).

All of the cysteine variants of ySgt2 successfully captured Bos1_BPA from Ssa1 similar to wild-type

suggesting that the cysteine mutations did not affect the structure or function of ySgt2.

For the distance experiment, each of the cysteine-mutant pairs was made in ySgt2-TPR-C

which lacks the dimerization domain. Each variant was coexpressed with an artificial TA client, a

cMyc-tagged BRIL (small, 4-helix bundle protein [51]) with a C-terminal TMD consisting of eight

leucines and three alanines, denoted 11[L8], and purified via nickel-affinity chromatography in

reducing buffer (Fig. S3*A*). All of the ySgt2 mutants bound the TA client and behaved similar to

the wild-type (cysteine-free) further suggesting the mutants did not perturb the native structure

(Fig. S3*B*). For disulfide crosslink formation, each eluate was oxidized and crosslinks were

identified by the visualization of a reducing-agent sensitive ~7.7kDa fragment in gel

electrophoresis (Fig. 3*B*). For both the wild-type construct and in N285C/G329C, where the pairs

are predicted from the Quark models to be too distant for disulfide bond formation, no higher

molecular weight band was observed. For the remaining pairs that are predicted to be close enough

for bond formation, the 7.7kDa fragment was observed in each case and is labile in reducing

conditions. Again, these results support the C_{cons} model derived from Quark.

With the four crosslinked pairs as distance constraints, new models were generated using

Robetta with a restraint on the corresponding pairs of C_β atoms less than 9Å (Fig. S4*A*). The

Robetta models from these runs are similar to the top scoring models from Quark (Fig. 3).

Satisfyingly, the pair of residues that do not form disulfide crosslinks are generally consistent (Fig.

S4*B*).

The improvement of the ySgt2 models predicted by Robetta with restraints included

encouraged us to generate models for hSgt2-C with constraints. For this, pairs were defined based

on sequence alignments of Sgt2 (Fig. 1*A*) and used as restraints. The resulting predictions had

architectures consistent with the equivalent regions predicted for ySgt2-C_{cons}, for example Robetta

4 (Fig. S4*C*, top). Although in general the predicted hSgt2 model is similar to that for ySgt2, the

region that corresponds to H2 occupies a position that precludes a clear hydrophobic groove. For

ySgt2, the longer N-terminal loop occupies the groove preventing the exposure of hydrophobics

to solvent (Fig. 3*C*, grey). For hSgt2, the shorter N-terminal loop may not be sufficient to similarly

occupy the groove and allowing for the clear hydrophobic hand seen for the ySgt2-C. To correct

for this, we replaced the sequence of the N-terminal loop of hSgt2-C with the ySgt2-C loop and

ran structure prediction with the pairwise distance restraints. This resulted in a model where the

loop occupies the groove and, when pruned away suggests the hydrophobic hand seen in yeast

(Fig. S4*C*, middle boxed). Of note, we also generated models of hSgt2-C using the most recent

Robetta method (transform-restrained) which produces new structures with a groove and similar

helical-hand architecture across the board (Fig. S4*C*, bottom).

We sought to further test the robustness of our model considering the intrinsic flexibility of

Sgt2-C by probing for disulfide bond formation with neighboring residues of one of our

crosslinking pairs. While the C_β-C_β distance puts these adjacent pairs at farther than 9Å, mutating

residues to cystines and measuring S-S distances across all possible pairs of rotamers provides a

wider interval on possible distances and, therefore, the likelihood a disulfide bond will form (Fig.

4*F*). Cysteine mutants were introduced to the residues adjacent to M289 and A319 in ySgt2-TPR-

C for four additional pairs: K288C/A319C, M290C/A319C, M289C/P318C, and M289C/L320C.

As described previously, these mutants were coexpressed with a TA substrate, in this case the

artificial BRIL-11[L8] which has a MBP-tag instead of a cMyc-tag. Complexes were purified by

amylose and then nickel affinity chromatography to ensure eluates contained only Sgt2-TPR-C

bound to substrate. Eluates were incubated in oxidizing conditions, quenched with 50mM NEM,

and digested with Glu-C protease. Again, a reductant sensitive band at 7.7kDa is observed for

each of these adjacent pairs. While the geometry of each of these C-C pairs might suggest against

disulfide bond formation, given the intrinsic flexibility of Sgt2-C, it is not surprising that each of

these pairs are able to form disulfide bonds. As before, disulfide bond formation was detected for

the M289C/A319C pair. In this new construct, we now see a small amount of disulfide bond

formation in the distant N285C/G329C pair, likely an effect of switching to the MBP tag.

Structural similarity of Sgt2-C domain to STI1 domains

Attempts to glean functional insight for Sgt2-C from BLAST searches did not reliably return

other families or non-Sgt2 homologs making functional comparisons difficult. A more extensive

profile-based search using hidden Markov models from the SMART database [52] identified a

similarity to domains in the yeast co-chaperone Sti1 (HOP in mammals). First called DP1 and DP2,

due to their prevalence of aspartates (D) and prolines (P), these domains have been shown to be

required for client-binding by Sti1 [53,54] and are termed 'STI1'-domains in bioinformatics

databases [52]. In yeast Sti1 and its human homolog HOP (combined will be referred to here as

Sti1), each of the two STI1 domains (DP1 and DP2) are preceded by Hsp70/90-binding TPR

domains, similar to the domain architecture of Sgt2. Deletion of the second, C-terminal STI1-

domain (DP2) from Sti1 *in vivo* is detrimental, impairing native activity of the glucocorticoid

receptor [53]. *In vitro*, removal of the DP2 domain from Sti1 results in the loss of recruitment of

the progesterone receptor to Hsp90 without interfering in Sti1-Hsp90 binding [55]. These results

implicate DP2 in binding of Sti1 clients. In addition, others have noted that, broadly, STI1-domains

may present a hydrophobic groove for binding the hydrophobic segments of a client [53,54].

Furthermore, the similar domain organizations (*i.e.* Sgt2 TPR-C, Sti1 TPR-STI1) and molecular

roles could imply an evolutionary relationship between these co-chaperones. Indeed, a multiple

sequence alignment of the Sgt2-C_{cons} with several yeast STI1 domains (Fig. 5*A*) reveals strong

conservation of structural features. H1-H5 of the predicted helical regions in C_{cons} align directly

with the structurally determined helices in the DP2 domain of Sti1; this includes complete

conservation of helix breaking prolines and close alignment of hydrophobic residues in the

amphipathic helices [53].

Based on the domain architecture and homology, a direct comparison between the STI1

domain and Sgt2-C_{cons} can be made. A structure of DP2 solved by solution NMR reveals that the

five amphipathic helices assemble to form a flexible helical-hand with a hydrophobic groove [53].

The lengths of the α-helices in this structure concur with those inferred from the alignment in Fig.

4*A*. Our molecular model of Sgt2-C_{cons} is strikingly similar to this DP2 structure (Fig. 5*B,C*). An

overlay of the DP2 structure and our molecular model demonstrates both Sgt2-C_{cons} and DP2 have

similar lengths and arrangements of their amphipathic helices (Fig. 5*D*). Consistent with our

observations of flexibility in Sgt2-C_{cons}, Sti1-DP2 generates few long-range NOEs between its

helices indicating that Sti1-DP2 also has a flexible architecture [53]. We consider this flexibility a

feature of these helical-hands for reversible and specific binding of a variety of clients.

Binding mode of TA clients to Sgt2

We examined the Sgt2-C_{cons} surface that putatively interacts with TA clients by constructing

hydrophobic-to-charge residue mutations that are expected to disrupt capture of TA clients by Sgt2.

Similar to the helix mutations in Fig. 2, the capture assay was employed to establish the relative

effects of individual mutations. A baseline was established based on the amount of the TA client

Sbh1 captured by wild-type Sgt2-TPR-C. In each experiment, Sbh1 was expressed at the same

level; therefore, differences in binding should directly reflect the affinity of Sgt2 mutants for clients.

In all cases, groove mutations from hydrophobic to aspartate led to a reduction in TA client binding

(Fig. 6). The effects are most dramatic with ySgt2 where each mutant significantly reduced binding

by 60% or more (Fig. 6*A*). While all hSgt2 individual mutants saw a significant loss in binding,

the results were more subtle with the strongest a ~36% reduction (M233D, Fig. 6*B*). Double

mutants were stronger with a significant decrease in binding relative to the individual mutants,

more reflective of the individual mutants in ySgt2. As seen before (Fig. 2*E&F*), we observe that

mutations toward the N-terminus of Sgt2-C have a stronger effect on binding than those later in

the sequence.

Sgt2-C domain binds clients with a hydrophobic segment ≥ 11 residues

With a molecular model for ySgt2-C$_{cons}$ and multiple lines of evidence for a hydrophobic

groove, we sought to better understand the specific requirements for TMD binding. We and others

have demonstrated that a monomeric C-domain from Sgt2 is sufficient for binding to TA clients

[13]. To study the minimal constraints on TA client binding, we chose to focus on a monomeric

construct of Sgt2 (Sgt2-TPR-C) binding to variable TMDs. TA clients were designed where the

overall (sum) and average (mean) hydrophobicity, length, and the distribution of hydrophobic

character were varied in the TMDs. These artificial TMDs, a Leu/Ala helical stretch followed by

a Trp, were constructed as C-terminal fusions to the soluble protein BRIL (Fig. 7*A*). The total and

mean hydrophobicity are controlled by varying the helix-length and the Leu/Ala ratio. For clarity,

we define a syntax for the various artificial TA clients to highlight the various properties under

consideration: hydrophobicity, length, and distribution. The generic notation is TMD-

length[number of leucines] which is represented, for example, as 18[L6] for a TMD of 18 amino

acids containing six leucines.

Our first goal with the artificial clients was to define the minimal length of a TMD to bind to

the C-domain. As described earlier, captures of his-tagged Sgt2-TPR-C with the various TA clients

were performed. We define a relative binding efficiency as the ratio of captured TA client by a

Sgt2-TPR-C normalized to the ratio of a captured wild-type TA client by Sgt2-TPR-C. In this case

we replaced the TMD in our artificial clients with the native TMD of Bos1 (Bos1$_{TMD}$). The

artificial client 18[L13] shows a comparable binding efficiency to Sgt2-TPR-C as that of Bos1$_{TMD}$

(Fig. 7*B*). From the helical wheel diagram of the TMD for Bos1, we noted that the hydrophobic

residues favored one face of the helix. We explored this 'hydrophobic face' by using model clients

that maintained this orientation while shortening the length and maintaining the average

hydrophobicity of 18[L13] (Fig. 7*B*). Shorter helices of 14 or 11 residues, 14[L10] and 11[L8],

also bound with similar affinity to Bos1. Helices shorter than 11 residues, 9[L6] and 7[L5], were

not able to bind Sgt2-TPR-C (Fig. 7*B*) establishing a minimal length of 11 residues for the helix

consistent with the dimensions of the groove predicted from the structural model (Fig. 3). In the

context of the full-length Sgt2, which exists as a dimer, an 11-residue cut-off suggests that two C-

domains could come together and bind to a single TA whose TMD lengths range from 18-24

residues.

Since a detected binding event occurs with TMDs of at least 11 amino acids, we decided to

probe this limitation further. The dependency of client hydrophobicity was tested by measuring

complex formation of Sgt2-TPR-C and artificial TA clients containing an 11 amino acid TMD with

increasing number of leucines (11[Lx]). As shown in Fig. 7*C*, increasing the number of leucines

monotonically enhances complex formation, echoing previous results [56]. hSgt2-TPR-C binds to

a wider spectrum of hydrophobic clients than ySgt2-TPR-C, which could mean it has a more

permissive hydrophobic binding groove, also reflected in the milder impact of Ala replacement

and Asp mutations in hSgt2-TPR-C to TA client binding (Fig. 2*C* and Fig. 6*A*).

331 Sgt2-C preferentially binds to TMDs with a hydrophobic face

Next, we address the properties within the TMD of TA clients responsible for Sgt2 binding.

In the case of ySgt2, it has been suggested that the co-chaperone binds to TMDs based on

hydrophobicity and helical propensity [56]. In our system, our artificial TMDs consist of only

alanines and leucines which have high helical propensities [57], and despite keeping the helical

propensity constant and in a range that favors Sgt2 binding, there is still variation in binding

efficiency. For the most part, varying the hydrophobicity of an artificial TA client acts as expected,

the more hydrophobic TMDs bind more efficiently to Sgt2 TPR-C (Fig. 7*C*). Our C$_{cons}$ model

suggests the hydrophobic groove of ySgt2-C protects a TMD with highly hydrophobic residues

clustered to one side (see Fig. 3*B*). To test this, various TMD pairs with the same hydrophobicity,

but different distributions of hydrophobic residues demonstrates TA clients with clustered leucines

have a higher relative binding efficiency than those with a more uniform distribution (Fig. 7*D*).

Helical wheel diagrams demonstrate the distribution of hydrophobic residues along the helix (*e.g.*

bottom Fig. 7*D*). The clustered leucines in the TMDs create a hydrophobic face which potentially

interacts with the hydrophobic groove formed by the Sgt2-C$_{cons}$ region, corresponding to the model

in Fig. 3*B*.

Discussion

Sgt2, the most upstream component of the GET pathway, plays a critical role in the targeting

of TA IMPs to their correct membranes. Its importance as the first confirmed selection step of ER

versus mitochondrial [56] destined TA IMPs necessitates a molecular model for TA client binding.

Previous work demonstrated a role for the C-domain of Sgt2 to bind to hydrophobic clients, yet

the exact binding domain remained to be determined. Through the combined use of biochemistry,

bioinformatics, and computational modeling, we conclusively identify the minimal client-binding

domain of Sgt2. Pulldown experiments revealed a minimal six α-helical region as the TMD binding

domain of Sgt2 that is also conserved in structure and sequence analyses. This domain

preferentially binds TMDs with hydrophobic residues organized onto one face of the helix. While

NMR spectroscopy displays an intrinsically disordered domain in the absence of substrate,

computational models of ySgt2-C consistently predict a structured helical hand in its conserved

region, while the rest of the domain remained flexible and varied between models. Structural

similarities between the ySgt2-C model and the STI1 domains DP1 and DP2 now place Sgt2

among a class of co-chaperones. Together, these results allow us to present a validated structural

model of the Sgt2 C-domain as a methionine-rich helical hand for grasping a hydrophobic helix

and to provide a mechanistic explanation for binding a TMD of at least 11 hydrophobic residues.

We confidently identify the C-domain of Sgt2 as containing a STI1 domain for client binding

through sequence alignments and structural homology. This places Sgt2 into a larger context of

conserved co-chaperones (Fig. 8*A*). In the co-chaperone family, the STI1 domains predominantly

follow HSP-binding TPR domains connected by a flexible linker, reminiscent of the domain

architecture of Sgt2. As noted above, for Sti1 these domains are critical for client-processing and

coordinated hand-off between Hsp70 and Hsp90 homologs [58] as well as coordinating the

simultaneous binding of two heat shock proteins. Both Sgt2 and the co-chaperone Hip coordinate

pairs of TPR and STI1 domains by forming stable dimers via their N-terminal dimerization

domains [59]. With evidence for a direct role of the carboxylate-clamp in the TPR domain of Sgt2

for client-binding now clear [21], one can speculate that the two TPR domains may facilitate TA

client entry into various pathways that use multiple heat shock proteins.

Computational modeling reveals a conserved region sufficient for TA client binding that

consists of a helical hand of five alpha-helices that form a hydrophobic groove to bind the client

TMD. The concept of TMD binding by a helical hand is reminiscent of other proteins involved in

membrane protein targeting. Like Sgt2, the signal recognition particle (SRP) contains a

methionine-rich domain that binds signal sequences and TMDs. While the helical order is inverted,

again five amphipathic helices form a hydrophobic groove that cradles the client signal peptide

[60] (Fig. 8*B*). Here once more, the domain has been observed to be flexible in the absence of

client [61,62] and, in the resting state, occupied by a region that includes a helix which must be

displaced [60]. Another helical-hand example recently shown to be involved in TA IMP targeting

is calmodulin where a crystal structure reveals two helical hands coordinating to clasp a TMD at

either end (Fig. 8*B*) [63]. Considering an average TMD of 18-20 amino acids (to span a ~40Å

bilayer), each half of calmodulin interacts with about 10 amino acids. This is in close

correspondence to the demonstrated minimal 11 amino acids for a TA client to bind to the

monomeric Sgt2-TPR-C. In the context of the full-length Sgt2, one can speculate that the Sgt2

dimer may utilize both C-domains to bind to a full TMD, similar to calmodulin. Cooperation of

the two Sgt2 C-domains in client-binding could elicit conformational changes in the complex that

could be recognized by downstream factors, such as additional interactions that increase the

affinity to Get5/Ubl4A.

Intriguingly, Sgt2-TPR-C preferentially binds to artificial clients with clustered leucines. If

the C-domain forms a hydrophobic groove as suggested by the computational model, it provides

an attractive explanation for this preference. In order to bind to the hydrophobic groove, a client

buries a portion of its TMD in the groove leaving the other face exposed. Clustering hydrophobic

residues contributes to the hydrophobic effect driving binding efficiency and protecting them from

the aqueous environment. Indeed, GET pathway substrates have been suggested to be more

hydrophobic TMDs than EMC substrates [64]. Of these, for the most hydrophobic substrates, like

Bos1, residues on both sides of the TMD could be protected by a pair of C-domains. Alternatively,

the unstructured N-terminal loop through H0 could act as a lid surrounding the circumference of

the client's TMD. Unstructured regions participating in substrate binding as well as capping a

hydrophobic groove have both suggested in the context of other domains, *e.g.* with Get3 [4]. The

role for this clustering of hydrophobic residues in TA client recognition and targeting merits further

investigation.

What is the benefit of the flexible helical-hand structure for hydrophobic helix binding? While

it remains an open question, it is notable that evolution has settled on similar simple solutions to

the complex problem of specific but temporary binding of hydrophobic helices. For all of the

domains mentioned above, the flexible helical-hands provide an extensive hydrophobic surface to

capture a client-helix—driven by the hydrophobic effect. Typically, such extensive interfaces are

between pairs of pre-ordered surfaces resulting in high affinities and slow off rates. These helical

hands are required to only engage temporarily, therefore the flexibility offsets the favorable free

energy of binding by charging an additional entropic cost for ordering a flexible structure in the

client-bound complex. The benefit for TA clients is a favorable transfer to downstream components

in the GET pathways as seen for ySgt2 [21] and hSgt2 [50]. The demonstration that TA transfer

from hSgt2 to Get3 is twice as fast as disassociation from hSgt2 into solution, perhaps interaction

with Get3 leads to conformational changes that further favor release [50].

While hSgt2 and ySgt2 share many properties, there are a number of differences between the

two homologs that may explain the different biochemical behavior. For the C_{cons}-domains, hSgt2

appears to be more ordered in the absence of client as the peaks in its NMR spectra are broader

(Fig. 1*E*). Comparing the domains at the sequence level, while the high glutamine content in the

C-domain is conserved it is higher in hSgt2 (8.8% versus 15.2%). The additional glutamines are

concentrated in the predicted longer H4 helix (Fig. 1*A*). The linker to the TPR domain is shorter

compared to ySgt2 while the loop between H3 and H4 is longer. Do these differences reflect

different roles? As noted, in every case the threshold for hydrophobicity of client-binding is lower

for hSgt2 than ySgt2 (Fig. 1*E*, 5, and 6) implying that the mammalian protein is more permissive

in client binding. The two C-domains have similar hydrophobicity, so this difference in binding

might be due to a lower entropic cost paid by having the hSgt2 C-domain more ordered in the

absence of client or the lack of an unstructured N-terminal loop.

The targeting of TA clients presents an intriguing and enigmatic problem for understanding

the biogenesis of IMPs. How subtle differences in each client modulates the interplay of hand-offs

that direct these proteins to the correct membrane remains to be understood. In this study, we focus

on a central player, Sgt2 and its client-binding domain. Through biochemistry and computational

analysis, we provide a structural model that adds more clarity to client discrimination.

Material and Methods

Plasmid constructs

MBP-Sbh1 full length, $ySgt2_{95-346}$ ($ySgt2$-TPR-C), $ySgt2_{222-346}$ ($ySgt2$-C), $ySgt2_{260-327}$ ($ySgt2$-C_{cons}), $ySgt2_{266-327}$ ($ySgt2$-ΔH0), $hSgt2_{87-313}$ ($hSgt2$-TPR-C), $hSgt2_{213-313}$ ($hSgt2$-C), $hSgt2_{219-300}$ ($hSgt2$-C_{cons}), and $hSgt2_{228-300}$ ($hSgt2$-ΔH0) were prepared as previously described [12,65]. Genes of $ySgt2$ or $hSgt2$ variants were amplified from constructed plasmids and then ligated into an pET33b-derived vector with a 17 residue N-terminal hexa-histidine tag and a tobacco etch virus (TEV) protease site. Single or multiple mutations on Sgt2 were constructed by site-direct mutagenesis. Artificial TA clients were constructed in a pACYC-Duet plasmid with a N-terminal cMyc tag, BRIL fusion protein [66], GSS linker, and a hydrophobic C-terminal tail consisting of leucines and alanines and ending with a tryptophan.

Protein expression and purification

All proteins were expressed in *Escherichia coli* NiCo21 (DE3) cells (New England BioLabs). To co-express multiple proteins, constructed plasmids were co-transformed as described [65]. Protein expression was induced by 0.3 mM IPTG at $OD_{600} \sim 0.7$ and harvested after 3 hours at 37°C. For structural analysis, cells were lysed through an M-110L Microfludizer Processor (Microfluidics) in lysis buffer (50 mM Tris, 300 mM NaCl, 25 mM imidazole supplemented with benzamidine, phenylmethylsulfonyl fluoride (PMSF), and 10 mM β-mercaptoethanol (BME), pH 7.5). For capture assays, cells were lysed by freeze-thawing 3 times with 0.1 mg/mL lysozyme. To generate endogenous proteolytic products of $ySgt2$-TPR-C for MS analysis, PMSF and benzamidine were excluded from the lysis buffer. His-tagged Sgt2 and his-tagged Sgt2/TA complexes were separated from the lysate by batch incubation with Ni-NTA resin at 4°C for 1hr. The resin was washed with 20 mM Tris, 150 mM NaCl, 25 mM imidazole, 10 mM BME, pH 7.5. The complexes of interest were eluted in 20 mM Tris, 150 mM NaCl, 300 mM imidazole, 10 mM BME, pH 7.5.

For structural analysis, the affinity tag was removed from complexes collected after the nickel elution by an overnight TEV digestion against lysis buffer followed by size-exclusion chromatography using a HiLoad 16/60 Superdex 75 prep grade column (GE Healthcare).

Measurement of Sgt2 protein concentration was carried out using the bicinchoninic acid (BCA) assay with bovine serum albumin (BSA) as standard (Pierce Chemical Co.). Samples for

NMR and CD analyses were concentrated to 10-15 mg/mL for storage at −80°C before
experiments.

For the *in vitro* transfer assay, plasmids encoding for the full-length ySgt2 cysteine mutants
were transformed into BL21 Star cells (Invitrogen). Cells were grown in 2x yeast-tryptone (2xYT)
media and induced with 0.1mM IPTG at an OD_{600} of 0.6 then harvested after 3 hours at 30°C by
centrifugation. Cells were lysed in 50mM Tris pH 8.0, 500mM NaCl, 10% glycerol, and 1x
BugBuster (Millipore Sigma), supplemented with protease inhibitors (4-(2-
aminoethyl)benzenesulfonyl fluoride hydrochloride (Roche), benzamidine, and BME). Full-length
his-tagged ySgt2 and cysteine mutants were separated from the lysate by batch incubation with
Ni-NTA resin (Qiagen) at 4°C for 1 hour. The resin was washed with 50mM Tris pH 8.0, 500mM
NaCl, 10% glycerol, and 25mM imidazole and then the protein was eluted in 50mM Tris pH 8.0,
500mM NaCl, 10% glycerol, and 300mM imidazole. For storage, protein was dialyzed in 25mM
K-HEPES pH 7.5, 150mM KOAc, and 20% glycerol at 4°C and then flash frozen in liquid nitrogen.
Purified Bos1 with p-benzoyl-l-phenylalanine (BPA) labeled at residue 230 (Bos1BPA) and yeast
Ssa1 were gifts from the lab of Shu-ou Shan (Caltech).

NMR Spectroscopy

482   15N-labeled proteins were generated from cells grown in auto-induction minimal media as
483 described [67] and purified in 20 mM phosphate buffer, pH 6.0 (for ySgt2-C, 10mM Tris, 100mM
484 NaCl, pH 7.5). The NMR measurements of 15N-labeled Sgt2-C proteins (~0.3-0.5 mM) were
485 collected using a Varian INOVA 600 MHz spectrometer at either 25°C (ySgt2-C) or 35°C (hSgt2-
486 C) with a triple resonance probe and processed with TopSpin™ 3.2 (Bruker Co.).

CD Spectroscopy

488   The CD spectra were recorded at 24°C with an Aviv 202 spectropolarimeter using a 1 mm
489 path length cuvette with 10 μM protein in 20 mM phosphate buffer, pH 7.0. The CD spectrum of
490 each sample was recorded as the average over three scans from 190/195 to 250 nm in 1 nm steps.
491 Each spectrum was then decomposed into its most probable secondary structure elements using
492 BeStSel [68].

Glu-C digestion of the double cysteine mutants on *ySgt2*-C

Complexes of the co-expressed wild type or double cysteine mutated His-*ySgt2*-TPR-C and

the artificial TA client, 11[L8], with either a cMyc or MBP tag were purified as the other His-Sgt2

complexes described above or initially purified via amylose affinity chromatography before nickel

chromatography explained earlier. The protein complexes were mixed with 0.2 mM CuSO4 and

0.4 mM 1,10-phenanthroline at 24°C for 20 min followed by 50 mM N-ethyl maleimide for 15

min. Sequencing-grade Glu-C protease (Sigma) was mixed with the protein samples at an

approximate ratio of 1:30 and the digestion was conducted at 37°C for 22 hours. Digested samples

were mixed with either non-reducing or reducing SDS-sample buffer, resolved via SDS-PAGE

using Mini-Protean® Tris-Tricine Precast Gels (10-20%, Bio-Rad), and visualized using

Coomassie Blue staining.

In vitro transfer assay of Bos1 from Ssa1 to *ySgt2*

The *in vitro* transfer assays were performed as in Chio et al. 2019 and Shao et al. 2017 [49,50].

Specifically, 39µM Bos1BPA (50mM HEPES, 300mM NaCl, 0.05% LDAO, 20% glycerol) was

diluted to a final concentration of 0.1µM and added to 4µM Ssa1 supplemented with 2mM ATP

(25mM HEPES pH7.5, 150mM KOAc). After one minute, 0.3µM of full-length ySgt2 or mutant

was added to the reaction. Samples were flash frozen after one minute and placed under a 365nm

UV lamp for 2 hours on dry ice to allow for BPA crosslinking.

Protein immunoblotting and detection

For western blots, protein samples were resolved via SDS-PAGE and then transferred onto

nitrocellulose membranes by the Trans-Blot® Turbo™ Transfer System (Bio-Rad). Membranes

were blocked in 5% non-fat dry milk and hybridized with antibodies in TBST buffer (50 mM Tris-

HCl pH 7.4, 150 mM NaCl, 0.1% Tween 20) for 1 hour of each step at 24°C. The primary

antibodies were used at the following dilutions: 1:1000 anti-penta-His mouse monoclonal (Qiagen),

1:5000 anti-cMyc mouse monoclonal (Sigma), and a 1:3000 anti-Strep II rabbit polyclonal

(Abcam). A secondary antibody conjugated either to alkaline phosphatase (Rockland, 1:8000) or

a 800nm fluorophore was employed, and the blotting signals were chemically visualized with

either the nitro-blue tetrazolium/5-bromo-4-chloro-3'-indolyphosphate (NBT/BCIP) chromogenic

assay (Sigma) or infrared scanner. All blots were photographed and quantified by image

densitometry using ImageJ [69] or ImageStudioLite (LI-COR Biosciences).

Quantification of Sgt2-TA complex formation

The densitometric analysis of MBP-Sbh1 capture by His-Sgt2-TPR-C quantified the intensity of the corresponding protein bands on a Coomassie Blue G-250 stained gel. The quantified signal ratios of MBP-Sbh1/His-Sgt2-TPR-C are normalized to the ratio obtained from the wild-type (WT). Expression level of MBP-Sbh1 was confirmed by immunoblotting the MBP signal in cell lysate. Average ratios and standard deviations were obtained from 3-4 independent experiments.

In artificial TA client experiments, both his-tagged Sgt2-TPR-C and cMyc-tagged artificial TA clients were quantified via immunoblotting signals. The complex efficiency of Sgt2-TPR-C with various TA clients was obtained by

$$E_{complex} = \frac{E_{TA}}{T_{TA}} \times \frac{1}{E_{capture}} \tag{1}$$

where E_{TA} is the signal intensity of an eluted TA client representing the amount of TA client co-purified with Sgt2-TPR-C and T_{TA} is the signal intensity of a TA client in total lysate which corresponds to the expression yield of that TA client. Identical volumes of elution and total lysate from different TA clients experiments were analyzed and quantified. In order to correct for possible variation the amount of Sgt2-TPR-C available for complex formation, $E_{capture}$ represents the relative amount of Sgt2-TPR-C present in the elution (E_{Sgt2}) compared to a pure Sgt2-TPR-C standard ($E_{purified,Sgt2}$).

$$E_{capture} = \frac{E_{Sgt2}}{E_{purified,Sgt2}}, \tag{2}$$

Each E_{TA} and T_{TA} value was obtained by blotting both simultaneously, *i.e.* adjacently on the same blotting paper. To facilitate comparison between TA clients, the Sgt2-TPR-C/TA client complex efficiency $E_{complex,TA}$ is normalized by Sgt2-TPR-C/Bos1 complex efficiency $E_{complex,Bos1}$.

$$\% \text{ Complex} = \frac{E_{complex,TA}}{E_{complex,Bos1}} \times 100 \tag{3}$$

Sequence alignments

An alignment of Sgt2-C domains was carried out as follows: all sequences with an annotated N-terminal Sgt2/A dimerization domain (PF16546 [70]), at least one TPR hit (PF00515.27, PF13176.5, PF07719.16, PF13176.5, PF13181.5), and at least 50 residues following the TPR domain were considered family members. Putative C-domains were inferred as all residues following the TPR domain, filtered at 90% sequence identity using CD-HIT [71], and then aligned using MAFFT G-INS-i [72]. Other attempts with a smaller set (therefore more divergent) of

sequences results in an ambiguity in the relative register of H0, H1, H2, and H3 when comparing
Sgt2 with SGTA.

Alignments of Sti1 (DP1/DP2) and STI1 domains were created by pulling all unique domain
structures with annotated STI1 domains from Uniprot. Where present, the human homolog was
selected and then aligned with PROMALS3D [73]. PROMALS3D provides a way of integrating
a variety of costs into the alignment procedure, including 3D structure, secondary structure
predictions, and known homologous positions.

All alignments were visualized using Jalview [74].

Molecular modeling

Putative models for ySgt2-C were generated with I-TASSER, PCONS, Quark, Robetta (*ab*
initio and transform-restrained modes), Phyre2, and RaptorX via their respective web servers
[48,75-78]. The highest scoring model from Quark was then chosen to identify putative TA client
binding sites by rigid-body docking of various transmembrane domains modelled as α-helices (3D-
HM [79]) into the ySgt2-C$_{cons}$ through the Zdock web server [80]. Pairwise distances were
calculated between C$_\beta$ atoms (the closer C$_\alpha$ proton on glycine) using mdtraj [81] . Based on our
disulfide crosslinks, new models were predicted using Robetta in *ab initio* mode specifying C$_\beta$-C$_\beta$
atom distance constraints bounded between 0 and 9 Å.

For hSgt2, using the same set of structure prediction servers above, we were only able to
produce a clear structural model using the Robetta transform-restrained mode. We were also unable
to generate a reliable model by directly using the ySgt2-C model as a template [82]. To crosslink
distance data from ySgt2 as restraints for hSgt2, pair positions were transferred from one protein
to the other via an alignment of Sgt2-C domains (excerpt in Fig. 1A) and ran Robetta *ab initio*.
Also, we grafted the N-terminal loop of ySgt2-C on hSgt2-C with the same set of restraints.

Images were rendered using PyMOL 2.3 (www.pymol.org).

Acknowledgements

We thank D. G. VanderVelde for assistance with NMR data collection; S. Mayo for providing
computing resources; S. Shan, H. J. Cho, Y. Liu, and members of the Clemons lab for support and
discussion. We thank J. Mock and A. M. Thinn for comments on the manuscript. This work was

supported by the National Institutes of Health (NIH) grants GM105385 and GM097572 (to WMC),

NIH/National Research Service Award Training Grant GM07616 (to SMS and MYF), and a

National Science Foundation Graduate Research fellowship Grant 1144469 (to SMS).

References

1. Pieper U, Schlessinger A, Kloppmann E, Chang GA, Chou JJ, Dumont ME, Fox BG, Fromme P, Hendrickson WA, Malkowski MG, et al. (2013) Coordinating the impact of structural genomics on the human α-helical transmembrane proteome. *Nat Struct Mol Biol* **20**: 135–138.

2. Aviram N, Schuldiner M (2017) Targeting and translocation of proteins to the endoplasmic reticulum at a glance. *J Cell Sci* **130**: 4079–4085.

3. Shao S, Hegde RS (2011) Membrane Protein Insertion at the Endoplasmic Reticulum. *Annu Rev Cell Dev Biol* **27**: 25–56.

4. Guna A, Hegde RS (2018) Transmembrane Domain Recognition during Membrane Protein Biogenesis and Quality Control. *Curr Biol* **28**: R498–R511.

5. Kutay U, Hartmann E, Rapoport TA (1993) A class of membrane proteins with a C-terminal anchor. *Trends Cell Biol* **3**: 72–75.

6. Hegde RS, Keenan RJ (2011) Tail-anchored membrane protein insertion into the endoplasmic reticulum. *Nat Rev Mol Cell Biol* **12**: 787–798.

7. Denic V (2012) A portrait of the GET pathway as a surprisingly complicated young man. *Trends Biochem Sci* **37**: 411–417.

8. Wattenberg B, Lithgow T (2001) Targeting of C-Terminal (Tail)-Anchored Proteins: Understanding how Cytoplasmic Activities are Anchored to Intracellular Membranes. *Traffic* **2**: 66–71.

9. Chartron JW, Clemons WM, Suloway CJM (2012) The complex process of GETting tail-anchored membrane proteins to the ER. *Curr Opin Struct Biol* **22**: 217–224.

10. Chio US, Cho H, Shan S-O (2017) Mechanisms of Tail-Anchored Membrane Protein Targeting and Insertion. *Annu Rev Cell Dev Biol* **33**: 417–438.

11. Shao S, Hegde RS (2011) A Calmodulin-Dependent Translocation Pathway for Small Secretory Proteins. *Cell* **147**: 1576–1588.

12. Chartron JW, Gonzalez GM, Clemons WM Jr. (2011) A Structural Model of the Sgt2 Protein and Its Interactions with Chaperones and the Get4/Get5 Complex. *J Biol Chem* **286**: 34325–34334.

13. Wang F, Brown EC, Mak G, Zhuang J, Denic V (2010) A Chaperone Cascade Sorts Proteins for Posttranslational Membrane Insertion into the Endoplasmic Reticulum. *Mol Cell* **40**: 159–171.

14. Simon AC, Simpson PJ, Goldstone RM, Krysztofinska EM, Murray JW, High S, Isaacson RL (2013) Structure of the Sgt2/Get5 complex provides insights into GET-mediated targeting of tail-anchored membrane proteins. *Proc Natl Acad Sci USA* **110**: 1327–1332.

15. Xu Y, Cai M, Yang Y, Huang L, Ye Y (2012) SGTA Recognizes a Noncanonical Ubiquitin-like Domain in the Bag6-Ubl4A-Trc35 Complex to Promote Endoplasmic Reticulum-Associated Degradation. *Cell Rep* **2**: 1633–1644.

16. Wunderley L, Leznicki P, Payapilly A, High S (2014) SGTA regulates the cytosolic quality control of hydrophobic substrates. *J Cell Sci* **127**: 4728–4739.

17. Leznicki P, High S (2012) SGTA antagonizes BAG6-mediated protein triage. *Proc Natl Acad Sci USA* **109**: 19214–19219.

629   18.   Gristick HB, Rao M, Chartron JW, Rome ME, Shan S-O, Clemons WM (2014) Crystal
630   structure of ATP-bound Get3–Get4–Get5 complex reveals regulation of Get3 by Get4.
631   *Nat Struct Mol Biol* **21**: 437–442.

632   19.   Mock J-Y, Chartron JW, Zaslaver M, Xu Y, Ye Y, Clemons WM Jr. (2015) Bag6
633   complex contains a minimal tail-anchor–targeting module and a mock BAG domain. *Proc*
634   *Natl Acad Sci USA* **112**: 106–111.

635   20.   Wang F, Whynot A, Tung M, Denic V (2011) The Mechanism of Tail-Anchored Protein
636   Insertion into the ER Membrane. *Mol Cell* **43**: 738–750.

637   21.   Cho H, Shan S-O (2018) Substrate relay in an Hsp70-cochaperone cascade safeguards
638   tail-anchored membrane protein targeting. *EMBO J* **37**:.

639   22.   Stefer S, Reitz S, Wang F, Wild K, Pang Y-Y, Schwarz D, Bomke J, Hein C, Löhr F,
640   Bernhard F, et al. (2011) Structural basis for tail-anchored membrane protein biogenesis
641   by the Get3-receptor complex. *Science* **333**: 758–762.

642   23.   Rome ME, Chio US, Rao M, Gristick H, Shan S-O (2014) Differential gradients of
643   interaction affinities drive efficient targeting and recycling in the GET pathway. *Proc Natl*
644   *Acad Sci USA* **111**: E4929–E4935.

645   24.   Vilardi F, Lorenz H, Dobberstein B (2011) WRB is the receptor for TRC40/Asna1-
646   mediated insertion of tail-anchored proteins into the ER membrane. *J Cell Sci* **124**: 1301–
647   1307.

648   25.   Yamamoto Y, Sakisaka T (2012) Molecular Machinery for Insertion of Tail-Anchored
649   Membrane Proteins into the Endoplasmic Reticulum Membrane in Mammalian Cells. *Mol*
650   *Cell* **48**: 387–397.

651   26.   Schuldiner M, Metz J, Schmid V, Denic V, Rakwalska M, Schmitt HD, Schwappach B,
652   Weissman JS (2008) The GET Complex Mediates Insertion of Tail-Anchored Proteins
653   into the ER Membrane. *Cell* **134**: 634–645.

654   27.   Kiktev DA, Patterson JC, Müller S, Bariar B, Pan T, Chernoff YO (2012) Regulation of
655   chaperone effects on a yeast prion by cochaperone Sgt2. *Mol Cell Biol* **32**: 4960–4970.

656   28.   Xu Y, Liu Y, Lee J-G, Ye Y (2013) A Ubiquitin-like Domain Recruits an Oligomeric
657   Chaperone to a Retrotranslocation Complex in Endoplasmic Reticulum-associated
658   Degradation. *J Biol Chem* **288**: 18068–18076.

659   29.   Rodrigo-Brenni MC, Gutierrez E, Hegde RS (2014) Cytosolic Quality Control of
660   Mislocalized Proteins Requires RNF126 Recruitment to Bag6. *Mol Cell* **55**: 227–237.

661   30.   Hessa T, Sharma A, Mariappan M, Eshleman HD, Gutierrez E, Hegde RS (2011) Protein
662   targeting and degradation are coupled for elimination of mislocalized proteins. *Nature*
663   **475**: 394–397.

664   31.   Dupzyk A, Williams JM, Bagchi P, Inoue T, Tsai B (2017) SGTA-Dependent Regulation
665   of Hsc70 Promotes Cytosol Entry of Simian Virus 40 from the Endoplasmic Reticulum. *J*
666   *Virol* **91**: 23.

667   32.   Long P, Samnakay P, Jenner P, Rose S (2012) A yeast two-hybrid screen reveals that
668   osteopontin associates with MAP1A and MAP1B in addition to other proteins linked to
669   microtubule stability, apoptosis and protein degradation in the human brain. *Eur J*
670   *Neurosci* **36**: 2733–2742.

671   33.   Trotta AP, Need EF, Selth LA, Chopra S, Pinnock CB, Leach DA, Coetzee GA, Butler
672   LM, Tilley WD, Buchanan G (2013) Knockdown of the cochaperone SGTA results in the
673   suppression of androgen and PI3K/Akt signaling and inhibition of prostate cancer cell
674   proliferation. *Int J Cancer* **133**: 2812–2823.

675   34.   Buchanan G, Ricciardelli C, Harris JM, Prescott J, Yu ZCL, Jia L, Butler LM, Marshall
676   VR, Scher HI, Gerald WL, et al. (2007) Control of Androgen Receptor Signaling in
677   Prostate Cancer by the Cochaperone Small Glutamine Rich Tetratricopeptide Repeat
678   Containing Protein. *Cancer Res* **67**: 10087–10096.
679   35.   Wang H, Zhang Q, Zhu D (2003) hSGT interacts with the N-terminal region of myostatin.
680   *Biochem Biophys Res Commun* **311**: 877–883.
681   36.   Chartron JW, VanderVelde DG, Clemons WM Jr. (2012) Structures of the Sgt2/SGTA
682   Dimerization Domain with the Get5/UBL4A UBL Domain Reveal an Interaction that
683   Forms a Conserved Dynamic Interface. *Cell Rep* **2**: 1620–1632.
684   37.   Liou S-T, Wang C (2005) Small glutamine-rich tetratricopeptide repeat-containing protein
685   is composed of three structural units with distinct functions. *Arch Biochem Biophys* **435**:
686   253–263.
687   38.   Cziepluch C, Kordes E, Poirey R, Grewenig A, Rommelaere J, Jauniaux JC (1998)
688   Identification of a novel cellular TPR-containing protein, SGT, that interacts with the
689   nonstructural protein NS1 of parvovirus H-1. *J Virol* **72**: 4149–4156.
690   39.   Callahan MA, Handley MA, Lee YH, Talbot KJ, Harper JW, Panganiban AT (1998)
691   Functional interaction of human immunodeficiency virus type 1 Vpu and Gag with a
692   novel member of the tetratricopeptide repeat protein family. *J Virol* **72**: 5189–5197.
693   40.   Winnefeld M, Grewenig A, Schn lzer M, Spring H, Knoch TA, Gan EC, Rommelaere J,
694   Cziepluch C (2006) Human SGT interacts with Bag-6/Bat-3/Scythe and cells with reduced
695   levels of either protein display persistence of few misaligned chromosomes and mitotic
696   arrest. *Exp Cell Res* **312**: 2500–2514.
697   41.   Darby JF, Krysztofinska EM, Simpson PJ, Simon AC, Leznicki P, Sriskandarajah N,
698   Bishop DS, Hale LR, Alfano C, Conte MR, et al. (2014) Solution Structure of the SGTA
699   Dimerisation Domain and Investigation of Its Interactions with the Ubiquitin-Like
700   Domains of BAG6 and UBL4A. *PLoS ONE* **9**: e113281–19.
701   42.   Dutta S, Tan Y-J (2008) Structural and functional characterization of human SGT and its
702   interaction with Vpu of the human immunodeficiency virus type 1. *Biochemistry* **47**:
703   10123–10131.
704   43.   Leznicki P, Korac-Prlic J, Kliza K, Husnjak K, Nyathi Y, Dikic I, High S (2015) Binding
705   of SGTA to Rpn13 selectively modulates protein quality control. *J Cell Sci* **128**: 3187–
706   3196.
707   44.   Waheed AA, MacDonald S, Khan M, Mounts M, Swiderski M, Xu Y, Ye Y, Freed EO
708   (2016) The Vpu-interacting Protein SGTA Regulates Expression of a Non- glycosylated
709   Tetherin Species. *Sci Rep* **6**: 1–14.
710   45.   Luo P, Baldwin RL (1997) Mechanism of helix induction by trifluoroethanol: a
711   framework for extrapolating the helix-forming properties of peptides from
712   trifluoroethanol/water mixtures back to water. *Biochemistry* **36**: 8413–8421.
713   46.   Dyson HJ, Wright PE (2004) Unfolded Proteins and Protein Folding Studied by NMR.
714   *Chem Rev* **104**: 3607–3622.
715   47.   Martínez-Lumbreras S, Krysztofinska EM, Thapaliya A, Spilotros A, Matak-Vinkovic D,
716   Salvadori E, Roboti P, Nyathi Y, Muench JH, Roessler MM, et al. (2018) Structural
717   complexity of the co-chaperone SGTA: a conserved C-terminal region is implicated in
718   dimerization and substrate quality control. *BMC Biol* **16**: 76.
719   48.   Xu D, Zhang Y (2012) Ab initio protein structure assembly using continuous structure
720   fragments and optimized knowledge-based force field. *Proteins* **80**: 1715–1735.

49. Chio US, Chung S, Weiss S, Shan S-O (2019) A Chaperone Lid Ensures Efficient and
Privileged Client Transfer during Tail-Anchored Protein Targeting. *Cell Rep* **26**: 37–
44.e37.
50. Shao S, Rodrigo-Brenni MC, Kivlen MH, Hegde RS (2017) Mechanistic basis for a
molecular triage reaction. *Science* **355**: 298–302.
51. Chun E, Thompson AA, Liu W, Roth CB, Griffith MT, Katritch V, Kunken J, Xu F,
Cherezov V, Hanson MA, et al. (2012) Fusion Partner Toolchest for the Stabilization and
Crystallization of G Protein-Coupled Receptors. *Structure* **20**: 967–976.
52. Letunic I, Doerks T, Bork P (2015) SMART: recent updates, new developments and status
in 2015. *Nucleic Acids Res* **43**: D257–D260.
53. Schmid AB, Lagleder S, Gräwert MA, Röhl A, Hagn F, Wandinger SK, Cox MB,
Demmer O, Richter K, Groll M, et al. (2012) The architecture of functional modules in the
Hsp90 co-chaperone Sti1/Hop. *EMBO J* **31**: 1506–1517.
54. Li Z, Hartl FU, Bracher A (2013) Structure and function of Hip, an attenuator of the
Hsp70 chaperone cycle. *Nat Struct Mol Biol* **20**: 929–935.
55. Nelson GM, Huffman H, Smith DF (2003) Comparison of the carboxy-terminal DP-repeat
region in teh co-chaperones Hop and Hip. *EMBO J* **8**: 125–133.
56. Rao M, Okreglak V, Chio US, Cho H, Walter P, Shan S-O (2016) Multiple selection
filters ensure accurate tail-anchored membrane protein targeting. *eLife* **5**: e21301–e21324.
57. Pace CN, Scholtz JM (1998) A Helix Propensity Scale Based on Experimental Studies of
Peptides and Proteins. *Biophys J* **75**: 422–427.
58. Röhl A, Wengler D, Madl T, Lagleder S, Tippel F, Herrmann M, Hendrix J, Richter K,
Hack G, Schmid AB, et al. (2015) Hsp90 regulates the dynamics of its cochaperone Sti1
and the transfer of Hsp70 between modules. *Nat Commun* **6**: 6655.
59. Coto ALS, Seraphim TV, Batista FAH, Dores-Silva PR, Barranco ABF, Teixeira FR,
Gava LM, Borges JC (2018) Structural and functional studies of the Leishmania
braziliensis SGT co-chaperone indicate that it shares structural features with HIP and can
interact with both Hsp90 and Hsp70 with similar affinities. *Int J Biol Macromol* **118**: 693–
706.
60. Voorhees RM, Hegde RS (2015) Structures of the scanning and engaged states of the
mammalian SRP-ribosome complex. *eLife* **4**: 1485–21.
61. Keenan RJ, Freymann DM, Walter P, Stroud RM (1998) Crystal Structure of the Signal
Sequence Binding Subunit of the Signal Recognition Particle. *Cell* **94**: 181–191.
62. Clemons WM, Gowda K, Black SD, Zwieb C, Ramakrishnan V (1999) Crystal structure
of the conserved subdomain of human protein SRP54M at 2.1 A resolution: evidence for
the mechanism of signal peptide binding. *J Mol Biol* **292**: 697–705.
63. Tidow H, Nissen P (2013) Structural diversity of calmodulin binding to its target sites.
FEBS J **280**: 5551–5565.
64. Guna A, Volkmar N, Christianson JC, Hegde RS (2018) The ER membrane protein
complex is a transmembrane domain insertase. *Science* **359**: 470–473.
65. Suloway CJ, Rome ME, Clemons WM (2011) Tail-anchor targeting by a Get3 tetramer:
the structure of an archaeal homologue. *EMBO J* **31**: 707–719.
66. Chu R, Takei J, Knowlton JR, Andrykovitch M, Pei W, Kajava AV, Steinbach PJ, Ji X,
Bai Y (2002) Redesign of a Four-helix Bundle Protein by Phage Display Coupled with
Proteolysis and Structural Characterization by NMR and X-ray Crystallography. *J Mol*
Biol **323**: 253–262.

67. Studier FW (2005) Protein production by auto-induction in high-density shaking cultures.
Protein Express Purif **41**: 207–234.
68. Micsonai A, Wien F, Kernya L, Lee Y-H, Goto Y, Réfrégiers M, Kardos J (2015)
Accurate secondary structure prediction and fold recognition for circular dichroism
spectroscopy. *Proc Natl Acad Sci USA* **112**: E3095–E3103.
69. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**: 671–675.
70. Finn RD, Coggill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M,
Qureshi M, Sangrador-Vegas A, et al. (2016) The Pfam protein families database: towards
a more sustainable future. *Nucleic Acids Res* **44**: D279–D285.
71. Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.
72. Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. *Mol Biol Evol* **30**: 772–780.
73. Pei J, Kim B-H, Grishin NV (2008) PROMALS3D: a tool for multiple protein sequence
and structure alignments. *Nucleic Acids Res* **36**: 2295–2300.
74. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ (2009) Jalview Version
2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**:
1189–1191.
75. Yang J, Yan R, Roy A, Xu D, Poisson J, Zhang Y (2015) The I-TASSER Suite: protein
structure and function prediction. *Nat Methods* **12**: 7–8.
76. Wallner B (2006) Identification of correct regions in protein models using structural,
alignment, and consensus information. *Prot Sci* **15**: 900–913.
77. Bradley P, Misura KM, Baker D (2005) Towards high-resolution de novo structure
prediction for small proteins. *Science* **309**: 1868–1871.
78. Yang J, Anishchenko I, Park H, Peng Z, Ovchinnikov S, Baker D (2019) Improved
protein structure prediction using predicted inter-residue orientations. *bioRxiv* **13**:
e1005324–e1005324.
79. Reißer S, Strandberg E, Steinbrecher T, Ulrich AS (2014) 3D Hydrophobic Moment
Vectors as a Tool to Characterize the Surface Polarity of Amphiphilic Peptides. *Biophys J*
106: 2385–2394.
80. Pierce BG, Wiehe K, Hwang H, Kim BH, Vreven T, Weng Z (2014) ZDOCK server:
interactive docking prediction of protein-protein complexes and symmetric multimers.
Bioinformatics **30**: 1771–1773.
81. McGibbon RT, Beauchamp KA, Harrigan MP, Klein C, Swails JM, Hernández CX,
Schwantes CR, Wang L-P, Lane TJ, Pande VS (2015) MDTraj: A Modern Open Library
for the Analysis of Molecular Dynamics Trajectories. *Biophys J* **109**: 1528–1532.
82. Webb B, Sali A (2002) *Comparative Protein Structure Modeling Using MODELLER*.
John Wiley & Sons, Inc., Hoboken, NJ, USA.
83. Drozdetskiy A, Cole C, Procter J, Barton GJ (2015) JPred4: a protein secondary structure
prediction server. *Nucleic Acids Res* **43**: W389–W394.
84. Schägger H (2006) Tricine–SDS-PAGE. *Nat Protoc* **1**: 16–22.
85. Kyte J, Doolittle RF (1982) A simple method for displaying the hydropathic character of a
protein. *J Mol Biol* **157**: 105–132.
86. Fernandez-Patron C, Hardy E, Sosa A, Seoane J, Castellanos L (1995) Double staining of
coomassie blue-stained polyacrylamide gels by imidazole-sodium dodecyl sulfate-zinc

reverse staining: sensitive detection of coomassie blue-undetected proteins. *Anal Biochem*
224: 263–269.

87. Bateman A, Martin MJ, O'Donovan C, Magrane M, Alpi E, Antunes R, Bely B, Bingley
M, Bonilla C, Britto R, et al. (2017) UniProt: the universal protein knowledgebase.
Nucleic Acids Res **45**: D158–D169.

Figures

Fig. 1. Structural characteristics of free Sgt2 C-domain. (*A*) *Top*, Schematic of the domain

organization of Sgt2. *Below,* representative sequences from a large-scale multiple sequence

alignment of the C domain: fungal Sgt2 from *S. cerevisiae*, *S. pombe*, and *C. thermophilum* and

metazoan Sgt2 from *C. savignyi*, *X. laevis*, and *H. sapiens*. Protease susceptible sites on ySgt2-C

identified by mass spectrometry are indicated by red arrowheads. Predicted helices of ySgt2 (blue)

and hSgt2 (orange) by Jpred [83] and/or structure prediction are shown. Blue/orange color scheme

for ySgt2/hSgt2 is used throughout the text. Residues noted in the text are highlighted by an

asterisk. (*B*) Overlay of size-exclusion chromatography traces of ySgt2-C (blue line), hSgt2-C

(orange line), ySgt2-TPR (blue dash), and hSgt2-TPR (orange dash). Traces are measured at 214

nm, baseline-corrected and normalized to the same peak height. (*C*) Far UV CD spectrum of 10

μM of purified ySgt2-C (blue) and hSgt2-C (orange) at RT with secondary structure decomposition

from BestSel [68]. (*D*) ^{1}H-^{15}N HSQC spectrum of ySgt2-C at 25°C. The displayed chemical shift

window encompasses all N-H resonances from both backbone and side chains. The range of

backbone amide protons, excluding possible side-chain NH_2 of Asn/Gln, is indicated by pairs of

red dashed lines. (*E*) As in *D* for hSgt2-C at 25°C.

Figure 1

Fig. 2. The minimal binding region of Sgt2 for TA client binding. (*A*) Diagram of the protein truncations tested for TA client binding that include the TPR-C domain, C-domain (C), C_{cons}, and $C_{cons} \Delta H0$ ($\Delta H0$) from ySgt2 and hSgt2. The residues corresponding to each domain are indicated, and grey blocks highlight the C_{cons} region. *(B)* Schematic of capture experiments of MBP-tagged Sbh1 (MBP-Sbh1) by Sgt2 variants. After co-expression, cell pellets are lysed and NTA-Ni$_{2+}$ is used to capture his-tagged Sgt2-TPR-C. (*C*) Tris-Tricine-SDS-PAGE gel [84] of co-expressed and purified MBP-Sbh1 and his-tagged Sgt2 truncations visualized with Coomassie Blue staining. (*D*) Helical wheel diagrams of predicted helices (see Fig. 1*A*) in the C_{cons} domain of ySgt2 and hSgt2. Residues are colored by the Kyte and Doolittle hydrophobicity scale [85]. (*E*) All of the hydrophobic residues (L, I, F, and M) in a predicted helix (H0, H1, etc.) are replaced with alanines and tested for the ability to capture MBP-Sbh1. Protein levels were quantified by Coomassie staining. Relative binding efficiency of MBP-Sbh1 by ySgt2 C-domain (ySgt2-C) variants was calculated relative to total amount of ySgt2-C captured (MBP-Sbh1/Sgt2-C) then normalized to the wild-type ySgt2-C. Experiments were performed 3-4 times and the standard deviations are presented. Total expression levels of the MBP-Sbh1 were similar across experiments as visualized by immunoblotting (IB) of the cell lysate. (*F)* As in *E* but for hSgt2.

Figure 2

Fig 3. A structural model for Sgt2-C$_{cons}$. *(A)* The top 10 models of the ySgt2-C$_{cons}$ generated by

the template-free algorithm Quark [48] are overlaid with the highest scoring model in solid.

Models are color-ramped from N- (blue) to C-terminus (red). *(B)* A model of ySgt2-C$_{cons}$ (surface

colored by Kyte-Doolittle hydrophobicity) bound to a TMD (purple helix) generated by rigid-body

docking through Zdock [80]. The darker purple corresponds to an 11 residue stretch. *(C)* The entire

ySgt2-C from the highest scoring model from Quark (C$_{cons}$ in rainbow with the rest in grey)

highlighting H0 and the rest of the flexible termini that vary considerably across models.

Figure 3

Fig 4. Validating the structural model with disulfide bond formation. Variants of His-ySgt2-TPR-C (WT or cysteine double mutants) were co-expressed with the artificial TA client, cMyc-BRIL-11[L8]. After lysis, ySgt2-TPR-C proteins were purified, oxidized, then digested by Glu-C protease and analyzed by gel either in non-reducing or reducing buffer. (*A*) Cα ribbon of ySgt2-C$_{cons}$ color-ramped with various pairs of cysteines highlighted. Scissors indicate protease cleavage sites resulting in fragments less than 3 kDa in size. (*B*) Tris-Glycine-SDS-PAGE gel visualized by imidazole-SDS-zinc stain [86]. For the WT (cys-free) no significant difference was found between samples in non-reducing vs. reducing conditions. All close residue pairs (A272/L327, I286/M323, M289/A319, and M289/N322) show peptide fragments (higher MW) sensitive to the reducing agent and indicate disulfide bond formation (indicated by arrow). A cysteine pair (N285/G329) predicted to be far apart by the model does not result in the higher MW species. (*C*) A schematic of the transfer of Bos1$_{BPA}$ from Ssa1 to full-length ySgt2 to demonstrate the double cysteine mutants are still functional. (*D*) A western blot visualizing cross-linked ySgt2-Bos1$_{BPA}$ complexes. All samples tested, WT, N285C/G329C, A272C/L327C, I286C/M323C, M289C/A319C, and M289C/N322C, had a higher molecular weight appear after the addition of ySgt2 which corresponds to the size of the cross-linked complex. As described in Cho et al., a band for cross-linked Ssa1-Bos1$_{BPA}$ complexes was not observed. (*E*) Cβ-Cβ distances between the residues mutated to cysteines based on various models predicted by the Quark, I-TASSER, PCONS, and Robetta. Cysteine pairs that are 9Å or less apart are colored in orange and are expected to be close enough to form disulfide bonds. Where all five pair distances are consistent with the experiment (4 near and 1 far), the row is shaded in grey. (*F*) Tris-Glycine SDS-PAGE gel probing the flexibility of ySgt2-C$_{cons}$. All new pairs (K288/A319, M290/A319, M289/P318, M289/L320) show peptide fragments sensitive to the reducing agent (indicated by arrow). The range of distances of the eight closest possible rotamer pairs is annotated below. The cysteine pair (N285/G329) shown to be far apart by the model does have a faint higher molecular weight band.

Figure 4

Fig. 5. Comparison of Sti1 domains and the Sgt2-C$_{cons}$ model. (*A*) Multiple sequence alignment

of Sgt2-C with STI1 domains (DP1, DP2) from STI1/Hop homologs. Helices are shown based on

the Sgt2-C$_{cons}$ model and the *Sc*Sti1-DP1/2 structures. Species for representative sequences are

from *S. cerevisiae (Scer)*, *S. pombe* (*Spom*), *C. thermophilum* (*Cthe*), *C. savignyi* (*Csav*), and *H.*

sapiens (*Hsap*). (*B*) Cα ribbon of *Sc*Sgt2-C$_{cons}$ color-ramped with large hydrophobic sidechains

shown as grey sticks (sulfurs in yellow). (*C*) Similar to *B* for the solution NMR structure of Sti1-

DP2$_{526-582}$ (PDBID: 2LLW) [53]. (*D*) Superposition of the Sgt2-C$_{cons}$ (blue) and Sti1-DP2$_{526-582}$

(red) drawn as cartoons.

Figure 5

Fig. 6. Effects on TA client binding of charge mutations to the putative hydrophobic groove of Sgt2-C$_{cons}$. For these experiments, individual point mutations are introduced into Sgt2-C and tested for their ability to capture Sbh1 quantified as in Figure 2D. (A) For ySgt2-C, a schematic and cartoon model are provided highlighting the helices and sites of individual point mutants both color-ramped for direct comparison. For the cartoon, the docked TMD is shown in purple. Binding of MBP-Sbh1 to his-tagged ySgt2-C and mutants were examined as in Figure 2D. Lanes for mutated residues are labeled in the same color as the schematic *(B)* Same analysis as in *A* for hSgt2-C. In addition, double point mutants are included.

Figure 6

Fig. 7. Minimal requirements for client recognition by Sgt2. (*A*) Schematic of model TA clients.

Quantification of complex formation is calculated and normalized to that of complexes with Bos1-

TMD, here defined as relative binding efficiency. (*B*) Complex formation of ySgt2 (blue) and hSgt2

(orange) with the TA client Bos1TMD and several artificial TA clients noted x[Ly], where x denotes

the length of the TMD and y denotes the number of leucines in the TMD. The helical wheel

diagrams of the TMD of TA clients here and for subsequent panels with leucines colored in dark

orange, alanines colored in pale orange, and tryptophans colored in grey. (*C*) Complex formation

of ySgt2-TPR-C and hSgt2-TPR-C with artificial TA clients with TMDs of length 11 and

increasing numbers of leucine. (*D*) Comparison of complex formation of ySgt2-TPR-C and hSgt2-

TPR-C with artificial TA clients of the same lengths and hydrophobicities but differences in the

distribution of leucines, i.e. clustered (solid line) vs distributed (dotted line).

Figure 7

Fig. 8. Various domain structures of STI1 and other helical-hand containing proteins. (*A*)
The domain architectures of proteins with a STI1 domain were obtained initially from InterPro
[87] and then adjusted as discussed in the text. Each domain within a protein is colored relative to
the key. (*B*) Structural comparison of various hydrophobic-binding helical-hand protein complexes.
For each figure only relevant domains are included. Upper row, color-ramped cartoon
representation with bound helices in purple. Lower row, accessible surface of each protein colored
by hydrophobicity again with docked helical clients in purple. In order, the predicted complex of
ySgt2-C_{cons} and *Sc*Sbh2-TMD, DP1 domain from yeast Sti1 with N-terminus containing H0 in
grey (*Sc*Sti1-DP1)(PDBID: 2LLV), human calmodulin (*Hs*CALM2) bound to a hydrophobic
domain of calcineurin (*Hs*PPP3CA) (PDBID: 2JZI), and M domain of SRP54 from *Oryctolagus*
cuniculus (*Oc*SRP54-M) and the signal sequence of human transferrin receptor (*Hs*TR-TMD)
(PDBID: 3JAJ).
933

A

Dimerization

TPR Motif HSP70/90 binding

Client Binding

ySgt2,hSgt2 — N-dimerization · 70 · TPR · 102 · 203 · STI1 · 267 · 331 · 346

Sti1/Hop, Sti1 — 127 · 184 · 223 · 348 · 359 · 461 · 484 · 538 · 543

Hip — 114 · 215 · 311 · 365 · 369

B

TA-protein targeting
ySgt2-C$_{cons}$
ScSbh1-TMD

Co-chaperone
ScSti1-DP1

C

Ca^{2+}/peptide binding
HsCALM2
HsPPP3CA

Co-translational targeting
OcSRP54-M
HsTR-TMD

Kyte & Doolittle
-4.5 (Arg)   4.5 (Ile)

Figure 8

Fig. S1. Biophysical characterization of the Sgt2-C$_{cons}$ domain. (*A*) CD spectra as in Fig. 1*C*

for the conserved C-terminal domains of ySgt2 (blue) and hSgt2 (orange). NMR spectra as in Fig.

1*D & E* for ySgt2-C$_{cons}$ (*B*, blue) and hSgt2-C$_{cons}$ (*C*, orange).

Fig. S2. Structural models across prediction methods. (*A*) Predictions from Quark, I-TASSER,

Pcons, Phyre2, RaptorX, and Robetta. Methods produce between 5 and 10 models. (*B*) Robetta

provides a residue-wise estimated error in Angstroms; this is shown below the corresponding

models with a grey bar indicating the C_{cons} region.

Figure S2

Fig. S3. Cysteine mutants are capable of binding to TA clients. (*A*) Schematic showing how
his-tagged ySgt2-TPR-C and double cysteine mutant constructs were coexpressed with the TA
client 11[L8], and complexes were purified by nickel affinity chromatography. (*B*) A SDS-PAGE
gel of the elution fractions demonstrates that 11[L8] was present in the elution suggesting double
cysteine mutations do not affect client binding. (*C*) An anti-cMyc western blot of the fractions
represented in the SDS-PAGE gel also demonstrates that 11[L8] was present in all elutions.
951

Figure S3

Fig. S4. Distance restraints lead to improved ySgt2-C and suggestive hSgt2-C models. (*A*)

Prediction of ySgt2-C using distances from in vitro crosslinking. (*B*) Cβ-Cβ distances between

residues probed by *in vitro* disulfide crosslinking for each ySgt2 model. Distances 9 Å or less are

colored orange. For models where all distances correspond (4 near and 1 far), the row is shaded

grey. (*C*) Models for hSgt2-C using restraints, adding a N-terminal loop, and via the new Robetta

TR method.

A ySgt2, using crosslink distances

	N285-G329	A272-L327	I286-M323	M289-A319	M289-N322	
1	18.6	4.6	6.8	7.3	12.4	C_β-C_β Distance (Å)
2	10.3	5.6	5.5	9.0	7.4	
3	12.7	6.7	4.8	5.0	8.0	
4	9.6	8.2	7.7	9.0	6.7	9 Å or less
5	8.5	8.0	7.2	7.2	5.1	

C

hSgt2, using crosslink inferred distances

hSgt2, using crosslink inferred distances with N-loop

hSgt2, Robetta TR

Figure S4