Title: Glycolysis upregulation is neuroprotective as a compensatory mechanism in ALS

Authors:

Ernesto Manzo[1], Ileana Lorenzini[4], Dianne Barrameda[1], Abigail G. O'Conner[1], Jordan M.

Barrows[1], Alexander Starr[4], Tina Kovalik[4], Benjamin E. Rabichow[4], Erik M. Lehmkuhl[1], Dakotah

D. Shreiner[1], Archi Joardar[1], Jean-Charles Liévens[5], Robert Bowser[4], Rita Sattler[4], Daniela C.

Zarnescu[1,2,3, *]

Affiliations:

1 Department of Molecular and Cellular Biology,

2 Department of Neuroscience

3 Department of Neurobiology, University of Arizona, Tucson, AZ 85721, USA

4 Neurology, Barrow Neurological Institute, Phoenix, AZ

5 Université de Montpellier, MMDN U1198, CC105, Place Eugène Bataillon, 34095 Montpellier

Cedex 5, France.

* corresponding author

Abstract

Amyotrophic Lateral Sclerosis (ALS), is a fatal neurodegenerative disorder, with TDP-43 inclusions as a major pathological hallmark. Using a *Drosophila* model of TDP-43 proteinopathy we found significant alterations in glucose metabolism including increased pyruvate, suggesting that modulating glycolysis may be neuroprotective. Indeed, a high sugar diet improves locomotor and lifespan defects caused by TDP-43 proteinopathy in motor neurons or glia, but not muscle, suggesting that metabolic dysregulation occurs in the nervous system. Overexpressing human glucose transporter GLUT-3 in motor neurons mitigates TDP-43 dependent defects in synaptic vesicle recycling and improves locomotion. Furthermore, *PFK* mRNA, a key indicator of glycolysis, is upregulated in flies and patient derived iPSC motor neurons with TDP-43 pathology. Surprisingly, *PFK* overexpression rescues TDP-43 induced locomotor deficits. These findings from multiple ALS models show that mechanistically, glycolysis is upregulated in degenerating motor neurons as a compensatory mechanism and suggest that increased glucose availability is protective.

Introduction

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease affecting upper and lower motor neurons resulting in progressive muscle weakness and eventual death (Hardiman et al., 2017). Cellular metabolic defects arise early, and contribute to the clinical manifestation of systemic defects that include weight loss, dyslipidemia, and hypermetabolism (reviewed in (Dupuis et al., 2011; Joardar et al., 2017). However, it is unclear how metabolic defects relate, or may contribute to motor neuron degeneration.

Observations from both patients and animal models have begun to elucidate the mechanisms that contribute to metabolic defects in ALS. Interestingly, the human brain relies on glucose as its major energy source, and consumes ~20% of the body's glucose derived energy (Mergenthaler et al., 2013). Glucose metabolism defects have been previously reported in the frontal lobe and the cortex of ALS patients (Ludolph et al., 1992), and in mutant SOD1 mouse models of ALS (Browne et al., 2006; Miyazaki et al., 2012). Increasing glucose availability has been shown to reduce protein misfolding and delay neuronal degeneration in *C. elegans* models of neurodegeneration (Tauffenberger et al., 2012). In a recent pilot study, a high caloric diet based on high carbohydrate content has been shown to be well tolerated by patients, and reduce serious adverse events in ALS patients (Wills et al., 2014b). Taken together, these findings suggest that although defective glucose metabolism is still poorly understood in ALS, there lies a great opportunity to better understand its relationship to disease and explore its potential as a therapeutic avenue.

Our lab has developed a *Drosophila* model of ALS based on overexpression of human TDP-43 that recapitulates multiple disease aspects including cytoplasmic aggregates, neuromuscular junction (NMJ) abnormalities, lifespan, and locomotor defects (Estes et al., 2011, 2013). Importantly, the large majority of ALS patients (>97%) harbor TDP-43 cytoplasmic aggregates regardless of etiology, highlighting the importance of understanding TDP-43 toxicity mechanisms (Ling et al., 2013; Neumann et al., 2006). Here, we report that in a *Drosophila* model of TDP-43 proteinopathy, glycolytic metabolites and genes are altered, and are consistent with increased glucose consumption. Moreover, key genes responsible for driving glucose metabolism

are upregulated in both patient derived induced pluripotent stem cell (iPSC) motor neurons and

postmortem patient spinal cord tissue. In a fly model of TDP-43 proteinopathy, increased dietary

glucose improves locomotor function and increases lifespan when TDP-43 is expressed in the

central nervous system but not muscles. Additionally, genetic over-expression of human glucose

transporters improves locomotor function, mitigates neuromuscular junction defects, and

improves lifespan in a variant dependent manner. Finally, motor neuron overexpression of *PFK*,

the rate limiting enzyme in glycolysis, improves locomotor function, suggesting that upregulation

of glycolysis is neuroprotective through a compensatory mechanism in ALS. Together, our results

show that increased glucose availability protects motor neurons and improves overall outcome in

models of TDP-43 proteinopathy.

Results

Glycolysis and pentose phosphate pathways are altered by TDP-43 expression in motor

neurons

To decipher metabolic alterations in ALS, we employed a *Drosophila* model of TDP-43

proteinopathy that recapitulates key features of the human disease including locomotor

dysfunction and reduced survival (Estes et al., 2011; Estes et al., 2013). We conducted

metabolomic profiling in whole third instar larvae expressing TDP-43 in motor neurons (D42>TDP-

43^{WT} or D42>TDP-43^{G298S}) and controls (D42>w^{1118}). These experiments showed significantly

increased phosphoenolpyruvate (PEP) and pyruvate in both TDP-43^{WT} and TDP-43^{G298S}

suggesting an increase in glucose metabolism (PEP, P_{value} = 0.034 and pyruvate, P_{value} = 0.007

for TDP-43^{WT}; PEP, P_{value} =0.002 and pyruvate, P_{value} = 0.009 for TDP-43^{G298S}; see Fig.1 and

Supplemental file 1). Interestingly, pyruvate does not appear increased when endogenous TDP-

43 is knocked down by RNAi in motor neurons suggesting that this alteration is caused by TDP-

43 proteinopathy (Figure 1 - supplemental 1). In larvae expressing TDP-43^{G298S}, we also found a

decrease in ribulose/xylulose 5-phosphate (P_{value} = 0.011) in conjunction with increased

sedoheptulose 7-phosphate (P_{value} = 0.008) suggesting that increased glycolytic input into the

Pentose Phosphate Pathway occurs in a variant dependent manner (see Fig. 1, Supplemental

file 1). Taken together these findings suggest that glucose metabolism is altered in the context of

TDP-43 proteinopathy, consistent with previous reports of metabolite alterations such as

increased pyruvate in plasma isolated from ALS patients (Lawton et al., 2012).

Key glycolysis and pentose phosphate pathway enzymes are upregulated by TDP-43

expression in motor neurons

An increase in glycolytic metabolites suggests a possible increase in glycolysis. To test this

hypothesis, we profiled the expression of multiple enzymes known to regulate flux through

glycolysis and the Pentose Phosphate Pathway. Phosphofructokinase-1 (*PFK1*; *PFK* in

Drosophila) levels are considered to be rate-limiting and control the rate of glycolysis, thus *PFK*

expression provides a reliable determinant of flux for glucose breakdown (Tanner et al., 2018).

To measure the levels of *PFK* in the context of TDP-43 proteinopathy we performed qPCR

in ventral nerve chords dissected from third instar larvae expressing TDP-43^{WT} or TDP-43^{G298S} in

motor neurons (D42>TDP-43) and w^{1118} controls. These experiments showed that *PFK* transcript

levels are increased in the context of TDP-43^{WT} or TDP-43^{G298S}, albeit statistical significance was

only achieved in the mutant TDP-43 (37% increase, P$_{value}$ = 0.06 for TDP-43^{WT} and 59% increase,

P$_{value}$ = 0.014 for TDP-43^{G298S}; see Fig. 2A). These results suggest that flux through glycolysis is

significantly increased in motor neurons in the context of TDP-43^{G298S}.

Finally, glucose-6-phosphate dehydrogenase (*G6PD*; *Zw* in *Drosophila*) was selected due

to its role as the commitment step for the pentose phosphate pathway (PPP). *Zw* transcript levels

are increased specifically in TDP-43^{G298S} (37% increase, P$_{value}$ = 0.012, see Fig. 2B) but not TDP^{WT},

consistent with increased glycolytic influx through the PPP in the context of mutant TDP-43.

PFK **transcript is upregulated in human tissue**

To confirm that the changes in our *Drosophila* model of ALS have clinical relevance, we assayed

the transcript levels of *PFK* in human tissues and patient derived iPSC motor neurons. Humans

have three isoforms of *PFK*, namely *PFKM*, *PFKL*, and *PFKP*. Although all are present in the

brain, *PFKL* is expressed at lower levels (Human Brain Transcriptome: http://hbatlas.org/) thus

we focused on *PFKP* and *PFKM*. For human tissues we used spinal cords isolated from ALS

patients with TDP-43 proteinopathy and controls lacking neurological defects (see Supplemental

file 2 for demographic information). qPCR experiments show that both *PFKP* and *PFKM* levels

are significantly increased in human spinal cords (*PFKP*: 80% increase, P_{value} <0.001; *PFKM*: 55%

increase, P_{value}=0.014, Fig. 2C). iPSC motor neurons were derived from a patient harboring the

TDP-43^{G298S} mutation and differentiated as previously described (Coyne et al., 2017). qPCR

profiling from iPSC derived motor neurons differentiated for 69-77 days (three independent

differentiations, see Supplemental file 3) showed a 58% increase in *PFKP* (P_{value}=0.010; see Fig.

2D) and a 45% increase in *PFKM* (P_{value}=0.036). These experiments show that glycolysis is also

upregulated in patient spinal cords and motor neurons consistent with our findings in the fly model.

Similarly, *G6PD* transcript levels are increased by 49% in spinal cords of ALS patients

(P_{value}=0.015; see Fig. 2C) and by 52% in patient derived motor neurons (P_{value}=0.012: see Fig.

2D).

TDP-43 proteinopathy alters the capacity of motor neurons to import glucose

Our findings that TDP-43 expression in motor neurons alters glycolysis could be explained by an

effect on glucose flux. To evaluate the levels of glucose in motor neurons, in the context of TDP-

43 proteinopathy, we used a genetically encoded, FRET based, glucose sensor, FLII12Pglu-

700μδ6 (Fehr et al., 2003; Takanaga et al., 2008). The sensor comprises a bacterial glucose-

sensing domain and a pair of cyan and yellow fluorescent proteins that upon glucose binding,

undergoes a conformational change and allows FRET to occur (Fig. 3A). This genetically encoded

sensor has been previously used in conjunction with the GAL-UAS system in *Drosophila* to detect

glucose flux and distribution in various cell types within the fly ventral nerve cord (Volkenhoff et

al., 2018).

Upon stimulation with glucose, neurons expressing the glucose sensor alone or together

with TDP-43 showed a small but significant increase in the FRET/CFP ratio (Figs. 3B, C). TDP-

43^{WT} expressing neurons showed no significant difference in the FRET/CFP signal compared to

controls (Figs. 3B, C, P_{value} <0.99). In contrast, TDP-43^{G298S} expressing neurons showed a

significant increase in the FRET/CFP signal compared to controls (Figs. 3B, C, P_{value} <0.001).

These results suggest that glucose flux is specifically enhanced in TDP-43^{G298S} expressing

neurons, and further substantiate our findings that glycolysis and the pentose phosphate pathway

are significantly increased in the context of TDP-43 proteinopathy.

A high glucose diet mitigates neuronal and glial induced TDP-43 toxicity

The alterations in glucose metabolism we have uncovered here could be a direct consequence

of TDP-43 proteinopathy, or alternatively, could reflect a coping mechanism. To distinguish

between these two scenarios, we fed TDP-43 expressing flies a high glucose diet. As we have

previously shown, the neuronal or glial expression of TDP-43^{WT} or TDP-43^{G298S} caused locomotor

defects as indicated by increased larval turning time compared to controls (Estes et al., 2011;

Estes et al., 2013). Of note, a high glucose diet increased the larval turning time of control larvae,

consistent with previous reports that high sugar intake induces insulin resistance and obesity in

flies (Musselman et al., 2011; Na et al., 2013). In contrast, a high glucose diet improved the larval

locomotor defect caused by either neuronal, or glial expression of TDP-43 (Fig. 4A and Fig. 4 –

figure supplement 1). Interestingly, we found that increasing the concentration of glucose five-fold

or higher rescued locomotor phenotypes caused by TDP-43 while no improvement was observed

when glucose is only increased by two-fold (Fig. 4 – figure supplement 1). Next, we examined the

effect of glucose on lifespan and found that the reduced survival caused by neuronal or glial

expression of TDP-43^{WT} or TDP-43^{G298S} was also significantly improved by feeding *Drosophila* a

high sugar (HS) versus a regular sugar (RS) diet (23.5 days for TDP-43^{WT} on RF and 53 days for

TDP-43^{WT} on HS, P_{value} = <0.0001; 14.5 days for TDP-43^{G298S} on RS and 41 days for TDP-43^{G298S}

HS, P_{value} = <0.0001 ; Fig. 4B, Fig. 4 – figure supplement 2). Together, these results suggest that

while a high sugar diet is detrimental to control flies, it improves key hallmarks of motor neuron

disease including locomotor dysfunction and reduced survival in ALS flies.

Overexpression of human glucose transporters mitigate TDP-43 dependent phenotypes in

neurons or glia

Our results on a high sugar diet suggest that increased glucose availability is protective in our fly

model of ALS. However, because feeding can cause systemic effects, and to further probe the

role of glucose in our *Drosophila* model of ALS*,* we co-expressed TDP-43 and the human glucose

transporter GLUT-3 in motor neurons or glia. GLUT-3 is the primary glucose transporter of

mammalian neurons, and has a high capacity for glucose (Ferreira et al., 2011). Co-expression

of GLUT-3 and TDP-43^{WT} or TDP-43^{G298S}, in either motor neurons or glia cells, improved the

locomotor deficits caused by TDP-43 expression alone (Fig. 4C). Moreover, GLUT-3 expression

in neurons improved the locomotor ability of adult flies expressing TDP-43^{WT} or TDP-43^{G298S} (as

measured by negative geotaxis assays, Fig. 4 – figure supplement 3). When using the standard

log-rank test for analyzing the survival data, lifespan was significantly increased when GLUT-3

was co-expressed with TDP-43^{G298S} in motor neurons, but not glial cells (18 days for TDP-43^{G298S}

and 30 days for TDP-43^{G298S}, P$_{value}$ = <0.0001). However, when using the Mantel-Cox test, which

weighs earlier deaths more significantly, we found that GLUT-3 overexpression increases lifespan

when TDP-43^{G298S} is expressed in glial cells (43.5 days vs 51.5 days, P$_{value}$ = 0.015). GLUT-3 had

no effect on lifespan when TDP-43^{WT} was expressed in either motor neurons or glia. Importantly,

GLUT3 overexpression had no effect on TDP-43 suggesting that its protective effect occurs

through a mechanism independent of TDP-43 levels (see Fig. 4 – figure supplement 4). Together,

these findings suggest that increased glucose availability is protective in both motor neurons and

glia but to a different extent that likely reflects different metabolic capabilities of the two cell types.

Although GLUT-3 is the predominant glucose transporter in the brain, recently, the human

glucose transporter GLUT-4, an insulin dependent glucose transporter expressed primarily in

muscle has been shown to localize to the plasma membrane to support synaptic activity (Ashrafi

et al., 2017). To test whether GLUT-4 also modulates TDP-43 proteinopathy, we co-expressed

GLUT-4 and TDP-43 in either motor neurons or glia. Similar to GLUT-3, co-overexpression of

GLUT-4 and TDP-43^{WT} or TDP-43^{G298S}, in either motor neurons or glial cells, improved the

locomotor deficits caused by TDP-43 expression alone (Fig. 4 – figure supplement 5). Notably,

while TDP-43 overexpression in muscle causes larval turning phenotypes, co-overexpression of

either GLUT-3 or GLUT-4 in muscle does not mitigate TDP-43 dependent locomotor deficits,

suggesting that the primary metabolic defect lies in the nervous system (Fig. 4 – figure
supplements 2 and 5).

TDP-43 dependent defects at the neuromuscular junction are rescued by GLUT-3
We previously described defects at the *Drosophila* neuromuscular junction (NMJ) caused by
neuronal expression of TDP-43. These defects include a reduction in synaptic bouton number
(Coyne et al., 2014; Estes et al., 2013) and a reduction in synaptic vesicle endocytosis as
indicated by FM1-43 dye uptake experiments (Coyne et al., 2017). To test whether increased
GLUT-3 could also rescue morphological and functional defects at the larval NMJ, we co-
expressed GLUT-3 and TDP-43 in motor neurons. Co-expression of GLUT-3 rescued NMJ size
as indicated by the total number of synaptic boutons in both TDP-43^{WT} (P_{value} = 0.004) and TDP-
43^{G298S} (P_{value} = 0.023) expressing larvae (Figs. 5A and 5C). Moreover, GLUT-3 overexpression
partially restores synaptic vesicle endocytosis deficits caused by TDP-43 (Figs. 5B and 5D, P_{value}
<0.001 for both TDP-43^{WT} and TDP-43^{G298S}). Taken together, these results show that GLUT-3
expression in motor neurons mitigates morphological and functional defects caused by TDP-43
at the NMJ and suggest that increased glucose availability in motor neurons is neuroprotective in
ALS.

Neuronal co-expression of *PFK* rescues TDP-43 induced toxicity
Increased *PFK* levels in conjunction with our findings that increased glucose availability is
protective suggest that increased glycolysis may be compensatory. To test this hypothesis, we
directly manipulated glycolysis genetically, either via *PFK* overexpression or RNAi knock-down,
in the context of TDP-43 proteinopathy. Overexpression of *PFK* alone in motor neurons negatively
altered locomotor activity suggesting that in normal neurons high glycolytic activity may be toxic
(Fig. 6A). However, the co-expression of *PFK* with TDP-43^{WT} or TDP-43^{G298S} rescued locomotor
defects to control levels (P_{value} = 0.003 for TDP-43^{WT} and P_{value} <0.001 for TDP-43^{G298S}, Fig. 6A).
In contrast, *PFK* RNAi knock-down had no significant effect on larval turning when expressed in
motor neurons alone (53% knock-down, P_{value}= 0.04, see Materials and Methods). However, when

PFK was knocked down by RNAi in the context of TDP-43 proteinopathy we found that it

significantly increased larval turning times for TDP-43^{WT} albeit it had no effect on TDP-43^{G298S}

(P_{value} < 0.001 for TDP-43^{WT} and P_{value} = 0.767 for TDP-43^{G298S}, Fig. 6B). One possible

interpretation for this variant dependent effect is that the metabolic deficit of TDP-43^{G298S} is so

severe that it may be difficult to exacerbate by further altering energy metabolism. Of note, a

second *PFK* RNAi line (y(1) sc[*] v(1) P{y[+t7.7] v[+t1.8]=TRiP.HMS01324}attP2) was tested, but

was lethal when expressed in motor neurons.

Similar to the experiments with motor neurons, the over-expression of *PFK* in glial cells

causes a locomotor defect, while the expression of two independent *PFK* RNAi lines has no effect

on locomotor activity as determined by larval turning assays (Fig. 6 – figure supplement 1). When

PFK is co-expressed in glial cells with TDP-43^{WT} there is a slight but not statistically significant

reduction in larval turning (P_{value} > 0.302) while it significantly mitigates TDP-43^{G298S} (P_{value} = 0.001,

Fig. 6 – figure supplement 1). Moreover, the knockdown of *PFK* in conjunction with TDP-43^{WT}

significantly impaired locomotor function when compared to TDP-43^{WT} expression alone (P_{value} <

0.001, Fig. 6 – figure supplement 1). *PFK* knock down did not affect TDP-43^{G298S} expressing flies

P_{value} > 0.057, Fig. 6 – figure supplement 1). Importantly, the protective effect of *PFK*

overexpression in motor neurons does not occur via reducing TDP-43 levels (Fig. 6 – figure

supplement 1). These results indicate that an increase in glycolysis, in either motor neurons or

glial cells, can ameliorate TDP-43 toxicity.

Compensatory upregulation of PFK transcript is not a consequence of sequestration in

TDP-43 insoluble complexes

Our findings of increased *PFK* mRNA levels and genetic interaction data with *PFK* support a

protective compensatory mechanism (see Fig. 7 for model). This could be in part explained by

the ribostasis hypothesis (Ramaswami et al., 2013) whereby *PFK* mRNA associates with TDP-43

in insoluble complexes and as a result, transcription is upregulated to compensate for the amount

of mRNA sequestered and unavailable for translation. This however, does not seem to be the

case, as soluble versus insoluble fractionation experiments do not support this hypothesis (Fig. 7

– figure supplement 1). Surprisingly, *PFK* mRNA appears to be more soluble in the context of

TDP-43^{WT} versus TDP-43^{G298S}, consistent with different TDP-43 variants utilizing distinct

pathomechanisms, as we have previously reported (Coyne et al., 2017; Estes et al., 2013).

Discussion

It is widely known that ALS patients display gross metabolic dysregulation described as

hypermetabolism (for a recent review, see (Joardar et al., 2017)). However, a major challenge in

the field has been to understand how these clinical observations relate to metabolic changes at

the cellular level. Here, we take advantage of cell type specific genetic tools available in

Drosophila to pinpoint how specific metabolic changes in neurons and glia relate to disease

progression. Using metabolic profiling we first found that the neuronal expression of TDP-43

increases the abundance of pyruvate, the end product of glycolysis. Notably, this alteration is

consistent with metabolite changes in plasma from ALS patients (Lawton et al., 2012) and suggest

an increase in glycolysis. Substantiating this scenario is transcriptional profiling of ALS spinal

cords and patient derived iPSC motor neurons showing that just as in flies, *PFK*, the rate limiting

step in glycolysis, is significantly upregulated (Fig. 2). Interestingly, metabolic profiling in flies also

suggests increased glycolytic input into the pentose phosphate pathway (Supplemental file 1),

which provides a mechanism for countering oxidative stress via increased NADPH production

(Kruger and von Schaewen, 2003). Supporting this possibility are our findings that G6PD, the rate

limiting enzyme of the pentose phosphate pathway is upregulated in fly and human ALS ventral

and spinal cords, respectively (Fig. 2). Oxidative stress is a well-established ALS manifestation

(Barber and Shaw, 2010) and the most recent FDA approved drug for treating ALS patients,

Radicava, is aimed at countering this aspect of the disease (Cruz, 2018). Collectively, molecular

and metabolic alterations identified in the fly validate in patient derived motor neurons and spinal

cords, albeit the magnitude of the changes found in human ALS tissues is often more dramatic

and, at some level, more comparable with mutant TDP-43 dependent phenotypes in flies. These

findings highlight the predictive power of the fly model and its relevance to studying disease

pathomechanisms in humans.

Using a genetically encoded glucose sensor we show that TDP-43^{G298S} expressing motor

neurons have increased capacity to import glucose (Fig. 3). This is consistent with increased

glycolysis and its end product, pyruvate. It is unclear why despite finding that pyruvate is also

increased in motor neurons overexpressing TDP-43^{WT}, we could not detect higher glucose uptake

than in controls, although this is consistent with PFK transcript levels trending up, yet not reaching

significance in our data set (Fig. 2). A possible explanation is that while our model is based on

TDP-43 expression specifically in motor neurons or glia, the metabolite profiling was performed

on whole larvae and captured non-cell autonomous changes, occurring in cells other than motor

neurons. Nevertheless, taken together our data show that there is a clear dysregulation of

glycolysis in TDP-43 expressing *Drosophila* and patient tissue samples. Alterations in glucose

metabolism may also be a feature of multiple ALS types. Indeed, pre-symptomatic SOD1 mutant

mice have been shown to have high glucose levels in spinal cords; however, glucose levels

gradually decrease as disease symptoms progress (Miyazaki et al., 2012). One question that

remains is whether alterations in glucose metabolism are compensatory or causative.

In an effort to address this question, we first increased glucose availability in *Drosophila*

by raising the dietary concentration of glucose (5-10 X higher than normal levels in standard fly

food), which mitigates locomotor defects and increases lifespan in ALS flies while making wild-

type larvae sluggish (Fig. 3). Interestingly, a high sugar diet only mitigates locomotor defects when

TDP-43 is expressed in motor neurons or glia but not muscles, indicating that the primary

metabolic defect lies in the nervous system and that muscles do not benefit from increased

glucose availability (Fig. 4 and Fig. 4 – figure supplements 2 and 5). This is somewhat surprising

given that muscles normally rely on glycolysis for ATP production but is also consistent with

findings that in ALS, muscles switch to primarily using lipids and not glucose for energy production

(Palamiuc et al., 2015).

This systemic approach, however, does not provide information on which cells specifically

benefit from increased glucose availability. To answer this question, we intervened genetically by

overexpressing human glucose transporters GLUT-3 or GLUT-4 in motor neurons, glia or muscles

(Figs. 4, 5 and Fig. 4 – figure supplements 2 and 5). These experiments showed that both GLUT-

3 and GLUT-4 rescued locomotor dysfunction of TDP-43 expressing larvae in the nervous system,

but not in muscles (Fig. 4 and Fig. 4 – figure supplements 2 and 5) confirming our findings from

high sugar feedings. Furthermore, our findings that overexpression of either GLUT-3 or GLUT-4

can help mitigate specific ALS-like phenotypes in flies are consistent with recent reports that both

GLUT-3 and GLUT-4 are required at synapses to regulate activity (Ashrafi et al., 2017; Ferreira

et al., 2011). Together, our dietary intervention and GLUT-3/4 overexpression experiments

indicate that increased glucose availability in motor neurons and glia confers protection in models

of TDP-43 proteinopathy, in a cell type dependent manner.

A possible explanation for why increased glucose availability improves defects caused by

TDP-43 proteinopathy is that cells are actively consuming glucose to increase glycolytic flux. A

recent study using vertebrate cell lines suggests that glycolytic flux depends on four key steps.

They include: 1) glucose uptake by glucose transporters, 2) hexokinase levels, 3) *PFK* levels (and

associated enzymes), and 4) lactate export, but not other enzymes or steps involved in glycolysis

(Tanner et al., 2018). The co-over-expression of *PFK* mitigates TDP-43 defects in flies, and is

consistent with our GLUT-3 over-expression experiments (Fig. 6). Surprisingly, *PFK* transcript

levels were higher in human ALS post-mortem tissues compared to controls, however protein

levels in total tissue homogenates were not significantly increased despite trending upwards (data

not shown). Nevertheless, our findings of increased *PFK* mRNA levels in flies and human tissues,

and the genetic interaction experiments support the notion that glycolysis is increased in ALS as

a compensatory mechanism (see Fig. 7 for model).

The fact that neurons would compensate deficits in cellular energetics through increasing

glycolysis makes sense in the light of well-established defects in mitochondrial function. TDP-43

has been previously shown to localize to the inner mitochondrial membrane of mitochondria

(Wang et al., 2016), and it has been shown to cause abnormally small mitochondria in *Drosophila*

(Khalil et al., 2017). Consistent with the importance of glycolysis in neurodegeneration, there are

reports of SNPs within *PFK* that are present in ALS patients but not in healthy controls (Rouillard

et al., 2016; Xie et al., 2014). Our findings that glycolysis is increased as a compensatory

mechanism and this confers neuroprotection is consistent with findings from a small clinical trial

showing that a high caloric high carbohydrate diet improves patient outcome (Wills et al., 2014a)

and also that patients with diabetes have a later onset of disease (Kioumourtzoglou et al., 2015).

While upregulation of glycolysis decreases the dependence of the organism for ATP derived from

mitochondria, it is also possible that increased glycolysis is improving organismal health through

a different mechanism. For example, pyruvate has been previously shown to protect mitochondria

from oxidative stress (Wang et al., 2007). Future studies should focus on the link between

glycolysis, oxidative stress and mitochondria in ALS.

Materials and Methods

***Drosophila* genetics**

Standard crosses were incubated at 25°C with a 12-hour alternating dark-light cycles. (i) w^{1118} (ii)

w^{1118}; UAS-TDP43^{WT}-YFP (iii) w^{1118}; UAS-TDP43^{G29S8}-YFP (iv) UAS-GLUT3 flies were crossed

with the GAL4 motor neuron driver D42. GLUT3 TDP-43 recombinant flies were generated using

standard genetic techniques. GLUT3 flies were described in (Besson et al., 2015).

TDP-43 was expressed in motor neurons and glia using the D42 GAL4 (Gustafson and

Boulianne, 1996) and repo GAL4 (Sepp et al., 2001) drivers, respectively. w^{1118} was crossed with

either the D42 or repo GAL4 drivers as a control. Endogenous TDP-43 was knocked down via

RNAi using y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01932}attP40 and y[1] v[1]; P{y[+t7.7]

v[+t1.8]=TRiP.HM05194}attP2.

To manipulate *PFK*, we used w^{1118}; P{w[+mC]=UAS-Pfk.T}3 for over expression, and y^1

sc* v^1; P{y[+t7.7] v[+t1.8]=TRiP.HMS01324}attP2 or y^1 sc* v^1; P{y[+t7.7] v[+t1.8]=TRiP.GL00298}-

attP2 (Line 2, viable only in glial expressing flies) for PFK knock down. *attP2* was used as a

genetic background control for PFK RNAi knock-down experiments. All lines were obtained from

the Bloomington Stock Center.

PFK RNAi efficiency was assayed by crossing *PFK* RNAi to D42 GAL4. Ventral nerve

cords were dissected and analyzed via qPCR (see below for primers). A Welch's t-test was used

to identify significance.

Fly food supplementation

Standard yeast/cornmeal/molasses food was heated and then allowed to cool to 55-60˚C. For high sugar supplementation, additional glucose was added to reach a final concentration of 357g/L (10X compared to regular food). Supplemented food was dispensed into vials and allowed to cool.

Locomotor assays

Larval turning assays were performed as previously described (Estes et al., 2011; Estes et al., 2013). Briefly, larvae are placed on a petri dish filled with solidified agar (2.5%) and grape juice. Third instar larvae are transferred from a vial onto the agar/juice surface and are allowed to acclimate. Using a clean paintbrush, larvae are turned ventral side up. The time it takes for larvae to turn themselves ventral side down and make a forward motion is recorded. Larval turning data were analyzed using Kruskal-Wallis multiple comparison tests (GraphPad Prism v8). Negative geotaxis assays on adult flies were performed by using TDP-43 expressing flies with a milder phenotype/lethality. TDP-43[WT] and TDP-43[G298S] flies were previously described in (Khalil et al., 2017). Briefly, TDP-43 and GLUT-3 were expressed in neurons with the pan neuronal driver, elav GAL4. Female flies of 11-day old were anesthetized with CO_2 and placed in a plastic column (diameter 1.3 cm) with a mark at 22 cm. After 20 min recovery, flies were gently tapped to the bottom of the column. Then the percentages of flies reaching the 22 cm mark or remaining at the bottom were counted after 60 sec. The test was repeated 3 times and results were averaged. The data are the mean of at least 6 trials and are expressed as mean ± SEM. Statistical comparisons were performed using one-way ANOVA followed by Tukey's multiple-comparison test.

Metabolite analysis

Metabolomic analysis was conducted by Metabolon, Inc as previously described (Joardar et al., 2014; Manzo et al., 2018). Briefly, 50–60 wandering third instar larvae (~50–60 mg) per sample were collected and flash-frozen in liquid nitrogen. A total of 5 replicates per genotype were collected. Samples were analyzed by ultrahigh performance liquid chromatography/mass

spectrometry (UHPLC/MS), or by gas chromatography/mass spectrometry (GC/MS). One-way

ANOVA was used to identify biochemicals that differed significantly between experimental groups.

Pyruvate measurements

Pyruvate was measured using the Pyruvate Assay Kit (Abcam-ab65342) as per manufacturer's

instructions. In brief, 20 3[rd] instar larvae per genotype were analyzed in three biological

replicates. Standard curve was generated kit provided reagents. Results are shown (Fig. 1 –

figure supplement 1) in nmoles/microliter.

RNA isolation, cDNA preparation and qPCR

cDNA protocol has been previously described (Manzo et al., 2018). Briefly, 5 ventral nerve cords

(VNCs) were isolated from third instar larvae per genotype using HL-3 saline buffer (70 mM NaCL,

5 mM KCl, 22 mM $MgCl_2$, 10 mM $NaHCO_3$, 5 mM trehalose, 115 mM sucrose, 5 mM HEPES, pH

7.3). RNA was extracted using the RNeasy RNA Extraction Kit (Qiagen) and stored at -80°C.

Genomic DNA (gDNA) was cleared using a DNase I (Thermo Fisher Scientific) digestion protocol.

Following gDNA clearing, reverse transcription was performed using the Maxima First Strand

cDNA Synthesis Kit for RT-qPCR (Thermo Fisher Scientific). The quality of the cDNA was tested

using GPDH primers (see below for sequence) and GoTaq endpoint polymerase chain reaction

(PCR) protocol (Promega). Quantitative PCR was performed using a SYBR Green protocol

(Applied Biosystems) and primers designed to span exon-exon junctions. Each reaction was

conducted in 3-6 biological replicates with three technical replicates each. ΔΔCT values are

normalized to GPDH (Pfaffl, 2001).The Kruskal-Wallis test was used to assess statistical

significance in GraphPad Prism v8. Protocols for human tissue RNA isolation and cDNA

preparation have been previously described (Bakkar et al., 2018). RT-PCR was performed using

TaqMan Gene Expression Assays following the TaqMan Fast Advanced Master Mix protocol

(Applied Biosystems) on the QuantStudio 6 Flex instrument (Applied Biosystems).

Primer sequences and Taqman probes

Drosophila Phosphofructokinase (PFK): Forward primer - TGGACGAGCTGGTCAAGAAC;

Reverse primer - CCACAAGAGCTAAATAGCCG

Drosophila G6PD (Zw): Forward primer - AAGCGCCGCAACTCTTTG; Reverse primer -

AGGGCGGTGTGATCTTCC

Drosophila GPDH: Forward primer - CCGCAGTGCTTGTTTTGCT*;* Reverse primer –

TATGGCCGAACCCCAGTTG

Human *Phosphofructokinase P* (PFKP): Assay ID Hs00242993_m1

Human *Phosphofructokinase M* (PFKM): Assay ID Hs00175997_m1

Human G6PD: Assay ID Hs00166169_m1

Human GAPDH: Assay ID Hs99999905_m1

Glucose sensor

Third instar larval brains where dissected in HL3-buffer (70 mM NaCl, 5 mM KCl, 20 mM MgCl2,

10 mM NaHCO3, 115 mM sucrose, 5 mM trehalose, 5 mM HEPES; pH 7.1). Of note, the glucose

sensor is not sensitive to trehalose (Volkenhoff et al., 2018). Larvae were pinned down on a

Sylgard plate using insect pins and cleaned so that the ventral nerve cord is clearly visible. A 40x

water immersion lens was used to image larvae individually. For glucose stimulation, HL3-buffer

was removed and replaced with 5mM glucose supplemented HL3 (70 mM NaCl, 5 mM KCl, 20

mM MgCl2, 10 mM NaHCO3, 115 mM sucrose, 5 mM glucose, 5mM HEPES, pH 7.1).

Fluorescent images were acquired using a Zeiss LSM880 NLO upright

multiphoton/confocal microscope. CFP was excited using a 405-nm laser, and images were

acquired in both the CFP (CFP 465–499 nm) and the FRET (535–695 nm) detection channels.

Data was analyzed using ImageJ with a drift correction plugin turboreg (Thevenaz et al.,

1998). Regions of interest (ROIs) were manually selected and the mean grey value of each ROI

was extracted. The mean grey values for the FRET and CFP channels was taken to calculate the

FRET ratio (F/C), which corresponds to the intracellular glucose concentration.

Each ROI was imaged every 10 seconds for 10 minutes (baseline) and then for another

10 minutes under high glucose HL-3 (stim; see Fig. 3 – figure supplement 1 for representative

examples). Each FRET ratio was normalized to the mean baseline value for each genotype. For

baseline vs stim comparisons (Fig. 3B), the normalized FRET ratio of each ROI was taken from

minutes 5-10 and minutes 15-20.

Western blots

TDP-43 was detected via its YFP tag using either the Clontech/Takara Living Colors GFP

monoclonal antibody (6326375) at 1:1,000 (for Fig. 4 – figure supplement 4) or the ThermoFisher

polyclonal antibody (A-11122) at 1:1,000 (for Fig. 6 – figure supplement 1). For normalization we

used either the Cell Signaling polyclonal rabbit anti β-Actin antibody (4967) at a 1:1,000 dilution

or the Millipore Sigma monoclonal antibody against beta tubulin (Mab3408) at 1:1,000. 5% milk

was used as blocking agent for all westerns.

FM1-43 experiments

Larvae are dissected in HL-3 buffer on a sylgard dish as previously described. FM1-43 dye uptake

assays were conducted using 4 μM FM1-43FX, 90 mM KCl, and 2 mM Ca2+ for stimulation.

Methods adapted from (Kuromi and Kidokoro, 2005; Verstreken et al., 2008). To determine dye

uptake signal, the background-corrected fluorescence for every synaptic bouton was calculated

by subtracting the mean background fluorescence of cell-free regions equal in size to the synaptic

bouton.

Bouton counts

Larval NMJs were dissected and fixed with 3.5% paraformaldehyde in PBS (pH 7.2: 130 mM NaCl,

5 mM Na_2HPO_4, 5mM NaH_2PO_4) for 30 minutes. Antibody dilutions are as follows: 1:300 mouse

anti-DCSP2 (DSHB), 1:500 anti-mouse Alexa Fluor 568 (Invitrogen), 1:200 Anti-HRP-CY5

conjugate (Invitrogen), and 1:40 Phalloidin-488 (Invitrogen). Blocking solution contained 2% BSA

and 5% Normal goat serum. Methods have been previously described in (Coyne et al., 2014).

Larval muscles 6 and 7, segment A3, were imaged using an LSM 880 confocal microscope

(Zeiss). NIH ImageJ was used to count bouton numbers and measure muscle area. Bouton count

was normalized to muscle area to account for size variation.

iPSC methods

Control iPSC and TDP-43 iPSC cells were differentiated to motor neurons as previously described

(Donnelly et al., 2013; Zhang et al., 2015).

Cellular Fractionations

To generate cellular fractions, 25 3rd instar wandering larvae were flash frozen, followed by

homogenization in Trizol^{TM}reagent using the Bullet Blender^{TM} Bead Lysis Kit Green in

conjunction with a Bullet Blender^{TM} Blue 24. Homogenates were then centrifuged at 2,000 x g

for 3 minutes. The supernatant was separated from both the fat layer and the beads then it was

further centrifuged at 25,000 g for 30 minutes. The resulting supernatant is the soluble fraction

and the insoluble fraction is generated by resuspending the pellet in urea buffer (30 mM Tris, 7

M Urea, 2 M Thiourea, 4% CHAPS, 1X Protease Inhibitor Cocktail (Roche), 0.5 mM PMSF, 40

units/μl RNAsin plus, pH 8.5). RNA was isolated from inputs and the cellular fractions utilizing

the manufactured described protocol for Trizol^{TM} reagent. cDNA was then generated utilizing

Thermoscientific^{TM} Maxima First Strand cDNA Synthesis Kit for RT-qPCR. RT-qPCR was

conducted on the StepOnePlus^{TM} RT-qPCR machine utilizing Sybr reagents.

Statistics

All statistical analyses for locomotor assays or qPCR comprising three or more data sets were

conducted using Kruskal-Wallis with uncorrected Dunn's test in Prism v8.0 (GraphPad). When

only two data sets were compared, either Welch's test or Wilcoxon signed-rank test were used.

Biological replicates for qPCR and metabolomics are defined as independent sample collections

per genotype, from independent genetic crosses. All qPCR experiments included three technical

replicates per biological replicate. Outliers were removed using ROUT method (Q=1%) in Prism

v8.0.

References cited:

Ashrafi, G., Wu, Z., Farrell, R.J., and Ryan, T.A. (2017). GLUT4 Mobilization Supports Energetic

Demands of Active Synapses. Neuron *93*, 606-615 e603.

Bakkar, N., Kovalik, T., Lorenzini, I., Spangler, S., Lacoste, A., Sponaugle, K., Ferrante, P.,

Argentinis, E., Sattler, R., and Bowser, R. (2018). Artificial intelligence in neurodegenerative

disease research: use of IBM Watson to identify additional RNA-binding proteins altered in

amyotrophic lateral sclerosis. Acta Neuropathol *135*, 227-247.

Barber, S.C., and Shaw, P.J. (2010). Oxidative stress in ALS: key role in motor neuron injury

and therapeutic target. Free Radic Biol Med *48*, 629-641.

Besson, M.T., Alegria, K., Garrido-Gerter, P., Barros, L.F., and Lievens, J.C. (2015). Enhanced

neuronal glucose transporter expression reveals metabolic choice in a HD Drosophila model.

PLoS One *10*, e0118765.

Browne, S.E., Yang, L., DiMauro, J.P., Fuller, S.W., Licata, S.C., and Beal, M.F. (2006).

Bioenergetic abnormalities in discrete cerebral motor pathways presage spinal cord pathology in

the G93A SOD1 mouse model of ALS. Neurobiol Dis *22*, 599-610.

Coyne, A.N., Lorenzini, I., Chou, C.C., Torvund, M., Rogers, R.S., Starr, A., Zaepfel, B.L., Levy,

J., Johannesmeyer, J., Schwartz, J.C.*, et al.* (2017). Post-transcriptional Inhibition of Hsc70-

4/HSPA8 Expression Leads to Synaptic Vesicle Cycling Defects in Multiple Models of ALS. Cell

Rep *21*, 110-125.

Coyne, A.N., Siddegowda, B.B., Estes, P.S., Johannesmeyer, J., Kovalik, T., Daniel, S.G.,

Pearson, A., Bowser, R., and Zarnescu, D.C. (2014). Futsch/MAP1B mRNA Is a Translational

Target of TDP-43 and Is Neuroprotective in a Drosophila Model of Amyotrophic Lateral

Sclerosis. J Neurosci *34*, 15962-15974.

Cruz, M.P. (2018). Edaravone (Radicava): A Novel Neuroprotective Agent for the Treatment of

Amyotrophic Lateral Sclerosis. P T *43*, 25-28.

Donnelly, C.J., Zhang, P.W., Pham, J.T., Haeusler, A.R., Mistry, N.A., Vidensky, S., Daley, E.L.,

Poth, E.M., Hoover, B., Fines, D.M., *et al.* (2013). RNA toxicity from the ALS/FTD C9ORF72

expansion is mitigated by antisense intervention. Neuron *80*, 415-428.

Dupuis, L., Pradat, P.F., Ludolph, A.C., and Loeffler, J.P. (2011). Energy metabolism in

amyotrophic lateral sclerosis. Lancet Neurol *10*, 75-82.

Estes, P.S., Boehringer, A., Zwick, R., Tang, J.E., Grigsby, B., and Zarnescu, D.C. (2011). Wild-

type and A315T mutant TDP-43 exert differential neurotoxicity in a Drosophila model of ALS.

Human molecular genetics *20*, 2308-2321.

Estes, P.S., Daniel, S.G., McCallum, A.P., Boehringer, A.V., Sukhina, A.S., Zwick, R.A., and

Zarnescu, D.C. (2013). Motor neurons and glia exhibit specific individualized responses to TDP-

43 expression in a Drosophila model of amyotrophic lateral sclerosis. Disease models &

mechanisms *6*, 721-733.

Fehr, M., Lalonde, S., Lager, I., Wolff, M.W., and Frommer, W.B. (2003). In vivo imaging of the

dynamics of glucose uptake in the cytosol of COS-7 cells by fluorescent nanosensors. J Biol

Chem *278*, 19127-19133.

Ferreira, J.M., Burnett, A.L., and Rameau, G.A. (2011). Activity-dependent regulation of surface

glucose transporter-3. J Neurosci *31*, 1991-1999.

Gustafson, K., and Boulianne, G.L. (1996). Distinct expression patterns detected within

individual tissues by the GAL4 enhancer trap technique. Genome *39*, 174-182.

Hardiman, O., Al-Chalabi, A., Chio, A., Corr, E.M., Logroscino, G., Robberecht, W., Shaw, P.J.,

Simmons, Z., and van den Berg, L.H. (2017). Amyotrophic lateral sclerosis. Nat Rev Dis Primers

3, 17071.

Joardar, A., Manzo, E., and Zarnescu, D.C. (2017). Metabolic Dysregulation in Amyotrophic

Lateral Sclerosis: Challenges and Opportunities. Curr Genet Med Rep *5*, 108-114.

Joardar, A., Menzl, J., Podolsky, T.C., Manzo, E., Estes, P.S., Ashford, S., and Zarnescu, D.C.

(2014). PPAR gamma activation is neuroprotective in a Drosophila model of ALS based on

TDP-43. Hum Mol Genet.

Khalil, B., Cabirol-Pol, M.J., Miguel, L., Whitworth, A.J., Lecourtois, M., and Lievens, J.C.

(2017). Enhancing Mitofusin/Marf ameliorates neuromuscular dysfunction in Drosophila models

of TDP-43 proteinopathies. Neurobiol Aging *54*, 71-83.

Kioumourtzoglou, M.A., Rotem, R.S., Seals, R.M., Gredal, O., Hansen, J., and Weisskopf, M.G.

(2015). Diabetes Mellitus, Obesity, and Diagnosis of Amyotrophic Lateral Sclerosis: A

Population-Based Study. JAMA Neurol *72*, 905-911.

Kruger, N.J., and von Schaewen, A. (2003). The oxidative pentose phosphate pathway:

structure and organisation. Curr Opin Plant Biol *6*, 236-246.

Kuromi, H., and Kidokoro, Y. (2005). Exocytosis and endocytosis of synaptic vesicles and

functional roles of vesicle pools: lessons from the Drosophila neuromuscular junction.

Neuroscientist *11*, 138-147.

Lawton, K.A., Cudkowicz, M.E., Brown, M.V., Alexander, D., Caffrey, R., Wulff, J.E., Bowser, R.,

Lawson, R., Jaffa, M., Milburn, M.V.*, et al.* (2012). Biochemical alterations associated with ALS.

Amyotrophic lateral sclerosis : official publication of the World Federation of Neurology

Research Group on Motor Neuron Diseases *13*, 110-118.

Ling, S.C., Polymenidou, M., and Cleveland, D.W. (2013). Converging mechanisms in ALS and

FTD: disrupted RNA and protein homeostasis. Neuron *79*, 416-438.

Ludolph, A.C., Langen, K.J., Regard, M., Herzog, H., Kemper, B., Kuwert, T., Bottger, I.G., and

Feinendegen, L. (1992). Frontal lobe function in amyotrophic lateral sclerosis: a

neuropsychologic and positron emission tomography study. Acta Neurol Scand *85*, 81-89.

Manzo, E., O'Conner, A.G., Barrows, J.M., Shreiner, D.D., Birchak, G.J., and Zarnescu, D.C.

(2018). Medium-Chain Fatty Acids, Beta-Hydroxybutyric Acid and Genetic Modulation of the

Carnitine Shuttle Are Protective in a Drosophila Model of ALS Based on TDP-43. Front Mol

Neurosci *11*, 182.

Mergenthaler, P., Lindauer, U., Dienel, G.A., and Meisel, A. (2013). Sugar for the brain: the role

of glucose in physiological and pathological brain function. Trends Neurosci *36*, 587-597.

Miyazaki, K., Masamoto, K., Morimoto, N., Kurata, T., Mimoto, T., Obata, T., Kanno, I., and Abe,

K. (2012). Early and progressive impairment of spinal blood flow-glucose metabolism coupling

in motor neuron degeneration of ALS model mice. J Cereb Blood Flow Metab *32*, 456-467.

Musselman, L.P., Fink, J.L., Narzinski, K., Ramachandran, P.V., Hathiramani, S.S., Cagan,

R.L., and Baranski, T.J. (2011). A high-sugar diet produces obesity and insulin resistance in

wild-type Drosophila. Dis Model Mech *4*, 842-849.

Na, J., Musselman, L.P., Pendse, J., Baranski, T.J., Bodmer, R., Ocorr, K., and Cagan, R.

(2013). A Drosophila model of high sugar diet-induced cardiomyopathy. PLoS Genet *9*,

e1003175.

Neumann, M., Sampathu, D.M., Kwong, L.K., Truax, A.C., Micsenyi, M.C., Chou, T.T., Bruce, J.,

Schuck, T., Grossman, M., Clark, C.M.*, et al.* (2006). Ubiquitinated TDP-43 in frontotemporal

lobar degeneration and amyotrophic lateral sclerosis. Science *314*, 130-133.

Palamiuc, L., Schlagowski, A., Ngo, S.T., Vernay, A., Dirrig-Grosch, S., Henriques, A., Boutillier,

A.L., Zoll, J., Echaniz-Laguna, A., Loeffler, J.P.*, et al.* (2015). A metabolic switch toward lipid

use in glycolytic muscle is an early pathologic event in a mouse model of amyotrophic lateral

sclerosis. EMBO Mol Med *7*, 526-546.

Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR.

Nucleic acids research *29*, e45.

Ramaswami, M., Taylor, J.P., and Parker, R. (2013). Altered ribostasis: RNA-protein granules in

degenerative disorders. Cell *154*, 727-736.

Rouillard, A.D., Gundersen, G.W., Fernandez, N.F., Wang, Z., Monteiro, C.D., McDermott,

M.G., and Ma'ayan, A. (2016). The harmonizome: a collection of processed datasets gathered

to serve and mine knowledge about genes and proteins. Database (Oxford) *2016*.

Sepp, K.J., Schulte, J., and Auld, V.J. (2001). Peripheral glia direct axon guidance across the

CNS/PNS transition zone. Dev Biol *238*, 47-63.

Takanaga, H., Chaudhuri, B., and Frommer, W.B. (2008). GLUT1 and GLUT9 as major

contributors to glucose influx in HepG2 cells identified by a high sensitivity intramolecular FRET

glucose sensor. Biochim Biophys Acta *1778*, 1091-1099.

Tanner, L.B., Goglia, A.G., Wei, M.H., Sehgal, T., Parsons, L.R., Park, J.O., White, E.,

Toettcher, J.E., and Rabinowitz, J.D. (2018). Four Key Steps Control Glycolytic Flux in

Mammalian Cells. Cell Syst *7*, 49-62 e48.

Tauffenberger, A., Vaccaro, A., Aulas, A., Vande Velde, C., and Parker, J.A. (2012). Glucose

delays age-dependent proteotoxicity. Aging Cell.

Thevenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to subpixel

registration based on intensity. IEEE Trans Image Process *7*, 27-41.

Verstreken, P., Ohyama, T., and Bellen, H.J. (2008). FM 1-43 labeling of synaptic vesicle pools

at the Drosophila neuromuscular junction. Methods Mol Biol *440*, 349-369.

Volkenhoff, A., Hirrlinger, J., Kappel, J.M., Klambt, C., and Schirmeier, S. (2018). Live imaging

using a FRET glucose sensor reveals glucose delivery to all cell types in the Drosophila brain. J

Insect Physiol *106*, 55-64.

Wang, W., Wang, L., Lu, J., Siedlak, S.L., Fujioka, H., Liang, J., Jiang, S., Ma, X., Jiang, Z., da

Rocha, E.L.*, et al.* (2016). The inhibition of TDP-43 mitochondrial localization blocks its neuronal

toxicity. Nat Med *22*, 869-878.

Wang, X., Perez, E., Liu, R., Yan, L.J., Mallet, R.T., and Yang, S.H. (2007). Pyruvate protects

mitochondria from oxidative stress in human neuroblastoma SK-N-SH cells. Brain Res *1132*, 1-

9.

Wills, A.M., Hubbard, J., Macklin, E.A., Glass, J., Tandan, R., Simpson, E.P., Brooks, B.,

Gelinas, D., Mitsumoto, H., Mozaffar, T.*, et al.* (2014a). Hypercaloric enteral nutrition in patients

with amyotrophic lateral sclerosis: a randomised, double-blind, placebo-controlled phase 2 trial.

Lancet *383*, 2065-2072.

Wills, A.M., Hubbard, J., Macklin, E.A., Glass, J., Tandan, R., Simpson, E.P., Brooks, B.,

Gelinas, D., Mitsumoto, H., Mozaffar, T.*, et al.* (2014b). Hypercaloric enteral nutrition in patients

with amyotrophic lateral sclerosis: a randomised, double-blind, placebo-controlled phase 2 trial.

Lancet *383*, 2065-2072.

Xie, T., Deng, L., Mei, P., Zhou, Y., Wang, B., Zhang, J., Lin, J., Wei, Y., Zhang, X., and Xu, R.

(2014). Genome-wide association study combining pathway analysis for typical sporadic

amyotrophic lateral sclerosis in Chinese Han populations. Neurobiol Aging *35*, 1778 e1779-

1778 e1723.

Zhang, P.W., Haidet-Phillips, A.M., Pham, J.T., Lee, Y., Huo, Y., Tienari, P.J., Maragakis, N.J.,

Sattler, R., and Rothstein, J.D. (2015). Generation of GFAP::GFP astrocyte reporter lines from

human adult fibroblast-derived iPS cells using zinc-finger nuclease technology. Glia.

Figure 1. **Glycolysis and pentose phosphate pathways are altered by TDP-43 expression**

in motor neurons. (A) Metabolite changes in glycolysis for whole larvae expressing TDP-43[WT]

or TDP-43[G298S] were analyzed using mass spectrometry (see Materials and Methods). Green and

red font represent metabolites that are significantly changed compared to controls (w[1118]), as

indicated. PEP and pyruvate were upregulated in both TDP-43[WT] and TDP-43[G298S] expressing

flies and is denoted by red asterisks. Changes in the pentose phosphate pathway metabolites are

specific to larvae expressing TDP-43[G298S]. (**B, C, D)** Significant changes in select metabolites

shown as box and whisker plots. Whiskers represent maximum and minimum values. Box edges

represent upper and lower quartiles. Median values are denoted by horizontal lines within each

box. One-way ANOVA was used to identify metabolites that differed significantly between

experimental groups (N=5).

 Figure 2. Glycolytic enzymes are transcriptionally upregulated. qPCR profiling of PFK (**A**;

677 N=5) and G6PD (**B**; N=5) from ventral nerve cords of *Drosophila*. Human PFKP, PFKM, or G6PD

678 mRNA levels were profiled in either spinal cords (**C**; N=8 control and 9 ALS cases) or human

679 iPSCs (**D**; N=2 differentiations). Kruskal-Wallis test was used to identify significance.

Figure 3. TDP expressing neurons have altered capacity to import glucose. FRET based

glucose sensor described in (Volkenhoff et al., 2018) was used to measure the glucose import

capacity. Glucose sensor schematic described in (**A**). Ex – Excitation; Em – Emission. **(B, C)**

TDP-43 expressing neurons and controls were imaged to detect CFP and FRET signal. 12-14

neurons were imaged every 10 seconds for 20 minutes. Values shown are the mean of 12-14

individual cells (ROI) from two ventral nerve cords **(B)**. Mean values for 5-10 minutes and 15-20

minutes time intervals were used to calculate the "baseline" and "stimulated" ("Stim") values

respectively **(C)**. Kruskal-Wallis test was used to calculate significance.

Figure 4. A high glucose diet rescues neuronal TDP-43 toxicity in flies. TDP-43^{WT} or ALS

associated TDP-43^{G298S} were expressed in MNs (using GAL4-UAS). **(A, B)** Larval turning and

lifespan assays for *Drosophila* fed a cornmeal based food containing either regular concentration

of sugar (RS) or a high sugar diet (HS:10x the standard amount of sugar). **(C, D)** Larval turning

and lifespan assays for *Drosophila* expressing GLUT-3 on its own or with TDP-43, as indicated.

At least 30 larvae were tested in larval turning assays and on average 20 adults were assayed

for survival. Kruskal-Wallis test and Log-rank (Mantel-Cox) test was used to determine statistical

significance for larval turning and survival curve respectively. * - $p < .05$, ** - $p<0.01$, *** - $p<0.001$.

Figure 5. TDP-43 dependent defects at the NMJ are rescued by GLUT-3. Third instar larvae

NMJ from segment A3, muscle 6/7 were immunostained for CSP and HRP **(A)** or analyzed for

their ability to endocytose FM1-43 dye upon stimulation with 90 mM KCl **(B)**. **(A, C)** Neuronal

TDP-43 expression in *Drosophila* neurons reduces the number of boutons (labeled with CSP and

HRP **A, C**) and reduces FM1-43 dye uptake **(B, D)**. These morphological **(A, C)** and functional

(B, D) deficits are rescued by co-expression of GLUT-3. N = 7-10 larvae. Kruskal-Wallis test was

used to identify significance.

Figure 6. Co-overexpression of PFK rescues TDP-43 induced locomotor defects. **(A)** TDP-43^{WT} or ALS associated TDP-43^{G298S} were expressed in MNs (using the GAL4-UAS system together with *Drosophila* UAS-PFK). **(B)** TDP-43^{WT} or ALS associated TDP-43^{G298S} were expressed in MNs (using the GAL4-UAS system together with *Drosophila* UAS-PFK^{RNAi}). N = 30 larvae. Kruskal-Wallis was used to determine statistical significance. * - P $_{value}$ < .05, ** - P $_{value}$ < 0.01, *** - P $_{value}$ < 0.001.

Figure 7. Proposed model showing PFK transcript levels increase in response to TDP-43 proteinopathy. (A) Neurons from non-diseased patients. **(B)** ALS neurons showing an increase in *PFK* transcript levels. SV – synaptic vesicle.

Supplemental Figures

Figure 1 – figure supplement 1

Pyruvate measurements in whole larvae expressing RNAi knock-down constructs for the endogenous *Drosophila* TDP-43 (TBPH) with the D42 GAL driver show no significant changes compared to w[1118] controls. P_{value}=0.092 (Student's t test).

Glucose sensor (GS)

GS-TDP^{G298S}

741 **Figure 3 – figure supplement 1**

742 Raw images of glucose sensor **(A)** or glucose sensor in the context of TDP-43^{G298S} **(B)**.

743 Sections shown are taken through the ventral nerve cord 7.5 min post mounting (baseline) and

744 7.5 min post stimulation (Glucose stim), as indicated. See Materials and Methods for details on

745 imaging and analyses. Scale bar as shown.

Figure 4 – figure supplement 1

Larval turning assays on regular cornmeal-molasses food supplemented with various amounts

of glucose (2X, 5X or 10X, as shown). Genotypes as indicated. N=30 larvae. Kruskal-Wallis test

was used to determine statistical significance.

Figure 4 – figure supplement 2

(i, ii) Larval turning assays **(i)** and lifespan assays **(ii)** on a high sugar diet (HS) compared to a

regular sugar diet (RF). Genotypes as indicated. **(iii and iv)** Larval turning assays **(iii)** and

lifespan assays **(iv)** for GLUT-3 overexpression alone or in conjunction with TDP-43. Genotypes

as indicated. **(v and vi)** Larval turning assays on HS versus RF **(v)** or in the context of GLUT-3

and TDP-43 overexpression in muscles **(vi)**. Kruskal-Wallis test was used for larval turning

assays and Log-rank or Mantel-Cox tests were used for lifespan assays to determine

significance.

Figure 4 – figure supplement 3

Negative geotaxis assay on adult flies expressing TDP-43^{WT} or TDP-43^{G298S} alone or in

conjunction with GLUT-3. The percent of flies that reach the top of the column are shown after

60 sec. Gene expression was specifically targeted in all neurons by using the elav GAL4 driver.

Statistical comparisons were performed using one-way ANOVA followed by Tukey's multiple-

comparison test.

Figure 4 – figure supplement 4

i. Western blot of ventral nerve cords probing for TDP-43-YFP and β-actin. D42 motor neuron

driver was used to express TDP-43 in motor neurons. ii. Quantification of 3 western blot

bioreplicates. Protein levels measured by GFP western blot (to detect TDP-YFP) are shown as a

ratio between GLUT-3-TDP^{WT}-YFP to TDP^{WT}-YFP alone and GLUT-3-TDP^{G298S}-YFP to TDP^{G298S}-

YFP alone.

Figure 4 – figure supplement 5

Larval turning assays for GLUT-4 and TDP-43 overexpression in motor neurons (i), glia (ii) or

muscles (iii). Genotypes as indicated. N=30 larvae. Kruskal-Wallis test was used to determine

statistical significance.

Supplemental Figure 4D. Negative geotaxis assay on adult flies expressing TDP-

43WT or TDP-43G298S alone or in conjunction with GLUT-3. The percent of flies that

reach the top (A) of the column are shown after 60 sec. Gene expression was

specifically targeted in all neurons by using the elav GAL4 driver. Statistical

comparisons were performed using one-way ANOVA followed by Tukey's multiple-

comparison test.

Figure 6 – figure supplement 1

Larval turning assays for PFK overexpression **(i)** or RNAi **(ii)** in the context of TDP-43 in glia. Genotypes as indicated. N=30 larvae. Kruskal-Wallis test was used to determine statistical significance. iii. Western blot of ventral nerve cords probing for TDP-43-YFP and tubulin. D42 motor neuron driver was used to express TDP-43 in motor neurons. ii. Quantification of 3 western blot bioreplicates. Protein levels measured by GFP western blot (to detect TDP-YFP) are shown as a ratio between PFK OE - TDP^{WT}-YFP to TDP^{WT}-YFP alone and PFK OE - TDP^{G298S}-YFP to TDP^{G298S}-YFP alone.

Figure 7 – figure supplement 1

Cellular fractionations from third instar larvae. TDP[WT] **(A)** or TDP[G298S] **(B)** were fractionated (see

supplemental materials and methods above) and PFK transcript levels in the urea fraction were

quantified and normalized to controls (w[1118]). Wilcoxon ranked sum test was used to perform

statistics.

	Biochemical name	$\dfrac{TDP^{WT}}{control}$	P_{value}	$\dfrac{TDP^{G298S}}{control}$	P_{value}
Glycolysis, Gluconeogenesis, and Pyruvate Metabolism	glucose	0.85	0.0003	0.95	0.2562
	glucose-6-phosphate (G6P)	1.13	0.697	1.37	0.1516
	glucose 1-phosphate	0.83	0.4723	0.87	0.5666
	fructose-6-phosphate	1.29	0.5165	1.35	0.235
	2-phosphoglycerate	0.77	0.9454	1.05	0.5316
	3-phosphoglycerate	1.03	0.9916	0.98	0.5479
	phosphoenolpyruvate (PEP)	3.29	0.0343	5.35	0.0016
	pyruvate	1.67	0.0072	1.6	0.0089
	lactate	0.9	0.2946	0.92	0.4443
	glycerate	1.05	0.5279	0.84	0.6078
Pentose Phosphate Pathway	6-phosphogluconate	0.62	0.0174	1.1	0.7614
	sedoheptulose-7-phosphate	1.16	0.5718	2.38	0.0084
	ribulose/xylulose 5-phosphate	0.79	0.211	0.58	0.0107
Pentose Metabolism	ribulose	1.13	0.1944	0.91	0.8221
	ribose	1.3	0.0543	0.9	0.7074
	ribitol	0.61	0.0433	0.58	0.0332
	ribonate	0.95	0.6298	1.03	0.7813
	xylonate	0.73	0.0615	1.11	0.7787
	xylose	0.9	0.6029	0.75	0.171
	xylitol	0.89	0.1002	0.85	0.0266
	threitol	1.4	0.4369	2.18	0.0355
	arabitol	1.09	0.5878	1.3	0.194
Glycogen Metabolism	maltopentaose	0.86	0.4789	1.23	0.3532
	maltotetraose	0.95	0.8841	0.98	0.8168
	maltotriose	1.02	0.9606	0.95	0.7228
	maltose	0.85	0.0094	0.97	0.6114
	isomaltose	0.83	0.2842	0.91	0.6496
Disaccharides and Oligosaccharides	trehalose	1.17	0.1078	1.04	0.6138
Fructose, Mannose and Galactose Metabolism	fructose	1.1	0.4677	1.05	0.7955
	sorbose	1.16	0.3647	1.11	0.4997
	sorbitol	0.8	0.0935	0.81	0.1236
	mannose	0.84	0.297	0.98	0.9304
	mannose-6-phosphate	1.15	0.6242	1.38	0.1272
	galactose 1-phosphate	0.78	0.3283	0.92	0.82
	galactonate	0.76	0.0366	0.95	0.8452
Nucleotide Sugar	UDP-glucuronate	0.82	0.4406	0.85	0.5114
	UDP-N-acetylglucosamine	0.76	0.0224	0.86	0.2222
Aminosugar Metabolism	glucosamine-6-phosphate	0.94	0.911	1.33	0.1329
	glucuronate	0.83	0.007	0.6	1.18E-09
	N-acetylglucosamine	0.76	0.0519	0.8	0.1516
	N-acetylglucosamine 6-phosphate	0.68	2.24E-05	0.84	0.0424

807
808

Supplemental file 1. Summary of carbohydrate metabolites in TDP-43[WT] and TDP-43[G298S]

compared to w[1118] controls. Altered metabolites in third instar larvae crossed with the motor

neuron driver D42 GAL4 were measured using gas or liquid chromatography followed by mass

spectrometry. Red and green colored cells indicate statistically significant changes ($P_{value} < 0.05$)

that are increased and decreased, respectively. Light red and light green colored cells indicate

upward or downward trends, respectively ($P_{value} < 0.1$).

Group/ID	Race	Gender	Age	PMI (hrs)
CON 1	W	M	22	6
CON 2	-	F	71	5
CON 3	-	M	72	7
CON 4	W	F	57	32
CON 6	W	F	53	4
CON 7	W	F	51	5
CON 8	W	F	57	11
CON 9	W	M	76	13
ALS 1	W	F	68	1
ALS 2	W	M	69	10
ALS 3	W	F	73	11
ALS 4	W	M	53	-
ALS 5	W	F	62	4
ALS 6	W	M	53	5
ALS 7	W	F	76	6
ALS 8	W	F	63	5
ALS 9	W	M	67	8

815 PMI = post-mortem interval; CON = Non-neurologic disease control

816 **Supplemental file 2.** Summary of demographic information for patient samples used to quantify

817 PFKP, PFKM and G6PD.

Cell line number	Passage number	Days in culture
ALS 1	39	69
ALS 2	35	70
ALS 3	36	77
Control 1	19	73
Control 2	28	74
Control 3	37	71
Control 4	21	70

820 **Supplemental file 3. Summary of iPSC MNs used to quantify PFKP, PFKM and G6PD.**

821 Patient cell lines used for qPCR analysis are shown.