MUTATION OF *MLH3* ENDONUCLEASE MOTIF REVEALS INTEGRATION BETWEEN CROSSOVER PATHWAYS IN MAMMALIAN MEIOSIS

Melissa Toledo[1,3,4], Xianfei Sun[1,4], Miguel A. Brieño-Enríquez[1,3, 5], Vandana Raghavan[2], Stephen Gray[1,3], Jeffrey Pea[1,3], Anita Venkatesh[2], Lekha Patel[1], Peter L. Borst[1,3], Eric Alani[2] and Paula E. Cohen[1,3,*].

Running Title: MLH3 endonuclease activity is critical for meiosis

Keywords: meiosis, germ cell, spermatocyte, MutL, MLH1, Crossover, Recombination

[1]Departments of Biomedical Sciences and [2]Molecular Biology and Genetics, and [3]The Center for Reproductive Genomics, Cornell University, Ithaca, NY

[4]Equal contributors

[5] Current address: Magee Womens Research Institute and University of Pittsburgh, Pittsburgh, PA

*Correspondence: paula.cohen@cornell.edu; (607) 280 2185

SUMMARY (40 words)

The MLH1-MLH3 complex is essential for crossing over in mammalian meiosis. We generated a mutation in mouse MLH3 that disrupts its conserved endonuclease domain and show that it disrupts crossing over in a manner distinct from the null, but also results in male infertility.

ABSTRACT

During meiotic prophase I, double strand breaks (DSBs) initiate homologous recombination leading to non-crossovers (NCOs) and crossovers (COs). In mouse, 10% of DSBs are designated to become COs, primarily through a pathway dependent on the MLH1-MLH3

10   heterodimer (MutLγ). Mlh3 contains an endonuclease domain that is critical for resolving COs in yeast. We generated a mouse *Mlh3^{DN}* allele harboring a mutation within this conserved domain that is predicted to generate a protein that is catalytically inert. *Mlh3$^{DN/DN}$* males, like fully null *Mlh3$^{-/-}$* males, have no spermatozoa and are infertile, yet spermatocytes have normal DSBs and undergo normal synapsis events in early prophase I. Unlike *Mlh3$^{-/-}$* males, however,

15   mutation of the endonuclease domain within MLH3 permits normal loading and frequency of MutLγ in pachynema. However, DSB repair and CO designation factors persist in *Mlh3$^{DN/DN}$* males, indicating a temporal delay in repair events. While *Mlh3$^{DN/DN}$* spermatocytes retain only 22% of wildtype chiasmata counts, this frequency is greater than observed in *Mlh3$^{-/-}$* males (10%), suggesting that the allele may be partially functional or that other pathways can

20   generate COs from these MutLγ-defined repair intermediates in *Mlh3$^{DN/DN}$* males, with evidence favoring the latter option. Double mutant mice that are homozygous for the *Mlh3$^{DN/DN}$* mutation along with a null allele of *Mus81*, show losses in chiasmata approaching levels observed in *Mlh3$^{-/-}$* males, suggesting that the MUS81-EME1-regulated crossover pathway accounts for some of the increased residual chiasmata observed in the *Mlh3$^{DN/DN}$* spermatocytes. These

25   results demonstrate that an intact MLH3 endonuclease domain is essential for most COs in

mammalian meiosis, and that an endonuclease-impaired MutLγ retains the ability to facilitate

the recruitment of other repair pathways, including MUS81-EME1.

INTRODUCTION

30   Meiosis is a specialized cell division process in which a diploid parental cell undergoes

one round of DNA replication followed by two rounds of division, resulting in up to four haploid

gametes. Successful halving of the genome during meiosis I depends on the tethering of

maternal and paternal homologous chromosomes during meiotic prophase I, and their

subsequent release at the first meiotic division. This tethering is ensured by homologous

35 recombination (leading to the formation of crossovers), by synapsis (the formation of a tripartite

proteinaceous structure, the synaptonemal complex, or SC) between homologous

chromosomes, and by cohesion (between replicated sister chromatids that ensures

appropriate tension on the metaphase I spindle) (Gray and Cohen 2016; Hunter 2015). Thus,

recombination and synapsis are hallmarks of prophase I, and are both essential for ensuring

40 homolog interactions leading to the formation of at least one crossover event per chromosome

pair. Moreover, the correct placement, frequency, and distribution of crossovers is critical for

ensuring appropriate disjunction at metaphase I and for maintaining genomic stability (Gray

and Cohen 2016; Bolcun-Filas and Handel 2018).

   Meiotic recombination begins with the introduction of a large number of programmed

45 double-strand breaks (DSBs), which are repaired as non-crossovers (NCOs) or crossovers

(COs). Evidence for distinct NCO versus CO pathways was obtained in *S. cerevisiae*, where it

was shown that the former occur earlier in meiotic prophase I, and subsequent work suggested

that they appeared primarily through synthesis-dependent strand annealing (SDSA)(McMahill

et al. 2007; Allers and Lichten 2001). In *M. musculus*, 90% of DSBs are repaired as NCOs,

50 presumably via SDSA or other pathways, while 10% of DSBs are repaired as COs (Holloway et

al. 2008; Guillon et al. 2005; Svetlanov et al. 2008; Cole et al. 2010).

COs can form via one of at least two distinct mechanisms (referred to as class I and class II), each of which is used in varying degrees in different eukaryotic organisms (de los Santos et al. 2003; Higgins et al. 2008; Holloway et al. 2008). The class I CO pathway is also known as the ZMM pathway, named after the major genes discovered in yeast that regulate this mechanism (Lynn et al. 2007; Jessop et al. 2006; Börner et al. 2004; Wang et al. 2009; Agarwal and Roeder 2000; Hollingsworth et al. 1995; Tsubouchi et al. 2006). Class II COs, on the other hand, do not involve the ZMM proteins, but instead appear to rely on the structure-specific endonuclease (SSN), MUS81/EME1 (Mus81/Mms4 in *S. cerevisiae*) (de los Santos et al. 2003; Higgins et al. 2008; Holloway et al. 2008). Class I COs differ from class II COs in that the former are regulated by interference, the process by which placement of one CO prevents the nearby localization of a second CO, thus resulting in appropriate spacing of these events across the genome (de Boer et al. 2006).

In the class I CO pathway, DSBs are processed and resected to form single-end invasion (SEI) intermediates. This is followed by displacement of single strand DNA (ssDNA) from the recipient homolog to produce a double Holliday junction (dHJ). The yeast ZMM proteins Msh4 and Msh5 form a complex known as MutSγ that associates with a subset of these intermediate structures (Novak et al. 2001; Nishant et al. 2010; Pochart et al. 1997). At least in yeast, this recruitment may be dependent on the STR complex, consisting of Sgs1 (BLM in mammals), Top3 and Rmi1 (Kaur et al. 2015; Tang et al. 2015). STR is proposed to act by disassembling the early recombination intermediates that would otherwise be processed through SSN-directed recombination pathways, thereby promoting either early NCO formation via SDSA, or CO formation through the capture of these recombination intermediates by the ZMM proteins, including MutSγ (Kaur et al. 2015). MutSγ is then thought to stabilize the dHJs, leading to the recruitment of a second MMR complex, MutLγ, consisting of the MutL homologs,

Mlh1 and Mlh3 (Hunter and Borts 1997; Wang et al. 1999). Similar to the situation in yeast, the mouse MutSγ complex associates with chromosome cores in zygonema at a frequency of approximately 150 foci per nucleus (Kneitz et al. 2000), recruiting the MutLγ complex in pachynema. However, unlike yeast, MutLγ associates with only a subset of MutSγ sites (~24-26 foci/nucleus), designating these events as class I COs (Kolas et al. 2005; Lipkin et al. 2002).

Though not formerly considered to be ZMM proteins, MLH1 and MLH3 are critical for class I CO events in numerous organisms (Al-Sweel et al. 2017; Woods et al. 1999; Wang et al. 1999; Lipkin et al. 2002; Nishant et al. 2008; Argueso et al. 2002; Rogacheva et al. 2014). In fact, the *M. musculus* MLH1-MLH3 heterodimer localizes to sites that are destined to become class I COs and the absence of either subunit in male spermatocytes leads to a dramatic decrease, but not complete absence, of chiasmata (the physical manifestation of a CO) (Kan et al. 2008; Kolas et al. 2005; Marcon and Moens 2003; Lipkin et al. 2002; Baker et al. 1996; Anderson et al. 1999). While MutLγ is known to be recruited to sites that are preloaded with MutSγ, recent studies have shown that *S. cerevisiae* MutLγ can bind to single and double-stranded DNA (ssDNA, dsDNA), as well as a variety of branched DNA structures (Rogacheva et al. 2014; Manhart et al. 2017; Claeys Bouuaert and Keeney 2017; Ranjha et al. 2014). How such binding properties relate to the *in vivo* functions of MutLγ remains unclear.

Class I CO formation in *M. musculus* is dependent on MLH3 (Lipkin et al. 2002; Svetlanov et al. 2008; Kan et al. 2008). Curiously, MLH3 is an essential binding partner for MLH1, but its recruitment precedes that of MLH1 (Kolas et al. 2005). Further analysis of MutLγ has shown that MLH3 contains a conserved metal binding motif, $DQHA(X)_2E(X)_4E$, originally discovered in the human MutL homolog, PMS2, and found to be required for human MutLα (hMLH1/hPMS2) endonuclease function (Kadyrov et al. 2006). This putative endonuclease

100   motif is highly conserved in eukaryotic homologs of human PMS2 and MLH3, but not in

homologs of human MLH1 and PMS1. The expectation for MLH3 is that this endonuclease

function might represent a "resolvase" activity for class I COs. Studies in *S. cerevisiae* have

shown that a single point mutation in the endonuclease motif of yeast Mlh3 (*mlh3-D523N*)

disrupts its endonucleolytic activity and results in meiotic crossover defects similar to full *mlh3*

105   (*mlh3Δ*) null mutants, yet does not affect the protein stability of Mlh3 or its interaction with Mlh1

(Nishant et al. 2008). Further analysis of the entire endonuclease domain in *S. cerevisiae*

revealed that mutation of any conserved residue results in a null or near-null phenotype with

respect to crossing over (Al-Sweel et al. 2017). Biochemical analysis reveals that the Mlh1-

mlh3D523N protein lacks the ability to nick closed circular double stranded DNA, indicating

110   loss of endonuclease activity (Rogacheva et al. 2014; Ranjha et al. 2014). Collectively, these

studies in *S. cerevisiae* suggest that MutLγ plays a direct role in resolving dHJs to generate

COs through its endonuclease activity.

To investigate the function of the putative endonuclease domain of MLH3 in mammalian

meiotic recombination, we generated a point mutant mouse (termed *Mlh3^{DN}*) in which the

115   endonucleolytic domain was disrupted at the orthologous residue to the D523N mutation in

yeast, allowing the overall structure of MLH3 to remain intact, as determined by the ability to

form a stable complex with MLH1. By generating a catalytically defective protein, we

hypothesized that the mutant MutLγ complex would remain structurally intact and thus might

reveal a functional interplay with other meiotic CO functions. For example, such a mutant might

120   reveal alternative CO resolvase pathways that could be recruited in the absence of the MutLγ

resolvase activity. We demonstrate that normal function of the MLH3 endonuclease domain is

required for resolution of DSB repair intermediates towards CO formation and thus for late

meiotic recombination events. *Mlh3$^{DN/DN}$* spermatocytes exhibit normal DSB formation and

early processing, and normal synapsis through early prophase I. *Mlh3*$^{DN/DN}$ spermatocytes

exhibit appropriate localization of MLH3 and MLH1 to the synaptonemal complex during

pachynema, along with pro-crossover factors, HEI10 and CDK2, phenotypes that are clearly

different to that observed in *Mlh3*$^{-/-}$ males. However, *Mlh3*$^{DN/DN}$ diakinesis-staged

spermatocytes show significantly fewer chiasmata compared to wild-type mice (WT), but

significantly more when compared to *Mlh3*$^{-/-}$ males, suggesting either that the MLH3^{DN} protein

retains partial endonucleolytic activity, or that the presence of the MutLγ complex, albeit altered

in its endonucleolytic capacity, can invoke other repair pathways to become active.

Interestingly, in line with this suggestion, we find that the RecQ helicase, BLM, is upregulated

throughout prophase I in *Mlh3*$^{DN/DN}$ spermatocytes, perhaps aiding the recruitment of other

repair proteins. To explore the increase in residual chiasmata observed at diakinesis in

Mlh3$^{DN/DN}$ males relative to that of *Mlh3*$^{-/-}$ males, we demonstrate that co-incident loss of the

class II CO pathway in *Mus81*$^{-/-}$*Mlh3*$^{DN/DN}$ double mutant males results in altered distribution of

MutLγ resulting in an increased proportion of synapsed autosomes bearing no MutLγ foci.

Furthermore, the proportion of chiasmata remaining in these double mutants is between that of

Mlh3$^{DN/DN}$ and *Mlh3*$^{-/-}$ males, suggesting that MUS81-EME1 accounts for a significant

proportion of these additional chiasmata. Collectively, our data show that the endonucleolytic

activity of MLH3 is important for normal processing of DSB repair intermediates through the

Class I pathway, and suggests that the presence of the MutLγ complex can facilitate the

recruitment of other repair pathways to achieve optimal CO frequency and distribution.

RESULTS

***Mlh3*$^{DN/DN}$ males are infertile**

To investigate the meiotic requirement for the presence of a functional endonuclease domain in mammalian MLH3, we generated a mouse line with a point mutation in a conserved endonuclease motif located in the *M. musculus* protein: DQHAAHERIRLE (Gueneau et al., 2013; Kadyrov et al., 2006). Specifically, we replaced the aspartic acid "D" in amino acid position 1185, with an asparagine "N" by changing GAC to AAC in the genomic sequence, termed MLH3^{DN} throughout. Extrapolating from an analogous mutation in the *S. cerevisiae* gene, this D-to-N replacement is predicted to disrupt the endonucleolytic function of MLH3 while maintaining the ability to interact stably with MLH1 ((Nishant et al., 2008); Figure S1). Mice were maintained on a C57Bl/6J background throughout the study.

Male *Mlh3$^{+/DN}$* mice were phenotypically similar to WT littermates and displayed full fertility. *Mlh3$^{DN/DN}$* males are also grossly normal when compared to WT littermates, survive into adulthood, and live normal lifespans. *Mlh3$^{DN/DN}$* males also exhibit normal mating behaviors as determined by observing a vaginal plug in WT females the morning after mating. However, breeding between multiple sets of *Mlh3$^{DN/DN}$* males and WT females was never observed to result in offspring over a four-year period.

Similar to the situation seen for *Mlh3$^{-/-}$* males (Lipkin et al., 2002), *Mlh3$^{DN/DN}$* males show complete infertility, demonstrated by significantly reduced testes size when compared to WT (Figure 1A, B; $p < 0.0001$) and the absence of spermatozoa in the epididymides (Figure 1C; $p < 0.0001$). Whereas histological cross-sections of testes stained with hemotoxylin and eosin from WT males showed the presence of meiotic and post-meiotic cells within the seminiferous epithelium, testis sections from *Mlh3$^{DN/DN}$* males were devoid of spermatids, but showed the presence of spermatogonia and spermatocytes (Figure 1D-G). In addition, metaphase I spermatocytes were observed in the tubular lumen of *Mlh3$^{DN/DN}$* mice (Figure 1G, black

170   arrows). Thus, mutation of the endonuclease domain of *Mlh3* in the mouse results in a sterility

phenotype grossly similar to that seen in *Mlh3*$^{-/-}$ mice.

Mlh3$^{DN/DN}$ spermatocytes, like those of _Mlh3_$^{-/-}$ mice, exhibit normal DSB formation and

synapsis

175   To investigate the progression of meiotic recombination, prophase I chromosome

spreads were prepared from WT, *Mlh3*$^{DN/DN}$, and *Mlh3*$^{-/-}$ adult males and stained for a variety of

markers involved in synapsis and recombination. Chromosome spreads were stained with

antibodies against γH2AX, the phosphorylated form of histone H2AX, as a marker of DSBs

(Viera et al. 2004; Mahadevaiah et al. 2001). In spermatocyte preparations from WT males,

180   γH2AX signal is abundant throughout the nucleus at leptonema, coincident with the induction of

several hundred DSBs (Mahadevaiah et al. 2001; Gray and Cohen 2016). The γH2AX signal

declines in zygonema as DSBs are processed for repair (Mahadevaiah et al. 2001; Hunter et

al. 2001). In pachynema and diplonema, γH2AX signal is absent from the autosomes, but

emerges throughout the sex body due to meiotic sex chromosome inactivation (MSCI)

185   ((Handel 2004); Figure S2A-D). Spermatocytes from both *Mlh3*$^{DN/DN}$ and *Mlh3*$^{-/-}$ males exhibit

the same γH$_2$AX signal and temporal dynamics as observed in WT spermatocytes, with

abundant staining in leptonema, slightly reduced signaling in zygonema, followed by the

absence of γH2AX signal on the autosomes of pachytene and diplotene spermatocytes, except

at the sex body (Figure S2F-I, K-N). We do not see specific persistent γH2AX signal on the

190   autosomes at pachynema in *Mlh3*$^{-/-}$ spermatocytes, as reported by other groups (Qiao et al.

2014), unless we markedly increase our imaging exposure time γH2AX (Figure S2E, J, O;

white arrows). Under these conditions, we see persistent foci of γH2AX in spermatocytes from

WT and from *Mlh3*$^{DN/DN}$ spermatocytes also. Thus, in our hands, we see no specific

persistence in autosomal γH2AX signal through pachynema in mice lacking MLH3 or harboring

195   a mutation within the endonuclease domain of MLH3.

Spermatocyte chromosome spreads from WT and *Mlh3^DN/DN^* males were stained with

antibodies against synaptonemal complex (SC) components, SYCP3 and SYCP1, marking the

axial/lateral elements and the central element, respectively. Prophase I progression in WT

spreads is characterized by the initial accumulation of SYCP3 signal in discrete dots along

200   chromosomes at leptonema, and these dots gradually coalesce into continuous filaments along

the chromosome cores in zygonema (Figure S2Q). At this time, SYCP1 appears in patches

along the SYCP3 signal, indicating that synapsis is occurring. By late zygonema, most of the

chromosome core is now labeled with SYCP1, and by pachynema synapsis is complete, as

demonstrated by complete overlap of the SYCP3/SYCP1 signals on the autosomes. For the

205   sex chromosomes, synapsis only occurs at the pseudoautosomal region (PAR). After meiotic

recombination occurs, the SC begins to degrade in diplonema, and the homologs are no longer

tethered to one another except at CO sites (Figure S2P-S).

Synapsis appears normal in *Mlh3^DN/DN^* spermatocytes with discrete accumulation of

SYCP3 on the chromosomes in leptonema, followed by continued accumulation of SYCP3

210   along the chromosomes as SYCP1 appears in patches in zygonema (Figure S2T, U).

Complete synapsis of the autosomes and the PAR is observed in pachynema with co-

localization of SYCP1 and SYCP3 (Figure S2V). Desynapsis is then observed in diplonema

with the degradation of the SC (Figure S2W). Thus, synapsis in *Mlh3^DN/DN^* spermatocytes

appears unaffected by loss of the endonuclease activity of MLH3, a result similar to that seen

215   for complete loss of MLH3 protein.

Mlh3^DN/DN^ spermatocytes show a persistence of RAD51 in pachytene cells

Early DSB repair events were monitored by examining and quantifying localization of the RecA strand exchange protein, RAD51, on chromosome cores of the autosomes throughout prophase I (Ashley et al. 1995). In WT mice, RAD51 localizes to chromosome cores of early and late zygotene cells as discrete foci at a high frequency (EZ and LZ, respectively; Figure 2A, G). Compared to WT littermates in early and late zygonema, RAD51 counts in spermatocytes from $Mlh3^{DN/DN}$ males were significantly elevated (Figure 2C,G; $p<0.001$ and $p<0.01$, respectively, by unpaired t-test with Welch's correction). However, while early zygotene RAD51 counts were indistinguishable in $Mlh3^{-/-}$ spermatocytes compared to both WT (Figure 2E,G), they were significantly lower than that seen at the equivalent stage in $Mlh3^{DN/DN}$ males ($p<0.001$ by unpaired t-test with Welch's correction). By late zygonema, the RAD51 counts in were significantly lower in $Mlh3^{-/-}$ spermatocytes compared to WT and $Mlh3^{DN/DN}$ animals ($p<0.001$ by unpaired t-test with Welch's correction).

By pachynema, RAD51 foci frequency in spermatocytes from WT decreased to very low numbers, as did that of $Mlh3^{-/-}$ males (Figures 2B,F,H; $p=0.55$ unpaired t-test). In contrast, focus counts in pachytene spermatocytes from $Mlh3^{DN/DN}$ males were significantly elevated (Figure 2D,H; $p<0.0001$). These observations suggest a persistence of DSB repair intermediates loaded with RecA homologs in $Mlh3^{DN/DN}$ spermatocytes in pachynema. We hypothesize, based on these observations, that RAD51 accumulation in zygonema is affected by loss or mutation of MLH3 protein, suggesting an early function for MutLγ in establishing appropriate DSB repair intermediates that is not confined to CO pathway fate. Further, we hypothesize that DSBs are repaired in a timely fashion in mice lacking MLH3 protein, perhaps through repair pathways that differ from those utilized in WT-derived spermatocytes. In contrast, DSBs in $Mlh3^{DN/DN}$ mutants are not repaired efficiently, resulting in a persistence of RAD51 foci on DSBs through to pachynema.

Mlh3^{DN/DN} pachytene spermatocytes show a hyper-accumulation and persistence of BLM

245   Bloom's syndrome mutated (BLM) is a mammalian RecQ DNA helicase whose _S. cerevisiae_ ortholog, Sgs1, was shown to promote the resolution of complex multi-chromatid joint molecule intermediates that may result from SEI events into both NCOs and COs (Tang et al. 2015; Kaur et al. 2015). During prophase I in WT male spermatocytes, BLM localizes to the chromosomal cores at a high frequency in zygonema and diminishes to a few foci in

250   pachynema (Moens et al. 2000; Walpita et al. 1999; Holloway et al. 2010). Recently, we showed that loss of MLH3 results in up-regulated BLM localization during prophase I, along with persistence of BLM on chromosome cores through late pachynema (Holloway et al. 2010). To determine if the disruption of the MLH3 endonuclease domain affects the localization of BLM in a similar fashion, we stained prophase I chromosome spreads with an antibody against

255   BLM. In zygonema, as previously reported, WT cells show the accumulation of BLM foci on the cores in high numbers, and this frequency is elevated in spermatocytes from both _Mlh3_^{DN/DN} and _Mlh3_^{-/-} spermatocytes (Figure 3A,B,E,F,I,J,M; p<0.0001 unpaired t-test). This is similar to that reported previously for _Mlh3_^{-/-} spermatocytes (Holloway et al. 2010). In early to mid-pachynema, BLM localization on chromosome cores persists in a small percentage of WT

260   spermatocytes, but the number of foci is very much reduced at this stage (Figure 3C,D,M). In contrast, all spermatocytes from _Mlh3_^{DN/DN} and _Mlh3_^{-/-} spermatocytes show persistent BLM focus localization along chromosome cores (Figure 3G,H,K,L) at a frequency that is elevated above that of WT spermatocytes (Figure 3M, p<0.0001 unpaired t-test). Moreover, the number of BLM foci in _Mlh3_^{DN/DN} spermatocytes is significantly elevated relative to that seen in _Mlh3_^{-/-}

265   spermatocytes (Figure 3M, p<0.05 unpaired t-test). By late pachynema to early diplonema this

difference was even greater, with BLM localization in *Mlh3$^{DN/DN}$* spermatocytes persisting, and

being lost in *Mlh3$^{-/-}$* spermatocytes by diplonema (Figure 3D,H,L). Thus, altered MLH3

endonuclease function, like complete loss of MLH3, leads to persistence of BLM helicase on

chromosome cores in late prophase I, but at an elevated frequency in *Mlh3$^{DN/DN}$* spermatocytes

270   relative to *Mlh3$^{-/-}$* spermatocytes.

Normal localization of crossover designation factor MSH4 *Mlh3$^{-/-}$* and *Mlh3$^{DN/DN}$* adult

male mice during prophase I.

      "Crossover designation" is defined as the process by which class I COs are selected

275   from an excess pool of DSB repair intermediates. In mouse, the 250+ DSBs are processed

through zygonema into various repair pathways, and only a subset of these will proceed

towards a class I CO fate (Gray and Cohen 2016). These sites become "licensed" for crossing

over through the accumulation of the MutS homolog heterodimer, MutSγ (MSH4 and MSH5;

(Edelmann et al. 1999; Kneitz et al. 2000)). The MutSγ complex then serves as an early pro-

280   crossover factor by recruiting the MutLγ complex to a select subset of sites and it is these sites

that will become "designated" as class I CO events. Notably, all of the 150+ MutSγ sites must

be repaired, either as a CO or an NCO, which means that approximately ~125 MutSγ sites

must leave the class I CO pathway and undergo repair through an alternate CO pathway or via

an NCO pathway, a situation that is unlike that seen in *S. cerevisiae* where the number of

285   MutSγ sites appear to correspond more closely to the number of CO events (Novak et al.

2001).

      To investigate how loss of MLH3 endonucleolytic function affects the paring down of

MutSγ sites through zygonema and pachynema, we explored MSH4 focus dynamics on

chromosome spreads throughout prophase I from WT, *Mlh3$^{DN/DN}$*, and *Mlh3$^{-/-}$* adult male mice

290 (Figure S3). We observed a similar paring down of MSH4 focus counts from early pachynema

(EP) to late pachynema (LP) in spermatocytes from WT, $Mlh3^{DN/DN}$, and $Mlh3^{-/-}$ adult males. In

all three cases, MSH4 foci appear on chromosome cores in zygonema (Figure

S3A,B,E,F,I,J,M), persisting at high levels in early pachynema (Figure S3C,G,K), and then

undergoing reduction to approximately 1-2 foci per chromosome in late pachynema (Figure

295 S3D,H,L,M). Quantification of MSH4 focus numbers in early and late pachytene spermatocytes

from WT, $Mlh3^{DN/DN}$, and $Mlh3^{-/-}$ adult male reveals the expected statistically significant decline

in these foci EP to LP (Figure S3M; $p < 0.0001$ Mann-Whitney U Test for all).

Normal localization of Class I Pro-CO factors in pachytene spermatocytes from

300 **$Mlh3^{DN/DN}$ adult male mice.**

To observe class I CO events in pachynema, we employed two well characterized

markers of these sites: the putative ubiquitin E3 ligase, Human Enhancer of Invasion-10

(HEI10), and cyclin-dependent kinase-2 (CDK2) (Ward et al. 2007; Qiao et al. 2014; Ashley et

al. 2001). In WT prophase I cells, CDK2 localizes to the telomeres (Figure 4A, yellow arrows)

305 as well as on the chromosome cores (Figure 4A, white arrows) during mid to late pachynema

and remains associated with SYCP3 signal through to diplonema (Ashley et al. 2001). The

localization of CDK2 along chromosome cores parallels the localization of MLH1 and MLH3,

and is associated with nascent class I CO events. In pachytene spermatocyte preparations

from $Mlh3^{DN/DN}$ mice, CDK2 signal remains associated with both the telomeres and

310 chromosome cores at a frequency and intensity that is reminiscent of that seen in WT

spermatocyte spreads (Figure 4B). This is in contrast to the situation in spermatocyte

preparations from $Mlh3^{-/-}$ males, in which CDK2 association with the telomere persists, but is

lost from nascent CO sites (Figure 4C).

HEI10 was recently shown to co-localize with MutLγ at sites of class I CO, and its

315   localization is dependent on Cyclin N-terminal Domain-containing-1 (CNTD1)(Holloway et al.

2014; Qiao et al. 2014). HEI10 is thought to play a key role in CO designation/maturation (Qiao

et al. 2014). As previously reported for WT cells at pachynema, HEI10 co-localizes with CDK2

and MutLγ (Figure 4D, pink arrows). Similar localization patterns and frequency were observed

for *Mlh3*[DN/DN] mice, with a frequency of one to two foci per chromosome (Figure 4E, pink

320   arrows), indicating normal recruitment of HEI10 on pachytene chromosome cores in *Mlh3*[DN/DN]

males. This is in contrast to the pattern of HEI10 staining in spermatocytes from *Mlh3*[-/-] mice,

where there is an increased accumulation of HEI10 foci (Figure 4F, pink arrows), as previously

reported (Qiao et al. 2014). Taken together, these observations demonstrate that loading of

HEI10 and CDK2 on class I CO designated sites is affected differently by mutation of *Mlh3:*

325   complete loss of MLH3 results in failure to load CDK2 and hyper-accumulation of HEI10, while

altered endonucleolytic activity of MLH3 results in normal loading of both CDK2 and HEI10.

Thus, the physical accumulation of MutLγ is required for normal loading of associated pro-

crossover maturation factors.

330   ***Mlh3*[DN/DN] pachytene staged spermatocytes exhibit normal localization of MLH3 and
MLH1.**

MutLγ represents the ultimate marker of DSB repair events that have adopted a class I

CO fate, and has been used as a CO proxy marker in many organisms (Lenzi et al. 2005;

Auton et al. 2013; Lipkin et al. 2002; Jean et al. 1999). We anticipated that the endonuclease

335   mutation in MLH3 would not affect localization of this complex. In WT spermatocytes, MLH3

localizes on the chromosomes during early pachynema, remaining associated with SYCP3

signal through to diplonema (Figure 5A). In pachytene spermatocyte preparations from

$Mlh3^{DN/DN}$ mice, MLH3 signal remains associated with the autosomal chromosome cores from early pachynema at a focus frequency that is statistically indistinguishable from that of WT

340 cells (Figure 5A-C, p=0.36 by unpaired t-test). MLH3 association with the PAR of the synapsed X and Y chromosomes was similarly unaffected in $Mlh3^{DN/DN}$ pachytene spermatocytes (data not shown). In addition, the timing of MLH3 appearance, in early pachynema and prior to that of MLH1, was normal in $Mlh3^{DN/DN}$ pachytene spermatocytes.

Localization of MLH1 was similarly explored in spermatocytes from $Mlh3^{DN/DN}$ mice. As

345 with MLH3, there was no difference in the timing of MLH1 accumulation on chromosome cores between WT and $Mlh3^{DN/DN}$ mice (Figure 5D-F). Moreover, when autosomal MLH1 foci were quantified, no statistical difference was observed in MLH1 focus frequency between WT and $Mlh3^{DN/DN}$ pachytene cells (Figure 5D-F, p= 0.2 by unpaired t-test). These data suggest that disruption of the endonuclease domain of MLH3 does not alter recruitment of MutLγ to

350 chromosomes in pachynema.

MLH1-MLH3-D1185N forms a stable heterodimer and displays similar chromatographic properties to MLH1-MLH3.

Mouse *Mlh1* and *Mlh3* were amplified from cDNA and cloned into pFastBac1 vectors as

355 described in the Methods. The MLH1-MLH3 and MLH1-MLH3-D1185N complexes were expressed from Sf9 cells infected with baculoviruses containing *MBP-Mlh1* and *His_{10}-Mlh3* or *His_{10}-Mlh3-D1185N* constructs (Figure 5G). Extracts from these cells were applied to a Ni-NTA column. Fractions containing induced proteins were pooled and then applied to an amylose column. Two major bands of molecular weights predicted for an MBP-MLH1-His_{10}-MLH3

360 complex were detected on SDS-PAGE after amylose chromatography (Figure 5H). These bands were further analyzed by mass spectrometry, and the results from this analysis

confirmed their identity (Figure 5I). Importantly, MLH1-MLH3 and MLH1-MLH3-D1185N eluted with an apparent 1:1 stoichiometry in both chromatography steps, indicating that the heterodimers were stable, and the protein yields of the two complexes after amylose

365   chromatography were similar (Figure 5H).

Mlh3$^{DN/DN}$ diakinesis staged spermatocytes exhibit significantly fewer chiasmata than WT spermatocytes, but elevated chiasmata compared to _Mlh3_$^{-/-}$ males.

Chiasmata are the physical manifestations of crossing over and, as such, can inform the

370   process of DSB repair via all pathways. Diakinesis-staged spermatocytes from WT and _Mlh3_$^{DN/DN}$ males were used to quantify chiasmata. WT cells exhibited a chiasmata frequency of 23.5 ±1.3 per nucleus (Figure 6A, D) whereas _Mlh3_$^{DN/DN}$ spermatocytes exhibited a dramatically reduced chiasmata count of 5.2 ± 1.7 chiasmata per nucleus (Figure 6B, D; p < 0.0001 by unpaired t-test). Chiasmata counts for _Mlh3_$^{-/-}$ males were even more dramatically

375   reduced at 2.8 ± 1.1 chiasmata per nucleus, a value that is significantly lower than both WT and _Mlh3_$^{DN/DN}$ spermatocytes (Figure 7C, D; p <0.0001 by unpaired t-test). Thus, complete loss of MLH3 protein leads to the loss of approximately 88% of chiasmata, while loss of endonuclease activity, but retention of MutLγ heterodimer results in only a 78% loss. Thus, the number of residual chiasmata observed in _Mlh3_$^{DN/DN}$ spermatocytes is higher than the

380   expected number of chiasmata achieved through the MUS81-EME1-driven class II CO pathway (assessed both cytoglogically and genetically; (Svetlanov et al. 2008; Kolas et al. 2005)).

Co-incident loss of *Mus81* on the *Mlh3^{DN/DN}* background results in altered MutLγ

distribution at pachynema and a partial reduction in the residual chiasmata observed in

Mlh3^{DN/DN} single null males

The increased residual chiasmata observed in *Mlh3^{DN/DN}* males compared to *Mlh3^{-/-}* animals

prompted us to ask whether the activity of the MUS81-EME1 heterodimer might be required for

these additional CO events. Previous studies in our lab showed that *Mus81^{-/-}* animals show

increased accumulation of MutLγ, resulting in normal chiasmata counts, suggesting that class I

CO events are up-regulated in the absence of the class II machinery (Holloway et al. 2008).

However, in the current study, co-incident mutation of one or both *Mlh3* alleles to the *Mlh3^{DN}*

variant on the *Mu81^{-/-}* mutant background yielded MLH1 focus counts that were not elevated

compared to those observed in *Mus81^{-/-}Mlh3^{+/+}* mice (Figure 6E, p<0.005 by unpaired t-test

with Welch's correction). Thus, the upregulation of MLH1 foci at pachynema requires both a

defective MUS81-EME1 dimer and a functional MutLγ heterodimer. Interestingly, pachytene

spermatocytes from *Mus81^{-/-}Mlh3^{+/DN}* and *Mus81^{-/-}Mlh3^{DN/DN}* males also show an abnormal

distribution of MLH1 foci across all autosomal pairs (Figure 6F). In *Mus81^{-/-}Mlh3^{+/+}* males, 68%

(21/31) cells show no synapsed autosomes without MLH1 foci (so-called "no exchange" or "E0"

chromosomes), while in *Mus81^{-/-}Mlh3^{+/DN}* males, this number dropped to 25% (5/20), with up to

3 E0 chromosomes per cell (Figure 6F). In *Mus81^{-/-}Mlh3^{DN/DN}* males, the number of synapsed

autosomes lacking a MLH1 focus was more significantly increased, with only 13% of cells

having no E0 chromosomes, and as many as 6 E0 bivalents being evident (Figure 6F). The

increasing frequency of MLH1-devoid autosomes in *Mlh3^{+/DN}* and *Mlh3^{DN/DN}* males on the

Mus81 null background was statistically significant in all pairwise comparisons (Figure 6F,

unpaired t-test with Welch's correction), perhaps indicating that interference is perturbed in

these double mutant animals. In yeast, and also probably in mice, CO interference requires a

functional MutSγ complex (Shinohara et al. 2008). However, MutSγ alone is not sufficient to ensure appropriate interference since $Mus81^{-/-}$ males exhibit disrupted interference despite appropriate MutSγ loading (Holloway et al. 2008).

Assessment of chiasmata counts in single (Figure 6D) and double mutants (Figure 6G) revealed the expected normal chiasmata frequency in spermatocytes from $Mus81^{-/-}Mlh3^{+/+}$ and $Mus81^{-/-}Mlh3^{+/DN}$ males, and the loss of most chiasmata in spermatocytes from $Mus81^{-/-}Mlh3^{DN/DN}$ males. However, whereas the loss of chiasmata structures in $Mlh3^{-/-}$ and $Mlh3^{DN/DN}$ single mutants was observed to be 88% and 78%, respectively (Figure 6D), the loss of chiasmata in $Mus81^{-/-}Mlh3^{DN/DN}$ males was 83%. The frequency of chiasmata in cells from these double mutants was statistically different from both $Mlh3$ single homozygous mutant animals (p<0.05, unpaired t-test with Welch's correction). Thus, loss of the class II pathway in addition to the mutation of the endonuclease domain of $Mlh3$ only partially reduces residual chiasmata counts to the levels observed in $Mlh3^{-/-}$ animals. Interestingly, despite reduced chiasmata, the overall number of bivalent structures observed in diakinesis preparations from $Mus81^{-/-}Mlh3^{DN/DN}$ males is the same as that seen in $Mlh3^{DN/DN}$ males, and is significantly elevated above that seen in $Mlh3^{-/-}$ single mutant males (Figure 6H). Taken together, these observations suggest three important features regarding class I/II interactions: (1) that the increased MutLγ localization observed in $Mus81^{-/-}$ males reported in previous publications (and confirmed here) is dependent on a fully functional MutLγ complex; (2) that a fully functional class I and class II machinery is required for appropriate distribution of MutLγ foci across the genome; and (3) that MUS81-EME1 activity only partially accounts for the increased residual chiasmata count observed in $Mlh3^{DN/DN}$ males.

DISCUSSION

Studies in *S. cerevisiae* and *M. musculus* have implicated MutLγ as the major resolvase of dHJs in the Class I CO pathway (Al-Sweel et al. 2017; Kadyrov et al. 2006; Lipkin et al. 2002; Manhart et al. 2017; Rogacheva et al. 2014; Ranjha et al. 2014; Nishant et al. 2008; Zakharyevich et al. 2012; Edelmann et al. 1996). The current study elucidates for the first time the importance of an intact endonuclease domain for the proper functioning of MLH3 during prophase I of mammalian meiosis. Our data reveal important similarities and differences in the mutant phenotypes to that observed with a nullizygous *Mlh3^-/-* allele. We generated a mouse lacking a functional endonuclease domain within MLH3, while allowing for heterodimer assembly and near-normal stability of the protein. We found that, as in *Mlh3^-/-* mutant males, *Mlh3^DN/DN* males are infertile, exhibit significantly smaller testes than their WT litter mates, and have no epididymal spermatozoa.

Looking specifically at prophase I events, we demonstrate that an intact endonuclease domain within MLH3 is not required for DSB or synaptonemal complex formation, similar to that seen in *Mlh3^-/-* males (Lipkin et al. 2002). However, there are distinct differences in RAD51 accumulation and persistence in spermatocytes from *Mlh3^-/-* and *Mlh3^DN/DN* males, suggesting that the effect of MutLγ loss on DSB repair processing is quite different from the presence of a defective MutLγ complex. Most importantly, while RAD51 is recruited in elevated numbers to chromosome cores of *Mlh3^DN/DN* males, it fails to be cleared effectively in pachynema, perhaps because the defective MutLγ complex blocks subsequent processing of DSB repair intermediates. Intriguingly, the significantly altered RAD51 accumulation in leptonema in both *Mlh3* mutants indicates a role for MutLγ prior to pachynema, far earlier than has been defined thus far. Indeed, analogous early pairing roles for MutLγ have been proposed for *M. Sordaria* and *S. cerevisiae* (Storlazzi et al. 2010; Claeys Bouuaert and Keeney 2017). In yeast, Al-Sweel *et al* constructed whole genome recombination maps for wild-type, endonuclease defective,

and null *mlh3* yeast mutants. Compared to wild-type, both the endonuclease defective and null mutants for *mlh3* showed increases in the number of NCO events, consistent with recombination intermediates being resolved through alternative recombination pathways (Al-Sweel et al. 2017). Thus in the case of yeast, loss of Mlh3 protein, or the production of an endonuclease defective protein, increases the frequency of other recombination outcomes, most notably including earlier NCO events.

Unexpectedly, spermatocytes from *Mlh3$^{DN/DN}$* males show more residual chiasmata at diakinesis than do *Mlh3$^{-/-}$* mice. Loss of either component of MutLγ is predicted to eradicate 90-95% of chiasmata, based on the established dogma that class I COs account for the majority, but not all, chiasmata in mammalian meiosis (Baker et al. 1996; Edelmann et al. 1996; Lipkin et al. 2002; Holloway et al. 2008; Svetlanov et al. 2008). Accordingly, in the absence of MLH1 or MLH3, we observed only 10% residual COs or chiasmata when assessed either genetically or cytogenetically. Thus, our observation of only a 78% loss of COs in *Mlh3$^{DN/DN}$* spermatocytes, leading to 22% residual chiasmata, is not in line with this expectation of a complete loss of all class I COs, and instead suggests that the endonucleolytic function of MLH3 does not account for the resolution of all class I COs and/or that other resolvases can replace MLH3 under certain conditions. Indeed, our previous analysis of *Mus81$^{-/-}$Mlh3$^{-/-}$* double mutant animals revealed a very small, but consistent, number of residual chiasmata, indicating the existence of other resolvase complexes (Holloway et al. 2008). Alternatively, it is feasible that the point mutation in the endonuclease domain does not completely eliminate endonucleolytic activity in the mouse, resulting in partial class I resolvase activity. However, we find this possibility highly unlikely due to the severity of the defect in endonucleolytic activity in the *S. cerevisiae* mlh3-D523N protein (Nishant et al. 2008; Rogacheva et al. 2014). Another explanation is that, in the absence of endonuclease activity, some class I CO events can be

480   processed by class II (or other) CO machinery following appropriate processing by DNA

helicases such as BLM, and the recruitment of one or more structure specific nucleases.

In *Mlh3*$^{-/-}$ (Holloway et al., 2010) and *Mlh3*$^{DN/DN}$ (current work) males, prophase I

spermatocytes show increased and persistent accumulation of BLM helicase. In *S. cerevisiae,*

loss of class I CO pathway components (for example, in *msh4/5* or *mlh1/3* mutants) is

485   suppressed by mutation of the BLM ortholog, *Sgs1,* highlighting the role of Sgs1 as an anti-

crossover factor, and suggesting that the loss of COs in these single MMR mutants is in part

due to the activity of Sgs1 (Oh et al. 2007; Jessop et al. 2006). In *msh4/5 sgs1* double mutant

strains, the restoration of COs occurs without any concomitant decrease in NCO events,

suggesting either that other CO pathways account for the non-class I COs, or that these DSBs

490   are repaired via inter-sister repair processes. In this sense, Sgs1 has been proposed to be

master orchestrator of recombination pathway choice (De Muyt et al. 2012), while the Sgs1-

Top3-Rmi1 complex as a whole can regulate CO formation both positively and negatively in

yeast (Kaur et al. 2015; Tang et al. 2015).

The situation we observe in *Mlh3*$^{-/-}$ and *Mlh3*$^{DN/DN}$ males with respect to BLM

495   persistence appears to be similar to that seen in yeast. What, then, is the function of BLM at

these dHJs that cannot now be processed by MLH3? Previous work in yeast provides a clue

as to the functional interaction between MutLγ and BLM: crossing over in the class I pathway

depends on MutSγ-driven designation on one side of the DSB, in part by suppressing Sgs1,

while Sgs1 also imposes an anticrossover bias on the other side of the DSB (Oh et al. 2007).

500   Collectively these two functions promote appropriate CO fates at designated DSB repair sites,

preventing the accumulation of unconventional CO/NCO variants through the use of the anti-

recombination activity of Sgs1. Extrapolating to the situation in spermatocytes from our two

Mlh3 mutant alleles, we can postulate that the loss of MLH3 protein entirely in *Mlh3*$^{-/-}$ males

results in a compensatory, but ineffective, increase in BLM that cannot overcome the failure to process class I COs sufficiently (a situation that is different to yeast). In the presence of intact, but catalytically inert MutLγ, on the other hand, the availability of additional BLM can then direct DSB repair in favor of other CO pathways in a similar fashion to the situation in yeast, where the engagement of Sgs1 promotes alternative repair mechanisms, primarily through the recruitment of structure specific nucleases, and the resolution of some dHJs through a class II (or other) CO pathway (De Muyt et al. 2012; Jessop et al. 2006; Jessop and Lichten 2008; Oh et al. 2007; Oh et al. 2008).

While we cannot assess the recruitment of the class II machinery to sites of DSB repair in prophase I in the mouse (due to the lack of available reagents), there is compelling evidence to favor this model, and to support the idea that MUS81-EME1 might participate in this CO resolution cross-talk. First, double mutants lacking *Mus81* and bearing a homozygous *Mlh3*[DN] allele show reduced chiasmata relative to *Mlh3*[DN/DN] males, indicating that MUS81 accounts for at least some of the increase in chiasmata above that of *Mlh3*[-/-] males. Secondly, *Mus81*[-/-] *Mlh3*[DN/DN] double mutant males do not show the increased frequency of MLH1 foci that are observed in *Mus81*[-/-] males, again indicating that the increased chiasmata observed in *Mlh3*[DN/DN] males is not due to upregulation of a weakly active MutLγ heterodimer, and again lending credence to the idea that the MLH3[DN] protein, like its yeast counterpart, is not effective for CO resolution. Thirdly, normal accumulation of MSH4 in *Mlh3*[DN/DN] spermatocytes recruits normal levels of pro-crossover factors, HEI10 and CDK2, suggesting that other CO resolution pathways must be engaged at the expense of class I pathways.

We provide evidence that additional COs are achieved in *Mlh3*[DN/DN] spermatocytes through the enhanced activity of the MUS81-EME1 endonuclease. However, this is clearly not the only route to achieve additional CO events, since we have shown previously

spermatocytes from *Mus81$^{-/-}$Mlh3$^{-/-}$* males retain a low number of chiasmata (Holloway et al. 2008). There are other resolvase complexes that might also generate COs in mammalian

530   meiosis, and these cannot be ruled out in the current analysis. Such resolvases in yeast

include SLX1-SLX4 and YEN1/GEN1(Zakharyevich et al. 2012; Arter et al. 2018; Muñoz-Galván et al. 2012). Indeed, the persistence of DSB repair intermediates into pachynema, along with the upregulated and persistent BLM might suggest that the defective MutLγ complex prevents accumulation of other such resolvase complexes. This might, in turn delay CO

535   maturation until later in prophase I when, for example, GEN1 can be invoked to resolve the CO (García-Luis and Machín 2014). Thus, we propose that the timing of MutLγ activity, and its clearance from nascent COs is an important factor in the recruitment of alternative CO processing machineries, but in a manner that is not dependent on its endonuclease activity.

Materials and Methods

Generation of mice and genotyping

PL253 targeting vector containing the *Mlh3-D1185N* point mutation in the potential

545 endonuclease domain and a loxP-neo-loxP cassette in intron 5-6 of *Mlh3* was incorporated into

an embryonic stem cell line. *Mlh3^{DN}* transgenic mice were crossed with a *Spo11-Cre* mouse

line to remove the *neo* cassette (Lyndaker et al., 2013), and then maintained on an inbred

background through backcrossing on to the C57Bl/6J line (Jackson Laboratory, Bar Harbor,

ME). Genotyping of WT, *Mlh3$^{+/DN}$*, and *Mlh3$^{DN/DN}$* mice was performed using the following PCR

550 primer pairs: forward (5'-AAGCCAAGTCTGCATGAGTA-3') and reverse (5'-

TAAATGTGCCACTGACTAAAT-3') followed by a restriction enzyme digestion with *Sau96I*

(New England Biolabs) at 37°C for 2-3 hours, which results in 439-bp and 263-bp fragments

from the WT allele and a 702-bp fragment from the mutant allele. Fertility tests were performed

by breeding *Mlh3$^{DN/DN}$* adult males with WT females. At least 3 males of each genotyped were

555 evaluated. Presence of a copulation plug the following morning counted as a successful mating

event. Pregnancy was confirmed by gentle palpation of the abdomen after gestation day 11 or

on delivery date of litters. Mice were housed and utilized under the guidance and approval of

the Cornell University Institutional Animal Care and Use Committee.

560 **Histology**

Testes from adult mice were fixed in Bouin's solution overnight at room temperature and

then washed 3 x 10 min with 70% ethanol at room temperature with agitation. Fixed and

paraffin-embedded testes were section at 5 μm. H&E staining was performed on Bouin's fixed

testes using standard methods. At least 6 males of each genotyped were evaluated.

Sperm counts

Caudal epididymides were removed from adult males and placed in pre-warmed 1X

PBS containing 4% bovine serum albumin. Sperm were released into solution by squeezing

epididymis with tweezers and incubated for 20 min at 32ºC/5% CO_2. After incubation, 20 μL of

570   sperm suspension was re-suspended in 480 μL of 10% formalin. Sperm counts were

performed with a hemocytometer. At least 10 males of each genotype were evaluated for

sperm counts and testis weights.

Prophase I chromosome analysis and immunofluorescence

575   Prophase I chromosome spreads from adult testes were prepared as previously

described (Kolas et al., 2005). For all experiments, at least 6 males of each genotyped were

evaluated. Chromosome slides were then washed in 0.4% Kodak Photo-Flo 200/1X PBS for 2

x 5 min, 0.4% Kodak Photo-Flo 200/dH$_2$O for 2 x 5 min, then air-dried for approximately 10 min

and stored in -80ºC or used immediately for staining. Primary antibodies used were: anti-

580   γH2AX (Millipore, NY, #05-636 1:10,000), anti-SYCP3 (Abcam, MA, #97672, 1:5000), anti-

SYCP1 (Abcam, MA, #15087, 1:1000), anti-RAD51 (Calbiochem, #PC130, 1:500), anti-BLM

(generous gift from Dr. Ramundo Freire; 1:100;), anti-CDK2 (Santa Cruz, TX, sc-163; 1:250),

anti-MLH3 ((Holloway et al., 2014); 1:1000;), anti-RNF212 (generous gift from Dr. Neil Hunter),

anti-MSH4 (Abcam, MA, #58666), anti-HEI10 (Anti-CCNB1IP1, Abcam, MA # 71977) and anti-

585   MLH1 (BD Biosciences Pharmingen, CA, #550838, 1:100). Secondary antibodies used were:

goat anti-mouse Alexa Fluor 488 (#62-6511), goat anti-mouse Alexa Fluor 555 (#A-10521),

goat anti-rabbit Alexa Fluor 488 (#65-6111), goat anti-rabbit Alexa Fluor 555 (#A-10520; all

Invitrogen, 1:2000).

Spermatocyte diakinesis spread preparations to observe chiasmata

Diakinesis chromosome spreads were prepared as previously described (Evans et al., 1964; Uroz et al., 2008) with slight modifications (Holloway et al., 2010). Slides were stained with 10% Giemsa for 10 mins, washed, air-dried and mounted with Permount.

Image acquisition

All chromosome spread slides were visualized using the Zeiss Imager Z1 microscope (Carl Zeiss, Inc.). Images were captured with a high-resolution microscopy camera AxioCam MRM (Carl Zeiss, Inc.) and processed with ZEN Software (version 2.0.0.0; Carl Zeiss, Inc.).

Cloning mouse *Mlh1* and *Mlh3*

cDNA was synthesized from total testis RNA from wildtype C57B/6J adult males using the SuperScript III Reverse Transcriptase Kit from ThermoFisher. *Mlh1* and *Mlh3* open reading frames were PCR amplified from cDNA using Expand High Fidelity DNA polymerase using primer pairs AO3365 (5'GCTAGCAGCTGATGCATATGGCGTTTGTAGCAGGAG) and AO3366 (5'TACCGCATGCTATGCATTAACACCGCTCAAAGACTTTG) for *Mlh1*, and AO3367 (5'ACGTCGACGAGCTCATATGCATCACCATCACCATCACCATCACCATCACATCAGGTGTC TATCAGATGAC) and AO3368 (5'CGAAAGCGGCCGCGATCATGGAGGCTCACAAGG) for *His$_{10}$-Mlh3*. Each fragment was cloned into the *Spe*1 site of pFastBac1 (ThermoFisher) using Gibson assembly PCR (NEB) to create pEAE393 (*Mlh1*) and pEAE397 (*His$_{10}$-Mlh3*). Constructs were verified by DNA sequencing with NCBI reference sequences NM_026810.2 and NM_175337.2 for *Mlh1* and *Mlh3,* respectively. These constructs were then modified as follows:

1. The MBP –TEV sequence was inserted at the N-terminus of *Mlh1* in pEAE393 to create pEAE395.

615 2. The *mlh3-D1185N* mutation was introduced into pEAE397 by Quick Change (Agilent) to create pEAE413.

Chromatography analysis of the mouse MLH1-MLH3-D1185N heterodimer from Baculovirus-infected Sf9 cells

620 *Sf9* cells were transfected with pEAE397 (His_{10}-*Mlh3*), pEAE413 (His_{10}-*Mlh3-D1185N*) and pEAE395 (*MBP-Mlh1*) using the Bac-to-Bac baculovirus infection system (Invitrogen). Fresh *Sf9* cells were co-infected with both viruses (containing *Mlh1* and *Mlh3* or *Mlh3-D1185N*). Cells were harvested 60 hours post infection, washed with phosphate buffered saline, and kept at -80°C until use.

625 Cell pellets from 250 ml of cells as thawed, resuspended in 60 ml hypotonic lysis buffer (20 mM HEPES-KOH pH 7.5, 5 mM NaCl, 1 mM $MgCl_2$, 1 mM PMSF and EDTA free protease inhibitor mixture from Roche and Thermo Scientific) and incubated for 15 min on ice. The suspension was adjusted to 250 mM NaCl, 15 mM imidazole, 10% glycerol, 2 mM ß-mercaptoethanol (BME), and clarified by centrifugation at 17,000 g for 20 min at 4°C. The

630 supernatant was mixed with 6 ml of 50% nickel-nitrolotriaceticacid-agarose (Ni-NTA) resin and allowed to bind for 2 hours or overnight followed by centrifugation to remove the unbound fraction. The resin was packed onto a column and washed with 7-10 column volumes of wash buffer (50 mM HEPES-KOH pH 7.5, 250 mM NaCl, 40 mM imidazole, 10% glycerol, 2 mM BME, 1 mM PMSF). Protein was eluted with 15 ml of 300 mM imidazole in 50 mM HEPES-

635 KOH pH 7.5, 250 mM NaCl, 40 mM imidazole, 10% glycerol, 2 mM BME and 1 mM PMSF. Elution fractions containing MLH1-MLH3, determined by SDS-PAGE, were pooled and loaded

onto 1 ml 100% amylose resin (NEB). The resin was washed with 10 column volumes of wash

buffer (50 mM HEPES-KOH pH 7.5, 250 mM NaCl, 10% glycerol, 2 mM BME, 1 mM PMSF)

and eluted with 6 ml wash buffer containing 10 mM maltose. Fractions containing MLH1-MLH3

640   were pooled and aliquots were flash frozen and stored in -80°C. The protein yield, following

amylose chromatography, was similar for wild-type and mutant complexes (approximately 120-

150 μg per 250 ml cells).

It is important to note that we were unable to detect a specific endonuclease activity for the

mouse MBP-MLH1-MLH3 complex, suggesting that the MBP tag interferes with MLH1-MLH3

645   functions. We were unable to test this directly because, despite numerous attempts, we were

unable to efficiently remove the MBP tag from MLH1 by treating MBP-MLH1-MLH3 with TEV

protease.

Mass-spectrometry of MLH1 and MLH3 bands from SDS-PAGE

650   SDS-PAGE bands following amylose chromatography predicted to contain MBP-MLH1

and His_{10}-MLH3 were excised and analyzed by the Cornell University Proteomics facility using

a Thermo LTQ Orbitrap Velos Mass Spectrometer.

Statistical methods and analysis

655   The majority of comparisons involved with unpaired parametric t-test with Welch's

correction or nonparametric Mann-Whitney U-test, depending on the data distribution. All

statistical analysis was performed with GraphPad Prism Version 7.00 for Mac, Graphpad

Software, La Jolla California USA, www.graphpad.com. P-values less than 0.05 were

considered statistically significant.

Acknowledgements

The authors of this work would like to thank Dr. Andrew Miller for necropsy analysis of WT, *Mlh3^{DN/DN}* and *Mlh3^{-/-}* mice (data not included), Dr. Ramundo Freire for providing anti-BLM antibody. We are grateful to members of the Cohen lab for their critical evaluation of this work. This work was supported by the National Institutes of Health (NIH) grant R01HD041012 to P.E. Cohen, and by graduate student funding through The Training Grant in Reproductive Genomics (PI Mark S. Roberson, T32HD052471) for M. Toledo. V. Raghavan and E. Alani were supported by funding to E. Alani from The NIH (grant GM53085). The authors declare no competing financial interests.

Author contribution

Melissa Toledo performed experiments, curated data, supervised undergraduates, and wrote the original draft; Xianfei Sun performed experiments, generated the mutant allele, and conceptualized the project; Miguel A. Brieño-Enríquez performed experiments, curated data, and reviewed/edited the manuscript; Vandana Raghavan performed experiments and analyzed data; Stephen Gray performed experiments, curated data, and reviewed/edited the manuscript; Jeffrey Pea performed experiments and analyzed data; Anita Venkatesh performed experiments ; Lekha Patel performed experiments; Peter L. Borst performed experiments; Eric Alani curated data, conceptualized experiments, analyzed data, and reviewed/edited the manuscript; Paula E. Cohen acquired funding, supervised the experiments, curated data, conceptualized experiments, wrote the manuscript, and reviewed/edited the manuscript.

Figure Legends

Figure 1. *Mlh3$^{DN/DN}$* males show a sterile phenotype. (A, B) *Mlh3$^{DN/DN}$* adult male testes are

significantly smaller when compared to WT littermates (*Mlh3$^{DN/DN}$* - 0.28% of total body weight

± 0.06, n = 13; WT - 0.77% ± 0.1, n = 15; p < 0.0001, unpaired t-test with Welch's correction)

690   and (C) have zero sperm in the epididymis (WT - 9.8 x 10^7 ± 4.4 sperm/mouse; p < 0.0001;

unpaired t-test with Welch's correction, n = 10 and 12 mice, respectively; error bars show

standard deviation). Hemotoxylin and eosin stained (D, E) WT testes show the presence of

meiotic and post-meiotic cells whereas (F, G) *Mlh3$^{DN/DN}$* are absent of spermatids and

spermatozoa. Higher magnification of WT and *Mlh3$^{DN/DN}$* testes sections are shown in E and G,

695   respectively. Black arrows in (G) indicate metaphase I spermatocytes in the *Mlh3$^{DN/DN}$*

seminiferous tubule lumen. Sg, spermatogonia; Sc, prophase I spermatocytes; St, postmeiotic

spermatids. Scale bar is 100 μm. Images were taken at 40X magnification.

Figure 2. *Mlh3$^{DN/DN}$* spermatocytes show a persistence of RAD51 foci in pachynema.

700   The localization of RAD51 (green) on synaptonemal complex protein SYCP3 (red) is observed

in (A, B) WT and (C, D) *Mlh3$^{DN/DN}$* male spermatocytes in zygonema and pachynema. (A, C, E,

G) In early zygonema, WT cells show high numbers of RAD51 associated with the

chromosome cores while *Mlh3$^{DN/DN}$* exhibit even higher RAD51 foci (WT mean $\pm$ standard

deviation = 142.1 ± 34.5 foci, *Mlh3$^{DN/DN}$* mean $\pm$ standard deviation = 188.8 ± 53.3 foci,

705   p<0.001 by unpaired t-test with Welch's correction). In late zygonema, *Mlh3$^{DN/DN}$*

spermatocytes continue to have higher RAD51 foci than WT (WT mean $\pm$ standard deviation =

193.0 ± 53.7 foci, *Mlh3$^{DN/DN}$* mean $\pm$ standard deviation = 225.3 ± 37.3 foci; p<0.01 by unpaired

t-test with Welch's correction). *Mlh3$^{-/-}$* late zygotene cells show significantly fewer RAD51 focus

counts (mean = 122.1 ± 39.9 foci) when compared to WT and *Mlh3*^{DN/DN} (both p < 0.0001 by

unpaired t-test with Welch's correction). In all cases, at least 3 mice were assessed for each

genotype and at least 20 cells per mouse (B, D, F, H) In pachynema, WT cells exhibit a

dramatic decrease of very few to no RAD51 associated with the autosomes while *Mlh3*^{DN/DN}

cells show persistent RAD51 (WT mean ± standard deviation = 2.8 ± 2.8 foci, *Mlh3*^{DN/DN} mean

± standard deviation = 6.1 ± 3.1 foci; p<0.0001, unpaired t-test). *Mlh3*^{-/-} pachytene cells exhibit

comparable RAD51 focus counts (mean ± standard deviation = 3.1 ± 2.1 foci) when compared

to WT (p = 0.55 by unpaired t-test), but significantly fewer than *Mlh3*^{DN/DN}

spermatocytes(p<0.0001 by unpaired t-test). Note that sex chromosome-associated RAD51

staining was excluded from counts at pachynema. For all chromosome imaging and foci

counts, at least three mice of each genotype were observed for each staining set.

Figure 3. *Mlh3*^{DN/DN} spermatocytes show a persistence of BLM in pachynema. Meiotic

spreads chromosomes from (A-D) WT, (E-H) *Mlh3*^{DN/DN}, and (I-L) *Mlh3*^{-/-} males showing the

localization of BLM (red) on synaptonemal complex protein SYCP3 (green) throughout the

progression of prophase I. (B, F, J, M) In zygonema, *Mlh3*^{DN/DN} and *Mlh3*^{-/-} cells show

significantly more BLM foci localized to the chromosome cores than WT (WT mean ± standard

deviation = 221.6 ± 35.1 foci, *Mlh3*^{DN/DN} mean ± standard deviation = 271.1 ± 34.2 foci, *Mlh3*^{-/-}

mean ± standard deviation = 265.5 ± 36.6 foci; p<0.0001 for each using unpaired t-test). (C, D,

K, M) In pachynema, BLM is no longer present on the chromosome cores of WT cells whereas

Mlh3^{DN/DN} and *Mlh3*^{-/-} cells show hyper-accumulation of BLM on the autosomes and the sex

body (WT mean ± standard deviation = 7.8 ± 4.7 foci, *Mlh3*^{DN/DN} mean ± standard deviation =

66.7 ± 23.2 foci, *Mlh3*^{-/-} mean ± standard deviation = 58.7 ± 17.8 foci; p<0.0001 for each using

unpaired t-test), which persists into diplonema. At zygonema, the frequency of BLM foci in

Mlh3$^{DN/DN}$ and *Mlh3$^{-/-}$* cells is not statistically different, whereas by pachynema, the number of

BLM foci remains significantly elevated in *Mlh3$^{DN/DN}$* spermatocytes, relative to that observed

735   in *Mlh3$^{-/-}$* cells (p<0.05, unpaired t-test). For all chromosome imaging and foci counts, at least

3 mice of each genotype were observed for each staining set.

Figure 4. Normal localization of CDK2 and HEI10 to nascent sites of Class I crossovers

in *Mlh3$^{DN/DN}$* spermatocytes. (A-C) CDK2 (red) localizes to the synaptonemal complex

740   SYCP3 (green) in WT (A) and *Mlh3$^{DN/DN}$* (B) pachytene spermatocytes, but not in *Mlh3$^{-/-}$* (C)

cells, except at the telomeres. Left-hand panels show merged red and green channels, in

which white arrows show examples of crossover associated CDK2, while yellow arrows show

examples of telomere associated CDK2. Right-hand panels show only the CDK2 signal in

white. (D-F) HEI10 (red) co-localizes with the synaptonemal complex SYCP3 (green) in

745   pachytene spermatocytes from WT (D), *Mlh3$^{DN/DN}$* (E), and *Mlh3$^{-/-}$* (F) males. Left-hand panels

show merged red and green channels, in which pink arrows show examples of crossover

associated HEI10. Right-hand panels show only the HEI10 signal in white. For all chromosome

imaging and foci counts, at least 3 mice of each genotype were observed for each staining set.

750   **Figure 5. *Mlh3$^{DN/DN}$* pachytene spermatocytes exhibit normal localization of MutLγ while**

the recombinant MLH3-D1185N protein can form complexes with MLH1. (A-C) MLH3

(green) localizes to SYCP3 (red) in WT and *Mlh3$^{DN/DN}$* pachytene spermatocytes with no

statistical difference in the number of MLH3 foci (WT = 21.7 ± 3.4 MLH3 foci n=131, *Mlh3$^{DN/DN}$*

= 21.3 ± 3.1 MLH3 foci n=122; p = 0.36 by unpaired t-test). (D-F) MLH1 (green) localizes to

755   SYCP3 (red) in WT and *Mlh3$^{DN/DN}$* pachytene spermatocytes also with no statistical difference

in the number of MLH1 foci (WT = 21.1 ± 2.6 MLH1 foci n=144, *Mlh3$^{DN/DN}$* = 20.2 ± 2.8 MLH1

foci n=142; p = 0.2 by unpaired t-test). Different colors in (C) and (F) indicate 3 sets of matched

littermates used, each color referring to the counts on a single animal (and color set from the

two genotypes being processed simultaneously); n.s = not significant; error bars show

standard deviation. (G-I) MLH1-mlh3-D1185N forms a stable heterodimer. (G) Schematic of

mouse *Mlh1* and *Mlh3* constructs (see Methods for details). (H) Representative purification of

MBP-MLH1-MLH3 (top) and MBP-MLH1-mlh3-D1185N (bottom) using Ni-NTA and amylose

resin chromatography as described in the Methods. Fractions were analyzed using SDS-

PAGE, stained by Coomassie brilliant blue. MLH1-MLH3 and MLH1-MLH3-D1185N were

eluted from amylose in the same fractions. The mass of molecular weight standards is

indicated on the left and the expected positions of MBP-MLH1 (127.5 KDa) and His$_{10}$-MLH3

(165 KDa) is indicated in the center. *Likely to be degradation products of MLH1-MLH3. (I)

Mass spectrometry analysis of the two major bands in SDS-PAGE detected after amylose

chromatography.

Figure 6. *Mlh3$^{DN/DN}$* **diakinesis staged spermatocytes show a reduced number of**

chiasmata (A-D), **while comparison of MLH1 focus frequency (E) and distribution (F)**

during pachynema, and chiasmata counts (G) and bivalent/univalent frequencies (H) in

Mus81$^{-/-}$ Mlh3$^{+/+}$, ***Mus81$^{-/-}$ Mlh3$^{+/DN}$***, **and** ***Mus81$^{-/-}$ Mlh3$^{DN/DN}$*** **males indicates partial**

involvement of class II CO pathway. (A-C) Diakinesis staged spermatocyte preparations

from WT, *Mlh3$^{DN/DN}$*, and *Mlh3$^{-/-}$* males stained with Giemsa showing chiasmata formation

between homologous chromosomes. (D) *Mlh3$^{DN/DN}$* cells exhibit significantly fewer chiasmata

when compared to WT (WT = 23.5 ± 1.3 chiasmata per nucleus, *Mlh3$^{DN/DN}$* = 5.2 ± 1.7; p <

0.0001 by unpaired t-test). *Mlh3$^{-/-}$* cells have significantly fewer chiasmata when compared to

WT and *Mlh3$^{DN/DN}$* (*Mlh3$^{-/-}$* = 2.8 ± 1.1 chiasmata per nucleus; p < 0.0001 by unpaired t-test).

All values are means $\pm$ standard deviation. (E-H). To explore the interplay between class I and class II CO events, we compared MutLγ frequency and distribution during pachynema between mice lacking *Mu81* with or without co-incident loss of a endonuclease-intact *Mlh3* allele (E, F). As expected, *Mus81$^{-/-}$Mlh3$^{+/+}$* males (E; filled octagons) show elevated MLH1 focus frequency relative to WT animals (Figure 5) and this is lost in mice bearing one or two copies of the *Mlh3^{DN}* allele (filled diamonds and down triangles, respectively). The reduced MLH1 focus count in *Mus81$^{-/-}$Mlh3$^{+/DN}$*, and *Mus81$^{-/-}$Mlh3$^{DN/DN}$* males is statistically significant (p<0.005, unpaired t-test with Welch's correction). (F) The distribution of MLH1 foci is also disrupted in *Mus81$^{-/-}$Mlh3$^{+/DN}$*, and *Mus81$^{-/-}$Mlh3$^{DN/DN}$* males, with increased numbers of synapsed autosomes showing no MLH1 foci, indicative of non-exchange (E0) chromosome pairs, compared to that seen in *Mus81$^{-/-}$Mlh3$^{+/+}$* males (statistical significance indicated on the graph; unpaired t-test with Welch's correction). (G, H) the outcome of class I and class II CO events was assessed by quantifying chiasmata (G) and intact bivalent pairs (H) in diakinesis preparations from mice of all three double mutant genotype combinations, compared to single mutants presented in panel C, above. (G) Chiasmata counts were similar to WT for *Mus81$^{-/-}$Mlh3$^{+/+}$* and *Mus81$^{-/-}$Mlh3$^{+/DN}$* males (filled octagons and diamonds, respectively), but were statistically significantly lower in *Mus81$^{-/-}$Mlh3$^{DN/DN}$* animals (filled down triangles). The frequency of chiasmata in *Mus81$^{-/-}$Mlh3$^{DN/DN}$* animals was significantly lower than that observed in *Mlh3$^{DN/DN}$* animals, but significantly higher than that observed in *Mlh3$^{-/-}$* animals (p<0.05, unpaired t-test with Welch's correction). However, the number of bivalent structures observed at diakinesis in *Mus81$^{-/-}$Mlh3$^{DN/DN}$* animals (panel H) was unchanged from that observed in *Mlh3$^{DN/DN}$* animals. In all cases, two animals were assessed for each genotype.

Figure S1. Amino acid sequence of MLH3 endonuclease domain and its conservation

across species. (A) Amino acid protein sequence of the *M. musculus, H. sapiens, S.*

cerevisiae, and A. thaliana MLH3 endonuclease domain, DQHA(X)$_2$E(X)$_4$E, shows the

conservation of this domain across these species. Asterisk refers to the conserved aspartic

acid (D) that was targeted for a point mutation and converted to asparagine (N) to generate the

Mlh3-DN mouse. (B) *Mus musculus* MLH3 is composed of a 1411 amino acid long sequence

that results in an ~158 kDa sized protein (UniProt, 2015) . MLH3 contains a globular N-terminal

domain (NTD; light gray) and C-terminal domain (CTD; light blue) connected by a flexible linker

arm (white). The NTD (light gray) contains ATP binding motifs (dark gray) that are conserved

across species. The CTD consists of the MLH1 interacting domain (light blue) and the

conserved endonuclease motif (dark blue). The aspartic acid (D) in the conserved

endonuclease motif in mouse, DQHAAHERIRLE, was converted to an asparagine (N; red) at

amino acid site 1185.

Figure S2. DSB formation and signaling as well as synapsis are normal in *Mlh3*$^{DN/DN}$

spermatocytes throughout prophase I. (A-O) *Mlh3*$^{DN/DN}$ prophase I cells exhibit normal DSB

formation and signaling as observed by γH2AX staining (green) on synaptonemal complex

protein SYCP3 (red) as compared to WT and *Mlh3*$^{-/-}$ cells. Images show abundant γH2AX

signal in leptonema, following by diminished signal in zygonema, with the absence of signal in

pachynema and diplonema, except at the sex body, due to MSCI. (E, J, O) Over exposure of

the γH2AX signal results in γH2AX foci or flares on the autosomes in WT, *Mlh3*$^{DN/DN}$, as well as

Mlh3$^{-/-}$ cells (white arrows), suggesting that this signal may not be representative of true un-

repaired DSBs. (P-W) *Mlh3*$^{DN/DN}$ prophase I cells have normal synapsis as observed by the

localization of synaptonemal complex protein SYCP1 (green) and SYCP3 (red) on the

chromosomes when compared to WT. SYCP3 forms as short patches along the chromosomes

in leptonema, extending into filaments in zygonema along with the appearance of SYCP1, full

830 synapsis with the co-localization of SYCP1 and SYCP3 are observed in pachynema, followed

by desynapsis in diplonema with the degradation of SYCP1.

Figure S3. Normal localization of crossover designation factor MSH4 in *Mlh3$^{DN/DN}$* and

Mlh3$^{-/-}$ spermatocytes. (A-M) MSH4 (green) localization on chromosome cores stained with

835 antibodies against SYCP3 (red) throughout prophase I (leptonema, zygonema, early

pachynema, and late pachynema). Diplonema and diakinesis are not shown. Representative

images of spermatocytes from WT (A-D), *Mlh3$^{DN/DN}$* (E-H), and *Mlh3$^{-/-}$* (I-L) adult males. Panel

M shows the quantitation of MSH4 foci in all three genotypes at early (EP) and late pachynema

(LP). A similar pattern of MSH4 foci accumulation and loss is observed for MSH4 as for

840 RNF212: accumulation of high numbers of foci in zygonema, diminishing to one or two foci per

chromosome in late pachynema. In all cases, ***$p < 0.0001$ by unpaired t-test with Welch's

correction. For all chromosome imaging and foci counts, at least three mice of each genotype

were observed for each staining set.

Bibliography

Agarwal, S. and Roeder, G.S. 2000. Zip3 provides a link between recombination enzymes and

850   synaptonemal complex proteins. *Cell* 102(2), pp. 245–255.

Allers, T. and Lichten, M. 2001. Intermediates of yeast meiotic recombination contain

heteroduplex DNA. *Molecular Cell* 8(1), pp. 225–231.

Anderson, L.K., Reeves, A., Webb, L.M. and Ashley, T. 1999. Distribution of crossing over on

mouse synaptonemal complexes using immunofluorescent localization of MLH1 protein.

855   *Genetics* 151, pp. 1569–79. Available at: http://www.ncbi.nlm.nih.gov/cgi-

bin/Entrez/referer?http://www.genetics.org/cgi/content/full/151/4/1569.

Argueso, J.L., Smith, D., Yi, J., Waase, M., Sarin, S. and Alani, E. 2002. Analysis of conditional

mutations in the Saccharomyces cerevisiae MLH1 gene in mismatch repair and in meiotic

crossing over. *Genetics* 160(3), pp. 909–921.

860   Arter, M., Hurtado-Nieves, V., Oke, A., Zhuge, T., Wettstein, R., Fung, J.C., Blanco, M.G. and

Matos, J. 2018. Regulated Crossing-Over Requires Inactivation of Yen1/GEN1 Resolvase

during Meiotic Prophase I. *Developmental Cell* 45(6), pp. 785–800.e6.

Ashley, T., Plug, A.W., Xu, J., Solari, A.J., Reddy, G., Golub, E.I. and Ward, D.C. 1995.

Dynamic changes in Rad51 distribution on chromatin during meiosis in male and female

865   vertebrates. *Chromosoma* 104, pp. 19–28.

Ashley, T., Walpita, D. and de Rooij, D.G. 2001. Localization of two mammalian cyclin

dependent kinases during mammalian meiosis. *Journal of Cell Science* 114(Pt 4), pp. 685–

693.

Auton, A., Rui Li, Y., Kidd, J., Oliveira, K., Nadel, J., Holloway, J.K., Hayward, J.J., Cohen, P.E., Greally, J.M., Wang, J., Bustamante, C.D. and Boyko, A.R. 2013. Genetic recombination is targeted towards gene promoter regions in dogs. *PLoS Genetics* 9(12), p. e1003984.

Baker, S.M., Plug, A.W., Prolla, T.A., Bronner, C.E., Harris, A.C., Yao, X., Christie, D.M., Monell, C., Arnheim, N., Bradley, A., Ashley, T. and Liskay, R.M. 1996. Involvement of mouse Mlh1 in DNA mismatch repair and meiotic crossing over. *Nature Genetics* 13(3), pp. 336–342.

De Boer, E., Stam, P., Dietrich, A.J.J., Pastink, A. and Heyting, C. 2006. Two levels of interference in mouse meiotic recombination. *Proceedings of the National Academy of Sciences of the United States of America* 103(25), pp. 9607–9612.

Bolcun-Filas, E. and Handel, M.A. 2018. Meiosis: the chromosomal foundation of reproduction. *Biology of Reproduction* 99(1), pp. 112–126.

Börner, G.V., Kleckner, N. and Hunter, N. 2004. Crossover/noncrossover differentiation, synaptonemal complex formation, and regulatory surveillance at the leptotene/zygotene transition of meiosis. *Cell* 117(1), pp. 29–45.

Claeys Bouuaert, C. and Keeney, S. 2017. Distinct DNA-binding surfaces in the ATPase and linker domains of MutLγ determine its substrate specificities and exert separable functions in meiotic recombination and mismatch repair. *PLoS Genetics* 13(5), p. e1006722.

Cole, F., Keeney, S. and Jasin, M. 2010. Comprehensive, fine-scale dissection of homologous recombination outcomes at a hot spot in mouse meiosis. *Molecular Cell* 39(5), pp. 700–710.

Edelmann, W., Cohen, P.E., Kane, M., Lau, K., Morrow, B., Bennett, S., Umar, A., Kunkel, T., Cattoretti, G., Chaganti, R., Pollard, J.W., Kolodner, R.D. and Kucherlapati, R. 1996. Meiotic pachytene arrest in MLH1-deficient mice. *Cell* 85(7), pp. 1125–1134.

Edelmann, W., Cohen, P.E., Kneitz, B., Winand, N., Lia, M., Heyer, J., Kolodner, R., Pollard, J.W. and Kucherlapati, R. 1999. Mammalian MutS homologue 5 is required for chromosome pairing in meiosis. *Nature Genetics* 21(1), pp. 123–127.

García-Luis, J. and Machín, F. 2014. Mus81-Mms4 and Yen1 resolve a novel anaphase bridge formed by noncanonical Holliday junctions. *Nature Communications* 5, p. 5652.

Gray, S. and Cohen, P.E. 2016. Control of meiotic crossovers: from double-strand break formation to designation. *Annual Review of Genetics* 50, pp. 175–210.

Guillon, H., Baudat, F., Grey, C., Liskay, R.M. and de Massy, B. 2005. Crossover and noncrossover pathways in mouse meiosis. *Molecular Cell* 20(4), pp. 563–573.

Handel, M.A. 2004. The XY body: a specialized meiotic chromatin domain. *Exp Cell Res* 296, pp. 57–63.

Higgins, J.D., Buckling, E.F., Franklin, F.C. and Jones, G.H. 2008. Expression and functional analysis of AtMUS81 in Arabidopsis meiosis reveals a role in the second pathway of crossing-over. *The Plant Journal: for Cell and Molecular Biology* 54(1), pp. 152–162.

Hollingsworth, N.M., Ponte, L. and Halsey, C. 1995. MSH5, a novel MutS homolog, facilitates meiotic reciprocal recombination between homologs in Saccharomyces cerevisiae but not mismatch repair. *Genes & Development* 9(14), pp. 1728–1739.

Holloway, J.K., Booth, J., Edelmann, W., McGowan, C.H. and Cohen, P.E. 2008. MUS81 generates a subset of MLH1-MLH3-independent crossovers in mammalian meiosis. *PLoS Genetics* 4(9), p. e1000186.

Holloway, J.K., Morelli, M.A., Borst, P.L. and Cohen, P.E. 2010. Mammalian BLM helicase is critical for integrating multiple pathways of meiotic recombination. *The Journal of Cell Biology*

188(6), pp. 779–789.

Holloway, J.K., Sun, X., Yokoo, R., Villeneuve, A.M. and Cohen, P.E. 2014. Mammalian CNTD1 is critical for meiotic crossover maturation and deselection of excess precrossover sites. *The Journal of Cell Biology* 205(5), pp. 633–641.

Hunter, N. 2015. Meiotic recombination: the essence of heredity. *Cold Spring Harbor Perspectives in Biology* 7(12).

Hunter, N., Borner, G.V., Lichten, M. and Kleckner, N. 2001. Gamma-H2AX illuminates meiosis. *Nat Genet* 27, pp. 236–8.

Hunter, N. and Borts, R.H. 1997. Mlh1 is unique among mismatch repair proteins in its ability to promote crossing-over during meiosis. *Genes & Development* 11(12), pp. 1573–1582.

Jean, M., Pelletier, J., Hilpert, M., Belzile, F. and Kunze, R. 1999. Isolation and characterization of AtMLH1, a MutL homologue from Arabidopsis thaliana. *Molecular & general genetics : MGG* 262(4-5), pp. 633–642.

Jessop, L. and Lichten, M. 2008. Mus81/Mms4 endonuclease and Sgs1 helicase collaborate to ensure proper recombination intermediate metabolism during meiosis. *Molecular Cell* 31(3), pp. 313–323.

Jessop, L., Rockmill, B., Roeder, G.S. and Lichten, M. 2006. Meiotic chromosome synapsis-promoting proteins antagonize the anti-crossover activity of sgs1. *PLoS Genetics* 2(9), p. e155.

Kadyrov, F.A., Dzantiev, L., Constantin, N. and Modrich, P. 2006. Endonucleolytic function of MutLalpha in human mismatch repair. *Cell* 126(2), pp. 297–308.

Kan, R., Sun, X., Kolas, N.K., Avdievich, E., Kneitz, B., Edelmann, W. and Cohen, P.E. 2008. Comparative analysis of meiotic progression in female mice bearing mutations in genes of the

935 DNA mismatch repair pathway. *Biology of Reproduction* 78(3), pp. 462–471.

Kaur, H., De Muyt, A. and Lichten, M. 2015. Top3-Rmi1 DNA single-strand decatenase is integral to the formation and resolution of meiotic recombination intermediates. *Molecular Cell* 57(4), pp. 583–594.

Kneitz, B., Cohen, P.E., Avdievich, E., Zhu, L., Kane, M.F., Hou, H., Kolodner, R.D.,
940 Kucherlapati, R., Pollard, J.W. and Edelmann, W. 2000. MutS homolog 4 localization to meiotic chromosomes is required for chromosome pairing during meiosis in male and female mice. *Genes & Development* 14(9), pp. 1085–1097.

Kolas, N.K., Svetlanov, A., Lenzi, M.L., Macaluso, F.P., Lipkin, S.M., Liskay, R.M., Greally, J., Edelmann, W. and Cohen, P.E. 2005. Localization of MMR proteins on meiotic chromosomes
945 in mice indicates distinct functions during prophase I. *The Journal of Cell Biology* 171(3), pp. 447–458.

Lenzi, M.L., Smith, J., Snowden, T., Kim, M., Fishel, R., Poulos, B.K. and Cohen, P.E. 2005. Extreme heterogeneity in the molecular events leading to the establishment of chiasmata during meiosis i in human oocytes. *American Journal of Human Genetics* 76(1), pp. 112–127.

950 Lipkin, S.M., Moens, P.B., Wang, V., Lenzi, M., Shanmugarajah, D., Gilgeous, A., Thomas, J., Cheng, J., Touchman, J.W., Green, E.D., Schwartzberg, P., Collins, F.S. and Cohen, P.E. 2002. Meiotic arrest and aneuploidy in MLH3-deficient mice. *Nature Genetics* 31(4), pp. 385–390.

Lynn, A., Soucek, R. and Börner, G.V. 2007. ZMM proteins during meiosis: crossover artists at
955 work. *Chromosome Research* 15(5), pp. 591–605.

Mahadevaiah, S.K., Turner, J.M., Baudat, F., Rogakou, E.P., de Boer, P., Blanco-Rodríguez,

J., Jasin, M., Keeney, S., Bonner, W.M. and Burgoyne, P.S. 2001. Recombinational DNA double-strand breaks in mice precede synapsis. *Nature Genetics* 27(3), pp. 271–276.

Manhart, C.M., Ni, X., White, M.A., Ortega, J., Surtees, J.A. and Alani, E. 2017. The mismatch repair and meiotic recombination endonuclease Mlh1-Mlh3 is activated by polymer formation and can cleave DNA substrates in trans. *PLoS Biology* 15(4), p. e2001164.

Marcon, E. and Moens, P. 2003. MLH1p and MLH3p localize to precociously induced chiasmata of okadaic-acid-treated mouse spermatocytes. *Genetics* 165(4), pp. 2283–2287.

McMahill, M.S., Sham, C.W. and Bishop, D.K. 2007. Synthesis-dependent strand annealing in meiosis. *PLoS Biology* 5(11), p. e299.

Moens, P.B., Freire, R., Tarsounas, M., Spyropoulos, B. and Jackson, S.P. 2000. Expression and nuclear localization of BLM, a chromosome stability protein mutated in Bloom's syndrome, suggest a role in recombination during meiotic prophase. *Journal of Cell Science* 113 (Pt 4), pp. 663–672.

Muñoz-Galván, S., Tous, C., Blanco, M.G., Schwartz, E.K., Ehmsen, K.T., West, S.C., Heyer, W.D. and Aguilera, A. 2012. Distinct roles of Mus81, Yen1, Slx1-Slx4, and Rad1 nucleases in the repair of replication-born double-strand breaks by sister chromatid exchange. *Molecular and Cellular Biology* 32(9), pp. 1592–1603.

De Muyt, A., Jessop, L., Kolar, E., Sourirajan, A., Chen, J., Dayani, Y. and Lichten, M. 2012. BLM helicase ortholog Sgs1 is a central regulator of meiotic recombination intermediate metabolism. *Molecular Cell* 46(1), pp. 43–53.

Nishant, K.T., Chen, C., Shinohara, M., Shinohara, A. and Alani, E. 2010. Genetic analysis of baker's yeast Msh4-Msh5 reveals a threshold crossover level for meiotic viability. *PLoS*

Genetics 6(8).

980   Nishant, K.T., Plys, A.J. and Alani, E. 2008. A mutation in the putative MLH3 endonuclease

domain confers a defect in both mismatch repair and meiosis in Saccharomyces cerevisiae.

Genetics 179(2), pp. 747–755.

Novak, J.E., Ross-Macdonald, P.B. and Roeder, G.S. 2001. The budding yeast Msh4 protein

functions in chromosome synapsis and the regulation of crossover distribution. *Genetics*

985   158(3), pp. 1013–1025.

Oh, S.D., Lao, J.P., Hwang, P.Y.-H., Taylor, A.F., Smith, G.R. and Hunter, N. 2007. BLM

ortholog, Sgs1, prevents aberrant crossing-over by suppressing formation of multichromatid

joint molecules. *Cell* 130(2), pp. 259–272.

Oh, S.D., Lao, J.P., Taylor, A.F., Smith, G.R. and Hunter, N. 2008. RecQ helicase, Sgs1, and

990   XPF family endonuclease, Mus81-Mms4, resolve aberrant joint molecules during meiotic

recombination. *Molecular Cell* 31(3), pp. 324–336.

Pochart, P., Woltering, D. and Hollingsworth, N.M. 1997. Conserved properties between

functionally distinct MutS homologs in yeast. *The Journal of Biological Chemistry* 272(48), pp.

30345–30349.

995   Qiao, H., Prasada Rao, H.B.D., Yang, Y., Fong, J.H., Cloutier, J.M., Deacon, D.C., Nagel, K.E.,

Swartz, R.K., Strong, E., Holloway, J.K., Cohen, P.E., Schimenti, J., Ward, J. and Hunter, N.

2014. Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic

recombination. *Nature Genetics* 46(2), pp. 194–199.

Ranjha, L., Anand, R. and Cejka, P. 2014. The Saccharomyces cerevisiae Mlh1-Mlh3

1000   heterodimer is an endonuclease that preferentially binds to Holliday junctions. *The Journal of*

Biological Chemistry 289(9), pp. 5674–5686.

Rogacheva, M.V., Manhart, C.M., Chen, C., Guarne, A., Surtees, J. and Alani, E. 2014. Mlh1-Mlh3, a meiotic crossover and DNA mismatch repair factor, is a Msh2-Msh3-stimulated endonuclease. *The Journal of Biological Chemistry* 289(9), pp. 5664–5673.

1005   De los Santos, T., Hunter, N., Lee, C., Larkin, B., Loidl, J. and Hollingsworth, N.M. 2003. The Mus81/Mms4 endonuclease acts independently of double-Holliday junction resolution to promote a distinct subset of crossovers during meiosis in budding yeast. *Genetics* 164(1), pp. 81–94.

Shinohara, M., Oh, S.D., Hunter, N. and Shinohara, A. 2008. Crossover assurance and

1010   crossover interference are distinctly regulated by the ZMM proteins during yeast meiosis. *Nature Genetics* 40(3), pp. 299–309.

Storlazzi, A., Gargano, S., Ruprich-Robert, G., Falque, M., David, M., Kleckner, N. and Zickler, D. 2010. Recombination proteins mediate meiotic spatial chromosome organization and pairing. *Cell* 141(1), pp. 94–106.

1015   Svetlanov, A., Baudat, F., Cohen, P.E. and de Massy, B. 2008. Distinct functions of MLH3 at recombination hot spots in the mouse. *Genetics* 178(4), pp. 1937–1945.

Al-Sweel, N., Raghavan, V., Dutta, A., Ajith, V.P., Di Vietro, L., Khondakar, N., Manhart, C.M., Surtees, J.A., Nishant, K.T. and Alani, E. 2017. mlh3 mutations in baker's yeast alter meiotic recombination outcomes by increasing noncrossover events genome-wide. *PLoS Genetics*

1020   13(8), p. e1006974.

Tang, S., Wu, M.K.Y., Zhang, R. and Hunter, N. 2015. Pervasive and essential roles of the Top3-Rmi1 decatenase orchestrate recombination and facilitate chromosome segregation in

meiosis. *Molecular Cell* 57(4), pp. 607–621.

Tsubouchi, T., Zhao, H. and Roeder, G.S. 2006. The meiosis-specific zip4 protein regulates crossover distribution by promoting synaptonemal complex formation together with zip2. *Developmental Cell* 10(6), pp. 809–819.

Viera, A., Santos, J.L., Page, J., Parra, M.T., Calvente, A., Cifuentes, M., Gómez, R., Lira, R., Suja, J.A. and Rufas, J.S. 2004. DNA double-strand breaks, recombination and synapsis: the timing of meiosis differs in grasshoppers and flies. *EMBO Reports* 5(4), pp. 385–391.

Walpita, D., Plug, A.W., Neff, N.F., German, J. and Ashley, T. 1999. Bloom's syndrome protein, BLM, colocalizes with replication protein A in meiotic prophase nuclei of mammalian spermatocytes. *Proceedings of the National Academy of Sciences of the United States of America* 96(10), pp. 5622–5627.

Wang, K., Tang, D., Wang, M., Lu, J., Yu, H., Liu, J., Qian, B., Gong, Z., Wang, X., Chen, J., Gu, M. and Cheng, Z. 2009. MER3 is required for normal meiotic crossover formation, but not for presynaptic alignment in rice. *Journal of Cell Science* 122(Pt 12), pp. 2055–2063.

Wang, T.F., Kleckner, N. and Hunter, N. 1999. Functional specificity of MutL homologs in yeast: evidence for three Mlh1-based heterocomplexes with distinct roles during meiosis in recombination and mismatch correction. *Proceedings of the National Academy of Sciences of the United States of America* 96(24), pp. 13914–13919.

Ward, J.O., Reinholdt, L.G., Motley, W.W., Niswander, L.M., Deacon, D.C., Griffin, L.B., Langlais, K.K., Backus, V.L., Schimenti, K.J., O'Brien, M.J., Eppig, J.J. and Schimenti, J.C. 2007. Mutation in mouse *Hei10*, an e3 ubiquitin ligase, disrupts meiotic crossing over. *PLoS Genetics* 3(8), p. e139.

1045   Woods, L.M., Hodges, C.A., Baart, E., Baker, S.M., Liskay, M. and Hunt, P.A. 1999.

Chromosomal influence on meiotic spindle assembly: abnormal meiosis I in female Mlh1

mutant mice. *The Journal of Cell Biology* 145(7), pp. 1395–1406.

Zakharyevich, K., Tang, S., Ma, Y. and Hunter, N. 2012. Delineation of joint molecule

resolution pathways in meiosis identifies a crossover-specific resolvase. *Cell* 149(2), pp. 334–

1050   347.

G *Mus musculus* MLH3 constructs

H Purification of MutLγ complex

I Mass Spectrometry identification of SDS gel bands

BAND	PROTEIN NAME	PROTEIN MW (KDa)	PEPTIDE NUMBER	PROTEIN SCORE	% PROTEIN COVERAGE	MOLAR %
Upper	His$_{10}$-MLH3	160	90	2077	64	95
Lower	MBP-MLH1	127.5	90	2413	79	94

E MLH1 Foci at pachynema

F Frequency of synapsed autosomes with no MLH1 foci in pachynema

G Chiasmata counts at diakinesis

H Bivalent and univalent chromosomes