1 **Phylogenomics reveals major diversification rate shifts in the evolution of silk moths and**
2 **relatives**
3

4 Hamilton CA[1,2*], St Laurent RA[1], Dexter, K[1], Kitching IJ[3], Breinholt JW[1,4], Zwick A[5], Timmermans
5 MJTN[6], Barber JR[7], Kawahara AY[1*]
6
Institutional Affiliations:
[1]Florida Museum of Natural History, University of Florida, Gainesville, FL 32611 USA
[2]Department of Entomology, Plant Pathology, & Nematology, University of Idaho, Moscow, ID
83844 USA
[3]Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK
[4]RAPiD Genomics, 747 SW 2nd Avenue #314, Gainesville, FL 32601. USA
[5]Australian National Insect Collection, CSIRO, Clunies Ross St, Acton, ACT 2601, Canberra,
Australia
[6]Department of Natural Sciences, Middlesex University, The Burroughs, London NW4 4BT, UK
[7]Department of Biological Sciences, Boise State University, Boise, ID 83725, USA
*Correspondence: hamiltonlab@uidaho.edu (CAH) or kawahara@flmnh.ufl.edu (AYK)

18
19
Abstract

The silkmoths and their relatives (Bombycoidea) are an ecologically and taxonomically
diverse superfamily that includes some of the most charismatic species of all the Lepidoptera.
Despite displaying some of the most spectacular forms and ecological traits among insects,
relatively little attention has been given to understanding their evolution and the drivers of
their diversity. We heavily sampled (both in taxa and loci) all major lineages of the
Bombycoidea, producing a well-supported phylogeny that identified important evolutionary
patterns (e.g., morphology, biogeography, and differences in speciation and extinction).
Importantly, analysis of diversification rates highlights the stark increases that exist within the
Sphingidae (hawkmoths) and Saturniidae (wild silkmoths). We postulate that these rate shifts
are due to differences in the intense selective pressures from insectivorous bats. The study also
introduces a new Bombycoidea-specific Anchored Hybrid Enrichment (AHE) probe set, a
modified DNA extraction protocol for Lepidoptera specimens from natural history collections,
and additional information on the existing AHE bioinformatics pipeline. Our research highlights
the flexibility of AHE to generate genomic data from a wide range of museum specimens, both
age and preservation method, and will allow researchers to tap into the wealth of biological
data residing in natural history collections around the globe.

37
38 **Keywords**
39 Anchored Hybrid Enrichment, Bombycoidea, Lepidoptera, natural history collections
40
Background
The Bombycoidea include some of the most charismatic moth species of all the
Lepidoptera. This ecologically diverse superfamily comprises ten families, 520 genera, and
6,092 species (1). Although widespread globally, the highest diversity of bombycoids occurs in

the tropics and subtropics. This diversity includes the most spectacular forms (i.e., range of
body sizes and wing shapes) and functions (i.e., mimicry, predator avoidance, flight capabilities,
and feeding strategies) in the Lepidoptera (2). Their large body sizes, ease of rearing in captivity
and utilization of larval silk has driven human interest in these moths.
A number of bombycoids have become important contributors to human culture,
originally as economically important species for either sericulture or as agricultural pests, but
more recently as model organisms for comparative studies of genetics, development, and
physiology (2). Additionally, many lineages play important roles as pollinators (3), (4), (5), (6),
(7), (8), (9), or as indicators in biodiversity and habitat quality assessments (10). Of the 10
families, three contain species that have been used as model organisms (Bombycidae,
Saturniidae, and Sphingidae). Unfortunately, the relationships among these three families have
remained largely elusive, with Bombycidae considered the sister lineage to either Saturniidae or
Sphingidae (11), (12), (13), (14), (15), (16), (17), (18); but see Breinholt and Kawahara (19).
Overall, relatively little attention has been given to the group with regards to understanding
their evolution.
Despite their charisma and intrigue, the lack a robust phylogeny based on broad taxon
sampling is dramatically affecting our ability to answer fundamental questions about the drivers
of their diversity. Monophyly of the Bombycoidea has been supported by six morphological
synapomorphies (20), but Zwick (21) determined that only two of these were systematically
informative: one poorly understood thoracic character (22), and one relating to forewing vein
arrangements (A. Zwick, unpublished). Recent molecular studies of bombycoid systematics (11),
(13), (21) have resulted in substantial differences from morphology-based phylogenetic
hypotheses (20), (22), (23). To date, nearly all molecular studies of the superfamily have
included fewer than 20 protein-coding genes for ≤50 species (11), (13), (16), (24). Although
these studies agreed on the monophyly of the superfamily, many relationships among families,
subfamilies, and tribes remained unclear and were characterized by weak branch support or
conflicting signal. Modern phylogenomics (e.g., based on Anchored Hybrid Enrichment) has
been shown to be effective to resolve relationships among Lepidoptera at multiple taxonomic
levels (25), (26), (27), (28), (29), including within the Bombycoidea (25), (30). However, studies
that utilize phylogenomics to resolve inter- and intra-familial relationships within the
Bombycoidea have been limited by sampling.
Perhaps the most interesting evolutionary aspect of this superfamily is their striking link
with bats, their main nocturnal predators (30), (31), (32), (33), (34). Bombycoids possess a
number of traits that appear to be anti-bat adaptations in response to echolocating bats, a
lineage thought to have arisen approximately 60 million years ago (35), (36), (37), (38), (39).
These nightly battles between bats and moths drove a predator/prey "arms race" that
produced anti-bat strategies such as hearing organs (40), (41), (42), (43), ultrasound producing
organs capable of jamming bat sonar (33), (44), evasive flight strategies (both in aerial
maneuvers and temporal partitioning of activity (45), and a variety of different anti-bat wing
morphologies (30), (46).
The most spectacular examples of these anti-bat strategies can be found in the
Saturniidae and Sphingidae – the two lineages that harbor the majority of described bombycoid
species (30), (32), (33), (47). The divergent life-history strategies of these two families (31), (48)
has likely played a major role in driving their diversity by exposing these lineages to differing

predatory selective forces. For example, the majority of hawkmoths feed as adults, seeking out
nectar resources during their relatively long lives (weeks to months) during which females
experience multiple mating events and retain the eggs internally for long periods of time to
allow egg maturation and host plant discovery (31). This ecological strategy is significantly
different from saturniids (16), (19), (49), which depend entirely upon the resources acquired
during the larval period. Adult saturniids possess reduced or non-functional mouthparts, lay
eggs almost immediately after mating, do not have ears or "jamming" organs, and exhibit
erratic evasive flight (31). These traits firmly establish the relative ecological roles of Saturniidae
and Sphingidae as an important natural experiment from which we can gather valuable
information regarding the evolution of predator/prey interactions in moths and their ultimate
effects on the diversification process in general.
To better answer questions regarding bombycoid evolution, we applied Anchored
Hybrid Enrichment (AHE) targeted-sequencing phylogenomics (50), sampling 571 loci in 117
taxa across Bombycoidea. In order to do this, we developed a new AHE probe set ("BOM1"),
redesigned from the "LEP1" probe set of Breinholt et al. (25), that more efficiently captures loci
across the Bombycoidea, allowing us to establish a robust evolutionary framework for future
research. This new probe set also includes loci that were not in the original "LEP1" (25) set, but
were part of existing Sanger-sequencing datasets (11), (13), (24), (51). Finally, we present a
modified DNA extraction protocol for use with historical natural history specimens, allowing
researchers to tap into the biological wealth of data residing in the many museum collections
around the globe.
110
Methods
The "BOM1" AHE probe set design
Anchored Hybrid Enrichment (AHE) is a targeted-sequencing methodology designed to
"capture" hundreds of unique orthologous loci (i.e., single copy, phylogenetically-informative
markers) from across the genome, for resolving both shallow and deep-level evolutionary
relationships (11), (13). Probes are designed to anchor in conserved regions that are flanked by
variable regions randomly spread throughout the genome. This approach creates a diverse set
of informative loci that include exons, introns, intergenic, and conserved regions of the
genome. Targeted-sequencing approaches like AHE provide mechanisms whereby different
researchers can confidently and effectively use the same loci for independent projects, allowing
for the combination of data across studies.
To build a more Bombycoidea-specific AHE probe set and phylogenomic dataset, we
began with the Lepidoptera Agilent Custom SureSelect Target Enrichment "LEP1" probe kit,
designed for 855 loci (21). We then modified this kit, henceforth the new "BOM1" kit, by
evaluating which loci were most phylogenetically-informative within the superfamily,
optimizing the set of probes to recover these loci, and including 24 previously-sequenced
"legacy" Sanger-sequenced loci, e.g., CO1 for species confirmation in BOLD (1), from (11), (13),
(24), and (51) (see Supp. Table 1 for loci names), as well as eight vision-related genes. The
probes for these vision-related genes are based phototransduction genes mined from eye or
head transcriptomes (unpublished; generated by AYK), and are included for future analyses to
investigate their evolution across the group.

Phylogenetically-informative loci were determined by examining the sequence variation
present in 56 bombycoid species from across the taxonomic breadth of the superfamily (55 AHE
samples, pulled from (25) or generated for this study, and the *Bombyx mori* reference genome
(52)). These samples were sequenced and processed using the "LEP1" probe kit and Breinholt
et al. (25) bioinformatics pipeline (53). Individual gene trees were inferred for each of the 855
loci, using Maximum Likelihood in RAxML v8.2 (54), under a GTRGAMMA model of evolution
with 100 non-parametric bootstraps for node support. For each tree, phylogenetic
informativeness was calculated using PhyDesign (55), (56), (57),
http://phydesign.townsend.yale.edu. Parsimony informative characters and number of
segregating sites were calculated using the R packages 'phyloch' (58) and 'ape' (59)
respectively. Additionally, each phylogeny was scrutinized by eye to determine the level of
informative branching (i.e., branch lengths and topological patterns). Loci deemed to be
phylogenetically uninformative, or those that were capturing poorly (<60% of sampled species
represented for a locus), were removed from consideration. The final "BOM1" probe kit
comprised 571 loci, 539 of which came from the original "LEP1" kit.
147
Taxon sampling
To build a "backbone" phylogeny of the Bombycoidea, we sampled all major lineages
(i.e., families, subfamilies, and tribes), with the exception of three rare, species-poor groups:
subfamily Munychryiinae (Anthelidae) and tribes Sataspedini and Monardini (Sphingidae), for
which representative samples were unavailable for DNA sequencing. In total, 115 ingroup
Bombycoidea species from 97 genera were included in the phylogenetic analysis, as well as two
lasiocampid outgroups (Supp. Table 2).
Specimens were obtained from fieldwork, historically preserved dry collections (Supp
Table 1), and molecular tissue collections. Field-collected specimens were stored in ≥95%
ethanol, RNAlater, or papered and dried with silica gel, and stored in freezers at -80°C. Genomic
DNA was extracted using OmniPrep Genomic DNA Extraction Kits (G-Biosciences, St. Louis, MO,
USA) and Dneasy Blood and Tissue Kits (Qiagen, Valencia, CA, USA). DNA concentration was
evaluated through agarose gel electrophoresis and fluorometery using a Qubit 2.0 (Invitrogen,
Thermo Fisher Scientific, Carlsbad, CA, USA). Library preparation, hybridization enrichment, and
Illumina HiSeq 2500 sequencing (PE100) was carried out by RAPiD Genomics (Gainesville, FL,
USA). All specimen wing vouchers and tissue storage methods follow Cho et al. (60). All ethanol,
RNAlater, freshly papered specimens, or previously extracted DNA were stored at -80°C in the
molecular tissue collection of the MGCL.
166
DNA extraction protocol for museum specimens
We evaluated the efficacy of obtaining DNA from historical museum specimens because
there is great interest in understanding the feasibility of this approach (28), (61), (62), (63). The
Lepidoptera specimens evaluated herein, were "field-pinned" (never rehydrated), "papered"
(stored in an envelope and kept dry since collected, thus not rehydrated or pinned), and
"traditionally-pinned" specimens (dried, rehydrated, and subsequently pinned) – the
historically most common method of Lepidoptera specimen storage. Collecting dates ranged
from 1987 to 2017 (Supp. Table 2). The samples themselves were not preserved for molecular
work and information about potential contaminants and/or extraction inhibitors (i.e.

fumigation compounds, other chemicals) was unavailable. In some cases there was little soft-
tissue to extract from within the abdomens, having degraded over time. Being that these
samples were not kept in molecular-grade conditions, many were contaminated with fungal
and bacterial growth, which could be identified visually by the spores left on the bodies or by
the smell of decay. These factors along with the amount of sclerotized tissue present and the
fact that the abdomens used needed to stay intact (not homogenized) for dissecting purposes,
made extracting good quality genomic DNA challenging.
The method, detailed in the Supplemental information, attempts to account for several
factors: the amount of degraded tissue, the presence of eggs, the relative fat content, and the
overall abdomen size. Many of the commercial DNA extraction kits on the market (including the
Omni Prep kit used in this study) recommend using 10mg-20mg of well preserved, soft tissue
for the extraction process. Given that the museum specimens used had been desiccated for
many years, a number of abdomens had little to no visible internal soft tissue left. We increased
the amount of proteinase K to lysis buffer ratio in hopes that the proteinase K would digest the
remaining DNA rich material in solution. The relative fat content of samples is an issue because
lipid-rich tissue can interfere with the digestion of the soft tissue as well as change the
chemistry of the DNA isolation buffers. The lipid-rich layer in the Chloroform phase separation
step made pipetting of the DNA rich supernatant difficult as well. Any specimen that appeared
to be "greasy" or seemed to have an oily film on the abdomen was not used for extraction.
Finally, the overall size of the abdomen dictated the amount of lysis buffer needed in order to
sufficiently submerge the abdomen and reach the available soft tissue, which then dictated the
volume of reagents needed for the remainder of the DNA isolation process. Lastly, and
importantly, the modified protocol will allow a user not only to obtain DNA, but also easily
prepare the genitalia for taxonomic or future morphological work using standard techniques.
Detailed explanation of the methods can be found in the Supplemental information.
201
Bioinformatics
The bioinformatics pipeline of Breinholt et al. (25) was used to clean and assemble raw
Illumina reads for each AHE locus. The pipeline uses a probe-baited iterative assembly that
extends beyond the "probe" region, checks for quality and cross contamination due to barcode
leakage, removes paralogs, and returns a set of aligned orthologs for each locus and taxon of
interest. To accomplish these tasks, the pipeline uses the *Bombyx mori* genome (52), and
"LEP1" or "BOM1" reference libraries – depending on the kit used for sequencing.
Loci for phylogenetic analysis were selected by applying a cutoff of ≥40% sampled taxa
recovery – meaning that for a locus to be included in the analysis, the locus had to be recovered
in at least 40% of the sampled taxa. The pipeline then evaluates density and entropy at each
site of a nucleotide sequence alignment. We elected to trim with entropy and density cutoffs
only in flanking regions, allowing the "probe" region (exon) to be converted into amino acid
sequences. For a site (outside of the "probe" region) to remain, that site must pass a 60%
density and 1.5 entropy cutoff, rejecting sites that fail these requirements. A higher first value
(60) increases the coverage cutoff (e.g., a site is kept if 60% of all taxa are represented at that
site). A higher second value (1.5) increases the entropy cutoff (i.e., entropy values represent the
amount of saturation at a site); sites with values higher than 1.5 possess higher saturation and
are thus deleted). AliView v. 1.18 (64) was used to check for frame shifts, recognize and remove

stop codons (if present), and edit sequencing errors or lone/dubious indels. Reading frames
were manually checked after "probe" region extraction prior to translation. Translation was
carried out with FASconCAT-G v1.02 (65). Because flanking sequences are generally "non-
coding" and sites have been deemed homologous (see 25), these flanking sequence regions
before and after the "probe" regions (exons) were separated from the exons, combined, and
treated together as an independent partition.
Of the 115 bombycoid and two outgroup specimens, 110 were sequenced directly using
the AHE target capture sequencing, of which 68 were sequenced using the "BOM1" kit and 42
using the "LEP1" kit. Seven specimens had their AHE loci "probe" regions mined from previously
sequenced transcriptomes or the *Bombyx mori* genome (Supp. Table 2). These specimens did
not have "flanking" data because of their transcriptome-data origin. All specimens were
processed using either the 'Bmori' (for "LEP1") or 'BOM1' (for "BOM1") reference libraries.
Previous scripts are available in Dryad (53). For reproducibility, instructions on how to use the
pipeline and additional scripts not used by Breinholt et al. (25) are provided in Dryad (available
upon publication).
235
Phylogenetics
Lepidoptera AHE probe sets comprise highly-conserved coding "probe" regions (i.e.,
exons) and more variable, generally non-coding "flanking" regions (e.g., introns or intergenic
regions) located on either side of the "probe" region (25). We evaluated both nucleotide and
amino acid datasets to examine phylogenetic signal and the role that saturation may play in the
data (see 19). Using the Breinholt et al. (25) pipeline and scripts (53), also included herein, three
datasets were built for phylogeny inference: 1) AA = an amino acid supermatrix composed of
translated "probe" region loci; 2) Pr+Fl = a "probe" + "flanking" supermatrix; and 3) ASTRAL =
the individual loci from the Pr+Fl supermatrix, used for individual gene tree inference and then
species tree estimation. Additional analyses (i.e., "probe" region-only, as nucleotides) were
investigated, but are not reported here because their outcomes did not differ from those
reported.
Concatenated supermatrices were assembled using FASconCAT-G v1.02 (65).
Phylogenetic inference was performed in a maximum likelihood (ML) framework using IQ-TREE
MPI multicore v1.5.3 (66). For nucleotide datasets, the '–m TEST' command was used in IQ-
TREE to perform a model of evolution search resembling that of jModelTest/ProtTest (66),
applying models to each partition (AA) or site (Pr+Fl) to determine the best model of amino acid
or nucleotide substitution. For all inferences, we performed 1000 random addition sequence
(RAS) replicates, and 1000 replicates each for both ultrafast bootstraps (UFBS) ('–bb' command)
and SH-aLRT tests ('-alrt' command). The SH-like approximate likelihood ratio test (SH-aLRT)
estimates branch support values that have been shown to be as conservative as the commonly
used non-parametric bootstrap values (67). SH-aLRT and bootstrap values tend to agree for
data sets with strong phylogenetic signal (i.e., datasets with loci that are "sufficiently" large and
tips that share "sufficiently" diverged sequences). Disagreements in branch support are thought
to arise as a consequence of small sample size, insufficient data, or saturated divergence levels
(see 68). We classified nodes as "robust" if they were recovered with support values of UFBS ≥
95and SH-aLRT ≥ 80 (67), (68).

Because concatenation can be misleading when there are high levels of incomplete
lineage sorting or deep coalescence (69), we assessed the impact of potential gene-tree
discordance (70), (71), (72) by inferring a phylogeny for each individual locus, using IQ-TREE
under the same parameters as above (i.e., "probe" and single partition of "flanking" loci were
modeled by site). Species tree estimation was performed in ASTRAL-III (73). ASTRAL is a
computationally efficient and statistically consistent (under the multi-species coalescent)
nonparametric method that takes input gene trees and estimates a highly accurate species
tree, even when there is a high level of incomplete lineage sorting (74). To evaluate support on
the species tree, we used the ASTRAL support values (ASV) – local posterior probabilities that
are more precise than evaluating bootstrap values across a set of input trees (75). ASTRAL
support values were determined to be "robust" if nodes were recovered with local posterior
probabilities ≥ 0.95. All pipeline steps and phylogenomic analyses were conducted on the
University of Florida HiPerGator HPC (http://www.hpc.ufl.edu/). All alignment FASTA files, loci
information, partition files, tree files, and other essential data files used for phylogenetic
inference are available as supplementary materials on Dryad (available upon publication).
278
Rogue taxon & outlier locus analyses
We investigated whether our molecular data included rogue taxa or outlier loci that
were potentially influencing our phylogenetic inferences. A rogue taxon analysis was carried
out using the online version of RogueNaRok (76), (77), http://rnr.h-its.org/, on the 1000
ultrafast bootstrap trees and the consensus tree from the Pr+Fl supermatrix. Outlier taxa and
loci analyses were carried out using Phylo-MCOA v.1.4 (PMCoA) (78) in R (on the 650 gene trees
used in the ASTRAL analysis). No rogue taxa or loci were found and therefore taxa were not
pruned in subsequent analyses.
287
Diversification rate analyses
As an initial investigation into why some bombycoid lineages are more diverse than
others, we examined and quantified how diversification rates (the interplay between speciation
and extinction) have changed over time. Instead of simply calculating species diversity per clade
and assuming extant diversity is a true indicator of increases in diversification rate, an approach
prone to biases, we applied BAMM (79) and 'BAMMtools' (80) to infer the number and location
of macroevolutionary rate shifts across our phylogeny, as well as to visualize the 95% credible
set of shift configurations. Because the Bombycoidea fossil record is poor (81), it is difficult to
infer a robust dated phylogeny and therefore we chose not to do so for this study. We took the
ML best tree and converted it into a relative-rate scaled ultrametric tree using the 'chronopl'
command in the R package 'ape' (59). To account for non-random missing speciation events, we
quantified the percentage of taxa sampled within each family and incorporated these in the
form of branch-specific sampling fractions. Sampling fractions were based on the updated
superfamily numbers calculated from Kitching and Rougiere et al. (1). Informed priors, based on
our sampling and phylogeny, were determined using 'setBAMMpriors' in BAMMtools. The
MCMC chain was run for 100 million generations, sampling every 1000 generations.
Convergence diagnostics was assessed using the R package 'coda' (82). The first 20% of runs
were discarded as burn-in.
306

Results

307
While there is no guaranteed method for extracting high quality Lepidoptera gDNA from
non-molecular grade specimens, our abdomen soaking approach proved relatively consistent
and yielded us enough target DNA to proceed with AHE sequencing. The datasets built for
phylogeny inference contained: 1) AA = 579 loci, 48,456 amino acid residues, modeled by locus;
2) Pr+Fl = 649 "probe" loci + one "flanking" locus (261,780 bp), each "probe" locus was
modeled by site, "flanking" data was maintained as a single partition and modeled by site; and
3) ASTRAL = 650 loci, each locus modeled by site.
Increasing the number of loci and species significantly improved our understanding of
Bombycoidea relationships. The inferred relationships are generally consistent across the three
phylogenetic inferences that we performed (AA, Pr+Fl, ASTRAL), with all major backbone
lineages robustly supported, and all bombycoid families *sensu* Zwick (21) and Zwick et al. (13)
recovered as monophyletic. Due to the methodological approach (i.e., the treatment of
different data types) and the more biologically realistic and parsimonious explanation of the
topology (see Systematics section in Supplemental), our preferred phylogeny was the "probe +
flanking" (Pr+Fl) dataset. All family-level placements of genera *sensu* Kitching and Rougiere et
al. (1) were supported, with the exception of *Arotros* Schaus, a genus long considered to be an
epiine bombycid (83). *Arotros* is clearly nested within Apatelodidae in our tree (Fig. 1; Supp. Fig.
1) and is hereby transferred to Apatelodidae (see supplemental text for additional details on
this transfer). The family-level rearrangements of Zwick (21) and Zwick et al. (13) were also
recovered in our phylogenetic results: the placement of taxa formerly classified in the
"Lemoniidae" (e.g., Lemaire & Minet (23)) are recovered within Brahmaeidae; the Apatelodidae
are distinct from the Bombycidae and Phiditiidae; and the broader concept of Endromidae,
which includes taxa formerly placed within the "Mirinidae" and Bombycidae (e.g., Lemaire &
Minet (23)), are recovered as a monophyletic group.
We find broad congruence with the major groupings designated by Zwick (21), though
internal relationships within these groups may not match previous phylogenies exactly.
Historically, the most problematic familial placement in the superfamily has been the
Bombycidae *sensu stricto*. Traditional Sanger-sequencing phylogenetics based on a handful of
gene regions, applied in (13), (18), and (21), placed this family either as sister to the Saturniidae
or to the Sphingidae (reviewed in 19), albeit without strong support for either. Our study clearly
places the Bombycidae as the sister lineage to the (Saturniidae + Sphingidae) clade, as seen in
both the AA and Pr+Fl datasets, but not in the ASTRAL inference – an outcome that mirrors
traditional studies that used low numbers of loci and lacked the phylogenetic signal to
confidently resolve relationships among these three families (Supp. Trees 1-3). The 'CAPOPEM'
group (Carthaeidae, Anthelidae, Phiditiidae, and Endromidae) is recovered in all three analyses,
though in the ASTRAL inference this group is nested within the 'SBS' group (Sphingidae,
Bombycidae and Saturniidae) - due to placement of the Bombycidae (Supp. Fig. 2). The relative
interfamilial relationships within the 'CAPOPEM' clade were not robustly supported in Zwick et
al. (13), and although our more refined taxon sampling and increased genetic coverage
confidently solidifies the group, the data still fail to fully resolve these relationships across
analyses (Supp. Fig. 2). Zwick et al. (13) recovered the Old World Endromidae as sister to the
Australian Anthelidae + (Australian Carthaeidae + Neotropical Phiditiidae). However, our Pr+Fl
analysis supports the Anthelidae as sister to Endromidae + (Carthaeidae + Phiditiidae) instead.

Effectively, Anthelidae and Endromidae swap places within 'CAPOPEM' when comparing Zwick
et al. (13) and our Pr+Fl phylogeny. Conversely, our AA results illustrate yet a different picture,
in which Phiditiidae are sister to Anthelidae + (Carthaeidae + Endromidae). This discordance is
important to note considering the disjunct geographic distribution of these families. Among
these families, only Phiditiidae is found in (and is endemic to) the New World, with each of the
other families being restricted to the Old World. Intrafamilial relationships in the 'SBS' and
'CAPOPEM' groups change depending on which of our datasets are used to infer the phylogeny
(Fig. 1, Supp. Figs. 1-3, Supp. Trees 1-3). Lastly, the 'BALE' group (Brahmaeidae, Apatelodidae,
and Eupterotidae) is recovered in all three analyses.
When we investigate intrafamilial relationships, the Bombycidae (disregarding the
taxonomically "incorrect" *Arotros*, mentioned above; see Supplemental text) possesses distinct
and monophyletic subfamilies Bombycinae (Old World) and Epiinae (New World; Fig. 1; Supp.
Fig. 1). Within the Eupterotidae, the subfamilies Striphnopteryginae and Janinae are
monophyletic, but others are not (Fig. 1; Supp. Fig. 1). Within the Saturniidae, the Pr+Fl and AA
topologies are similar to Regier et al. (51) (Fig. 1 & Supp. Fig. 3). The Oxyteninae are sister to
the rest of the family, followed by the Cercophaninae. Arsenurinae is the sister lineage to the
Hemileucinae and the Ceratocampinae, while Salassinae is the sister lineage to the Saturniinae.
A major subfamily relationship difference lies with the Agliinae, sister to the Salassinae +
Saturniinae (Pr+Fl) or sister to the Arsenurinae (AA, also (13) and (51)). The ASTRAL inference is
quite different, placing the Ceratocampinae as the sister lineage to a clade containing the
Agliinae, Arsenurinae + Hemileucinae, and a Saturniinae + Salassinae clade (Supp. Fig. 3),
though with low support. Within the Sphingidae, our topologies (both Pr+Fl & AA) are quite
different from Zwick et al. (13). Instead of the Smerinthinae being paraphyletic, the
Smerinthinae are monophyletic and the Sphinginae are polyphyletic (Fig. 1; Supp. Fig. 1). This
outcome is robustly supported in the Pr+Fl and ASTRAL phylogenies, though internal structuring
is slightly different. In the AA phylogeny, while the placement of the Macroglossinae, Langiinae
and Sphinginae are robustly supported (and the same topology as in Pr+Fl), low internal
support obfuscates relationships within the Smerinthinae.
Importantly, a number of subfamilies and genera are not monophyletic, based on our
sampling and the most current taxonomic definition (see Kitching and Rougerie et al. (1)) (Fig.
1; Supp. Fig. 1). Within Eupterotidae, the Eupterotinae and the "Ganisa group" are not
monophyletic. The Eupterotinae are rendered paraphyletic due to the placements of
Panacelinae and Striphnopteryginae. The Striphnopteryginae genus *Phiala* is paraphyletic with
respect to *Lichenopteryx*, and the "Ganisa group" is polyphyletic due to the traditional inclusion
of the genus *Neopreptos* (84). Within the Anthelidae, the genus *Anthela* is paraphyletic due to
the placement of *Nataxa* and *Pterolocera*, a finding in agreement with (21) . And lastly, within
the Sphingidae, Sphinginae is polyphyletic due to the placement of *Pentateucha* as sister to the
Langiinae, and the Smerinthinae genus *Polyptychus* is paraphyletic.
To quantify whether certain bombycoid lineages exhibit differing diversification rates,
we evaluated how the interplay between speciation and extinction has changed over relative
time rather than using just the simple sums of described species in each family. At the family
scale, a major diversification rate shift occurs along the lineage leading to the Saturniidae and
Sphingidae (Fig. 1). Unfortunately, due to the poor Bombycoidea fossil record and the poor
ability of computational approaches to estimate extinction rates, we are not able to discern

whether the differences that exist are due to increases or decreases in speciation, although the
raw numbers, based on (1), do agree with the outcomes found herein: Sphingidae (1,602 spp.);
Saturniidae (3,454 spp.); Bombycidae (202 spp.); Endromidae (70 spp.); Phiditiidae (23 spp.);
Carthaeidae (1 sp.); Anthelidae (94 spp.); Eupterotidae (396 spp.); Brahmaeidae (68 spp.);
Apatelotidae (182 spp.).
400
Discussion
Within the past decade, only a handful of studies have attempted to use molecular
sequence data to understand the phylogenetic relationships of the Bombycoidea (11), (13), (21)
and none corroborated the earlier subjective morphology-based hypotheses (20), (23). So, to
establish a foundational backbone for future evolutionary, comparative, and taxonomic studies,
we sampled exemplars from all major lineages in the superfamily and used Anchored Hybrid
Enrichment (AHE) phylogenomics to provide a robust phylogeny of the superfamily based on
the largest taxonomic and molecular sampling to date.
To achieve dense taxonomic coverage, we included many samples from natural history
collections, and modified previous DNA extraction methodology to obtain DNA for high-
throughput sequencing. We also redesigned the AHE probe set developed by Breinholt et al.
(25) and modified it to more efficiently recover phylogenetically-informative loci within the
Bombycoidea, as well as adding "legacy" Sanger-sequenced loci for increasing the usability of
the data (e.g., integration of "legacy" datasets and CO1 for species identification). To assist
future usage of this and other AHE probe sets, we expanded the command log instructions of
Breinholt et al. (25) to include step-by-step instructions for running the bioinformatics pipeline.
Our phylogeny largely reinforces the phylogenetic results of earlier molecular works,
although with topological differences within families (i.e., some subfamilies and genera are not
recovered as monophyletic in our study). Topological discordances with trees published in
previous works are likely to be the product of increased locus sampling, which provided
significantly more phylogenetic information. For example, our results provide a well-supported
placement of the historically problematic family Bombycidae *sensu stricto*, as the sister lineage
to Saturniidae + Sphingidae. An earlier study (19) had provided some evidence in support of this
relationship, but its sampling was very limited. Additionally, it is quite likely that for the deep to
mid-timescale nodes, the AA phylogeny is the best hypothesis for Bombycoidea relationships.
Our study highlights how morphological convergence in Bombycoidea has confused our
understanding of their evolution. This convergence is sometimes exceptional. For example, the
bombycid genus *Rotunda*, endemic to the Old World, and the Apatelodidae *Arotros*, endemic to
the New World, are widely separated on the phylogenetic tree (Fig. 1; Supp. Fig. 1) and
geographically, yet are astonishingly similar in body size, wing shape, and phenotypic
appearance (see Systematics section in Supplemental Information). Additionally, when we
examine the 'CAPOPEM' group closely, we see a biogeographical distribution that, based on the
AA and ASTRAL topologies, appears to be a more biologically realistic and parsimonious
evolutionary history due to the placement of the Neotropical Phiditiidae as sister to the
remaining and Old World 'CAPOPEM' families (Supp. Fig. 2).
One of the most surprising results of the present study is the dramatic shift in
diversification rates within both the Sphingidae and Saturniidae compared to other bombycoid
lineages. These differences in "survivability" may be explained by their very different life-

history strategies, morphology (e.g., body sizes and wing shapes), flight speed and
maneuverability (31), as well as very different anti-bat strategies (30), (32), (33), (47). Although
we did not conduct a divergence time estimation analysis in this study, the origin of the 'SBS'
group has been postulated to be around 50 mya (85), with hawkmoths originating soon
thereafter (33). Insectivorous bats are thought to have originated roughly 60 mya, and the
diversification of Sphingidae and Saturniidae around that time suggests that the incredible
taxonomic diversity within these two families is in part due to bat-related selection pressures
resulting in diverse anti-bat traits, traits that other bombycoid lineages lack.
However, the hawkmoth and silk moth evolutionary story is certainly more complicated
than simply reflecting their interactions with bats. As evolution is often complex, there are
undoubtedly a suite of factors playing significant roles in their diversity of form and function.
Although we hypothesize that the differences in diversification rates are correlated with bat
predation, it is possible that these rate shifts are due to other factors, such as ecological
specialization or shifts in host plant usage, both as larvae and adults. Amassing and collating
behavioral and ecological datasets for the tips of the Bombycoidea Tree of Life for
macroevolutionary comparative investigations is essential to furthering our understanding of
this diverse, global superfamily, as well as understanding how bats, ecological traits, and/or
biogeographical history may or may not have shaped their diversity.
457

Acknowledgements
We thank our collaborators Seth Bybee, Deborah Glass, Yash Sondhi, Jamie Theobald, and
Emmanuel Touissant. We are indebted to those individuals that provided specimens for this
project, contributed help in identifying specimens, or in other ways: Samantha Epstein, John
Heppner, Geena Hill, Nick Homziak, Jacqueline Miller, Charles Mitter, Kim Mitter, Lary Reeves,
Andrei Sourakov, and Andrew Warren.
464

465 **Funding**
466 This work was partly supported by NSF grants [IOS-1121807, JRB], [IOS-1121739, AYK], [DBI-
467 1349345, AYK], [DEB-1557007, AYK], and [PRFB-1612862, CAH]; a NERC grant [NE/P003915/1,
468 IJK]; National Geographic Society grants [CRE 9944-16, JRB]; and postdoctoral support from the
469 Florida Museum of Natural History (to CAH).
470

471 **Author Contributions**
472 CAH, AYK, JWB, IJK, MT, AZ designed the study. CAH and JWB analyzed data. CAH, AYK, RAS
473 wrote the paper. CAH, AYK, JRB, KD, IJK, RAS, AZ collected, identified, and processed specimens.
474 All authors discussed results and contributed to the writing of the manuscript.
475

Supplemental Information
The datasets supporting this article have been uploaded as part of the supplementary
material or can be found in Dryad, (available upon publication). Supplemental Information
includes details of the DNA extraction protocol using museum specimens, three figures, one
table, detailed systematic information on the Bombycoidea families, bioinformatics pipeline
scripts, detailed instructions on running the pipeline, all alignment FASTA files (nucleotides,
amino acids, supermatrices, and individual loci), partition files, tree files, and other essential

data files used for phylogenetic inference and analysis. Previous Breinholt et al. bioinformatics
scripts are available in Dryad (53).

Figures

Figure 1. Maximum likelihood tree of Bombycoidea, based on 650 AHE loci. All nodes are
supported by ≥95% bootstrap values unless otherwise noted. Branch color indicates the
estimated diversification rate, with warmer colors representing lineages with higher rates.
Major taxonomic groups such as families and subfamilies are labeled. Photographs represent
lineages sampled in the phylogeny. Species-level diversity, based on Kitching and Rougerie et al.
(1), are also noted next to Family.

References

1.   Kitching I, Rougerie R, Zwick A, St Laurent R, Ballesteros Mejia L, Kawahara A, et al. A
global checklist of the Bombycoidea (Insecta: Lepidoptera). BDJ. 2018 Feb
12;6(1):e22236–13.

2.   Roe AD, Weller SJ, Baixeras J, Brown J, Cummings MP, Davis DR, et al. Evolutionary
framework for Lepidoptera model systems. In: Goldsmith M, Marec F, editors. Molecular
Biology and Genetics of the Lepidoptera. 2010. pp. 1–24.

3.   Gregory D. Hawkmoth pollination in the genus *Oenothera*. Aliso A Journal of Systematic
and Evolutionary Botany. 1963;5(3):357–84.

4.   Haber WA, Frankie GW. A Tropical Hawkmoth Community- Costa Rican Dry Forest
Sphingidae. Biotropica. 1989 Dec 9;21(2):155–72.

5.   Nilsson LA, Jonsson L, Rason L, Randrianjohany E. Monophily and pollination mechanisms
in *Angraecum arachnites* Schltr. (Orchidaceae) in a guild of long-tongued hawk-moths
(Sphingidae) in Madagascar. Biological Journal of the Linnean Society. 1985;26(1):1–19.

6.   Nilsson LA, Rabakonandrianina E, Pettersson B. Exact tracking of pollen transfer and
mating in plants. Nature. 1992;360(6405):666–8.

7.   Wasserthal LT. The pollinators of the Malagasy star orchids *Angraecum sesquipedale*, *A.*
sororium and *A. compactum* and the evolution of extremely long spurs by pollinator shift.
Botanica Acta. 1997 Oct;110(5):343–59.

8.   Nilsson LA. Deep flowers for long tongues. Trends In Ecology & Evolution. 1998
Jul;13(7):259–60.

9.   Raguso RA, Willis MA. Synergy between visual and olfactory cues in nectar feeding by
naïve hawkmoths, *Manduca sexta*. Animal Behaviour. 2002 Nov 1;64(5):685–95.

10.   Beck J, Kitching IJ, Linsenmair KE. Effects of habitat disturbance can be subtle yet
significant: biodiversity of hawkmoth-assemblages (Lepidoptera: Sphingidae) in

Southeast-Asia. Biodiversity and Conservation. 2nd ed. Kluwer Academic Publishers;
2006 Jan;15(1):465–86.

11.   Regier JC, Cook CP, Mitter C, Hussey A. A phylogenetic study of the "bombycoid
complex" (Lepidoptera) using five protein-coding nuclear genes, with comments on the
problem of macrolepidopteran phylogeny. Systematic Entomology. 2008;33:175–89.

12.   Meusemann K, Reumont von BM, Simon S, Roeding F, Strauss S, Kück P, et al. A
phylogenomic approach to resolve the Arthropod Tree of Life. Mol Biol Evol. 2010 Oct
15;27(11):2451–64.

13.   Zwick A, Regier JC, Mitter C, Cummings MP. Increased gene sampling yields robust
support for higher-level clades within Bombycoidea (Lepidoptera). Systematic
Entomology. 2011;36(1):31–43.

14.   Kim MJ, Kang AR, Jeong HC, Kim K-G, Kim I. Reconstructing intraordinal relationships in
Lepidoptera using mitochondrial genome data with the description of two newly
sequenced lycaenids, *Spindasis takanonis* and *Protantigius superans* (Lepidoptera:
Lycaenidae). Mol Phylogenet Evol. Elsevier Inc; 2011 Nov 1;61(2):436–45.

15.   Simon S, Narechania A, Desalle R, Hadrys H. Insect phylogenomics: exploring the source
of incongruence using new transcriptomic data. Genome Biol Evol. Oxford University
Press; 2012 Dec 1;4(12):1295–309.

16.   Regier JC, Mitter C, Zwick A, Bazinet AL, Cummings MP, Kawahara AY, et al. A large-scale,
higher-level, molecular phylogenetic study of the insect order Lepidoptera (moths and
butterflies). Moreau CS, editor. PLoS ONE. 2013 Mar 12;8(3):e58568–23.

17.   Timmermans MJTN, Lees DC, Simonsen TJ. Towards a mitogenomic phylogeny of
Lepidoptera. Mol Phylogenet Evol. Elsevier Inc; 2014 Oct 1;79(C):169–78.

18.   Wang X, Chen Z-M, Gu X-S, Wang M, Huang G-H, Zwick A. Phylogenetic relationships
among Bombycidae s.l. (Lepidoptera) based on analyses of complete mitochondrial
genomes. Systematic Entomology. John Wiley & Sons, Ltd; 2018 Nov 20;20:537–9.

19.   Breinholt JW, Kawahara AY. Phylotranscriptomics: saturated third codon positions
radically influence the estimation of trees based on next-gen data. Genome Biol Evol.
2013 Nov 1;5(11):2082–92.

20.   Minet J. The Bombycoidea: Phylogeny and higher classification (Lepidoptera, Glossata).
Entomologica Scandinavica. 1994;25(1):63–88.

21.   Zwick A. Molecular phylogeny of Anthelidae and other bombycoid taxa (Lepidoptera:
Bombycoidea). Systematic Entomology. 2008 Jan 7;33:190–209.

22.   Minet J. Tentative reconstruction of the ditrysian phylogeny (Lepidoptera, Glossata).
553   Entomologica Scandinavica. 1991;22(1):69–95.

554   23.   Lemaire C, Minet J. 18. The Bombycoidea and their relatives. In: Kristensen NP, editor.
555   Lepidoptera, Moths and Butterflies vol 1 Evolution, Systematics, and Biogeography
556   Handbook of Zoology. Walter de Gruyter, Berlin; 1998. pp. 322–53.

557   24.   Regier JC, Zwick A, Cummings MP, Kawahara AY, Cho S, Weller S, et al. Toward
558   reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia):
559   an initial molecular study. BMC Evolutionary Biology. BioMed Central; 2009;9(1):280–21.

560   25.   Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. Resolving
561   relationships among the megadiverse butterflies and moths with a novel pipeline for
562   anchored phylogenomics. Syst Biol. 2018;67(1):78–93.

563   26.   Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A
564   comprehensive and dated phylogenomic analysis of butterflies. CURBIO. Elsevier Ltd;
565   2018 Mar 5;28(5):770–5.

566   27.   Johns CA, Toussaint EFA, Breinholt JW, Kawahara AY. Origin and macroevolution of
567   micro-moths on sunken Hawaiian Islands. Proc Biol Sci. The Royal Society; 2018 Aug
568   29;285(1885):20181047–10.

569   28.   St Laurent RA, Hamilton CA, Kawahara AY. Museum specimens provide phylogenomic
570   data to resolve relationships of sack-bearer moths (Lepidoptera, Mimallonoidea,
571   Mimallonidae). Systematic Entomology. Wiley/Blackwell (10.1111); 2018 May
572   31;43(4):729–61.

573   29.   Toussaint EFA, Breinholt JW, Earl C, Warren AD, Brower AVZ, Yago M, et al. Anchored
574   phylogenomics illuminates the skipper butterfly tree of life. BMC Evolutionary Biology.
575   BMC Evolutionary Biology; 2018 Jun 16;18(101):1–11.

576   30.   Rubin JJ, Hamilton CA, McClure CJW, Chadwell BA, Kawahara AY, Barber JR. The
577   evolution of anti-bat sensory illusions in moths. Science Advances. American Association
578   for the Advancement of Science; 2018 Jul;4(7).

579   31.   Janzen D. Two ways to be a tropical big moth: Santa Rosa saturniids and sphingids.
580   Oxford surveys in evolutionary biology. 1984;1(85):140.

581   32.   Barber JR, Kawahara AY. Hawkmoths produce anti-bat ultrasound. Biology Lett. 2013
582   May 8;9(4):20130161–1.

583   33.   Kawahara AY, Barber JR. Tempo and mode of antibat ultrasound production and sonar
584   jamming in the diverse hawkmoth radiation. Proceedings of the National Academy of
585   Sciences. 2015 May 19;112(20):6407–12.

34. Kawahara AY, Plotkin D, Hamilton CA, Gough H, Laurent RS, Owens HL, et al. Diel
behavior in moths and butterflies: a synthesis of data illuminates the evolution of
temporal activity. Organisms Diversity & Evolution; 2017 Dec 4;:1–15.

35. Jones KE, Bininda-Emonds ORP, Gittleman JL. Bats, clocks, and rocks: diversification
patterns in Chiroptera. Evolution. 2005 Oct;59(10):2243–55.

36. Teeling EC, Springer MS, Madsen O, Bates P, O'Brien SJ, Murphy WJ. A molecular
phylogeny for bats illuminates biogeography and the fossil record. Science. American
Association for the Advancement of Science; 2005 Jan 28;307(5709):580–4.

37. Miller-Butterworth CM, Murphy WJ, O'Brien SJ, Jacobs DS, Springer MS, Teeling EC. A
family matter: conclusive resolution of the taxonomic position of the long-fingered bats,
Miniopterus. Mol Biol Evol. 2007 Jul;24(7):1553–61.

38. Shi JJ, Rabosky DL. Speciation dynamics during the global radiation of extant bats.
Evolution. 2015 Jun;69(6):1528–45.

39. Lei M, Dong D. Phylogenomic analyses of bat subordinal relationships based on
transcriptome data. Nature Publishing Group. 2016 Jun 13;6(1):27726.

40. Roeder KD, Treat AE, Vandeberg JS. Auditory sense in certain hawkmoths. Science.
1968;159:331–3.

41. Spangler HG. Moth hearing, defense, and communication. Annual Review of Entomology.
Annual Reviews 4139 El Camino Way, PO Box 10139, Palo Alto, CA 94303-0139, USA;
1988;33(1):59–81.

42. Göpfert MC, Surlykke A, Wasserthal LT. Tympanal and atympanal "mouth-ears" in
hawkmoths (Sphingidae). P Roy Soc B-Biol Sci. The Royal Society; 2002 Jan
7;269(1486):89–95.

43. Hofstede ter HM, Goerlitz HR, Ratcliffe JM, Holderied MW, Surlykke A. The simple ears of
noctuoid moths are tuned to the calls of their sympatric bat community. J Exp Biol. The
Company of Biologists Ltd; 2013 Nov 1;216(Pt 21):3954–62.

44. Barber JR, Conner WE. Acoustic mimicry in a predator-prey interaction. P Natl Acad Sci
Usa. National Academy of Sciences; 2007;104(22):9331–4.

45. Lamarre GPA, Mendoza I, Rougerie R, Decaëns T, Hérault B, Bénéluz F. Stay out (almost)
all night: contrasting responses in flight activity among tropical moth assemblages.
Neotrop Entomol. 2015 Feb 10;44(2):109–15.

46. Zhong M, Hill GM, Gomez JP, the DPTJO, 2016. Quantifying wing shape and size of
saturniid moths with geometric morphometrics. Journal of the Lepidopterists' Society.
2016;70(2):1–9.

47.   Barber JR, Leavell BC, Keener AL, Breinholt JW, Chadwell BA, McClure CJW, et al. Moth
621   tails divert bat attack: evolution of acoustic deflection. Proceedings of the National
622   Academy of Sciences. National Acad Sciences; 2015 Mar 3;112(9):2812–6.

623   48.   Tammaru T, Haukioja E. Capital breeders and income breeders among Lepidoptera:
624   consequences to population dynamics. Oikos. 1996;77(3):561–4.

625   49.   Kawahara AY, Breinholt JW. Phylogenomics provides strong evidence for relationships of
626   butterflies and moths. P Roy Soc B-Biol Sci. 2014 Jun 18;281(1788):20140970–0.

627   50.   Lemmon AR, Emme SA, Lemmon EM. Anchored Hybrid Enrichment for massively high-
628   throughput phylogenomics. Syst Biol. 2012 Aug 11;61(5):727–44.

629   51.   Regier JC, Grant MC, Mitter C, Cook CP, Peigler RS, Rougerie R. Phylogenetic relationships
630   of wild silkmoths (Lepidoptera: Saturniidae) inferred from four protein-coding nuclear
631   genes. Systematic Entomology. 2008 Feb 4;33(2):219–28.

632   52.   Xia Q, Zhou Z, Lu C, Cheng D, Dai F, Bin Li, et al. A draft sequence for the genome of the
633   domesticated silkworm (*Bombyx mori*). Science. American Association for the
634   Advancement of Science; 2004 Dec 10;306(5703):1937–40.

635   53.   Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. Data from:
636   Resolving relationships among the megadiverse butterflies and moths with a novel
637   pipeline for Anchored Phylogenomics. Dryad Digital Repository; 2017.

638   54.   Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
639   phylogenies. Bioinformatics. 2014.

640   55.   Mayrose I, Graur D, Ben-Tal N, Pupko T. Comparison of site-specific rate-inference
641   methods for protein sequences: empirical Bayesian methods are superior. Mol Biol Evol.
642   2004 May 21;21(9):1781–91.

643   56.   Pond SLK, Frost SDW, Muse SV. HyPhy: hypothesis testing using phylogenies. Springer,
644   New York, NY; 2005. pp. 125–81.

645   57.   López-Giráldez F, Townsend JP. PhyDesign: an online application for profiling
646   phylogenetic informativeness. BMC Evolutionary Biology. BioMed Central Ltd; 2011 May
647   31;11(1):152.

648   58.   Heibl C. PHYLOCH: R language tree plotting tools and interfaces to diverse phylogenetic
649   software packages. 2013.

650   59.   Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R
651   language. Bioinformatics. 2004 Jan 20;20(2):289–90.

60.   Cho S, Epstein SW, Mitter K, Hamilton CA, Plotkin D, Mitter C, et al. Preserving and
vouchering butterflies and moths for large-scale museum-based molecular research.
PeerJ. 2016;4(4):e2160.

61.   Blaimer BB, Lloyd MW, Guillory WX, Brady SG. Sequence capture and phylogenetic utility
of genomic Ultraconserved Elements obtained from pinned insect specimens. O'Grady P,
editor. PLoS ONE. Public Library of Science; 2016 Aug 24;11(8):e0161531–20.

62.   Mikheyev AS, Zwick A, Magrath MJL, Grau ML, Qiu L, Su YN, et al. Museum genomics
confirms that the Lord Howe Island stick insect survived extinction. CURBIO. Elsevier Ltd;
2017 Oct 23;27(20):3157–3161.e4.

63.   Ruane S, Austin CC. Phylogenomics using formalin-fixed and 100+ year-old intractable
natural history specimens. Molecular Ecology Resources. Wiley/Blackwell (10.1111);
2017 Feb 13;17(5):1003–8.

64.   Larsson A. AliView: a fast and lightweight alignment viewer and editor for large datasets.
Bioinformatics. 2014.

65.   Kück P, Longo GC. FASconCAT-G: extensive functions for multiple sequence alignment
preparations concerning phylogenetic studies. Frontiers in Zoology. 2014;11(81).

66.   Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015
Jan;32(1):268–74.

67.   Minh BQ, Nguyen MAT, Haeseler von A. Ultrafast approximation for phylogenetic
bootstrap. Mol Biol Evol. 2013 Apr 15;30(5):1188–95.

68.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms
and methods to estimate Maximum-Likelihood phylogenies: assessing the performance
of PhyML 3.0. Syst Biol. 2010 May;59(3):307–21.

69.   Mendes FK, Hahn MW. Why concatenation fails near the anomaly zone. Syst Biol. 2017
Jul 11;67(1):158–69.

70.   Maddison W. Gene trees in species trees. Syst Biol. 1997 Sep 1;46(3):523–36.

71.   Slowinski J, Page R. How should species phylogenies be inferred from sequence data?
Syst Biol. 1999;48(4):814–25.

72.   Edwards SV. Is a new and general theory of molecular systematics emerging? Evolution.
Wiley Online Library; 2009;63(1):1–19.

73.   Zhang C, Sayyari E, Mirarab S. ASTRAL-III: increased scalability and impacts of contracting
low support branches. In: RECOMB International Workshop on Comparative Genomics.
2017. pp. 53–75.

74.   Mirarab S, Warnow T. ASTRAL-II: coalescent-based species tree estimation with many
hundreds of taxa and thousands of genes. Bioinformatics. 2015 Jun 9;31:i44–i52.

75.   Sayyari E, Mirarab S. Anchoring quartet-based phylogenetic distances and applications to
species tree reconstruction. BMC Genomics. BioMed Central; 2016 Nov 11;17(10):101–
13.

76.   Aberer AJ, Stamatakis A. A simple and accurate method for rogue taxon identification.
2011. pp. 118–22.

77.   Aberer AJ, Krompass D, Stamatakis A. Pruning rogue taxa improves phylogenetic
accuracy: an efficient algorithm and webservice. Syst Biol. 2012 Nov 8;62(1):162–6.

78.   de Vienne DM, Ollier S, Aguileta G. Phylo-MCOA: a fast and efficient method to detect
outlier genes and species in phylogenomics using multiple co-inertia analysis. Mol Biol
Evol. 2012 May 15;29(6):1587–98.

79.   Rabosky DL. Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. Kolokotronis S-O, editor. PLoS ONE. 2014 Feb
26;9(2):e89543.

80.   Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, et al. BAMMtools: an R
package for the analysis of evolutionary dynamics on phylogenetic trees. Kembel S,
editor. Methods in Ecology and Evolution. 2014 May 10;5(7):701–7.

81.   Sohn JC, Labandeira CC, Davis DR, Mitter C. An annotated catalog of fossil and subfossil
Lepidoptera (Insecta: Holometabola) of the world. Zootaxa. 2012.

82.   Plummer M, Best N, Cowles K, Vines K. CODA: convergence diagnosis and output analysis
for MCMC. R News. 2006;6(1):7–11.

83.   Schaus W. Familie Bombycidae (Epiinae). In: Seitz A, editor. Die Gross-Schmetterlinge der
Erde. Kernen, Stuttgart; 1929. pp. 675–711. (Die amerikanischen Spinner und
Schwärmer).

84.   Nässig WA, Oberprieler RG. An annotated catalogue of the genera of Eupterotidae
(Insecta, Lepidoptera, Bombycoidea). Senckenbergiana biologica. 2008;88(1):53–80.

85.   Misof B, Liu S, Meusemann K, Peters RS, Donath A. Phylogenomics resolves the timing
and pattern of insect evolution. Science. 2014;346(6210):763–7.

Andriasa contraria
Falcatula falcata
Pseudoclanis kenyae
Polyptychus digitatus
Polyptychus andosa
Marumba juvencus
Clanis undulosa
Mimas tiliae
Smerinthus jamaicensis
Coequosa triangularis
Parum colligata
Callambulyx amanda
Cypa decolor
Ambulyx canescens

Smerinthinae

46.7/91

Sphingidae
1,602 spp.

Sphinx chersis
Dolbina tancrei
Hopliocnema brachycera

Sphinginae

Langia zenzeroides
Langiinae
Pentateucha curiosa
Sphinginae
74.8/92
Macroglossum sylvia
Hemaris thysbe
Pseudosphinx tetrio

Macroglossinae

Bunaea alcinoe
Eudaemonia agriphontes
Micragone agathylla
Saturnia pavonia
Rothschildia lebeau

Saturniinae

23.6/85
Salassa sp.
Salassinae
Aglia japonica
Agliinae
Polythysana cinerascens
Hemileucinae
Hemileuca magnifica

Saturniidae
3,454 spp.

Anisota pellucida
Eacles imperialis
Ceratocampinae
70.1/92
Almeidaia aidae
Arsenura armida
Arsenurinae
Cercophana venusta
Janiodes ecuadoriensis
Cercophaninae
Oxytenis naemia
Oxyteninae

'SBS'

Epia lunilinea
Epia muscosa
74.1/94
Colla micacea
New genus new species
Colla sp.
Quentalia lividia
Quentalia melchthalia
Anticla ortygia
Anticla antica

Epiinae

Bombycidae
202 spp.

Penicilifera sp.
Ernolatia moorei
Triuncina brunnea
Trilocha sp.
Racinoa sp.
Gunda ochracea
Bombyx mori
Rotunda rotundapex

Bombycinae

Oberthueria formosibia
Oberthueria caeca
Mustilizans dierli
Mustilia glabrata
Comparmustilia sphingiformis
Pseudandraca flavamaculata
Andraca trilochoides
Mirina christophi
Endromis versicolora
Prismosticta fenestrata

Endromidae *70 spp.*

39.6/86
Phiditia sp.
Phiditia sp.
Sorocaba sp.

Phiditiidae *23 spp.*

Carthaea saturnioides
Carthaeidae *1 sp.*

48.2/91
Anthela asterias
Anthela ocellata
Nataxa flavescens
Pterolocera amplicornis
Anthela varia
Chelepteryx collesi

Anthelidae *94 spp.*

'CAPOPEM'

Cotana serranotata
Eupterotinae
Panacela lewinae
Panacelinae
Eupterote sp. Malaysia
Eupterotinae
Phiala sp.
Lichenopteryx despecta
Phiala wichgrafi
Striphnopteryginae
Sphingonatha asclepiades
Eupterotinae
Neopreptos sp.
"Ganisa group"

72.2/92
Melanothrix sp. Phillipines
Melanothrix sp. Malaysia
Ganisa similis
Pseudoganisa currani
Pseudojana sp.

"Ganisa group"

Eupterotidae *396 spp.*

Jana strigina
Jana preciosa
Rhodopteriana sp.
Hoplojana sp.
Hilbrides sp.
88.4/98
Camerunia sp.
Stenoglene uniformis
Acrojana scutaea

Janinae

Bombycoidea

Brahmaea japonica
Brahmaea hearseyi
Brahmaea europaea
Dactyloceras sp.
Spiramiopsis comma

Brahmaeidae *68 spp.*

Lemonia philopalus
Ephoria (Zanola) liliana
48.7/95
Zanola sp.
Zanola sp.
Ephoria marginalis

'BALE'

44.9/92
Olceclostera amoria
Olceclostera seraphica
Falcatelodes anava
Bombycidae per Kitching et al., (2018)
Arotros striata *
Apatelodes pithala
Apatelodes firmiana
Prothysana sp.
Drepatelodes sp.

Apatelotidae *182 spp.*

Drepatelodes quadrilineata
Trabala hantu
Artace sp.

Lasiocampidae

Lasiocampoidea

Smerinthus jamaicensis

Sphinx asella

Eudaemonia argiphontes

Arsenura sp.

Bombyx mandarina

Endromis versicolor

Carthaea saturnioides

Phiditia sp.

Anthela sp.

Jana eurymas

Brahmaea paukstadtorum

Apatelodes torrefacta

19
16
12
9.1