Phylogenomics resolves major relationships and reveals significant diversification rate shifts

in the evolution of silk moths and relatives

Hamilton CA[1,2*], St Laurent RA[1], Dexter, K[1], Kitching IJ[3], Breinholt JW[1,4], Zwick A[5], Timmermans

MJTN[6], Barber JR[7], Kawahara AY[1*]

[1]Florida Museum of Natural History, University of Florida, Gainesville, FL 32611 USA

[2]Department of Entomology, Plant Pathology, & Nematology, University of Idaho, Moscow, ID

83844 USA

[3]Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

[4]RAPiD Genomics, 747 SW 2nd Avenue #314, Gainesville, FL 32601. USA

[5]Australian National Insect Collection, CSIRO, Clunies Ross St, Acton, ACT 2601, Canberra,

Australia

[6]Department of Natural Sciences, Middlesex University, The Burroughs, London NW4 4BT, UK

[7]Department of Biological Sciences, Boise State University, Boise, ID 83725, USA

[*]Corresponding authors: hamiltonlab@uidaho.edu (CAH) or kawahara@flmnh.ufl.edu (AYK)

Abstract

Background

The silkmoths and their relatives constitute the ecologically and taxonomically diverse

superfamily Bombycoidea, which includes some of the most charismatic species of Lepidoptera.

Despite displaying some of the most spectacular forms and ecological traits among insects,

relatively little attention has been given to understanding their evolution and the drivers of

their diversity.

Results

To begin to address this problem, we created a new Bombycoidea-specific Anchored Hybrid

Enrichment (AHE) probe set and sampled up to 571 loci for 117 taxa across all major lineages of

the Bombycoidea, producing a well-supported phylogeny. The tree was overall consistent with

prior morphological and molecular studies, although some taxa (e.g., *Arotros* Schaus) were

misplaced in the Bombycidae and here formally transferred to Apatelodidae. We identified

important evolutionary patterns (e.g., morphology, biogeography, and differences in speciation

and extinction), and our analysis of diversification rates highlights the stark increases that exist

within the Sphingidae (hawkmoths) and Saturniidae (wild silkmoths).

Conclusions

We postulate that these rate shifts are due to the well-documented bat-moth "arms race" and

differences in selective pressures from insectivorous bats. The study establishes a backbone for

future evolutionary, comparative, and taxonomic studies, and presents a modified DNA

extraction protocol that allows Lepidoptera specimens to be readily sequenced from pinned

natural history collections, succeeding in samples up to 30 years old. Our research highlights

the flexibility of AHE to generate genomic data from a wide range of museum specimens, both

age and preservation method, and will allow researchers to tap into the wealth of biological

data residing in natural history collections around the globe.

Keywords

Anchored Hybrid Enrichment, targeted sequence capture, phylogenomics, Bombycoidea,

Lepidoptera, natural history collections

Background

The Bombycoidea include some of the most charismatic moths among all Lepidoptera.

This ecologically diverse superfamily comprises ten families, 520 genera, and 6,092 species [1].

Although widespread globally, the highest diversity of bombycoids occurs in the tropics and

subtropics. This diversity includes the most spectacular forms (i.e., range of body sizes and wing

shapes) and functions (i.e., mimicry, predator avoidance, flight capabilities, and feeding

strategies) in the Lepidoptera [2].

A number of bombycoids have become important contributors to human culture,

originally as economically important species for sericulture or as agricultural pests, but more

recently as model organisms for comparative studies of genetics, development, and physiology

[2]. Additionally, many lineages play important roles as pollinators ([3], [4], [5], [6], [7], [8], [9]),

or as indicators in biodiversity and habitat quality assessments [10]. Of the 10 families, three

contain species that have been used as model organisms (Bombycidae, Saturniidae, and

Sphingidae). Unfortunately, relationships among bombycoid families, and especially these three

families, have remained largely elusive. For example, Bombycidae have been considered the

sister lineage to either Saturniidae or Sphingidae ([11], [12], [13], [14], [15], [16], [17], [18]); but

see Breinholt and Kawahara [19]. Overall, relatively little attention has been given to the group

with regards to understanding their evolution.

Despite their charisma and intrigue, the lack of a robust phylogeny based on broad and

dense taxon sampling across the Bombycoidea is dramatically affecting our ability to answer

fundamental questions about the drivers of their diversity. Monophyly of the Bombycoidea has

been supported by six morphological synapomorphies [15], but Zwick [20] determined that only

two of these were systematically informative: one poorly understood thoracic character [21],

and one relating to the arrangement of the forewing vein (A. Zwick, unpublished). Recent

molecular studies of bombycoid systematics ([11], [13], [20]) have resulted in substantial

differences in terms of relationships from morphology-based phylogenetic hypotheses ([15],

[21], [22]). To date, nearly all molecular studies of bombycoids have included fewer than 20

protein-coding genes for ≤50 species (e.g., [11], [13], [23], [24]). Although these studies agreed

on the monophyly of the superfamily, many relationships among families, subfamilies, and

tribes remained unclear and were characterized by weak branch support or conflicting signal.

Modern phylogenomics (e.g., based on Anchored Hybrid Enrichment) has been shown to be

effective to resolve relationships among Lepidoptera at multiple taxonomic levels ([16], [25],

[26], [27], [28]) including the Bombycoidea ([16], [29]). However, those studies that utilize

phylogenomics to resolve inter- and intra-familial relationships within the Bombycoidea have

been limited by taxon sampling.

To better answer questions regarding bombycoid evolution, we applied Anchored

Hybrid Enrichment (AHE) targeted-sequencing phylogenomics [30]. We developed a new,

bombycoid-specific, AHE probe set (here called "BOM1"), redesigned from the "LEP1" probe set

of Breinholt et al. [16]. Our new probe set captures "legacy" Sanger sequencing-based loci that

were part of existing bombycoid molecular datasets ([11], [13], [24], [31]), enabling the merging

of older published datasets with those generated from the BOM1 probe set. To further improve

the ability to generate a dataset with greater taxon and locus sampling, we developed a new

method for extracting DNA from pinned natural history specimens, and show that our

extraction approach is successful in obtaining DNA sequence data for phylogenomics. In total,

our dataset resulted in 571 loci for 117 species across Bombycoidea.

We also use this opportunity to utilize the phylogeny to examine patterns of

diversification in the superfamily. Bombycoidea are well-known to have multiple different

ecological life-history strategies, especially for the Saturniidae and Sphingidae – two lineages

that harbor the majority of described bombycoid species ([29], [32], [33], [34]). The divergent

life-history strategies of these two families ([35], [36]) has likely played a major role in driving

their diversity. For example, the majority of hawkmoths feed as adults, seeking out nectar

resources during their relatively long lives (weeks to months). During this time, females

experience multiple mating events and retain the eggs internally for long periods to allow egg

maturation and host plant discovery [35]. This ecological strategy is significantly different from

saturniids ([23], [19], [37]), which depend entirely upon the resources acquired during the larval

period. Adult saturniids possess reduced or non-functional mouthparts and lay eggs almost

immediately after mating. Furthermore, these lineages possess a number of different traits

that appear to be anti-bat adaptations in response to echolocating bats, a lineage thought to

have arisen approximately 60 million years ago ([38], [39], [40], [41], [42]). Hawkmoths,

especially those in the Sphinginae and Macroglossinae, are strong fliers thought to have

evolved hearing organs and ultrasound producing organs capable of jamming bat sonar ([33],

[43]). Saturniids, in contrast, lack hearing organs and exhibit erratic evasive flight [35] and

hindwing tails that deflect bat echoes ([29], [34]). These traits firmly establish the relative

ecological roles of Saturniidae and Sphingidae as an important natural experiment from which

we can gather valuable information regarding the evolution of predator/prey interactions in

moths and their ultimate effects on the diversification process in general.

Methods

The "BOM1" AHE probe set design

Anchored Hybrid Enrichment (AHE) is a targeted-sequencing methodology designed to

capture hundreds of unique orthologous loci (i.e., single copy, phylogenetically-informative

markers) from across the genome, for resolving both shallow and deep-level evolutionary

relationships ([11], [13]). Probes are designed to anchor in conserved regions that are flanked

by variable regions randomly spread throughout the genome. This approach creates a diverse

set of informative loci that include exons, introns, intergenic, and conserved regions of the

genome. Targeted-sequencing approaches, like AHE and UCE (Ultraconserved Elements),

provide mechanisms whereby different researchers can confidently and effectively use the

same loci for independent projects, allowing for the combination of data across studies.

Breinholt et al. [16] constructed a Lepidoptera Agilent Custom SureSelect Target

Enrichment "LEP1" probe kit, designed for 855 loci. However, this probe set is not specific to

Bombycoidea, and does not include some of the traditional loci that have been used in

phylogenetics of Bombycoidea. In order to build a more Bombycoidea-specific AHE probe set

and phylogenomic dataset, we began by modifying the LEP1 kit, evaluating which loci were

most phylogenetically-informative within the superfamily, optimizing the set of probes to

recover these loci, and including 24 previously-sequenced Sanger-sequenced loci, such as e.g.,

CO1, CAD, DDC, period, wingless and others from [11], [13], [24], and [31], as well as eight

vision-related genes (see Supp. Table 1 for locus names). The probes for these vision-related

genes are based phototransduction genes mined from eye or head transcriptomes

(unpublished; generated by AYK), and are included for future analyses to investigate their

evolution across the superfamily.

To determine the informative loci for BOM1, phylogenetically-informative loci were

identified by examining the sequence variation of 56 bombycoid species from across the

taxonomic breadth of the superfamily (55 AHE samples, from either Breinholt et al. [16] or

generated for this study, plus loci mined from the *Bombyx mori* reference genome [44]). These

samples were sequenced and processed using the LEP1 probe kit and Breinholt et al. [16]

bioinformatics pipeline [45]. Individual gene trees were generated for each of the 855 loci,

using Maximum Likelihood in RAxML v8.2 [46], under a GTRGAMMA model of evolution with

100 non-parametric bootstraps for node support. For each tree, phylogenetic informativeness

was calculated using PhyDesign ([47], [48], [49]). Parsimony informative characters and the

number of segregating sites were calculated using the R packages 'phyloch' [50] and 'ape' [51]

respectively. Additionally, each phylogeny was scrutinized visually to determine whether

branch lengths and topological patterns appeared realistic (i.e., no significant outliers present).

Loci deemed to be phylogenetically uninformative, or those that were capturing poorly (<60%

of sampled species represented for a locus), were excluded from the probe set. The final BOM1

probe kit comprised 571 loci, 539 of which came from the original LEP1 kit.

Taxon sampling

To build a backbone phylogeny of the Bombycoidea, we sampled all major lineages (i.e.,

families, subfamilies, and tribes), with the exception of three rare, species-poor groups:

subfamily Munychryiinae (Anthelidae) and tribes Sataspedini and Monardini (Sphingidae), for

which representative samples were unavailable for DNA sequencing. Sampled lineages were

chosen because: 1) they were appropriate representatives of the taxonomic group needed for

the analysis (i.e., good morphological and evolutionary representative of a tribe); and 2) they

were accessible for use in phylogenomics. In total, 115 ingroup Bombycoidea species from 97

genera were included in the phylogenetic analysis, as well as two Lasiocampidae outgroups –

the sister lineage to the bombycoids (Supp. Table 2).

Specimens were obtained from fieldwork, historically preserved dry collections (Supp.

Table 1), and molecular tissue collections. Field-collected specimens were stored in ≥95%

ethanol, RNAlater, or papered and dried with silica gel. Genomic DNA was extracted using

OmniPrep Genomic DNA Extraction Kits (G-Biosciences, St. Louis, MO, USA) and DNeasy Blood

and Tissue Kits (Qiagen, Valencia, CA, USA). DNA concentration was evaluated through agarose

gel electrophoresis and fluorometry using a Qubit 2.0 (Invitrogen, Thermo Fisher Scientific,

Carlsbad, CA, USA). Library preparation, hybridization enrichment, and Illumina HiSeq 2500

sequencing (PE100) was carried out at RAPiD Genomics (Gainesville, FL, USA). Specimen wing

vouchering and tissue storage methods follow Cho et al. [52]. All DNA extracts and specimens

preserved in ethanol, RNAlater, or those freshly papered were stored at -80°C at the Florida

Museum of Natural History, McGuire Center of Lepidoptera and Biodiversity (MGCL).

DNA extraction protocol for museum specimens

We evaluated the efficacy of obtaining DNA from historical museum specimens because

there is great interest in understanding the feasibility of this approach for use in phylogenetics

and systematics ([27], [53], [54], [55]). The Lepidoptera specimens evaluated herein, were

"field-pinned" (never rehydrated), "papered" (stored in an envelope and kept dry since

collected, thus not rehydrated or pinned), and "traditionally-pinned" specimens (dried,

rehydrated, and subsequently pinned) – the historically most common method of Lepidoptera

specimen storage. Collecting dates ranged from 1987 to 2017 (Supp. Table 2). Samples were

not initially intended to be preserved for molecular sequencing and information about potential

contaminants and/or extraction inhibitors (i.e. fumigation compounds, other chemicals) was

unavailable. In some cases there was little soft-tissue to extract from within the abdomens,

having degraded over time. Being that these samples were not kept in molecular-grade

conditions, many were contaminated with fungal and bacterial growth, which could be

identified visually by the spores left on the bodies or by the smell of decay. These factors along

with the amount of sclerotized tissue present and the fact that the abdomens used needed to

stay intact (not homogenized) for dissecting purposes, made extracting good quality genomic

DNA challenging.

Our extraction method, detailed in the Supplemental information, attempts to account

for several factors: the amount of degraded tissue, the presence of eggs, the relative fat

content, and the overall abdomen size. Many commercial DNA extraction kits on the market

(including the Omni Prep kit used in this study) recommend using 10 mg-20 mg of well

preserved, soft tissue for the extraction process. Given that the museum specimens used had

been desiccated for many years, a number of abdomens had little to no visible internal soft

tissue remaining. To digest the remaining material in solution, we increased the ratio of

proteinase K to lysis buffer. Companies that produce DNA extractions kits know, as can be seen

by their specific extraction kits, the relative fat content of samples is an issue because lipid-rich

tissue can interfere with the digestion of the soft tissue as well as change the chemistry of the

DNA isolation buffers. Specimens that appeared to be "greasy" or seemed to have an oily film

on the abdomen were not used for extraction. The overall size of the abdomen was used to

estimate the amount of lysis buffer needed in order to sufficiently submerge the abdomen and

reach the available soft tissue. The amount of buffer also reflected the volume of reagents

needed for the remainder of the DNA isolation process. Lastly, the modified protocol also

allows a user to easily prepare the genitalia for taxonomic or morphological taxonomic work as

these structures remain undamaged.

Bioinformatics

The bioinformatics pipeline of Breinholt et al. [16] was used to clean and assemble raw

Illumina reads for each AHE locus. The pipeline uses a probe-baited iterative assembly that

extends beyond the probe region, checks for quality and cross contamination due to barcode

leakage, removes paralogs, and returns a set of aligned orthologs for each locus and taxon of

interest. To accomplish these tasks, the pipeline uses the *Bombyx mori* genome [44], and an

AHE reference library, which in this study was the BOM1 reference library.

Loci for phylogenetic analysis were selected by applying a cutoff of ≥40% sampled taxa

recovery (i.e., for a locus to be included in the analysis, the locus had to be recovered in at least

40% of the sampled taxa). The pipeline evaluates density and entropy at each site of a

nucleotide sequence alignment. We elected to trim with entropy and density cutoffs only in

"flanking" regions, allowing the "probe" region (exon) to be converted into amino acid

sequences. For a site (outside of the probe region) to remain, that site must pass a 60% density

and 1.5 entropy cutoff, rejecting sites that fail these requirements. A higher first value (60)

increases the coverage cutoff (e.g., a site is kept if 60% of all taxa are represented at that site).

A higher second value (1.5) increases the entropy cutoff (i.e., entropy values represent the

amount of saturation at a site; sites with values higher than 1.5 possess higher saturation and

are thus deleted). AliView v1.18 [56] was used to translate to amino acids, check for frame

shifts, recognize and remove stop codons, and edit sequencing errors or lone/dubious indels.

Because flanking sequences are generally non-coding and sites have been deemed homologous

(see [16]), these flanking sequences, before and after the probe regions, were separated from

the exons, then combined and treated together as an independent partition. Due to the

filtering steps in the bioinformatics pipeline (i.e., site orthology, and density and saturation

evaluation), the flanking partition can be viewed as a SNP supermatrix, where each site is

homologous, but uninformative sites, saturated sites, or sites with large amounts missing data

have been removed.

Of the 115 bombycoid and two outgroup specimens, 110 were sequenced directly using

AHE target capture sequencing, of which 68 were sequenced using the BOM1 and 42 using the

LEP1 kit. Seven specimens had their AHE loci probe regions mined from previously sequenced

transcriptomes or the *B. mori* genome (Supp. Table 2). These specimens did not have flanking

data because of nature of transcriptome data. All specimens were processed using either the

'Bmori' (for LEP1) or 'BOM1' (for BOM1) reference libraries. Previous Breinholt et al., [16] a

scripts are available in Dryad [45]. Instructions on how to use the pipeline and additional scripts

that were not part of Breinholt et al. [16] are provided in Dryad (will update upon acceptance).

Phylogenetics

Lepidoptera AHE probe sets comprise highly-conserved coding probe regions (i.e.,

exons) and more variable, generally non-coding flanking regions (e.g., introns or intergenic

regions) located on either side of the probe region [16]. We evaluated both nucleotide and

amino acid datasets to examine phylogenetic signal and the role that saturation may play in the

data (see [19]). With the Breinholt et al. [16] pipeline and scripts [45], three datasets were built

for phylogeny inference: 1) AA = an amino acid supermatrix composed of translated probe

region loci; 2) Pr+Fl = a probe + flanking supermatrix; and 3) ASTRAL = the individual loci from

the Pr+Fl supermatrix, used for individual gene tree inference and then species tree estimation

to evaluate the potential effects of deep coalescence. Additional analyses (i.e., probe region-

only, as nucleotides) were investigated, but are not reported here because their outcomes did

not differ from those reported.

Concatenated supermatrices were assembled using FASconCAT-G v1.02 [57].

Phylogenetic inference was performed in a maximum likelihood (ML) framework using IQ-TREE

MPI multicore v1.5.3 [58]. For both nucleotide and amino acid datasets, the '–m TEST'

command was used in IQ-TREE to perform a search for the most appropriate model of amino

acid or nucleotide substitution. For all inferences, we performed 1000 random addition

sequence (RAS) replicates, and 1000 replicates each for both ultrafast bootstraps (UFBS) ('–bb'

command) and SH-aLRT tests ('-alrt' command). The SH-like approximate likelihood ratio test

(SH-aLRT) estimates branch support values that have been shown to be as conservative as the

commonly used non-parametric bootstrap values [59]. SH-aLRT and bootstrap values tend to

agree for data sets with strong phylogenetic signal (i.e., datasets with loci that are sufficiently

large in number of bases, and tips that share sufficient divergence between sequences).

Disagreements in branch support are thought to arise as a consequence of small sample size,

insufficient data, or saturated divergence levels (see [60]). We classified nodes as "robust" if

they were recovered with support values of UFBS ≥ 95 and SH-aLRT ≥ 80 ([59], [60]).

Because concatenation can be misleading when there are high levels of incomplete

lineage sorting or deep coalescence [61], we assessed the impact of potential gene-tree

discordance ([62], [63], [64]) by inferring a phylogeny for each individual locus, using IQ-TREE

under the same parameters as above (i.e., probe and single partition of flanking loci were

modeled by site). Species tree estimation was performed in ASTRAL-III [65]. ASTRAL is a

computationally efficient and statistically consistent (under the multi-species coalescent)

nonparametric method that takes input gene trees and estimates a highly accurate species

tree, even when there is a high level of incomplete lineage sorting (or deep coalescence) [66].

The use of ASTRAL is also an informative "data exploration" exercise with phylogenomic

datasets, providing valuable information regarding the level of general tree discordance across

your set of gene trees, and the potential presence of incomplete lineage sorting/deep

coalescence that should be investigated further. To evaluate node support on the species tree,

we used the ASTRAL support values (ASV) – local posterior probabilities that are more precise

than evaluating bootstrap values across a set of input trees [67]. ASTRAL support values were

determined to be "robust" if nodes were recovered with local posterior probabilities ≥ 0.95. All

pipeline steps and phylogenomic analyses were conducted on the University of Florida

HiPerGator HPC (http://www.hpc.ufl.edu/). All alignment FASTA files, loci information, partition

files, tree files, and other essential data files used for phylogenetic inference are available as

supplementary materials on Dryad (will update upon acceptance).

Rogue taxon & outlier locus analyses

We investigated whether our molecular data included rogue taxa or outlier loci that

were potentially influencing our phylogenetic results. A rogue taxon analysis was carried out

using the online version of RogueNaRok ([68], [69]), http://rnr.h-its.org/, on the 1000 ultrafast

bootstrap trees and the consensus tree from the Pr+Fl supermatrix. Outlier taxa and loci

analyses were carried out using Phylo-MCOA v.1.4 (PMCoA) [70] in R (on the 650 gene trees

used in the ASTRAL analysis). No rogue taxa or loci were found, and therefore we did not prune

any taxa or loci from subsequent analyses.

Diversification rate analyses

As an initial investigation into why some bombycoid lineages are more diverse than

others, we examined and quantified how diversification rates (the interplay between speciation

and extinction) have changed over time. Simply calculating species diversity per clade and

assuming extant diversity is a true indicator of increases in diversification rate could produce

significant biases in one's interpretations due to some charismatic lineages receiving more

taxonomic effort than their "boring" sister lineages (see [71], [72], [73], [74]). We therefore

applied BAMM [75] and 'BAMMtools' [76] to infer the number and location of

macroevolutionary rate shifts across our phylogeny, and visualize the 95% credible set of shift

configurations.

The fossil record of Lepidoptera, especially of the Bombycoidea, is poor [77]. Prior

lepidopteran studies that have included fossils in dating analyses have been scrutinized for

incorrect fossil identification or placement on phylogeny [78]. Because Lepidoptera fossils are

limited and characters are difficult to discern, we decided not to conduct a dating analysis for

this study. Instead, the ML best tree was converted into a relative-rate scaled ultrametric tree

using the 'chronopl' command in the R package 'ape' [51]. This approach produces a tree

whose branches are scaled to evolutionary rates, not a dated tree, and provides a way to

understand evolutionary changes over relative "time" of the group being investigated.

For the first time, quantifiable rates of diversification were calculated for the

Bombycoidea. This is important, because whether a taxonomic group possesses more described

species than another related lineage, does not mean they have diversified "more". A larger

number of described species could simply be due to the taxonomic effort, a well-known bias in

bombycoids, with the Sphingidae and Saturniidae representing the charismatic groups that

most bombycoid taxonomists have historically worked. To account for non-random missing

speciation events, we quantified the percentage of taxa sampled within each family and

incorporated these in the form of branch-specific sampling fractions. Sampling fractions were

based on the updated superfamily numbers calculated from Kitching et al. [1]. Informed priors,

based on our sampling and phylogeny, were determined using 'setBAMMpriors' in BAMMtools.

The MCMC chain was run for 100 million generations, sampling every 1000 generations.

Convergence diagnostics was assessed using the R package 'coda' [79]. The first 20% of runs

were discarded as burn-in.

Results

Our abdomen soaking approach proved relatively consistent with high enough yield of

target DNA to proceed with AHE sequencing. Datasets constructed for phylogeny inference

contained the following number of loci, sequence length, and model specifications: 1) AA = 579

loci, 48,456 amino acid residues, modeled by locus; 2) Pr+Fl = 649 probe loci + one flanking

locus (261,780 bp), each probe locus was modeled by site, flanking data was maintained as a

single partition and modeled by site; and 3) ASTRAL = 650 loci, each locus modeled by site.

Differences in loci number between the AA dataset and the Pr+Fl dataset are the result of loci

being removed from the AA phylogenetic inference due to a lack of variation at the amino acid

level for those loci (i.e., the probe regions were highly conserved with no variation across taxa).

Increasing the number of loci and taxon sampling significantly improved our

understanding of Bombycoidea relationships. The inferred relationships are generally

consistent across the three phylogenetic inferences that we performed (AA, Pr+Fl, ASTRAL),

with all major backbone relationships robustly supported, and all bombycoid families *sensu*

Zwick [20] and Zwick et al. [13] were recovered as monophyletic. Due to the methodological

approach (i.e., the treatment of different data types) and the more biologically realistic and

parsimonious explanation of the topology (see Systematics section in Supplemental Materials),

our preferred phylogeny is the tree generated from the "probe + flanking" (Pr+Fl) dataset. All

family-level placements of genera *sensu* Kitching et al. [1] were supported, with the exception

of *Arotros* Schaus, a genus long considered to be an epiine bombycid [80]. *Arotros* is clearly

nested within Apatelodidae in our tree (Fig. 1; Supp. Fig. 1) and is hereby transferred to

Apatelodidae (see supplemental text for additional details on this taxonomic change). The

family-level rearrangements of Zwick [20] and Zwick et al. [13] were also recovered in our

phylogenetic results: the placement of taxa formerly classified in the "Lemoniidae" (e.g.,

Lemaire & Minet [22]) are recovered within Brahmaeidae; the Apatelodidae are distinct from

the Bombycidae and Phiditiidae; and the broader concept of Endromidae, which includes taxa

formerly placed within the "Mirinidae" and Bombycidae (e.g., Lemaire & Minet [22]), are

recovered as monophyletic.

We find broad congruence with the major groupings designated by Zwick [20], though

internal relationships within these groups did not exactly match previously published trees.

Historically, the most problematic familial placement in the superfamily has been the

Bombycidae *sensu stricto*. Phylogenetic studies that were based on a handful of gene regions

(e.g., [13], [18], [20]), placed this family either as sister to the Saturniidae or to the Sphingidae

(reviewed in [19]), albeit without strong support for either. Our study clearly places the

Bombycidae as the sister lineage to the Saturniidae + Sphingidae (the 'SBS' group – coined by

Zwick et al. [13]), as seen in trees from the AA and Pr+Fl datasets, but not in the ASTRAL tree

(Supp. Trees 1-3) – an outcome that mirrors traditional Sanger sequencing studies based on few

loci, where individual gene trees can lack the phylogenetic signal of supermatrices. The

'CAPOPEM' group (Carthaeidae, Anthelidae, Phiditiidae, and Endromidae – coined by Regier et

al. [11]) is recovered in all three analyses, although in the ASTRAL inference this group is nested

within the clade containing the Sphingidae, Bombycidae and Saturniidae (Supp. Fig. 2).

Interfamilial relationships within the CAPOPEM clade were not robustly supported in Regier et

al. [11] or Zwick et al. [13], but our AHE-based trees confidently solidify relationships within this

group. Zwick et al. [13] recovered the Old World Endromidae as sister to the Australian

Anthelidae + (Australian Carthaeidae + Neotropical Phiditiidae), however, our Pr+Fl analysis

supports the Anthelidae as sister to Endromidae + (Carthaeidae + Phiditiidae) instead.

Effectively, Anthelidae and Endromidae swap places within CAPOPEM when comparing Zwick et

al. [13] and our Pr+Fl phylogeny. Conversely, our AA results illustrate yet a different picture, in

which Phiditiidae are sister to Anthelidae + (Carthaeidae + Endromidae), mirroring Regier et al.

[11]. This discordance is important to note considering the disjunct geographic distribution of

these families. Among these families, only Phiditiidae is found in (and endemic to) the New

World, with each of the other families being restricted to the Old World. Lastly, while

intrafamilial relationships in the SBS and CAPOPEM groups change depending on which of our

datasets are used to infer the phylogeny (Fig. 1, Supp. Figs. 1-3, Supp. Trees 1-3), the 'BALE'

group (Brahmaeidae, Apatelodidae, and Eupterotidae – coined by Regier et al. [11]) is

recovered in all three analyses.

When we investigate intrafamilial relationships, the Bombycidae (excluding *Arotros*,

mentioned above; see Supplemental text) possesses distinct and monophyletic subfamilies

Bombycinae (Old World) and Epiinae (New World; Fig. 1; Supp. Fig. 1). Within the Eupterotidae,

the Striphnopteryginae and Janinae are monophyletic, but others are not (Fig. 1; Supp. Fig. 1).

Within the Saturniidae, the Pr+Fl and AA topologies are similar to Regier et al. [31] (Fig. 1 &

Supp. Fig. 3), where the Oxyteninae are sister to the rest of the family, followed by the

Cercophaninae. Arsenurinae is the sister lineage to the Hemileucinae and the Ceratocampinae,

while Salassinae is the sister lineage to the Saturniinae. A major difference in subfamily

relationships is the placement of Agliinae, which was either sister to the Salassinae +

Saturniinae (Pr+Fl) or to the Arsenurinae (AA, also [13] and [31]). The ASTRAL inference is quite

different, placing the Ceratocampinae as the sister lineage to a clade containing the Agliinae,

Arsenurinae + Hemileucinae, and a Saturniinae + Salassinae clade (Supp. Fig. 3), albeit with low

branch support. Within the Sphingidae, our topologies (both Pr+Fl and AA) are largely

congruent, but differ slightly from Kawahara et al. [81], Zwick et al. [13], and Kawahara and

Barber [33]. Smerinthinae, which now excludes *Langia* (Kitching et al [1]) is monophyletic and

the Sphinginae are polyphyletic (Fig. 1; Supp. Fig. 1). This outcome is robustly supported in the

Pr+Fl and ASTRAL trees, although internal relationships slightly differ between the two trees.

The placement of the Macroglossinae, Langiinae, and Sphinginae are robustly supported in the

AA and Pr+FI tree, but low internal node support obfuscates relationships within Smerinthinae.

Among the other families, a number of subfamilies and genera, as defined by the most

current classification of the Bombycoidea [1], are not monophyletic (Fig. 1; Supp. Fig. 1). Within

Eupterotidae, the Eupterotinae and the "Ganisa group" are not monophyletic. The Eupterotinae

are rendered paraphyletic due to the placements of Panacelinae and Striphnopteryginae. The

Striphnopteryginae genus *Phiala* is paraphyletic with respect to *Lichenopteryx*, and the Ganisa

group is polyphyletic due to the traditional inclusion of the genus *Neopreptos* [82]. Within the

Anthelidae, *Anthela* is paraphyletic due to the placement of *Nataxa* and *Pterolocera*, a finding

congruent with Zwick [20]. Within Sphingidae, Sphinginae is polyphyletic due to the placement

of *Pentateucha* as sister to the Langiinae, and the Smerinthinae genus *Polyptychus* is

paraphyletic.

A number of factors can lead to the appearance of a taxonomic group being more

"diverse" than other sister lineages. For example, when simply looking at numbers of described

species, taxonomic bias in interest and effort could substantially affect our understanding. This

is why it is essential to formally test diversification rates. While there are more described

species of Saturniidae and Sphingidae than the rest of the Bombycoidea, that doesn't mean

those two families are necessarily more evolutionarily diverse. For the first time, actual

quantifiable rates of diversification were calculated for the Bombycoidea, by evaluating how

the interplay between speciation and extinction has changed over relative time. At the family

level, a major diversification rate shift occurs along the lineage leading to the Saturniidae and

Sphingidae (Fig. 1). The effective sample size of the log-likelihood was 8475.169 and the

effective sample size of the number of shift events present in each sample was 18141.4. The

95% credible set of rate shift configurations sampled with BAMM can be seen in Supp. Fig. 4.

Unfortunately, due to the poor Bombycoidea fossil record and the limited computational

approaches currently available to adequately estimate extinction rates, we are unable to

discern whether the differences that exist are due to increases or decreases in speciation,

although species diversity numbers, based on the classification of Kitching et al. [1] agree with

our diversification results. The number of described species for each of the ten bombycoid

families, according to Kitching et al. [1] is: Anthelidae (94 spp.); Apatelotidae (182 spp.);

Bombycidae (202 spp.); Brahmaeidae (68 spp.); Carthaeidae (1 sp.); Endromidae (70 spp.);

Eupterotidae (396 spp.); Phiditiidae (23 spp.); Saturniidae (3,454 spp.); Sphingidae (1,602 spp.).

Discussion

Within the past decade, a few studies using molecular sequence data have attempted to

resolve the phylogenetic relationships of the Bombycoidea ([11], [13], [20]). None have

corroborated the earlier morphology-based hypotheses ([15], [22]). In order to establish a

backbone for future evolutionary, comparative, and taxonomic studies, we sampled exemplars

from all major lineages in the superfamily and used Anchored Hybrid Enrichment (AHE)

phylogenomics to provide a robust phylogeny of the superfamily based on the largest

taxonomic and molecular sampling to date.

To achieve dense taxonomic coverage, we included many samples from pinned

specimens in natural history collections. We modified previous DNA extraction protocols to

increase DNA yield for high-throughput sequencing and modified the AHE probe set developed

by Breinholt et al. [16] to more efficiently recover phylogenetically-informative loci within the

Bombycoidea. To allow more flexibility in the use of the data (e.g., integration of these samples

with "legacy" datasets and CO1 for species identification), we added a selection of traditional

Sanger-sequenced loci. To assist future usage of the BOM1 and other AHE probe sets, we

expanded the bioinformatic protocol of Breinholt et al. [16] to include step-by-step instructions

for running the bioinformatics pipeline.

Our phylogeny largely reinforces the results of earlier molecular works, although

phylogenomics finally brings into focus those long problematic relationships, while also

identifying important topological where some subfamilies and genera are not recovered as

monophyletic. When comparing with trees published in previous works, topological

discordances are likely the product of increased locus sampling, which provided significantly

more phylogenetic information, as well as morphological homoplasy or convergence that likely

obscured the true placement of certain taxonomic groups. For example, our results provide a

well-supported placement of the historically troublesome family Bombycidae *sensu stricto*, as

the sister lineage to Saturniidae + Sphingidae. An earlier phylogenomic study [19] had provided

some evidence in support of this relationship, but its taxon sampling was very limited.

Our study highlights how morphological convergence in Bombycoidea has confused our

understanding of their evolution. For example, the bombycid genus *Rotunda*, endemic to the

Old World, and the apatelodid genus *Arotros*, endemic to the New World, are clearly unrelated

(Fig. 1; Supp. Fig. 1). However, they are both astonishingly similar in body size, wing shape, and

phenotypic appearance (see Supplemental Information: Systematics). Such results imply that

there may be adaptive advantages to evolving particular wing shape and size, as found in other

bombycoid lineages such as in the Saturniidae [29]. Furthermore, within the CAPOPEM clade,

we see a biogeographical distribution that, based on the AA and ASTRAL topologies, appears to

be a more biologically realistic and parsimonious explanation (than the Pr+Fl topology) of the

evolutionary history due to the placement of the Neotropical Phiditiidae as sister to the

remaining and Old World CAPOPEM families (Supp. Fig. 2). In the future, increased taxon

sampling will likely help bring better resolution to both the backbone and internal bombycoid

relationships. These types of findings, wherein different dataset types answer different

questions, highlight the importance of evaluating phylogenetic data in different ways because

phylogenetic signal could be hiding in phylogenomic datasets.

From an evolutionary viewpoint, one of the most interesting results came from the first

attempt to quantify the diversification rates across the Bombycoidea, in particular, the dramatic

shift in diversification rates leading to the Sphingidae and Saturniidae lineages. While maybe

not "surprising", because of the number of species described, this had never been quantified

before in this group. We know that simple numbers of described species does not mean

necessarily that they have diversified "more" than other closely related lineages. A number of

factors could cause one taxonomic group to appear as if there are more species than another

related lineage. Perhaps the most important of these is simply due to the taxonomic effort that

has historically been applied to a group, a bias that can certainly be found in the bombycoids,

with Sphingidae and Saturniidae representing highly charismatic groups where most bombycoid

taxonomists have worked.

These findings are fascinating because the Sphingidae and Saturniidae have contrasting

life-history strategies, such as larvae feeding on different kinds of food plants, adults having the

ability nectar source [35], and anti-bat strategies ([29], [32], [33], [34]) which are reflected in

their morphology (e.g., body sizes and wing shapes) and behavior, including flight speed and

maneuverability [35]. In an "arms race", as has been shown between moths and bats, bat

predation selectively removes unfit lineages from the environment, thus increasing the speed

of evolution of these surviving lineages. As new traits evolve that can be used to effectively

evade the predator, subsequent release from the predatory pressure provides the

opportunities to diversify ecologically and behaviorally. Although we did not conduct a

divergence time estimation analysis in the present study, the origin of the SBS group has been

postulated to be approximately 50 mya [83], with Sphingidae originating soon thereafter [33].

Insectivorous bats are thought to have originated roughly 60 mya, and the diversification of

Sphingidae and Saturniidae around that time suggests that the incredible taxonomic diversity

within these two families could be in part due to bat-related selection pressures resulting in

diverse anti-bat traits.

The hawkmoth and silk moth evolutionary story is certainly more complex than simply

reflecting their interactions with bats. Although we postulate that the differences in

diversification rates are correlated with bat predation, it is possible that these rate shifts are

due to other factors, such as ecological specialization or shifts in host plant usage, both as

larvae and adults. Amassing and collating behavioral and ecological datasets for the tips of the

Bombycoidea Tree of Life for macroevolutionary comparative investigations is essential to

furthering our understanding of this diverse, global superfamily, as well as understanding how

bats, ecological traits, and/or biogeographical history may or may not have shaped their

diversity. At this time, large trait datasets do not exist for these groups, but are currently being

worked on to be included in future trait-dependent diversification analyses with much more

complete sampling, at the genus and species level across these families, to truly explore the

drivers of Bombycoidea diversity. This research establishes some hypotheses to be further

tested when more complete sampling of the Bombycoidea has been completed, and robust

trait datasets have been collected.

Conclusions

Our study finally brings into focus long problematic bombycoid relationships and

establishes a backbone for future evolutionary, comparative, and taxonomic studies. Our

modified DNA extraction protocol allows Lepidoptera specimens to be readily sequenced from

pinned natural history collections, and highlights the flexibility of AHE to generate genomic data

from a wide range of museum specimens, both age and preservation method. By allowing

researchers to tap into the wealth of biological data residing in natural history collections

around the globe, these types of methodologies (e.g., DNA from museum specimens and

targeted sequencing capture) will provide the opportunities for us to continually add to our

understanding of Lepidoptera and Bombycoidea evolution, as well as refinine our

understanding of relationships across the Tree of Life.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets supporting the results of this article are available in Dryad Data Repository

(will update upon acceptance with the unique persistent identifier and hyperlink to datasets in

http:// format). Supplemental Information includes four figures, one table, plus the full

museum specimen DNA extraction protocol, systematic information on Bombycoidea families,

bioinformatics pipeline scripts and instructions, alignment FASTA files (nucleotides, amino

acids, supermatrices, and individual loci), partition files, tree files, and other essential data files

that were inputted or outputted from the phylogenetic inference and analysis. Previous

Breinholt et al., [16] bioinformatics scripts are available in Dryad [45].

Competing interests

The authors declare that they have no competing interests.

Authors' Information

ORCID ID - Hamilton: 0000-0001-7263-0755; St Laurent: 0000-0001-6439-5249; Dexter: 0000-

0002-5136-6083; Kitching: 0000-0003-4738-5967; Breinholt: 0000-0002-3867-2430; Zwick:

0000-0002-7532-1752; Timmermans: 0000-0002-5024-9053; Barber: 0000-0003-3084-2973;

Kawahara: 0000-0002-3724-4610

Funding

This work was partly supported by NSF grants [IOS-1121807, JRB], [IOS-1121739, AYK], [DBI-

1349345, AYK], [DBI-1601369, AYK], [DEB-1557007, AYK], and [PRFB-1612862, CAH]; a NERC

grant [NE/P003915/1, IJK]; National Geographic Society grants [CRE 9944-16, JRB]; and

postdoctoral support from the Florida Museum of Natural History (to CAH).

Authors' Contributions

CAH, AYK, JWB, IJK, MT, AZ designed the study. CAH and JWB analyzed data. CAH, AYK, RAS

wrote the paper. CAH, AYK, JRB, KD, IJK, RAS, AZ collected, identified, and processed specimens.

All authors discussed results and contributed to the writing of the manuscript.

Acknowledgements

We thank Seth Bybee, Deborah Glass, Yash Sondhi, Jamie Theobald, and Emmanuel Toussaint.

We are indebted to Samantha Epstein, John Heppner, Geena Hill, Nicholas Homziak, Jacqueline

Miller, Charles Mitter, Kim Mitter, Lary Reeves, Andrei Sourakov, and Andrew Warren, who

provided specimens for this project, contributed help in identifying or organizing specimens, or

assisted the project in other ways (e.g., lab work or intellectually). The authors would also like

to thank the peer-review process for improving the manuscript. The authors acknowledge

University of Florida Research Computing for providing computational resources and support

that contributed to the research results reported in this publication. URL:

http://researchcomputing.ufl.edu

References

1. Kitching I, Rougerie R, Zwick A, St Laurent R, Ballesteros Mejia L, Kawahara A, et al. A global

checklist of the Bombycoidea (Insecta: Lepidoptera). BDJ. 2018;6:e22236–13.

2. Roe AD, Weller SJ, Baixeras J, Brown J, Cummings MP, Davis DR, et al. Evolutionary

framework for Lepidoptera model systems. In: Goldsmith M, Marec F, editors. Molecular

Biology and Genetics of the Lepidoptera. 2010. pp. 1–24.

3. Gregory D. Hawkmoth pollination in the genus *Oenothera*. Aliso A Journal of Systematic and

Evolutionary Botany. 1963;5:357–84.

4. Haber WA, Frankie GW. A Tropical Hawkmoth Community- Costa Rican Dry Forest

Sphingidae. Biotropica. 1989;21:155–72.

5. Nilsson LA, Jonsson L, Rason L, Randrianjohany E. Monophily and pollination mechanisms in

Angraecum arachnites Schltr. (Orchidaceae) in a guild of long-tongued hawk-moths

(Sphingidae) in Madagascar. Biological Journal of the Linnean Society. 1985;26:1–19.

6. Nilsson LA, Rabakonandrianina E, Pettersson B. Exact tracking of pollen transfer and mating

in plants. Nature. 1992;360:666–8.

7. Wasserthal LT. The pollinators of the Malagasy star orchids *Angraecum sesquipedale*, *A.*

sororium and *A. compactum* and the evolution of extremely long spurs by pollinator shift.

Botanica Acta. 1997;110:343–59.

8. Nilsson LA. Deep flowers for long tongues. Trends In Ecology & Evolution. 1998;13:259–60.

9. Raguso RA, Willis MA. Synergy between visual and olfactory cues in nectar feeding by naïve

hawkmoths, *Manduca sexta*. Animal Behaviour. 2002;64:685–95.

10. Beck J, Kitching IJ, Linsenmair KE. Effects of habitat disturbance can be subtle yet significant:

biodiversity of hawkmoth-assemblages (Lepidoptera: Sphingidae) in Southeast-Asia.

Biodiversity and Conservation. 2006;15:465–86.

11. Regier JC, Cook CP, Mitter C, Hussey A. A phylogenetic study of the "bombycoid complex"

(Lepidoptera) using five protein-coding nuclear genes, with comments on the problem of

macrolepidopteran phylogeny. Systematic Entomology. 2008;33:175–89.

12. Meusemann K, Reumont von BM, Simon S, Roeding F, Strauss S, Kück P, et al. A

phylogenomic approach to resolve the Arthropod Tree of Life. Mol Biol Evol. 2010;27:2451–64.

13. Zwick A, Regier JC, Mitter C, Cummings MP. Increased gene sampling yields robust support

for higher-level clades within Bombycoidea (Lepidoptera). Systematic Entomology. 2011;36:31–

43.

14. Kim MJ, Kang AR, Jeong HC, Kim K-G, Kim I. Reconstructing intraordinal relationships in

Lepidoptera using mitochondrial genome data with the description of two newly sequenced

lycaenids, *Spindasis takanonis* and *Protantigius superans* (Lepidoptera: Lycaenidae). Mol

Phylogenet Evol. 2011;61:436–45.

15. Minet J. The Bombycoidea: Phylogeny and higher classification (Lepidoptera, Glossata).

Entomologica Scandinavica. 1994;25:63–88.

16. Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. Resolving relationships

among the megadiverse butterflies and moths with a novel pipeline for anchored

phylogenomics. Syst Biol. 2018;67:78–93.

17. Timmermans MJTN, Lees DC, Simonsen TJ. Towards a mitogenomic phylogeny of

Lepidoptera. Mol Phylogenet Evol. 2014;79:169–78.

18. Wang X, Chen Z-M, Gu X-S, Wang M, Huang G-H, Zwick A. Phylogenetic relationships among

Bombycidae s.l. (Lepidoptera) based on analyses of complete mitochondrial genomes.

Systematic Entomology. 2018;20:537–9.

19. Breinholt JW, Kawahara AY. Phylotranscriptomics: saturated third codon positions radically

influence the estimation of trees based on next-gen data. Genome Biol Evol. 2013;5:2082–92.

20. Zwick A. Molecular phylogeny of Anthelidae and other bombycoid taxa (Lepidoptera:

Bombycoidea). Systematic Entomology. 2008;33:190–209.

21. Minet J. Tentative reconstruction of the ditrysian phylogeny (Lepidoptera, Glossata).

Entomologica Scandinavica. 1991;22:69–95.

22. Lemaire C, Minet J. 18. The Bombycoidea and their relatives. In: Kristensen NP, editor.

Lepidoptera, Moths and Butterflies. vol. 1. Evolution, Systematics, and Biogeography. Handbook

of Zoology. Walter de Gruyter, Berlin; 1998. pp. 322–53.

23. Regier JC, Mitter C, Zwick A, Bazinet AL, Cummings MP, Kawahara AY, et al. A large-scale,

higher-level, molecular phylogenetic study of the insect order Lepidoptera (moths and

butterflies). PLoS ONE. 2013;8:e58568–23.

24. Regier JC, Zwick A, Cummings MP, Kawahara AY, Cho S, Weller S, et al. Toward

reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia): an

initial molecular study. BMC Evolutionary Biology. 2009;9:280–21.

25. Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A

comprehensive and dated phylogenomic analysis of butterflies. CURBIO. 2018;28:770–5.

26. Johns CA, Toussaint EFA, Breinholt JW, Kawahara AY. Origin and macroevolution of micro-

moths on sunken Hawaiian Islands. Proc. Biol. Sci. 2018;285:20181047–10.

27. St Laurent RA, Hamilton CA, Kawahara AY. Museum specimens provide phylogenomic data

to resolve relationships of sack-bearer moths (Lepidoptera, Mimallonoidea, Mimallonidae).

Systematic Entomology. 2018;43:729–61.

28. Toussaint EFA, Breinholt JW, Earl C, Warren AD, Brower AVZ, Yago M, et al. Anchored

phylogenomics illuminates the skipper butterfly tree of life. BMC Evolutionary Biology.

2018;18:1–11.

29. Rubin JJ, Hamilton CA, McClure CJW, Chadwell BA, Kawahara AY, Barber JR. The evolution of

anti-bat sensory illusions in moths. Science Advances. 2018;4.

30. Lemmon AR, Emme SA, Lemmon EM. Anchored Hybrid Enrichment for massively high-

throughput phylogenomics. Syst Biol. 2012;61:727–44.

31. Regier JC, Grant MC, Mitter C, Cook CP, Peigler RS, Rougerie R. Phylogenetic relationships of

wild silkmoths (Lepidoptera: Saturniidae) inferred from four protein-coding nuclear genes.

Systematic Entomology. 2008;33:219–28.

32. Barber JR, Kawahara AY. Hawkmoths produce anti-bat ultrasound. Biology Lett.

2013;9:20130161–1.

33. Kawahara AY, Barber JR. Tempo and mode of antibat ultrasound production and sonar

jamming in the diverse hawkmoth radiation. Proceedings of the National Academy of Sciences.

2015;112:6407–12.

34. Barber JR, Leavell BC, Keener AL, Breinholt JW, Chadwell BA, McClure CJW, et al. Moth tails

divert bat attack: evolution of acoustic deflection. Proceedings of the National Academy of

Sciences. 2015;112:2812–6.

35. Janzen D. Two ways to be a tropical big moth: Santa Rosa saturniids and sphingids. Oxford

surveys in evolutionary biology. 1984;1:140.

36. Tammaru T, Haukioja E. Capital breeders and income breeders among Lepidoptera:

consequences to population dynamics. Oikos. 1996;77:561–4.

37. Kawahara AY, Breinholt JW. Phylogenomics provides strong evidence for relationships of

butterflies and moths. P Roy Soc B-Biol Sci. 2014;281:20140970–0.

38. Jones KE, Bininda-Emonds ORP, Gittleman JL. Bats, clocks, and rocks: diversification patterns

in Chiroptera. Evolution. 2005;59:2243–55.

39. Teeling EC, Springer MS, Madsen O, Bates P, O'Brien SJ, Murphy WJ. A molecular phylogeny

for bats illuminates biogeography and the fossil record. Science. 2005;307:580–4.

40. Miller-Butterworth CM, Murphy WJ, O'Brien SJ, Jacobs DS, Springer MS, Teeling EC. A family

matter: conclusive resolution of the taxonomic position of the long-fingered bats, *Miniopterus*.

Mol Biol Evol. 2007;24:1553–61.

41. Shi JJ, Rabosky DL. Speciation dynamics during the global radiation of extant bats. Evolution.

2015;69:1528–45.

42. Lei M, Dong D. Phylogenomic analyses of bat subordinal relationships based on

transcriptome data. Scientific Reports. 2016;6:27726.

43. Barber JR, Conner WE. Acoustic mimicry in a predator-prey interaction. Proceedings of the

National Academy of Sciences. 2007;104:9331–4.

44. Xia Q, Zhou Z, Lu C, Cheng D, Dai F, Bin Li, et al. A draft sequence for the genome of the

domesticated silkworm (*Bombyx mori*). Science. 2004;306:1937–40.

45. Breinholt JW, Earl C, Lemmon AR, Lemmon EM, Xiao L, Kawahara AY. Data from: Resolving

relationships among the megadiverse butterflies and moths with a novel pipeline for Anchored

Phylogenomics. Dryad Digital Repository; 2017.

46. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014.

47. Mayrose I, Graur D, Ben-Tal N, Pupko T. Comparison of site-specific rate-inference methods

for protein sequences: empirical Bayesian methods are superior. Mol Biol Evol. 2004;21:1781–

91.

48. Pond SLK, Frost SDW, Muse SV. HyPhy: hypothesis testing using phylogenies. Springer, New

York, NY; 2005. pp. 125–81.

49. López-Giráldez F, Townsend JP. PhyDesign: an online application for profiling phylogenetic

informativeness. BMC Evolutionary Biology. 2011;11:152.

50. Heibl C. PHYLOCH: R language tree plotting tools and interfaces to diverse phylogenetic

software packages. 2013.

51. Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R language.

Bioinformatics. 2004;20:289–90.

52. Cho S, Epstein SW, Mitter K, Hamilton CA, Plotkin D, Mitter C, et al. Preserving and

vouchering butterflies and moths for large-scale museum-based molecular research. PeerJ.

2016;4:e2160.

53. Blaimer BB, Lloyd MW, Guillory WX, Brady SG. Sequence capture and phylogenetic utility of

genomic Ultraconserved Elements obtained from pinned insect specimens. PLoS ONE.

2016;11:e0161531–20.

54. Mikheyev AS, Zwick A, Magrath MJL, Grau ML, Qiu L, Su YN, et al. Museum genomics

confirms that the Lord Howe Island stick insect survived extinction. CURBIO. 2017;27:3157–

3161.e4.

55. Ruane S, Austin CC. Phylogenomics using formalin-fixed and 100+ year-old intractable

natural history specimens. Molecular Ecology Resources. 2017;17:1003–8.

56. Larsson A. AliView: a fast and lightweight alignment viewer and editor for large datasets.

Bioinformatics. 2014.

57. Kück P, Longo GC. FASconCAT-G: extensive functions for multiple sequence alignment

preparations concerning phylogenetic studies. Frontiers in Zoology. 2014;11.

58. Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ. IQ-TREE: a fast and effective stochastic

algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32:268–74.

59. Minh BQ, Nguyen MAT, Haeseler von A. Ultrafast approximation for phylogenetic bootstrap.

Mol Biol Evol. 2013;30:1188–95.

60. Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and

methods to estimate Maximum-Likelihood phylogenies: assessing the performance of PhyML

3.0. Syst Biol. 2010;59:307–21.

61. Mendes FK, Hahn MW. Why concatenation fails near the anomaly zone. Syst Biol.

2017;67:158–69.

62. Maddison W. Gene trees in species trees. Syst Biol. 1997;46:523–36.

63. Slowinski J, Page R. How should species phylogenies be inferred from sequence data? Syst

Biol. 1999;48:814–25.

64. Edwards SV. Is a new and general theory of molecular systematics emerging? Evolution.

2009;63:1–19.

65. Zhang C, Sayyari E, Mirarab S. ASTRAL-III: increased scalability and impacts of contracting

low support branches. RECOMB International Workshop on Comparative Genomics. 2017. pp.

53–75.

66. Mirarab S, Warnow T. ASTRAL-II: coalescent-based species tree estimation with many

hundreds of taxa and thousands of genes. Bioinformatics. 2015;31:i44–i52.

67. Sayyari E, Mirarab S. Anchoring quartet-based phylogenetic distances and applications to

species tree reconstruction. BMC Genomics. BioMed Central; 2016;17:101–13.

68. Aberer AJ, Stamatakis A. A simple and accurate method for rogue taxon identification. IEEE

International Conference on Bioinformatics and Biomedicine BIBM. 2011. pp. 118–22.

69. Aberer AJ, Krompass D, Stamatakis A. Pruning rogue taxa improves phylogenetic accuracy:

an efficient algorithm and webservice. Syst Biol. 2012;62:162–6.

70. de Vienne DM, Ollier S, Aguileta G. Phylo-MCOA: a fast and efficient method to detect

outlier genes and species in phylogenomics using multiple co-inertia analysis. Mol Biol Evol.

2012;29:1587–98.

71. Dial KP, Marzluff JM. Nonrandom Diversification Within Taxonomic Assemblages.

Systematic Zoology. 1989;38:26–37.

72. Doyle JA, Donoghue MJ. Phylogenies and Angiosperm Diversification. Paleobiology.

1993;19:141–67.

73. Sanderson MJ, Donoghue MJ. Shifts in Diversification Rate with the Origin of Angiosperms.

Science. 1994;264:1590–3.

74. Sanderson MJ, Donoghue MJ. Reconstructing shifts in diversification rates on phylogenetic

trees. Trends In Ecology & Evolution. 1996;11:15–20.

75. Rabosky DL. Automatic detection of key innovations, rate shifts, and diversity-dependence

on phylogenetic trees. PLoS ONE. 2014;9:e89543.

76. Rabosky DL, Grundler M, Anderson C, Title P, Shi JJ, Brown JW, et al. BAMMtools: an R

package for the analysis of evolutionary dynamics on phylogenetic trees. Methods in Ecology

and Evolution. 2014;5:701–7.

77. Sohn JC, Labandeira CC, Davis DR, Mitter C. An annotated catalog of fossil and subfossil

Lepidoptera (Insecta: Holometabola) of the world. Zootaxa. 2012,1-132.

78. de Jong R. Fossil butterflies, calibration points and the molecular clock (Lepidoptera:

Papilionoidea). Zootaxa. 2017;4270:1–63.

79. Plummer M, Best N, Cowles K, Vines K. CODA: convergence diagnosis and output analysis

for MCMC. R News. 2006;6:7–11.

80. Schaus W. Familie Bombycidae (Epiinae). In: Seitz A, editor. Die Gross-Schmetterlinge der

Erde. Kernen, Stuttgart; 1929. pp. 675–711.

81. Kawahara AY, Mignault AA, Regier JC, Kitching IJ, Mitter C. Phylogeny and Biogeography of

Hawkmoths (Lepidoptera: Sphingidae): Evidence from Five Nuclear Genes. Knapp M, editor.

PLoS ONE. 2009;4:e5719.

82. Nässig WA, Oberprieler RG. An annotated catalogue of the genera of Eupterotidae (Insecta,

Lepidoptera, Bombycoidea). Senckenbergiana biologica. 2008;88:53–80.

83. Misof B, Liu S, Meusemann K, Peters RS, Donath A. Phylogenomics resolves the timing and

pattern of insect evolution. Science. 2014;346:763–7.

Figures

Figure 1. Maximum likelihood tree of Bombycoidea, based on 650 AHE loci. All nodes are

supported by ≥95% UFBoot and ≥80% SH-aLRT values unless otherwise noted. Branch color

indicates the estimated diversification rate, with warmer colors representing lineages with

higher rates. Major taxonomic groups such as families and subfamilies are labeled. Photographs

represent species in lineages sampled in the phylogeny. Species diversity, based on Kitching et

al. [1], are noted next to families.