**Siliplant1 (Slp1) protein precipitates silica in sorghum silica cells**

**Santosh Kumar[1,a], Nurit Adiram-Filiba[2], Shula Blum[1], Javier Arturo Sanchez-Lopez[3],**

**Oren Tzfadia[4], Ayelet Omid[5], Hanne Volpin[5], Yael Heifetz[3], Gil Goobes[2] and Rivka**

**Elbaum[1]**

(S.K. ORCID ID: 0000-0001-8261-5620)

[1]Robert H Smith Faculty of Agriculture, Food and Environment, Robert H Smith Institute of

Plant Sciences and Genetics in Agriculture, Hebrew University of Jerusalem, Rehovot

7610001, Israel; [2]Department of Chemistry, Bar-Ilan University, Ramat Gan 5290002, Israel;

[3]Department of Entomology, Robert H Smith Faculty of Agriculture, Food and Environment,

Hebrew University of Jerusalem, Rehovot 7610001, Israel; [4]Bioinformatics and Systems

Biology, VIB/Ghent University, Gent B-9052, Belgium; [5]Danziger Innovations Limited,

Mishmar Hashiva 5029700, Israel

[a]Present address: Department of Plant and Environmental Sciences, Weizmann Institute of

Science, Rehovot 7610001, Israel

Address for correspondence:

*Santosh Kumar*

*Tel: +972-544378508*

*Email: skg.research@gmail.com*

*Rivka Elbaum*

*Tel: +972-8-9489335*

*Email: rivka.elbaum@mail.huji.ac.il*

| Total word count (excluding summary, references and legends): | **6230** | No. of figures: | 7 (all in colour) |
|---|---|---|---|
| Summary: | **200** | No. of Tables: | nil |
| Introduction: | **926** | No of Supporting Information files: | 3 (1) Supplementary Text and figures |

| | | | (2) Video S1 |
|---|---|---|---|
| | | | (3) Video S2 |
| Materials and Methods: | **2103** | | |
| Results: | **2072** | | |
| Discussion: | **986** | | |
| Accession number | **48** | | |
| Acknowledgements: | **95** | | |

**Summary**

- Silicon is absorbed by plant roots as silicic acid. The acid moves with the transpiration stream to the shoot, and mineralizes as silica. In grasses, leaf epidermal cells called silica cells deposit silica in most of their volume by unknown mechanism.

- Using bioinformatics tools, we identified a previously uncharacterized protein in sorghum (*Sorghum bicolor*), which we named Siliplant1 (Slp1). Silica precipitation activity *in vitro*, expression profile, and activity in precipitating biosilica *in vivo* were characterized.

- Slp1 is a basic protein with seven repeat units rich in proline, lysine, and glutamic acid. A short peptide, repeating five times in the protein precipitated silica *in vitro* at a biologically relevant silicic acid concentration. Raman and NMR spectroscopies showed that the peptide attached the silica through lysine amine groups, forming a mineral-peptide open structure. We found Slp1 expression in immature leaf and inflorescence tissues. In the immature leaf active silicification zone, Slp1 was localized to the cytoplasm or near cell boundaries of silica cells. It was packed in vesicles and secreted to the paramural space. Transient overexpression of Slp1 in sorghum resulted in ectopic silica deposition in all leaf epidermal cell types.

- Our results show that Slp1 precipitates silica in sorghum silica cells.

**Keywords:** active silicification zone (ASZ), biomineralization, phytolith, silica cell, silicification, silicon, Siliplant1 (Slp1), *Sorghum bicolor*

## Introduction

Grasses are well known for their high silica ($SiO_2 \cdot nH_2O$) content, reaching up to 10% of their dry weight (Hodson *et al.*, 2005). Silicon is available to plants as mono-silicic acid [$Si(OH)_4$] whose concentration in soil solution usually varies between 0.1 to 0.6 mM (Epstein, 1994). Grass roots actively uptakes silicic acid from soil (Ma *et al.*, 2006, 2007) of which more than 90% is loaded into the xylem. A cooperative uptake of silicic acid by two root transporters leads to Si supersaturation in the grass xylem sap (Sakurai et al., 2015). Si concentration may reach 5-7.5 mM, which is 3 to 4 times higher than its saturation concentration (Ma *et al.*, 2002; Casey *et al.*, 2003; Mitani & Ma, 2005). Silicic acid molecules then move with water inside the plant body to reach the shoots where they are unloaded from the xylem (Yamaji *et al.*, 2008). Finally, silicic acid polymerises as solid biogenic silica at several locations inside the plant body. The most prominent sites of silica deposition in grasses are the walls of leaf epidermal cells, abaxial epidermal cells of the inflorescence bracts (glumes and lemma) and root endodermal cells. The silicification mechanism across the cell types is not uniform. Models of deposition suggest either spontaneous formation as a result of evapo-transpirational loss of water, or a tightly controlled process (Kumar *et al.*, 2017b). Uptake of silicic acid and its deposition are also affected by plant mechanical damage (McLarnon *et al.*, 2017), and possibly other unknown physiological factors affecting root silicon transporters (Talukdar *et al.*, 2019).

One of the cell types most frequently silicified in grass leaves is silica cells (Motomura *et al.*, 2000; Kumar & Elbaum, 2018). Silica cells are specialized epidermal cells occurring mostly as silica-cork cell pairs, both above and below the leaf longitudinal veins (Kaufman *et al.*, 1985), on internode epidermis (Kaufman *et al.*, 1969) and on the abaxial epidermis of glumes (Hodson *et al.*, 1985). We earlier estimated that the whole process of leaf silica cell silicification is completed within 10 hours of cell division (Kumar & Elbaum, 2018). Within this time period, almost the entire cell lumen is filled with solid silica (Hodson *et al.*, 1985). Silica cell silicification is immediately followed by cell death (Kumar & Elbaum, 2018) which ceases the silicification process (Markovich *et al.*, 2015; Kumar *et al.*, 2017a; Kumar & Elbaum, 2018). Silicification in silica cells is thus different than silicification in macro-hairs and abaxial epidermal cells in lemma where silica deposition takes place in a course of weeks on the thickened cell wall material (Hodson *et al.*, 1984; Perry *et al.*, 1984), or in glume prickle hairs and papillae where silica is deposited in the empty lumen of these cells even after cell death (Hodson *et al.*, 1985). The template on which silica cells deposit silica is unknown. Cell

wall polysaccharides such as mixed-linkage glucan (Fry *et al.*, 2008; Kido *et al.*, 2015) and

callose (Law & Exley, 2011; Brugiére & Exley, 2017; Kulich *et al.*, 2018) have been suggested

as a template for silicification in plants. However, lumen silica structure of *Triticum durum*

silica cells has continuously distributed organic matter with the N/C ratio indicative of amino

acids (Alexandre *et al.*, 2015). This suggests protein(s) as the templating organic matrix for the

silicification process in silica cells.

Silicification is widespread among living beings from unicellular microbes to highly evolved

multicellular organisms (Perry, 2003). Several bio-silica associated proteins have been

reported, for example, silaffins (Kröger *et al.*, 1999; Poulsen & Kröger, 2004), silacidins

(Wenzl *et al.*, 2008) and silicanin-1 (Kotzsch *et al.*, 2017) from diatoms; and silicateins

(Shimizu *et al.*, 1998) and glassin (Shimizu *et al.*, 2015) from sponges. Among plants, a short

peptide derived from an inducible proline-rich protein precipitates silica *in vitro*. The protein,

involved in systemic acquired resistance in cucumber, may precipitate silica locally at

attempted fungal penetration sites (Kauss *et al.*, 2003). All the above-mentioned protein groups

do not share sequence homology.

To study silica deposition in silica cells, we used sorghum (*Sorghum bicolor*), a member of the

grass (Poaceae) family. Sorghum is categorized as an active silica accumulator (Hodson *et al.*,

2005; Coskun *et al.*, 2018). In grasses, leaves appear successively (Skinner & Nelson, 1995).

Even within an individual leaf, there is a gradient of cell maturation. The leaf epidermal cell

division zone is confined to the base of the leaf and the newly divided cells displace the

maturing cells away from the leaf base. Hence, leaf epidermal cells that are close to the leaf

apex are older than the cells close to the base (Skinner & Nelson, 1995).

We earlier found that silica cell silicification is confined to elongating leaves, in a well-defined

active silicification zone (ASZ) (Kumar *et al.*, 2017a; Kumar & Elbaum, 2018). The

mineralization initiates in the paramural space of viable silica cells, producing a thick silica

cell wall, and restricting the cytoplasmic space to smaller and smaller volumes (Kumar *et al.*,

2017a; Kumar & Elbaum, 2018). During this fast process, the cell maintains cell-to-cell

connectivity to the neighbouring cells through plasmodesmata (Kumar *et al.*, 2017a). As the

silicic acid resides in the apoplast in supersaturation, a crucial stage would be the initiation of

controlled silica deposition. A possible way would be by adding to the cell wall a

biomineralizing protein in appropriate time and place. Here, we report on a previously

uncharacterized protein that is expressed in silica cells, exported to the cell wall during their

silicification, and induces precipitation of silica *in planta*. Hence, we named this protein as

Siliplant1 (Slp1).

## Materials and Methods

### Plant material, growth conditions and tissue nomenclature

Seeds of *Sorghum bicolor* (L.) Moench (line BTx623) were surface sterilized and grown in soil

as reported previously (Kumar & Elbaum, 2018). Unless indicated otherwise, we used sorghum

seedlings of about 2 weeks of age for our studies. Immature, silicifying leaves ($L_{IS}$) in our

present study is analogous to leaf-2, as reported in our earlier studies (Kumar & Elbaum, 2017,

2018; Kumar *et al.*, 2017a). The immature leaf was cut into five equal segments. The middle

segment is most active in terms of silica cell silicification (Kumar *et al.*, 2017a; Kumar &

Elbaum, 2018) and was named active silicification zone (ASZ). The segment just older than

ASZ (towards leaf tip) was named ASZ+1, while the segment just younger than ASZ (towards

leaf base) was ASZ-1. The youngest segment (at leaf base) was named ASZ-2. Mature leaves

were cut into 10 equal segments and only the eighth segment from the leaf-base was used for

all the experiments.

### Sequence-based analyses of Slp1

Secondary structure prediction was done using the program GOR4 (Combet *et al*., 2000).

Intrinsically disordered tertiary structure was predicted using IUPred (Dosztányi *et al*.,

2005a,b). The prediction type we chose was 'long disorder'. Residues with score greater than

0.5 were regarded disordered. SignalP was used to predict signal peptide in Slp1 (Petersen *et*

*al.*, 2011). TargetP was used to predict whether the protein is secretory in nature (Emanuelsson

*et al.*, 2000). The organism group selected was 'plant' and the program was run with default

cutoff selection.

### Silica precipitation using Peptide-1 and Peptide-3

We tested the peptide sequence HKKPVPPKPKPEPK (Peptide-1) which appears 5 times in

Slp1 primary sequence for its silica precipitating activity *in vitro*, and compared it to a mutated

peptide, where all lysine groups were replaced by alanine (HAAPVPPAPAPEPA, Peptide-3).

We freshly prepared 1 M silicic acid solution by mixing 150 µl of tetramethyl orthosilicate to

850 µl of 1 mM HCl for four minutes under gentle stirring.

For small-scale silica precipitation at 90.9 mM silicic acid, 5 µl of 1 M silicic acid solution was

mixed with 50 µl of peptide solution (1.5 or 2.0 mg ml$^{-1}$ in 0.1 M potassium phosphate buffer,

pH 7.0) and incubated for 5 minutes under gentle shaking at room temperature. For silica

precipitation at 5 mM silicic acid, 55 µl of freshly prepared 1 M silicic acid solution was diluted

in 945 µl of 0.1 M potassium phosphate buffer (pH 7.0), of which 10 µl was mixed with 100

µl of Peptide-1 solution (1.5 mg ml$^{-1}$ in 0.1 M potassium phosphate buffer, pH 7.0) and

incubated for 30 min under gentle shaking condition. The same reactions lacking any peptide

served as control. Sediment was collected by centrifugation at 14000 g for 5 minutes. The

supernatant was thrown, and the pellet was washed thrice with 1 ml $H_2O$.

For large-scale precipitating of circa 100 mg of silica we incubated 30 mg of peptide (1.5 mg

ml$^{-1}$ in 0.1 M potassium phosphate buffer, pH 7.0) with 2 ml of freshly prepared 1 M silicic

acid solution for 30 min at room temperature under gentle shaking condition. The precipitate

with Peptide-1 was collected by centrifugation at 5000 X g for 5 minutes and washed thrice by

$H_2O$. The precipitate with Peptide-3 was collected by centrifugation at 5000 X g for 20 min

and washed thrice with $H_2O$. Both the precipitates were dried at 60 ℃.

**Scanning electron microscopy (SEM)**

The pellet of the small-scale reaction was suspended in $H_2O$ and 1 µl of the suspension was

smeared on carbon tape, air dried and coated with iridium. The coated sample was observed in

Jeol JSM IT100 (Peabody, MA, USA) scanning electron microscope (SEM).

**High resolution transmission electron microscopy (HRTEM)**

Powder from peptide-1 large-scale precipitation was resuspended in double distilled water and

spread on a TEM grid. HRTEM measurements were carried out on a JEM 2100 JEOL

microscope. The samples were imaged at an electron-beam accelerating voltage of 200 kV.

The images were collected to characterize the silica internal structure.

**Raman spectroscopic observation of Peptide-1 and silica precipitated with Peptide-1**

Lyophilized peptide solution was reconstituted in double distilled water (50 mg ml$^{-1}$) and 1 µl

of this solution was put on a steel slide and dried at 37 ℃ for 30 minutes before measuring the

Raman spectra.

The small-scale precipitation of 90.9 mM silicic acid with peptide-1 was dried at 60 ℃ until

attaining a constant dried weight. The dried powder was mounted on a steel slide. Raman

spectra of the samples were collected using InVia microspectrometer (Renishaw, New Mills,

UK) equipped with polarized 532 nm laser (45 mW max. intensity, 4 µm$^2$ beam) excitation

under a 50X air objective lens (NA=0.75). Spectra analysis was done in WiRE3 (Renishaw),
including smoothing, background subtraction, and peak picking.

**Solid state NMR spectroscopy**

Solid state NMR spectra of the peptide-1 large-scale precipitation were recorded on a Bruker
11.7 T Avance III spectrometer (Billerica, MS, USA) equipped with a 4 mm VTN CPMAS
probe at a spinning rate of 10 kHz. The $^{29}$Si cross polarization experiment was done with 8192
scans and recycle delay of 6 sec and the $^{29}$Si direct excitation experiment was recorded with
2560 scans and recycle delay of 60 sec. Analysis of the $^{29}$Si spectrum was done using the
DMFIT program (Massiot *et al.*, 2002). $^{13}$C cross polarization was done with 2048 scans using
recycle delay of 5 sec.

**Tissue-specific expression of Slp1 in sorghum**

Polyclonal antibodies against Peptide-1 was raised in rabbit (GenScript, NJ, USA). Crude
proteins from root tissues, whole of immature silicifying leaves, mature leaves and immature
inflorescence (expected to emerge from the flag leaf of mature plants within about a week)
were extracted in the extraction buffer (0.1 M potassium phosphate buffer, pH 7.0; 1% protease
inhibitor cocktail, Sigma and 0.1% 2-Mercaptoethanol). Forty-eight µg of the extracted protein
from each sample was loaded in separate wells and run on 15% polyacrylamide separating gel
under denaturing condition. The gel was blotted onto nitrocellulose membrane and the
membrane was put in blocking buffer (5% milk powder in Tris buffer saline, 0.1% Tween 20)
for one hour. We incubated the membrane with purified polyclonal antibody against Peptide-1
(1 µg ml$^{-1}$) at 4 °C for overnight. The membrane was washed and incubated with secondary
antibody (anti-rabbit IgG mouse monoclonal antibody, Genscirpt, NJ, USA) conjugated with
Horseradish peroxidase. Chemiluminescence was developed (PerkinElmer, Akron, US) and
the membrane was scanned for one second exposure time.

**Transcript abundance of *Slp1***

To check transcription of *Slp1* in immature leaf, mature leaf and immature inflorescence, RNA
was isolated from these tissues and cDNA was synthesized using 1 µg of total RNA after on
column-DNase treatment (Zymoresearch, CA, USA). Equal volume of cDNA was PCR
amplified using the primers specific for *Slp1* (Slp1-RT-F and Slp1-RT-R) and ubiquitin-
conjugating enzyme (UbCE; Sb09g023560) as internal control (Shakoor *et al.*, 2014;
Markovich *et al.*, 2015). The primer pair used to amplify *UbCE* was UbCE-RT-F and UbCE-
RT-R. Nucleotide sequences of primers are given in Supporting Information Table S1.

To measure relative transcript abundance of *Slp1* in ASZ-2, ASZ-1, ASZ, ASZ+1 and the youngest mature leaf, three independent sorghum seedlings with immature leaf length between 13-16 cm were used. The youngest mature leaves were used as a measure of background transcription level of *Slp1* as silica cells in these regions are dead and silicification is no longer taking place in them (Kumar & Elbaum, 2018). RNA isolation and cDNA synthesis were carried out as written above. Reactions were performed using SYBR mix (Invitrogen, MA, USA) in 7300 Real Time PCR machine (Applied Biosystems, MA, USA) as explained before (Markovich *et al.*, 2015) except that the primer concentrations were kept at 150 nM each. A melt curve analysis at the end of the PCR cycle was carried out to ensure that only one PCR product was formed. *Slp1* and *UbCE* transcripts were amplified using primers as mentioned above. In addition, we also amplified transcripts of an RNA recognition motif-containing protein (*RRM*; Sb07g027950) as another internal control (Shakoor *et al.*, 2014) using the primers RRM-RT-F and RRM-RT-R (Table S1).

Using *UbCE* as internal control, the relative transcript abundance of *Slp1* was calculated according to the formula of (Pfaffl, 2001). The relative transcript level of *RRM* is also reported, showing that the transcript level of the two house-keeping genes do not change significantly in the tested tissues (Supporting Information Fig. S1).

**Immunolocalization of Slp1 in sorghum leaves**

Leaf tissues from ASZ and mature leaves were cut into about 1 mm X 1 mm pieces and fixed in 4% para-formaldehyde (w/v) containing 0.05% triton X-100 using vacuum infiltration. Blocking was done in PBS buffer containing 0.1% BSA. The tissue sections were incubated with purified polyclonal antibody against Peptide-1 (10 µg ml$^{-1}$) for one hour at room temperature. After washing, the tissue sections were incubated with secondary antibody (goat anti-rabbit IgG) tagged with Alexa flour 488. The tissue sections were further washed and incubated for 10 min in propidium iodide (5 µg ml$^{-1}$) solution to stain the cell wall red. Tissue sections undergone the same treatment (i) with pre-immune serum, or (ii) without the primary or the secondary; or without both the antibodies worked as control. The tissue segments were mounted on microscopic slide and observed under Leica SP8 inverted confocal laser scanning microscope with solid-state laser (Wetzlar, Germany). The excitation wavelength was 488 nm while the emission filter range was 500-550 nm for Alexa flour 488. Propidium iodide was excited at 514 nm and the emission filter range was 598-634 nm.

**Transient overexpression of Slp1 in tobacco (*Nicotiana benthamiana* Domin)**

pBIN-19 plasmid was modified by inserting CaMV35S promoter in between HindIII and SalI

restriction sites, whereas GFP followed by NOS terminator was inserted in between XbaI and

EcoRI restriction sites. The resulting plasmid (pBIN-GFP) was used as control plasmid to drive

the expression the GFP under the control of CaMV35S promoter. Primers Slp1-SalI-F and

Slp1-XbaI-R (Table S1) were used to amplify the full length of Slp1 without the stop codon

and flanked with the restriction sites SalI and XbaI, respectively. The PCR product was

digested with SalI and XbaI and ligated to the SalI and XbaI digested pBIN-GFP plasmid. The

expression cassette consisted of CaMV35S promoter-Slp1-GFP-NoS terminator. We

immobilized the plasmids into *Agrobacterium tumefaciens* (strain EHA105) and infiltrated

*Nicotiana benthamiana* source leaves on the abaxial side with slight modifications from the

reported protocol (Li, 2011). Our resuspension solution consisted of 50 mM MES buffer, 2

mM $NaH_2PO_4$ supplemented with 200 µM acetosyringone. After 48 hours, leaf pieces were

observed under laser scanning confocal microscope (excitation: 488 nm; emission filter range:

500-550 nm).

**Transient overexpression of Slp1 in sorghum**

We created a construct exploiting the maize dwarf mosaic virus (MDMV) genome

(https://patentscope.wipo.int/search/en/detail.jsf?docId=WO2016125143). We fused the

expression cassette containing a CaMV35S promoter driving the expression of Slp1 and *nos*-

terminator in between *Age*I and *Apa*I restriction sites to the MDMV genome as follows. We

amplified full length of Slp1 except the start and stop codons using the primers Slp1-AgeI-F

and Slp1-ApaI-R (Table S1). After digestion of the PCR product with AgeI and ApaI restriction

enzymes, we ligated the digested PCR product with AgeI and ApaI digested MDMV-GUS

plasmid. The resulting plasmid was named MDMV-Slp1. The plasmids were coated on gold

microparticles (1 µm diameter, Bio-Rad, California, USA) and bombarded on about one week

old sorghum seedlings according to the protocol of (Jose-Estanyol, 2013) with the differences

explained below. Sorghum seeds were surface sterilized, germinated and then grown with their

roots immersed in tap water for up to seven days before they were bombarded. Just before

bombardment, the seedling roots were quickly taken out of water, and a bunch of five seedlings

were arranged and loosely stuck in the center of a petri dish and bombarded with the plasmid

coated gold particles using 1100 psi rupture discs. After bombardment, the seedlings roots were

wrapped in moist tissue paper and put in a beaker with tap water for 24 hours, after which the

bombarded seedlings were transferred to soil. The seedlings were grown for about three weeks

and the phenotypes in the leaves showing viral symptoms were observed in the SEM. RNA

was isolated from the young leaves of the infected plants and cDNA was synthesized. Primers

derived from the MDMV-Slp1 plasmid (MDMV-Slp1-F and MDMV-Slp1-R, Table S1) were

used to amplify Slp1 with viral flanking regions, or primers from viral protein coat (MDMV-

Coat-F and MDMV-Coat-R, Table S1) were used to ascertain the viral infection in the control

plants (infected by MDMV-GUS plasmid). We conducted this experiment twice, each time

starting with 15 independent biological replicates and analysing altogether 10 plants showing

strong viral symptoms.

**Results**

**Screening for putative silicification protein(s)**

In order to find candidate silicification regulating factors, we analysed a publicly available data

set which reports the influence of silicon treatments on gene expression in wheat (GEO NCBI

dataset GSE12936) (Chain *et al.*, 2009), as both sorghum and wheat belong to Poaceae (grass)

family. We searched for genes that co-express with wheat silicon transporters (*TaLsi1*, *TaLsi2*,

*TaLsi6*) irrespective of the plant pathogenic infection status, using Comparative Co-Expression

Network Construction and Visualization (CoExpNetViz) online tool (Tzfadia *et al.*, 2016).

Once co-expressed genes were identified, their probe sequences were extracted and used to

BLAST against the sorghum genome. The BLAST yielded a list of 18 sorghum orthologues

genes (e-value > 100 was used as orthology cutoff) (Table S2). The primary sequence of all

the initially screened proteins were analysed for the theoretical isoelectric point (pI) using

ProtParam tool in ExPASy server (https://web.expasy.org/protparam/); and the presence of

predicted signal peptide (SignalP server). Since positively charged amino acids have been

shown involved in biological silicification, we discarded all the proteins with predicted pI value

less than 7. We also rejected the proteins that lacked signal peptide as a silicifying protein must

be secreted outside the silica cell membrane in order for silicification to take place in the

paramural space. Proteins were further screened for the presence of internal repeat units as

found in several silica precipitating proteins (Kröger *et al.*, 1999; Kauss *et al.*, 2003; Shimizu

*et al.*, 2015), abundance of histidine, glutamic acid (Shimizu *et al.*, 2015), lysine (Kröger *et al.*,

1999; Kauss *et al.*, 2003) or serine, glycine-rich motifs, and frequency of proline-lysine and

proline-glutamic acid residues (Harrison, 1996). Based on our selection criteria, we identified

a protein in sorghum (Sb01g025970) that had seven repeat units rich in lysine, similar to

silaffins from *Cylindrotheca fusiformis*, a diatom species (Kröger *et al.*, 1999); histidine-

aspartic acid rich regions similar to glassin from a marine sponge, *Euplectella* (Shimizu *et al.*,

2015), and several proline residues as in a proline-rich protein (PRP1) from cucumber (Kauss

*et al.*, 2003). We functionally characterized this protein and named it Siliplant1 (Slp1).

**Molecular architecture of Sorghum Slp1**

Sorghum Slp1 comprises of 524 amino acids with a predicted N-terminal signal sequence (Fig.

1). The signal sequence is predictably cleaved between amino acid positions 24 and 25. Further

sequence characterization using TargetP predicted Slp1 to be a secretory protein with high

probability. BLAST-based homology search revealed that Slp1 is a single copy gene in

sorghum and belongs to a family of proline-rich proteins with unknown function. The transcript

of this protein is highly abundant in young leaves and inflorescences (MOROKOSHI, © 2018

RIKEN, query Sobic.001G265900). Slp1 is rich in proline, lysine, and histidine, respectively

comprising 20%, 13%, and 11% of total amino acids. The theoretical pI value of the protein is

9.28. Structure predictions using GOR4 (Combet *et al.*, 2000) suggest that about 80% of the

protein is random coil and 14% is alpha helix. IUPred suggested that Slp1 has intrinsically

disordered tertiary structure (Dosztányi *et al.*, 2005a,b). Slp1 has repeat units (R1 to R7) rich

in proline, lysine, and glutamic acid (P, K, E-rich domain) with the consensus sequence

KKPXPXKPKPXPKPXPXPX. Near a wide range of physiological pH, close to half of this

domain is positively charged (lysine-rich) and the rest of the domain is negatively charged

(aspartic acid-rich). Five of the repeat units (except R1 and R3) have a histidine and aspartic

acid (H, D) rich domain with the consensus sequence DXFHKXHDYHXXXXHFH

immediately preceding the P, K, E-rich domain. This aspartic acid rich domain is negatively

charged under physiological pH. In five repeats (except R2 and R4) there is a third domain

following the P, K, E-rich domain, rich in proline, threonine and tyrosine (P,T,Y-rich) with the

consensus sequence YHXPXPTYXSPTPIYHPPX. Before the start of the repeat units (amino

acids 1-148), there is a region rich in alanine, serine, glycine and valine (comprising together

45% of the region). At the end of R1, R3, and R5, we found an RXL domain. This domain acts

as recognition site for unknown proteases that cleave at the C-terminus of the leucine residue

of the RXL domain in many bio-silica associated proteins in diatoms (Kröger *et al.*, 1999;

Wenzl *et al.*, 2008; Scheffel *et al.*, 2011; Kotzsch *et al.*, 2017). Despite these common features,

Slp1 does not show sequence similarity with any protein known to be involved in bio-

mineralization.

**Silica precipitated *in vitro* by Peptide-1 derived from Slp1 sequence**

In order to test the possible activity of Slp1 as a silica mineralizer we synthesized the peptide sequence HKKPVPPKPKPEPK (Peptide-1) which appears 5 times in the P, K, E-rich domain of Slp1 primary sequence. When metastable (90.9 mM) silicic acid solution was added to Peptide-1 solution (final peptide concentration 1.82 or 1.36 mg ml$^{-1}$) at neutral pH, the reaction mixture became cloudy within seconds, and very fine, white sand-like powder precipitated. In contrast, a mutant peptide, where the lysine residues in Peptide-1 were replaced by alanine (HAAPVPPAPAPEPA; Peptide-3), did not precipitate powder-like silica at 90.9 mM silicic acid. Gel-like material formed when the reaction mixture was centrifuged (Fig 2a). Scanning electron micrographs of the powder sediment, forming within 5 minutes revealed spheres of about 0.5 microns in diameter (Fig. 2b). Silica was also precipitated by Peptide-1 (1.36 mg ml$^{-1}$) at silicic acid concentration of 5 mM (pH=7.0), which is in the lower concentration range found in grass leaves (Casey *et al.*, 2003). The precipitation was invisible to the naked eyes even after 30 minutes of incubation. However, SEM examination of the sediment showed that there indeed was silica precipitation and the diameter of an individual silica nanosphere was about 250 nm (Fig. 2c). In control solutions (without peptide) at 90.9 and 5 mM silicic acid, precipitation was not observed by SEM. High resolution transmission electron micrographs (HRTEM) of particles of 0.5 microns revealed molecular scale dark and bright regions, possibly resulting from the contrasting electron density of the peptide and poly-siloxane chains. The dark and bright patterns create a short range periodic order a few nanometers long and 1 nm thick.

**Raman and NMR spectroscopic characterization of Peptide-1 – silica precipitate**

The precipitated silica with Peptide-1 at 90.9 mM silicic acid was characterized by Raman and magic angle spinning (MAS) solid-state nuclear magnetic resonance (ss-NMR) spectroscopies (Fig. 3). Raman spectrum of the sediment showed that the mineral is silica, recognized by the peaks at 489 cm$^{-1}$ and 997 cm$^{-1}$ (Aksan *et al.*, 2016). These peaks are missing in the Raman spectrum of Peptide-1 (Fig. 3a). Further analysis indicated chemical interactions between the silica surface and the amino group of the lysine side chain (Supporting Information Notes S1).

By conducting large scale silica precipitation assay with Peptide-1 at 90.9 mM silicic acid, we could collect 113 mg of dried silica, sufficient for NMR analyses. Confirmation of the HRTEM and Raman findings that the peptide was complexed with the silica was given through NMR $^{1}$H-$^{13}$C cross polarization measurements (Fig. 3b). The spectrum shows narrow peaks (full width at half maximum (FWHM) was 282 Hz), as compared with spectra of a diatom peptide complexed with silica (FWHM of 489 Hz) (Geiger *et al.*, 2016). The narrow peaks allowed us

to identify many of the amino acid sidechain carbons, with peaks shifted to a higher field by about 2 ppm. Such shifts, caused by the silica shielding the magnetic field that is felt by the peptide, indicate of close proximity of the mineral to the peptide.

In order to study the structure of the mineral we collected NMR direct polarization signal from Si atoms (Fig. 3c). $^{29}$Si NMR can detect the number of groups of -OSi and -OH bound to a central Si atom. A Qn band is defined as $Si-(OSi)_nOH_{4-n}$ (Engelhardt & Michel, 1987). The relative intensities of the Q4:Q3:Q2 peaks in the spectrum were 62:33:5, indicating that there are about 5 bulk (Q4) Si atoms for every 3 surface (Q3+Q2) atoms. To examine further the silicon atoms at the surface we excited them through near-by hydrogen atoms, by cross polarization measurement (Fig. 3c). This measurement revealed that in addition to surface Q3 and Q2 we detected some Q4 siloxane species bound to surface oxygen. These Si atoms were excited by protons from nearby silanols and water, indicating their closeness to the surface of the silica particle.

**Expression pattern and subcellular localization of Slp1**

Our sequence-based predictions and *in vitro* results suggested a role for Slp1 in silica deposition. To investigate whether the expression profiles of Slp1 correlate with silica deposition times and locations, we raised an antibody against Peptide-1. Using Western hybridization, we detected several bands in immature leaves and inflorescence. Slp1 expression was not detected in roots and mature leaves (Fig. 4a). In correlation with the protein expression profile, we detected RNA transcription of *Slp1* in immature leaves and inflorescences, but not in mature leaves (Fig. 4b). This profile matches regions of active silica deposition, which occurs only in young leaves but not in mature leaves. RNA transcript profile of *Slp1* along immature, silicifying leaf tissues showed that *Slp1* is strongly transcribed in the leaf base, reaching the highest levels just before the active silicification zone (ASZ-1). The transcription levels dropped by a factor of about 15 in the ASZ, and further reduced in older tissues of immature leaves (Fig. 4c). In the youngest mature leaf tissue, we found that *Slp1* is transcribed at a basic, low level constituting the background transcription level in these leaves. Background transcription level was about 1/19,000 that of the maximal transcript in ASZ-1.

To further correlate Slp1 with silica deposition, we tested its distribution in leaf epidermal cells using antibody against Peptide-1. In the ASZ, the antibody bound to silica cells (Fig. 5a,b). We noticed the antibody signal in either the cytoplasmic volume or near the cell periphery of silica cells (Video S1). Further image analysis suggested that the cytoplasmic immunofluorescence

signal originated from vesicles (Fig. 5c). In mature leaves, where silica cells are silicified and dead, the antibody hybridized only to the boundary of silica cells (Fig. 5d,e). No vesicles were identified by image analysis (Fig. 5f). Immunolocalization with the pre-immune serum, or the reactions lacking any one or both the antibodies as control did not fluoresce (Fig. S2).

The presence of N-terminal signal peptide in Slp1 predicted it to be secretory in nature. Hence, to verify this prediction, we made translational fusion of green fluorescent protein (GFP) to the C-terminal of full length Slp1 and transiently overexpressed it in tobacco. Similar to our observations in the silica cells of the ASZ, the cells expressing the fusion protein fluoresced in discrete packets distributed throughout the cytoplasm (Fig. 6a,b). In addition, we could identify packets fusing to the cell membrane as well as diffused fluorescence at cell boundaries (Fig. 6c,d, Video S2). In control tobacco plants that overexpressed GFP without Slp1, the fluorescence was uniform in the cytoplasm and nucleus, and lower between cells (Fig. 6e,f). Mock infiltration of tobacco leaves using MES buffer did not fluoresce (Fig. 6g,h). Thus, our results show that native Slp1 is packed in vesicles inside silica cells and suggest that silica cells release Slp1 packages to the paramural space. Since the apoplasm of grass leaves is super-saturated with silicic acid, secretion of the Slp1 to the apoplast may lead to immediate silica precipitation.

**Functional characterization of Slp1 *in planta***

To elucidate the role of Slp1 *in planta*, we transiently overexpressed it in sorghum, using a construct derived from maize dwarf mosaic virus. Compared with the wild type plants, control sorghum plants infected with virus lacking the Slp1 sequence showed infection lesions and viral symptoms (Fig. S3) but no unusual silica deposition (Fig. 7a-f). In contrast, the Slp1 overexpressing plants had ectopic silica deposition in patches, especially close to the veins and where viral symptoms were observed (Fig. 7g-i). Silica was also deposited in cells which are not usually silicified in sorghum leaves, like guard cells, cork cells and long cells (Fig. 7j-l). SEM in the back-scattered electron mode suggested the silicification intensity to be of a level similar to that in silica cells. Similar observations were recorded in ten independent biological replicates. Energy-dispersive X-Ray spectroscopic signal for silicon from the heavily silicified patches in the Slp1 overexpressing leaves was much higher than in the wild type and control leaves. In correlation, the locations of high Si contained also high concentrations of oxygen, low concentrations of carbon, and similar background levels for other elements.

**Discussion**

Silicification in plants may be passive or biologically controlled depending upon the site of
deposition (Kumar *et al.*, 2017b). Out of several cell types that deposit silica in grasses, silica
cells are unique as more than 95% of cells silicify in young, elongating leaves. Their
mineralization requires metabolic activity and is independent of transpiration (Sangster and
Parry 1971). The fast silica precipitation over a few hours (Lawton, 1980) in addition to the
formation of the mineral at the cell wall, as opposed to the cytoplasm (Kumar *et al.*, 2017a;
Kumar & Elbaum, 2018), point to a factor exported from the cells that induces biogenic silica
formation. Our work shows that silica cells express and export the protein Slp1 to the apoplast,
timed with silicification in the paramural space (Kumar *et al.*, 2017a; Kumar & Elbaum, 2018).
It is obvious that silica deposition depends on the presence of silicic acid which is absorbed by
roots from the soil. Typical silicic acid concentrations measured in the xylem sap are 5-7.5 mM
for wheat (Casey *et al.*, 2003), 5-25 mM for rice (Ma *et al.*, 2002; Mitani & Ma, 2005), and 4,
7-12 mM in sorghum seedlings and mature plants, respectively (our unpublished data). Since
Slp1 overexpression caused deposition in all epidermis cell types, we can conclude that it
actively precipitates silica at *in planta* silicic acid concentrations. Furthermore, the fact the
silica formed ectopically in stomata and long cells indicates that Slp1 is sufficient to cause
precipitation in those cells. Based on our results we conclude that Slp1 in its native form
actively deposits silica *in planta*.

**Expression patterns are in correlation with silica deposition in silica cells**
Slp1 expression was not detected in roots where silica cells do not form. Furthermore, we could
not detect Slp1 expression in mature leaf tissues, in correlation with the absence of viable,
active silica cells (Kumar & Elbaum, 2018). This suggests that silica deposition in live silica
cells is dependent on Slp1, while silica deposition in cell walls and possibly other locations is
governed by other means such as specific cell wall polymers (Fry *et al.*, 2008; Law & Exley,
2011; Kido *et al.*, 2015; Brugiére & Exley, 2017; Kulich *et al.*, 2018). Our report on Slp1
expression in immature inflorescence is consistent with published transcriptomic data, showing
*Slp1* transcript both before and after inflorescence emergence (Table S5 of Davidson et al.
2012). We propose that some cell types in the inflorescence employ Slp1 to produce silica
without transpiration. These may be glume abaxial epidermal cells that silicify before
inflorescence emergence and macro-hairs and the long cells that are located on the abaxial
epidermis of lemma, which lacks stomata and thus the transpiration stream (Hodson *et al.*,
1984; Hodson, 2016; Kumar *et al.*, 2017b).

**Possible post-translational modifications**

Some indication for Slp1 post-translational modifications may be the time gap between its
highest transcription level in the ASZ-1 tissue, and the highest silicification activity in the ASZ
(Kumar *et al.*, 2017a; Kumar & Elbaum, 2018). One such possible modification may be
glycosylation (Elbaum *et al.*, 2009). Processed Slp1 molecules are then packed inside vesicles
(Lawton, 1980) and stored in the cytoplasm for later release (Alberts *et al.*, 2002) at the time
of silicification. Modifications may be tissue and species specific. The difference in the size of
the Western hybridization signal between immature leaf and inflorescence that we observed
may be attributed to differential processing of Slp1 in these two tissues. Slp1 has three RXL
domains in the primary structure. RXL domains are proteolytically cleaved in many diatom
biosilica associated proteins (Kröger *et al.*, 1999; Wenzl *et al.*, 2008; Scheffel *et al.*, 2011;
Kotzsch *et al.*, 2016, 2017). It would be interesting to study if Slp1 undergoes alternative
processing, similar to *Thalassiosira pseudonana* (a diatom species) silaffin (Poulsen & Kröger,
2004).

**Slp1 precipitates together with silicic acid creating highly porous biosilica**

Binding of anti-Peptide-1 antibody in the boundary of dead silica cells of mature leaves
suggests that as Slp1 templates the silica precipitation, it is caught inside the silica structure.
Silicanin-1, a biosilica associated protein from *T. pseudonana* gets embedded inside biosilica
structure and is accessible to anti-silicanin-1 antibody (Kotzsch *et al.*, 2017). Similarly, anti-
Peptide-1 antibody may also have access to the epitope remains on the surface of the deposited
silica in silica cells. Our *In-vitro* experiments support such entrapment, with significant shifts
in the NMR and Raman peaks, indicating close proximity between the peptide and the mineral.
The periodic short-range order, as seen in HRTEM, suggests a very intimate interaction of the
forming mineral and the peptide, similarly to a peptide derived from Silaffin3, a silica
precipitating protein from diatom (Iline-Vul *et al.*, 2018). The mineral that we produced *in vitro*
contains many surface Si atoms, forming an open mesoporous structure. Similarly, native plant
silica may form a permeable matrix that will allow movement of solutes. Diffusion of silicic
acid through the mineral is required in order for silica cells to fully silicify, because the
mineralization front is at the paramural space (Kumar & Elbaum, 2017, 2018; Kumar *et al.*,
2017a).

**Conclusions**

Over-expression and localization studies of Slp1 show that this protein is involved in
silicification in sorghum leaf silica cells. Slp1 has unique amino acid composition, charge
distribution and probably post-translational modifications necessary for its activity. Soon after

cell division, silica cells start their preparation for silicification. Slp1 is transcribed, translated,

and post-translationally modified, packed inside vesicles and stored in the cytoplasm until the

cell is ready to silicify. The vesicles fuse to the membrane and release their content in the

paramural space to come in contact with silicic acid. This immediately precipitates open-

structured silica that allows the diffusion of more silicic acid from the apoplastic space. The

rapid formation of the mineral explains the difficulty associated with finding silicification in

an intermediate state. The expression pattern, localization and modifications of Slp1 in the

inflorescence bracts need to be studied in detail.

**Accession number:** The nucleotide sequence of *Sorghum bicolor* Slp1 has been submitted to

NCBI with the GenBank accession number- MH558953. GenBank accession number of the

pBIN19 plasmid is U09365.1. The MDMV-GUS plasmid that we used in the current study is

covered by a published patent number WO2016125143

(https://patentscope.wipo.int/search/en/detail.jsf?docId=WO2016125143).

**Acknowledgements**

S.K. was a recipient of post-doctoral fellowship from Planning and Budgeting Committee

(PBC), Council of Higher Education, Israel. This research was funded by the Israel Science

Foundation grant 534/14. Partial support to the Centre for Scientific Imaging, Hebrew

University (Rehovot campus) by a grant from USAID (AID-ASHA-G 1400005) to the

American Friends of The Hebrew University and The Robert H Smith Faculty of Agriculture,

Food and Environment is gratefully acknowledged. The authors thank. Shmuel Wolf for the

pBIN-GFP plasmid, and Nerya Zexer and Victor M. Rodriguez Zancajo for measuring silicic

acid concentration in sorghum sap.

**Author Contributions**

S.K. and R.E. planned and designed the research; S.K. identified Slp1 from the list of putative

silicification related proteins and performed most of the experiments; N.A.-F. did the NMR

spectroscopy and analysed the data with G.G.; S.B. performed parts of the experiments; S.K.

and J.A.S.-L. performed immunolocalization experiment under the supervision of Y.H.; O.T.

analysed the microarray data and prepared the list of Si responsive genes in sorghum; A.O.

prepared the MDMV construct under the supervision of H.V.; S.K. and R.E. analyzed the

results and wrote the paper. All the authors commented and approved the final version of the

manuscript for publication.

**References**

**Aksan A, Hubel A, Reategui E**. **2016**. Patent no. US 9492271B2. [WWW document] URL
https://patents.google.com/patent/US9492271 [accessed 04 June 2019] Silica-based
composite ocular device and methods.

**Alberts B, Johnson A, Lewin J, Raff M, Roberts K, Walter P**. **2002**. *Molecular Biology of*
*the Cell*. New York: Garland Science.

**Alexandre A, Basile-Doelsch I, Delhaye T, Borshneck D, Mazur JC, Reyerson P, Santos**
**GM**. **2015**. New highlights of phytolith structure and occluded carbon location: 3-D X-ray
microscopy and NanoSIMS results. *Biogeosciences* **12**: 863–873.

**Brugiére T, Exley C**. **2017**. Callose-associated silica deposition in Arabidopsis. *Journal of*
*Trace Elements in Medicine and Biology* **39**: 86–90.

**Casey WH, Kinrade SD, Knight CTG, Rains DW, Epstein E**. **2003**. Aqueous silicate
complexes in wheat, *Triticum aestivum* L. *Plant, Cell and Environment* **27**: 51–54.

**Chain F, Côté-Beaulieu C, Belzile F, Menzies JG, Bélanger RR**. **2009**. A comprehensive
transcriptomic analysis of the effect of silicon on wheat plants under control and pathogen
stress conditions. *Molecular Plant-Microbe Interactions* **22**: 1323–1330.

**Combet C, Blanchet C, Geourjon C, Deléage G**. **2000**. NPA@: Network Protein Sequence
Analysis. *Trends in Biochemical Sciences* **25**: 147–150.

**Coskun D, Deshmukh R, Sonah H, Menzies JG, Reynolds O, Ma JF, Kronzucker HJ,**
**Bélanger RR**. **2018**. The controversies of silicon's role in plant biology. *New Phytologist*.

**Davidson RM, Gowda M, Moghe G, Lin H, Vaillancourt B, Shiu S-H, Jiang N, Robin**
**Buell C**. **2012**. Comparative transcriptomics of three Poaceae species reveals patterns of gene
expression evolution. *The Plant journal* **71**: 492–502.

**Dosztányi Z, Csizmok V, Tompa P, Simon I**. **2005a**. IUPred: web server for the prediction
of intrinsically unstructured regions of proteins based on estimated energy content.
*Bioinformatics* **21**: 3433–3434.

**Dosztányi Z, Csizmók V, Tompa P, Simon I**. **2005b**. The pairwise energy content estimated
from amino acid composition discriminates between folded and intrinsically unstructured
proteins. *Journal of Molecular Biology* **347**: 827–839.

**Elbaum R, Melamed-Bessudo C, Tuross N, Levy AA, Weiner S**. **2009**. New methods to
isolate organic materials from silicified phytoliths reveal fragmented glycoproteins but no
DNA. *Quaternary International* **193**: 11–19.

**Emanuelsson O, Nielsen H, Brunak S, Von Heijne G**. **2000**. Predicting subcellular
localization of proteins based on their N-terminal amino acid sequence. *Journal of Molecular*
*Biology* **300**: 1005–1016.

**Engelhardt G, Michel D**. **1987**. *High resolution solid state NMR of silicates and zeolites*.
Chichester: John Wiley & Sons.

**Epstein E**. **1994**. The anomaly of silicon in plant biology. *Proceedings of the National*
*Academy of Sciences of the United States of America* **91**: 11–17.

**Fry SC, Nesselrode BHWA, Miller JG, Mewburn BR**. **2008**. Mixed-linkage (1→3,1→4)-
*β*-D-glucan is a major hemicellulose of *Equisetum* (horsetail) cell walls. *New Phytologist*
**179**: 104–115.

**Geiger Y, Gottlieb HE, Akbey Ü, Oschkinat H, Goobes G**. **2016**. Studying the conformation of a silaffin-derived pentalysine peptide embedded in bioinspired silica using solution and dynamic nuclear polarization magic-angle spinning NMR. *Journal of the American Chemical Society* **138**: 5561–5567.

**Harrison CC**. **1996**. Evidence for intramineral macromolecules containing protein from plant silicas. *Phytochemistry* **41**: 37–42.

**Hodson MJ**. **2016**. The development of phytoliths in plants and its influence on their chemistry and isotopic composition. Implications for palaeoecology and archaeology. *Journal of Archaeological Science* **68**: 62–69.

**Hodson MJ, Sangster AG, Parry DW**. **1984**. An ultrastructural study on the development of silicified tissues in the lemma of *Phalaris canariensis* L. *Proceedings of the Royal Society of London B* **222**: 413–425.

**Hodson MJ, Sangster AG, Parry DW**. **1985**. An ultrastructural study on the developmental phases and silicification of the glumes of *Phalaris canariensis* L. *Annals of Botany* **55**: 649–665.

**Hodson MJ, White PJ, Mead A, Broadley MR**. **2005**. Phylogenetic variation in the silicon composition of plants. *Annals of Botany* **96**: 1027–1046.

**Iline-Vul T, Adiram-Filiba N, Matlahov I, Geiger Y, Abayev M, Keinan-Adamsky K, Akbey U, Oschkinat H, Goobes G**. **2018**. Understanding the roles of functional peptides in designing apatite and silica nanomaterials biomimetically using NMR techniques. *Current Opinion in Colloid and Interface Science* **33**: 44–52.

**Jose-Estanyol M**. **2013**. Maize embryo transient transformation by particle bombardment. *Bio-protocol* **3**: e865.

**Kaufman PB, Bigelow WC, Petering LB, Drogosz FB**. **1969**. Silica in developing epidermal cells of avena internodes: electron microprobe analysis. *Science* **166**: 1015–1017.

**Kaufman PB, Dayanandan P, Franklin CI**. **1985**. Structure and function of silica bodies in the epidermal system of grass shoots. *Annals of Botany* **55**: 487–507.

**Kauss H, Seehaus K, Franke R, Gilbert S, Dietrich RA, Kröger N**. **2003**. Silica deposition by a strongly cationic proline-rich protein from systemically resistant cucumber plants. *The Plant Journal* **33**: 87–95.

**Kido N, Yokoyama R, Yamamoto T, Furukawa J, Iwai H, Satoh S, Nishitani K**. **2015**. The matrix polysaccharide (1;3,1;4)-2-$\beta$-glucan is involved in silicon-dependent strengthening of rice cell wall. *Plant and Cell Physiology* **56**: 268–276.

**Kotzsch A, Gröger P, Pawolski D, Bomans PHH, Sommerdijk NAJM, Schlierf M, Kröger N**. **2017**. Silicanin-1 is a conserved diatom membrane protein involved in silica biomineralization. *BMC Biology* **15**: 65.

**Kotzsch A, Pawolski D, Milentyev A, Shevchenko A, Scheffel A, Poulsen N, Shevchenko A, Kröger N**. **2016**. Biochemical composition and assembly of biosilica-associated insoluble organic matrices from the diatom *Thalassiosira pseudonana*. *Journal of Biological Chemistry* **291**: 4982–4997.

**Kröger N, Deutzmann R, Sumper M**. **1999**. Polycationic peptides from diatom biosilica that direct silica nanosphere formation. *Science* **286**: 1129–1132.

**Kulich I, Vojtíková Z, Sabol P, Ortmannová J, Neděla V, Tihlaříková E, Žárský V**.
**2018**. Exocyst subunit EXO70H4 has a specific role in callose synthase secretion and silica
accumulation. *Plant Physiology*: pp.01693.2017.

**Kumar S, Elbaum R**. **2017**. Estimation of silica cell silicification level in grass leaves using
in situ charring method. *Bio-protocol* **7**: e2607.

**Kumar S, Elbaum R**. **2018**. Interplay between silica deposition and viability during the life
span of sorghum silica cells. *New Phytologist* **217**: 1137–1145.

**Kumar S, Milstein Y, Brami Y, Elbaum M, Elbaum R**. **2017a**. Mechanism of silica
deposition in sorghum silica cells. *New Phytologist* **213**: 791–798.

**Kumar S, Soukup M, Elbaum R**. **2017b**. Silicification in grasses: variation between
different cell types. *Frontiers in Plant Science* **8**: 438.

**Law C, Exley C**. **2011**. New insight into silica deposition in horsetail (Equisetum arvense).
*BMC Plant Biology* **11**: 112.

**Lawton JR**. **1980**. Observations on the structure of epidermal cells, particularly the cork and
silica cells, from the flowering stem internode of *Lolium temulentum* L. (Gramineae).
*Botanical Journal of the Linnean Society* **80**: 161–177.

**Li X**. **2011**. Infiltration of *Nicotiana benthamiana* protocol for transient expression via
*Agrobacterium*. *Bio-protocol* **1**: e95.

**Ma JF, Tamai K, Ichii M, Wu GF**. **2002**. A rice mutant defective in Si uptake. *Plant*
*Physiology* **130**: 2111–2117.

**Ma JF, Tamai K, Yamaji N, Mitani N, Konishi S, Katsuhara M, Ishiguro M, Murata Y,**
**Yano M**. **2006**. A silicon transporter in rice. *Nature* **440**: 688–691.

**Ma JF, Yamaji N, Mitani N, Tamai K, Konishi S, Fujiwara T, Katsuhara M, Yano M**.
**2007**. An efflux transporter of silicon in rice. *Nature* **448**: 209–212.

**Markovich O, Kumar S, Cohen D, Addadi S, Fridman E, Elbaum R**. **2015**. Silicification
in leaves of sorghum mutant with low silicon accumulation. *Silicon* **doi:** 10.1007/s12633-
015-9348-x.

**Massiot D, Fayon F, Capron M, King I, Le Calvé S, Alonso B, Durand J-O, Bujoli B,**
**Gan Z, Hoatson G**. **2002**. Modelling one- and two-dimensional solid-state NMR spectra.
*Magnetic Resonance in Chemistry* **40**: 70–76.

**Mitani N, Ma JF**. **2005**. Uptake system of silicon in different plant species. *Journal of*
*experimental botany* **56**: 1255–1261.

**Motomura H, Fujii T, Suzuki M**. **2000**. Distribution of silicified cells in the leaf blades of
*Pleioblastus chino* (Franchet et Savatier) Makino (Bambusoideae). *Annals of Botany* **85**:
751–757.

**Perry CC**. **2003**. Silicification: The processes by which organisms capture and mineralize
silica. *Reviews in Mineralogy and Geochemistry* **54**: 291–327.

**Perry CC, Mann S, Williams RJP**. **1984**. Structural and analytical studies of the silicified
macrohairs from the lemma of the grass *Phalaris canariensis* L. *Proceedings of the Royal*
*Society of London B* **222**: 427–438.

**Petersen TN, Brunak S, von Heijne G, Nielsen H**. **2011**. SignalP 4.0: discriminating signal
peptides from transmembrane regions. *Nature Methods* **8**: 785–786.

**Pfaffl MW**. **2001**. A new mathematical model for relative quantification in real-time RT-
PCR. *Nucleic acids research* **29**: 2002–2007.

**Poulsen N, Kröger N**. **2004**. Silica morphogenesis by alternative processing of silaffins in
the diatom *Thalassiosira pseudonana*. *The Journal of Biological Chemistry* **279**: 42993–
42999.

**Scheffel A, Poulsen N, Shian S, Kröger N**. **2011**. Nanopatterned protein microrings from a
diatom that direct silica morphogenesis. *Proceedings of the National Academy of Sciences of*
*the United States of America* **108**: 3175–3180.

**Shakoor N, Nair R, Crasta O, Morris G, Feltus A, Kresovich S**. **2014**. A *Sorghum bicolor*
expression atlas reveals dynamic genotype-specific expression profiles for vegetative tissues
of grain, sweet and bioenergy sorghums. *BMC plant biology* **14**: 35.

**Shimizu K, Amano T, Bari MR, Weaver JC, Arima J, Mori N**. **2015**. Glassin, a histidine-
rich protein from the siliceous skeletal system of the marine sponge *Euplectella*, directs silica
polycondensation. *Proceedings of the National Academy of Sciences of the United States of*
*America* **112**: 11449–11454.

**Shimizu K, Cha J, Stucky GD, Morse DE**. **1998**. Silicatein α: Cathepsin L-like protein in
sponge biosilica. *Proceedings of the National Academy of Sciences of the United States of*
*America* **95**: 6234–6238.

**Skinner RH, Nelson CJ**. **1995**. Elongation of the grass leaf and its relationship to the
phyllochron. *Crop Science* **35**: 4–10.

**Wenzl S, Hett R, Richthammer P, Sumper M**. **2008**. Silacidins: Highly acidic
phosphopeptides from diatom shells assist in silica precipitation in vitro. *Angewandte Chemie*
**47**: 1729–1732.

**Yamaji N, Mitatni N, Ma JF**. **2008**. A transporter regulating silicon distribution in rice
shoots. *The Plant Cell* **20**: 1381–1389.

**Supplementary Information**

**Fig. S1** Transcript level of *RRM* in relation to *UbCE* as internal control gene, showing that the

transcript level of the two house-keeping genes do not change significantly in the tested tissues.

**Fig. S2** Immunolocalization control reactions using pre-immune serum; or lacking either one

or both the antibodies.

**Fig. S3** Maize dwarf mosaic virus (MDMV) infected sorghum plants showing symptoms.

**Table S1** List of primers used in the present study.

**Table S2** List of the wheat (genes identified that show significantly differential transcription

upon silicon treatment verses the non-treated plants, and their *Sorghum bicolor* homologues.

**Video S1** Confocal microscopy video clip showing Slp1 localization in the cytoplasmic space

and near the cell boundary of silica cells.

**Video S2** Confocal microscopy video clip showing the vesicles packed Slp1 fusing to the cell

membrane.

**Notes S1** Raman analysis

**Fig. 1** Primary sequence of *Sorghum bicolor* Slp1, showing the N-terminal signal sequence and the seven repeats (R1-R7). The predicted signal sequence is shown in italics. The arrow denotes the predicted signal sequence cleavage site. All the repeats (R1-R7) have a proline, lysine and glutamic acid rich (P, K, E-rich) domain forming a zwitterionic center. In addition, a histidine-aspartic acid rich (H, D-rich) domain, negatively charged near physiological pH precedes the P, K, E-rich domain in repeats 2, 4, 5, 6 and 7. Whereas, a proline, threonine and tyrosine rich (P, T, Y-rich) domain follows the P, K, E-rich domain in repeats 1, 3, 5, 6 and 7. At the end of repeats 1, 3 and 5, there is an RXL domain (shaded) which serves as cleavage site for unknown proteases in many bio-silica associated proteins in diatoms. The underlined sequence is Peptide-1 which appears five times in the primary sequence and was used for raising antibodies and for silica precipitation assays. The consensus sequence for the domains is given in boxes below each domain, where X denotes any amino acid.

**Fig. 2** Imaging of silica precipitation by peptides derived from *Sorghum bicolor* Slp1 sequence.

(a) Sand-like powder sediment produced from metastable (90.9 mM) silicic acid solution with

Peptide-1, while a gel-like material formed after several rounds of centrifugation of silicic acid

solution with Peptide-3. (b) Scanning electron micrograph of the powder sediment formed by

Peptide-1 at 90.9 mM silicic acid. (c) Scanning electron micrograph of the silica sediment

formed by Peptide-1 at 5 mM silicic acid. (d) High-resolution transmission electron microscopy

(HRTEM) of one particle imaged in panel (b) reveals a mesoporous structure on a nanometric

scale. Bar, 5 nm. (e) Dark and bright granulation have short-range order, as marked by arrows

in the inset panel. Bars of 1 μm in panels (b, c), 60 nm μm in (d), 5 nm in (e), and 2 nm in (e-

inset).

**Fig. 3** Characterization of the peptide-silica sediment using Raman and NMR spectroscopy. (a) Raman spectra of Peptide-1 (red line) and the sediment formed with Peptide-1 (black line). The spectra show that the mineral is silica, with altered surface, and that the peptide is bound to the mineral through the lysine aliphatic amine and proline ring residues and the -COO⁻ C-terminal group. See Note S1 for peak assignment. (b) Spectrum of the peptide-silica sediment, measured by magic angle spinning (MAS) solid-state nuclear magnetic resonance (ss-NMR) $^1$H-$^{13}$C cross polarization. The NMR signals, typical to aliphatic bonds in the amino acid sidechain, are shifted to a high field by about 2 ppm. These shifts reflect a shielding effect of the silica, suggesting that the sidechains are bound to the mineral. (c) Spectra of $^{29}$Si showing peaks of Si-(OSi)$_2$(OH)$_2$ (Q2); Si-(OSi)$_3$(OH) (Q3); and Si-(OSi)$_4$ (Q4). Direct polarization (black line) samples all the Si atoms in the sample, while $^1$H-cross polarization (blue line) samples Si atoms in proximity to protons.

**Fig. 4** Expression profile of *Sorghum bicolor* Slp1. (a) Western blot of crude protein extract from roots (R), immature silicifying leaves ($L_{IS}$), mature leaves ($L_M$), and immature inflorescences (ImI), detected by an antibody against Peptide-1. Slp1 was expressed in developing leaves and inflorescence, but not in roots and mature leaves. Multiple sized bands in the inflorescence may indicate that Slp1 is processed differently in immature leaf and inflorescence. (b) RNA transcription of *Slp1* in immature silicifying leaves ($L_{IS}$), mature leaves ($L_M$), and immature inflorescences (ImI). In accordance with protein translation, transcription was detected only in the immature tissues. *UbCE* (ubiquitin-conjugating enzyme) was used as internal control gene. (c) RNA transcript profile of *Slp1* along immature silicifying leaf tissues. Maximal transcript was found in ASZ-1, which lies just below the Active Silicification Zone (ASZ). Error bars indicate standard deviation (n=3).

**Fig. 5** Immunolocalization of *Sorghum bicolor* Slp1 in the active silicification zone (ASZ) of young leaves (a-c), and near the tips of fully silicified mature leaves (d-f). In sorghum leaves, silica cells usually pair with cork cells (indicated by thunder sign) and exist as long chains of alternating silica and cork cells; but near the leaf tip, cork cells are absent and silica cells fuse to form long polylobate bodies. (a) Immunolocalization (green fluorescence) shows that Slp1 is localized to silica cells of sorghum leaf active silicification zone (ASZ), inside the cytoplasm (arrows) or near the cell periphery (arrowhead). (b) Fluorescence image in (a) merged with the corresponding brightfield image. (c) The anti-peptide-1 antibody fluorescence was processed to select for punctuated regions (pseudocoloured yellow) in the ASZ. The Slp1 appears in packets in the cytoplasm and cell boundary. (d) The anti-peptide-1 antibody (green fluorescence) binds to the edges of silicified silica cells (arrows) in mature leaves. (e) Fluorescence image in (d) merged with the corresponding brightfield image. (f) Processed fluorescence of mature leaf image demonstrates that peptide-1 (pseudocoloured yellow) is embedded inside the polymerized silica. Red fluorescence is from propidium iodide, blue is background autofluorescence of the cell walls. Bar 25 µm in panels (a, b, d, e) and 4 µm in panels (c) and (f).

**Fig. 6** Fluorescence confocal microscopy showing the secretory nature of *Sorghum bicolor* Slp1. (a) Slp1 fused with GFP was transiently overexpressed in tobacco leaf using *Agrobacterium tumefaciens*. The green fluorescence, marking the location of Sil, was found in packets inside the cytoplasm (arrows), while diffused green fluorescence can also be seen along the margin of the cell. White rectangle marks a region enlarged in panel (c). (b) Fluorescent image in (a) merged with the corresponding brightfield image. (c) Segmenting the green fluorescence to diffuse (green) and punctate (pseudocoloured yellow) regions shows packets fusing to the cell membrane (arrows) as well as diffused fluorescence at cell boundaries. (d) Fluorescent image in (c) merged with the corresponding brightfield image (e) Control plants expressing GFP without Slp1 showing green uniform fluorescence of the cytoplasm and nucleus (arrowheads). Arrows indicate the cytoplasm of adjacent cells, demonstrating the low fluorescence between cells. (f) Fluorescent image in (e) merged with the corresponding brightfield image. (g) Mock infiltration of tobacco leaves using MES buffer did not fluoresce. (h) Fluorescent image in (h) merged with corresponding brightfield image. Bars in panels (a, b, e, f, g, h) are 20 µm, and in (c, d) are 10 µm.

**Fig. 7** Overexpression of *Sorghum bicolor* Slp1 in sorghum. Arrowheads indicate silica cells, arrows- stomata, thunder signs- cork cells, stars- long cellsand tick mark indicate viral lesions. (a,d,g,j) Scanning electron microscope (SEM) image in back-scattered electron (BSE) mode. Silicified cells appear brighter compared with the background. (b,e,h,k) Energy-dispersive X-ray spectra (EDS) of the corresponding SEM image showing the comparative elemental composition of the scanned area. (c,f,i,l) EDS map for Si-signal form the corresponding SEM image. Maps are pseudocoloured. Black colour means no signal and increasing intensity of the colour denotes higher density of Si atoms. (a) Wild type sorghum mature leaf scanned by SEM under low magnification showing the general silification pattern in a mature leaf. Inset: epidermal cell types are indicated in a higher magnification image. (b) EDS of the image in (a). (c) EDS map for Si-signal of the image in (a). (d) SEM image of control plants showing silicification only at usual locations. Dust particle with high Si content is seen on the bottom

right.  (f) EDS map for Si of the image in (d). (g) SEM image of ectopic silica deposition in a viral lesion in Slp1 overexpressing plant. (h) EDS of the image in (g). (i) EDS map for Si of the image in (g). (j) SEM image of Slp1 overexpressing plants showing high intensity silica deposition in cells that do not usually silicify in wild type plants. All epidermal cell types can be seen silicified. (k) EDS of the image in (j). (l) EDS map for Si of the image in (j).  Bars represent 200 µm in panels (a; d and g); and 50 µm in panel (j).