<u>Title: Many unreported crop pests and pathogens are probably already present</u>

Running title: Crop pest and pathogen distributions

Daniel P. Bebber[1*], Elsa Field[2], Gui Heng[3], Peter Mortimer[3], Timothy Holmes[4], Sarah J. Gurr[1]

[1] Department of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, UK

[2] Department of Plant Sciences, University of Oxford, South Parks Road, Oxford, OX1 3RB, UK

[3] Kunming Institute of Botany, Chinese Academy of Sciences, 132 Lanhei Road, Kunming 650201,

People's Republic of China

[4] CABI, Nosworthy Way, Wallingford, OX10 8DE, UK

[*] Corresponding author, email d.bebber@exeter.ac.uk, tel. +44 1392 725851

Summary

Biotic invasions threaten global biodiversity and ecosystem function. Such incursions present

challenges to agriculture where invasive pest species cause significant production losses require major

economic investment to control and can cause significant production losses. Pest Risk Analysis (PRA)

is key to prioritizing agricultural biosecurity efforts, but is hampered by incomplete knowledge of

current crop pest and pathogen distributions. Here we develop predictive models of current pest

distributions and test these models using new observations at sub-national resolution. We apply

generalized linear models (GLM) to estimate presence probabilities for 1739 crop pests in the CABI

pest distribution database. We test model predictions for 100 unobserved pest occurrences in the

People's Republic of China (PRC), against observations of these pests abstracted from the Chinese

literature. This resource has hitherto been omitted from databases on global pest distributions. Finally,

we predict occurrences of all unobserved pests globally. Presence probability increases with host

presence, presence in neighbouring regions, per capita GDP, and global prevalence. Presence

probability decreases with mean distance from coast and known host number per pest. The models were

good predictors of pest presence in Provinces of the PRC, with area under the ROC curve (AUC) values

of 0.75 – 0.76. Large numbers of currently unobserved, but probably present pests (defined here as

unreported pests with a predicted presence probability > 0.75), are predicted in China, India, southern

Brazil and some countries of the former USSR. Our results shows that GLMs can predict presences of

pseudo-absent pests at sub-national resolution. The Chinese scientific literature has been largely

inaccessible to Western academia but contains important information that can support PRA. Prior

studies have often assumed that unreported pests in a global distribution database represents a true

absence. Our analysis provides a method for quantifying pseudo-absences to enable improved PRA and

species distribution modelling.

Keywords

biogeography, crop pathogens, crop pests, Generalized Linear Model, observational bias, Pest Risk

Analysis, pseudo-absence, species distribution model.

Introduction

The spread of invasive species is homogenizing the biosphere, with wide-ranging implications for natural ecosystems (Baiser *et al.*, 2012; Santini *et al.*, 2013) and agriculture (Fisher *et al.*, 2012; Bebber *et al.*, 2014a; Bebber, 2015). The number of first observations of crop pests and pathogens has accelerated in recent years, driven primarily by global trade (Ding *et al.*, 2008; Bacon *et al.*, 2014), but also potentially by climate change and our improving ability to monitor and identify threats (Bebber *et al.*, 2014a; Bebber, 2015). Here, we use the term 'pest' to describe any herbivorous arthropod, pathogenic microbe or virus known to attack agricultural crops. Emerging pests can be extremely damaging to agricultural production and the economy, causing both pre-harvest and post-harvest losses (Bebber & Gurr, 2015; Paini *et al.*, 2016; Savary *et al.*, 2017). Recently, for example, sub-Saharan Africa has suffered from the virulent Ug99 strain of the wheat stem rust fungus (*Puccinia graminis tritici*) (Patpour *et al.*, 2015), the newly-evolved Maize Lethal Necrosis viral syndrome (Wangai *et al.*, 2012), arrival of the Asian citrus psyllid (*Diaphorina citri*) which vectors citrus greening disease (Shimwela *et al.*, 2016), and the appearance of Tropical Race 4 of *Fusarium oxysporum f. sp. cubense* attacking Cavendish bananas (Ordonez *et al.*, 2015). Central America, Europe, East Africa and Australia have been identified as hotspots of new pest invasions, with maize, bananas, citrus and potato as the crops most likely to be affected (Bebber, 2015). Outbreaks of resident pests due to favourable weather conditions, virulence evolution, or crop management factors, add to the burden on farmers. For example, a major outbreak of coffee leaf rust (*Hemileia vastatrix*) in Latin America, likely to have been triggered by a failure in disease management, is reported to have caused large-scale unemployment and social upheaval in recent years (Avelino *et al.*, 2015).

Despite the expanding ranges of many pests, complete occupation of their potential ranges has not yet occurred (Bebber *et al.*, 2014a) and so there remains a strong impetus for implementation of biosecurity measures at international borders (Fears *et al.*, 2014; Flood & Day, 2016; MacLeod *et al.*, 2016). Control of spread within countries is extremely difficult because of unhindered transport of plants and soils (Ward, 2016), and biosecurity measures focus on quarantine and inspections at borders (MacLeod *et al.*, 2016). A key component of international phytosanitary action is Pest Risk Analysis (PRA), a suite of methods that allow countries to prioritize protective measures against those pests most likely to arrive and cause serious economic damage (Robinet *et al.*, 2012; Baker *et al.*, 2014). PRA involves assessment of the likelihood of pest arrival, the likelihood of establishment, the potential economic impact if uncontrolled, and the prospect of successful control or eradication (Baker *et al.*, 2014). To date, PRA has primarily been based upon expert opinion regarding the likelihood of arrival and potential impact of individual pests. For example, the UK's recently-established Plant Health Risk Register (PHRR) (Baker *et al.*, 2014) employs simple climate-matching (based on known pest distributions) and host availability to assign qualitative risks of invasion and impact, but not quantitative predictive models. Examples of registered pests include the Oleander aphid *Aphis nerii* which has been assigned

78   very low likelihoods of arrival and establishment, and would cause negligible damage if it did, whereas
79   the Zebra chip phytoplasma *Candidatus liberibacter solanacearum* is thought moderately likely to
80   arrive and would have a very serious impact if it did (DEFRA, 2018).
81
While quantitative PRA protocols have been recently developed recently by the European Food Safety
Authority (Jeger *et al.*, 2018), examples of quantitative PRA are rare in international phytosanitary
legislation and practice. This contrasts with the long and vibrant history of research in predictive species
distribution modelling (SDM) for pests (Elith & Leathwick, 2009; Sutherst, 2014). The geographic
distributions of species are non-random, determined by their biotic environment (e.g. hosts or prey), the
abiotic environment (e.g. climate, edaphic factors), and migration (dispersal to suitable habitat)
(Soberón & Peterson, 2005; Soberón, 2007; Soberón & Nakamura, 2009). Thus, pest invasion risk is,
in theory, quantifiable. Numerous modelling approaches are now available to predict the likely
distributions and impacts of pests (Elith & Leathwick, 2009; Venette *et al.*, 2010; Robinet *et al.*, 2012),
ranging from process-based, or mechanistic models, to statistical, or correlative approaches (Dormann
et al., 2012). Regional and global databases on known pest distributions are commonly used to
parameterize these models, either providing direct estimates of pests' ecological niches (Venette *et al.*,
2010; Kriticos, 2012), or indirectly via shared geographic ranges (Paini *et al.*, 2010, 2016; Eschen *et*
al., 2014).
96
One seldom-acknowledged issue with pest distribution data in global databases is geographic bias in
the likelihood that a pest will be detected, correctly identified, reported and recorded (Pyšek *et al.*,
2008). Analysis of one of the most widely studied global pest distribution databases suggests that
hundreds of pests already present in many developing countries have not been reported (Bebber *et al.*,
2014b). The total number of observed pests in an administrative area (country, or administrative
division for larger countries) can be largely explained by scientific capacity and agricultural production.
Under a scenario of globally high scientific and technical capacity (i.e. where all countries have US-
level per capita GDP and research expenditure), analysis predicts that many countries across the
developing world would report hundreds more pests. This suggests that a large fraction of the *current*
agricultural pest burden is unreported and unknown, and that even the best global databases suffer from
severe observational bias. This has potentially serious consequences for both plant biosecurity activities
and for research based upon these databases. Such observational bias may have implications for SDM
methods that infer environmental tolerances from observed distributions. Scientific capacity, economic
development, and the ability to detect, identify and report pests, are strongly correlated with latitude, as
is climate (Bebber *et al.*, 2014b). Under-reporting of pests at low latitudes will therefore bias estimation
of climate tolerances, as occurrence is underreported in warmer regions. Reducing this observational
bias by strengthening pest identification efforts in the developing world is therefore critical in
improving scientific understanding of pest distributions, and in PRA.

The People's Republic of China (henceforth referred to as China) has been predicted to harbour the largest number of pests (Bebber *et al.*, 2014b). China produces the largest quantity of crops by tonnage globally, and has the greatest diversity of production. Both factors are strong determinants of recorded pest numbers (Bebber *et al.*, 2014b). Yet, the actual recorded number of pests in China is much smaller than expected (Bebber *et al.*, 2014b). For many countries, under-reporting of agricultural pests is likely to be purely a function of the lack of institutional capacity to detect, identify, and report incidences in the scientific and 'grey' literature used by CABI to populate the distribution database. For China, there is potentially an interesting alternative. The Chinese literature was, until the reforms of 1978, largely inaccessible to Western academia. Even post-reform and the opening of China instigated by Deng Xiaoping, Chinese-language publications are not commonly accessed by English-speaking researchers. A scientifically-important translation of the Chinese literature is the reporting of the anti-malarial compound artemisinin (Klayman, 1985). The Chinese research literature, having developed isolated from the Western literature, therefore provides a potentially independent data source for testing models of pest distributions.

Here, we develop statistical models of global pest presence using a database of known pest occurrences and confront the predictions of individual pest or pathogen presence in China's provinces with observations from the Chinese literature. In addition, we compare models in which pest absences are treated as true absences with models in which absences are weighted according to estimates of scientific and technical capacity of a given country to report plant health risks, to investigate the effect of observational bias and pseudoabsences in pest distribution modelling. We then apply our distribution models globally to all unreported pests in all regions, to give predicted probabilities of presence. Finally, we list those pests that are probably present, but as yet unreported, around the world.

Materials and Methods

We obtained pest distribution data from the CABI Knowledge Bank database in January 2014 with permission (CABI, Wallingford, UK). The database comprised 91,030 records of the observed distributions of 1901 agricultural pests by administrative division of each country, e.g. US States. In total, 384 geographical units were included in the model, comprising 221 countries plus sub-national divisions for Australia (7), Brazil (28), Canada (13), China (31), India (33), and the USA (51). Geographical regions such as Bouvet Island which were smaller than a single pixel (5 arc minute resolution, or approximately 100 sq. km) of the gridded crop distribution database we employed were excluded from the analysis. Host crop spatial distributions for 175 crops at 5 arc minute resolution were obtained from the EarthStat database (http://www.earthstat.org/; Monfreda *et al.*, 2008). Known plant hosts of each pest were taken from the CABI Knowledge Bank, and the host genera matched to the genera in the list of 175 crops. Pests without known hosts in this list of 175 crops were excluded from

the analysis. Pests from taxonomic groups with fewer than 50 species (e.g. Acari, Gastropoda and

various other insect taxa) were also excluded from the analysis. This left a total of 1739 pests comprising

124 species, subspecies and pathotypes of Bacteria, 106 Diptera, 215 Coleoptera, 398 Fungi, 233

Hemiptera, 248 Lepidoptera, 99 Nematoda, 61 Oomycota and 209 viruses. Assigning reported

presences for each pest to each geographical region gave a dataset of 667,776 presences or absences for

each pest-region pair. In total, there were 81,821 presences (12.2 per cent of the total) in the final dataset.

We developed Generalized Linear Models (GLM), using the *glm* function (*MASS* package) in R v.3.4.0

with logit link for binomial data (R Development Core Team, 2017), for the presence or (pseudo-)

absence of each pest in each region. Model predictors were as follows: log-transformed *per capita* GDP

for the country as a whole in 2016 (World Bank data, http://data.worldbank.org/); log-transformed total

number of crop host genera for the pest (CABI Knowledge Bank, obtained with permission); log-

transformed area (ha) of the pest's host crop distribution (summing planted areas of all known host

crops in each geographical region); log-transformed host crop area (ha) of neighbouring (i.e. with land

border) regions which have reported the pest as present (set to zero if no neighbours have reported the

pest); log-transformed total fraction of regions globally that have reported the pest; and log transformed

distance (km) of crop area to the coast (calculated as the distance of the centroid of the crop area

distribution from the nearest coastline). Log transformations were applied to distribute the predictor

variable values more evenly across the sample space. Briefly, the rationale for these predictors was that

GDP is a proxy for historical trade (Pyšek *et al.*, 2010) and observational capacity (Bebber *et al.*,

2014b), host area indicates the available habitat for each pest, host number indicates the degree of biotic

generalism of the pest, neighbouring-region presence indicates the potential for spread across a land

border, fraction of regions reporting presence indicates global ubiquity and environmental generalism,

and distance to coast indicates proximity to international shipping ports (Chapman *et al.*, 2017).

We developed two pest distribution models. The 'unweighted' model included geographical and

biological predictors and treated all unobserved pests as absent from a region. The 'weighted' model

treated unobserved pests as potentially pseudo-absent, using a function of the scientific and technical

capacity of each country (Bebber *et al.*, 2014b). Presences were taken as being correct and

unambiguous, and given a weighting of unity. Absences were weighted by the logarithm of the

agricultural and biological sciences publication output of each country from 1996 – 2016 (Scimago

Lab, 2017), normalized to the logarithm of the output of the USA (the world's most scientifically

productive country), such that the absence weight $w_0 = \log(s)/\log(s_{USA})$. Thus, pests unreported from

scientifically advanced nations were assumed not to be present (or, present at undetectable population

density), while pests unreported from developing nations were less informative of absence. China, with

the second largest research output, had $w_0 = 0.93$, suggesting that non-reporting of a pest should be

relatively strong evidence of its physical absence. However, we hypothesized that non-reporting in the

CABI databases could be due to lack of translation from the Chinese literature, therefore we set w_0 to
zero for China, effectively omitting these pseudo-absences from the analysis. The weighted and
unweighted models were compared with a null model assuming constant presence probability using
Likelihood Ratio Tests.
193
To validate the models we predicted the probability of presence for a random sample of 100 as-yet
unobserved pests in all Chinese Provinces, but excluding Taiwan. The Chinese literature was searched
for observations of these unobserved pests in China. We used the text mining methodology designed
by CABI for their Plantwise Knowledge Bank. The following rules were followed to locate pest records
in the Chinese literature:

- Include only papers that are primarily about distribution data, not those where distribution is
mentioned, but where something else is the primary focus. If this is unclear do not process the
paper.
- Mine only the primary literature (including Masters and Doctoral theses), not meta-analyses,
reviews, or non-peer reviewed ("grey") literature.
- Pest and host names must be preferred scientific names, following the CAB Thesaurus
(www.cabi.org/cabthesaurus/) and the Plant List (http://www.theplantlist.org/).
- Record country and location information given in the paper, including latitude/longitude. CABI
uses five levels for location, from the largest scale (i.e., provincial) to the smallest (i.e.,
village/town).
- Record date of observation/collection (entering each year separately) and date of publication.
Can be left blank if not given, or use the date of receipt in the diagnostic laboratory as a
surrogate for date of collection.
- Record pest status – present/not found. Only record absence if pest absence is specifically stated
in the paper.
- Record pest status using only the status terms defined by CABI, and only if used in the paper
e.g. "widespread", "restricted" "soil only" "greenhouse only" (see CABI guidelines for
complete list).
- Record if the paper was a first record of that pest or not and details of this (e.g. "first record in
<country/location>", "first record on <host species name>")
- Only enter data where the pest/pathogen has been clearly identified, not just symptoms seen.
- Record only natural infections, not artificial inoculants.
221

Combinations of pests and locations were submitted to several search engines. The priority of search
engines was: Baidu (www.baidu.com), China National Knowledge Infrastructure (CNKI,
http://www.cnki.net), Chongqing VIP Information Company (CQVIP, http://lib.cqvip.com/), and
Wangfang Data (http://www.wanfangdata.com.cn). Baidu is the most popular Chinese internet search

engine. CNKI is led by Tsinghua University, and supported by ministries of the Chinese Government.

CQVIP, formerly known as Database Research Center under the Chongqing Branch of the Institute of

Scientific & Technical Information of China (CB-ISTIC), was China's first Chinese journal database

research institution. Wanfang Data is an affiliate of the Chinese Ministry of Science & Technology, and

provides access to a wide range of database resources.

Publication titles were searched first, followed by full text interrogation. The first 50 search results were

scanned before dismissing a search term. The first search term combination was pest name and location

(Province). If this yielded no result, then pest name and various distribution terms were tried. These

distribution terms were: "catalogues" OR "checklists" OR "distribution" OR "inventories" OR "new

records" OR "surveys" OR "geographical distribution" OR "new geographic records" OR "new host

records". Searches included local names in Chinese where these were known or could be identified

from the literature, preferred scientific names, and non-preferred scientific names from CAB Thesaurus

(https://www.cabi.org/cabthesaurus/). Searches continued until one piece of literature was found for

that pest in that region, that fitted all of the requirements for CABI text mining.

If a pest was not found in any of these searches, it was assumed to be absent from the literature and thus

effectively absent from the region. We cannot prove, however, that a pest is present at very low

population density and has not yet been detected (Crooks, 2005).

Modelled probabilities of reported pest presence in the global dataset, P_G, were obtained from the

predictor variables for each pest-region combination, for each GLM (*predict* function in R). We then

compared P_G with the observed presence-absence data for our Chinese sample data using logistic

regressions (*glm* function in R) and Receiver-Operator Characteristic (ROC) curves (*pROC* library for

R). The logistic regression coefficients *c* and *m* determine the probability of pest presence in the Chinese

sample as $P_C = 1/(1 + \exp(-(c + mP_G)))$. ROC curves describe the relationship between the true positive

rate (sensitivity, the fraction of presences correctly identified as presences) and false positive rate (1 –

specificity, where specificity is the fraction of absences correctly classified as absences) as the threshold

for a binary classifier is decreased from one (classifying any presence probability less than one as

absent) to zero (classifying any positive probability as present). A good predictor will have a high true

positive rate and low false positive rate for any classification threshold, whereas a poor predictor will

have roughly equal true and false positive rates (i.e., be uninformative). The Area Under Curve (AUC)

for the ROC curves gives the probability that, for a random pair of presence and absence observations,

the presence probability will be greater for the presence than the absence (Jiménez-Valverde, 2012).

Models with good discrimination ability should have AUC significantly greater than half.

For illustration, we identified probably present pests (PPP) as those for which are currently unreported from a particular region, but for which $P_G > 0.75$ in our weighted model. This threshold was chosen based on the Kent scale which suggests a probability of 0.75 as an event that would generally be described as 'probable' (Kent, 1994). This is an arbitrary definition but allows us to suggest some of the pests that PRA and phytosanitary activities may want to focus on.

Results

Globally, P_G increased significantly with presence in neighbouring regions, the area of host crops, the global prevalence of the pest and per capita GDP in both models (Table 1). P_G declined with mean distance from the coast and known host crop genera per pest. The models explained similar fractions of the deviance, and had very similar ROC curves with AUC around 88 per cent (Table 1). P_G was always higher for the weighted model, because absences were down-weighted (i.e. fewer true zeros), but predictions for the two models were very highly correlated (r = 0.98). The models found the highest P_G for Hemiptera and Lepidoptera, and lowest for Nematoda, Bacteria and Acari, compared with other taxonomic groupings.

Table 1. GLMs for global pest presence. The unweighted model treated unobserved pests as true absences. The weighted model weighted pseudo-absences as a function of country scientific capacity. The unweighted model had AIC = 339872, AUC = 0.88, Nagelkerke R^2 = 0.40, McFadden R^2 = 0.32. The weighted model had AIC = 308171, AUC = 0.88, Nagelkerke R^2 = 0.37, McFadden R^2 = 0.31. CoastDist is distance of crop centroid from the coast (km), GDP is per capita GDP (US\$), Hosts is reported number of host crop genera, HostArea is harvested area of known host crops, NeigArea is harvested area of host crops in neighbouring regions that have reported the pest, and Prevalence is the fraction of all regions that have reported the pest.

	Unweighted model				Weighted model			
	Mean	SE	Z	Pr(>|Z|)	Mean	SE	Z	Pr(>|Z|)
Acari (Intercept)	-3.67	0.051	-72.3	0.000	-0.897	0.055	-16.3	0.000
+ Bacteria	-0.091	0.032	-2.9	0.004	-0.073	0.034	-2.2	0.014
+ Coleoptera	0.036	0.030	1.2	0.240	0.039	0.032	1.2	0.180
+ Diptera	0.092	0.034	2.7	0.006	0.104	0.036	2.9	0.026
+ Fungi	0.027	0.028	1.0	0.337	0.033	0.030	1.1	0.380
+ Hemiptera	0.167	0.029	5.7	0.000	0.150	0.031	4.8	0.000
+ Lepidoptera	0.145	0.029	4.9	0.000	0.134	0.032	4.2	0.000
+ Nematoda	-0.150	0.033	-4.5	0.000	-0.143	0.035	-4.1	0.000
+ Oomycota	0.046	0.034	1.4	0.176	0.061	0.037	1.7	0.151
+ Virus	0.047	0.030	1.6	0.120	0.067	0.033	2.1	0.137

log(CoastDist + 1)	-0.176	0.004	-49.0	0.000	-0.222	0.004	-57.7	0.000
log(GDP + 1)	0.295	0.004	81.5	0.000	0.086	0.004	22.3	0.000
log(Hosts + 1)	-0.300	0.004	-71.4	0.000	-0.297	0.005	-65.6	0.000
log(HostArea + 1)	0.171	0.001	123.1	0.000	0.159	0.001	108.1	0.000
log(NeigArea + 1)	0.140	0.001	181.2	0.000	0.142	0.001	173.5	0.000
log(Prevalence)	0.842	0.007	124.3	0.000	0.867	0.007	121.3	0.000

We validated the models with reports of pests abstracted from the Chinese literature.

For illustration, we defined a 'probably present pest' (PPP) as one unreported from a region, but with $P_G > 0.75$ (using the weighted model). Overall, only 4702 of 585955 (0.8 per cent) of all unreported pest-region combinations fell into this class (Supplementary Table S1). The number of PPPs per pest category was greatest for Fungi (2052) and Hemiptera (859). Overall, 86 per cent of unreported pest-region combinations were predicted to be unlikely ($P_G < 0.25$). China, India, the USA and Eastern Europe had the largest numbers of predicted PPPs, along with other parts of East Asia and Southern Brazil (Figure 1). The top ten PPPs by number of global regions were *Cochliobolus heterostrophus* (Ascomycota: Pleosporales, a pathogen of maize), *Rhopalosiphum padi* (Arthropoda: Hemiptera, cereal pest), *Gibberella fujikuroi* (Ascomycota: Hypocreales, rice pathogen), *Sitophilus zeamais* (Arthropoda: Coleoptera), maize and rice pest), *Schizaphis graminum* (Arthropoda: Hemiptera, pest of Poaceae cereals), *Setosphaeria turcica* (Ascomycota: Pleosporales, maize pathogen), *Aphis spiraecola* (Arthropoda: Hemiptera, wide host range), *Nezara viridula* (Arthropoda: Hemiptera, legume pest), *Acyrthosiphon pisum* (Arthropoda: Hemiptera, legume pest) and *Rhopalosiphum maidis* (Arthropoda: Hemiptera, pest of maize and other crops).

Figure 1. Total number of probably present pests (PPP) in all countries and sub-national regions.

Total numbers of recorded pests in China's Provinces and municipalities increased from northern and central regions to southern and coast regions (Figure 2a), except for the central province of Gansu which had 826 reported pests. There is no obvious reason why numbers would be so large in Gansu. Here,

agricultural production is moderate, and there are no particular academic centres which could account
for observational bias. Hence, the Gansu values appear to be an artefact of the CABI database. The
smallest numbers of recorded pests were in the mountainous provinces of Qinghai (0) and Xizang
Zizhiqu (Tibet) (73), the central provinces of Ningxia (48), and the municipalities of Chongqing (24),
Tianjin (3), Beijing (50) and Shanghai (55). Total numbers were largest in the coastal provinces of
Guangdong (301), Zeijiang (294), Jiangsu (293), Fujian (263), and also in the southern provinces of
Yunnan (291) and Sichuan (259).
316

317
318   Figure 2. a) Total number of pests recorded in the CABI pest distribution database by China Province
319   (excluding Taiwan). Hatched region is Gansu, see text for details. b) Total number of probably present
320   pests (PPP) in China Provinces.
321

We validated our models using published pest observations from the Chinese literature. Both models
were significant predictors of pest presence/absence for 100 randomly-sampled pest-Province
combinations, of which 27 were found to be present (Figure 3, Supplementary Table S2). For the
unweighted model, the coefficients of the logistic function were $c = -1.73 \pm 0.34$ and $m = 3.52 \pm 1.25$
(likelihood ratio test *vs* null model, p = 0.0043). For the weighted model, the coefficients were -1.90 $\pm$
0.38 and 3.19 $\pm$ 0.96 (likelihood ratio test, p = 0.0006). The predictive power of the models was also
tested using ROC curves, demonstrating significant discriminant ability with AUC of 0.76 (95 per cent
Confidence Interval 0.66 – 0.86) for the unweighted model, and AUC 0.75 (0.64 – 0.86) for the
weighted model (Figure 3). Our analysis revealed gaps in the CABI database, which is commonly used
for analyses of global pest distributions. Taking one important potato pest, late blight *Phytophthora*
infestans (Oomycota), as an example, high presence probabilities (> 0.75) were predicted for ten
provinces listed as not reporting this pest in the CABI database. However, this pathogen has been
reported present throughout the potato-growing regions of China, including Guangdong (Guo *et al.*,
2010).
336

Figure 3. Model prediction tests. Observed presence/absence of 100 pest-province combinations vs. P_G from a) unweighted model and b) weighted model. Curves and shaded areas show mean and 95% CI for logistic regression fits. Tick marks show observed data. Grey diagonals show identity relationship. ROC curves for c) unweighted and d) weighted models. Error bars show 95% CI for specificity and sensitivity derived from 2000 bootstrap replications.

For China, the total number of PPPs increased from west to east (Figure 2b), and was greatest in the north eastern provinces of Jilin (59), Heilongjiang (58), and Inner Mongolia (58), the eastern provinces of Shandong (60) and Anhui (61), well as the ports of Shanghai (71) and Tianjin (51). The eastern provinces of Xizang Zizhiqu (Tibet) (1), Qinhai (1), Gansu (0) and Ningxia (2) had the lowest numbers, along with the island of Hainan (0) (Figure 3). The total number of PPPs in China was 827, the majority being Fungi (332) and Hemiptera (175). The top ten most-common PPPs in China were (in decreasing order) *Gibberella fujikuroi* (Ascomycota: Hypocreales, rice pathogen), *Aphis spiraecola* (Arthropoda: Hemiptera, generalist), *Delia platura* (Arthropoda: Diptera, pest of legumes), *Acyrthosiphon pisum* (Arthropoda: Hemiptera, legume pest), *Rhopalosiphum padi* (Arthropoda: Hemiptera, cereal pest), *Schizaphis graminum* (Arthropoda: Hemiptera, pest of Poaceae), *Curvularia* sp. (Fungi: Ascomycota, generalist pathogen), *Rhopalosiphum maidis* (Arthropoda: Hemiptera, pest of maize and other crops), *Agrotis ipsilon* (Arthropoda: Lepidoptera, generalist pest), *Lasiodiplodia theobromae* (Ascomycota: Botryosphaeriales, generalist pathogen). Thus, many of the most common PPPs in China were also common globally.

Discussion

The Chinese literature provided strong and significant support for the predictions of pest distribution models based upon host distribution, pest prevalence, and other socioeconomic factors. China's growing economy is expected to lead to large influxes of invasive species, including pests, in coming years (Ding *et al.*, 2008). Analysis of temporal trends in CABI pest observations show a relatively smooth increase in pests from 1950-2000, but the pattern for China is more complex, with a slow increase from 1950 until the late 1970s, a step increase, and then a more rapid growth in pest numbers from 1980 onwards (Bebber *et al.*, 2014a). One potential determinant of this sudden acceleration is the strong support for science and technology given by Deng Xiaoping in 1978, which lead to an increase in funding and academic freedom following the anti-intellectualism of the Cultural Revolution. China now ranks second only to the USA in annual R&D expenditure (IMF, 2013) and scientific output (Scimago Lab, 2017).

We identified a number of pests that were very likely to be present, and the majority of these PPPs were globally distributed and had wide host ranges. Their distributions commonly spanned wide latitudinal ranges, indicating broad climatic tolerances. *C. heterostrophus*, or Southern Leaf Spot, is primarily known as a pathogen of maize but has a wide host range. It has a wide geographic distribution both latitudinally and across continents, resulting in a high likelihood of occurrence in other regions where hosts are present. For example, *C. heterostrophus* is currently recorded only in eastern regions of North America, where most maize is grown. The lack of reported observations in the western regions of North America may be due to the fact that maize, the major host, is uncommon, and hence the disease currently has little impact. *C. sativus*, causing root and foot rot, also has a very wide geographic distribution, but an even wider host range. It is reported from Texas, Oklahoma, Mississippi, Illinois and Tennessee, but not from neighbouring Arkansas or Missouri. Hence, the high presence probability in these States. A similar pattern is seen for the maize pathogen *S. turcica.* Another global species, *R. maidis*, the green corn aphid, is reported across Europe and in Russia, but, like many other pests, not from the former Soviet states of Ukraine, Belarus, Lithuania, Latvia and Estonia. It is plausible that reporting from these nations was less likely when they were part of the USSR.

Predictors like host availability, presence in neighbouring territories and global prevalence were expected to have positive relations with presence probability. The negative relation with distance from coast is likely to be related to import via shipping ports (Huang *et al.*, 2012; Liebhold *et al.*, 2013), and supports the observation that islands report more pests than countries with land borders (Bebber *et al.*, 2014b). Detailed modelling of individual pest climate responses (Bregaglio *et al.*, 2012; Kriticos *et al.*, 2013) for such a large number of pests was beyond the scope of this study. Implicitly, we can assume that the presence of the host crop indicates that the climate is suitable for the pest (Paini *et al.*, 2016), though we acknowledge that this is not necessarily the case (Berzitis *et al.*, 2014). The negative

relationship with number of host genera per pest might suggest that host specialists are more likely to
invade and establish than host generalists, once host availability has been taken into account. For the
practical purposes of PRA, our models provide reliable probability estimates for the presence of
unreported pests at subnational resolution, and we have provided a global list of the unreported pests
whose presence is most likely (Table S2).

We addressed the issue of pseudo-absences in the CABI data by statistically weighting missing pest
observations in proportion to the scientific output of the reporting nation, since scientific output had
been confirmed as a strong determinant of total reported pest numbers (Bebber *et al.*, 2014b). Often,
unreported pests are treated as true absences in pest risk analyses (Paini *et al.*, 2016). The positive
relation of GDP with presence probability supports our hypothesis that wealthy countries are more
likely to detect and report pests (Bebber *et al.*, 2014b). Once observational bias is controlled for using
scientific capacity-based weighting, per capita GDP becomes a weaker determinant. Our weighted
model has similar overall explanatory power to our unweighted model. Nevertheless, the issue of
observational biases related to country-level socioeconomic variation has been raised several times for
various classes of organism (Jones *et al.*, 2008; Pyšek *et al.*, 2008; Westphal *et al.*, 2008; Boakes *et al.*,
2010; Bebber *et al.*, 2013, 2014b), and we therefore recommend the application of appropriate statistical
controls when analysing datasets produced from reports of species presences (as opposed to
distributional datasets derived from rigorous sampling protocols).

Our SDM was statistical, fitting response functions for various predictors to the probability of pest
presence. Many SDM approaches exist, from highly mechanistic models based on pest biology and
ecology (Bregaglio *et al.*, 2012; Skelsey *et al.*, 2016) to purely statistical models that utilize only
patterns in known distributions (Paini *et al.*, 2010). The rarity of quantitative model input into PRAs is
partly due to the scarcity of empirical data available on pest biology and epidemiology required to
parameterize mechanistic models, and so key biological parameters are often inferred from known
distributions (Robinet *et al.*, 2012). This is particularly the case for newly emergent pathogens for which
experimental investigations have not yet been conducted. The European Food Safety Authority (EFSA)
has developed quantitative PRA guidelines that recommend modelling approaches and data sources for
assessing invasion and establishment risk (Jeger *et al.*, 2018), and application of these methods was
attempted for *Diaporthe vaccinii*, a pest of blueberries (Jeger *et al.*, 2017). However, most of the
epidemiological data required for this pest was unavailable, and the risk assessment was thus based on
expert opinion or data from related pests (Jeger *et al.*, 2017). Epidemiological parameters can be poorly
constrained even for long-established pests. For example, coffee leaf rust fungus (*Hemileia vastatrix*)
has affected coffee production for more than a century, but a recent infection model relied upon
temperature response functions derived from the single available study published three decades
previously (Bebber *et al.*, 2016). Initiatives such as the EU-funded PRATIQUE project (2008-11) have

attempted to fill this knowledge gap and enable modelling by collating available ecophysiological data

for insect pests (Baker, 2012). While the advantages and disadvantages of the many different pest

distribution and impact models continue to be researched and debated (Venette *et al.*, 2010; Dormann

et al., 2012; Robinet *et al.*, 2012; Sutherst, 2014), it is clear that practical application of these methods

in PRA remains limited.

SDM for pests has direct policy implications for PRA and plant biosecurity. PRA is guided by

International Standards for Phytosanitary Measures (ISPM), which are part of the International Plant

Protection Convention (IPPC) (MacLeod *et al.*, 2010). ISPMs tend to rely on expert judgement for

PRA, rather than quantitative modelling to support decision making. ISPM No. 21 "Pest Risk Analysis

for Regulated Non-Quarantine Pests", endorsed in 2004, mentions use of pest and host life-cycle and

epidemiological information, but not quantitative modelling (FAO, 2004). Individual PRAs similarly

employ a qualitative approach. For example, the Australian Government's PRA for *Drosophila suzukii*

references only a single unpublished report on SDM for this species, conducted for North America.

Probabilities of *D. suzukii* spread within Australia are qualitatively assessed by comparison with

observations in other countries (Department of Agriculture, Fisheries and Forestry, 2013). The

European and Mediterranean Plant Protection Organization (EPPO) PRAs occasionally include model

results. For example, a climate matching for the bacterium *Xanthomonas axonopodis* pv. *allii* was

undertaken using the CLIMEX model, to identify areas at risk within the EPPO region (EPPO, 2008).

However, as discussed previously, appropriate empirical studies are rare (Jeger *et al.*, 2017). Our results

contribute to the quantification of risk within PRA by providing probabilistic estimates for the presence

of hundreds of unreported pests around the world, thereby improving understanding of the threats to

global agriculture. With growing evidence that pest ranges are shifting poleward in response to global

climate change (Bebber *et al.*, 2013), our poor knowledge of pest distributions, particularly in the

developing world, is troubling, both because of the burden these organisms place on farmers who have

little access to detection and control technologies, and because invasions of temperate regions are likely

to occur from warmer regions. Improved targeting of phytosanitary measures through quantitative PRA

is therefore vital to crop protection.

Acknowledgements

We thank the British Society for Plant Pathology for an Undergraduate Vacation Bursary for Elsa Field.

We thank Le Yei and Li Huili (Kunming Institute of Botany) for assistance in text mining the Chinese

literature and Ho Wai Yim for translations. CABI is an international intergovernmental organisation

and we gratefully acknowledge the core financial support from our member countries (and lead

agencies) including the United Kingdom (Department for International Development), China (Chinese

Ministry of Agriculture), Australia (Australian Centre for International Agricultural Research), Canada

(Agriculture and Agri-Food Canada), Netherlands (Directorate-General for International Cooperation),

and Switzerland (Swiss Agency for Development and Cooperation). See https://www.cabi.org/about-
cabi/who-we-work-with/key-donors/ for full details.
472

473 **Author contribution**

474 DB conducted the analyses and wrote the manuscript. EF and GH searched the Chinese literature. TH
475 assisted with CABI data acquisition. All authors contributed ideas and edited the manuscript.
476

477 **Data accessibility**

478 Pest distribution data are available with permission from CABI, Nosworthy Way, Wallingford, OX10
479 8DE, UK
480 Sources for other datasets used in the analysis are given in the text.
481
482

References

Avelino, J., Cristancho, M., Georgiou, S., Imbach, P., Aguilar, L., Bornemann, G., Läderach, P., Anzueto, F., Hruska, A.J. & Morales, C. (2015) The coffee rust crises in Colombia and Central America (2008–2013): impacts, plausible causes and proposed solutions. *Food Security*, **7**, 303–321.

Bacon, S.J., Aebi, A., Calanca, P. & Bacher, S. (2014) Quarantine arthropod invasions in Europe: the role of climate, hosts and propagule pressure. *Diversity and Distributions*, **20**, 84–94.

Baiser, B., Olden, J.D., Record, S., Lockwood, J.L. & McKinney, M.L. (2012) Pattern and process of biotic homogenization in the New Pangaea. *Proceedings of the Royal Society B: Biological Sciences*, **279**, 4772–4777.

Baker, R.H.A. (2012) An introduction to the PRATIQUE Research Project. *EPPO Bulletin*, **42**, 1–2.

Baker, R.H.A., Anderson, H., Bishop, S., MacLeod, A., Parkinson, N. & Tuffen, M.G. (2014) The UK Plant Health Risk Register: a tool for prioritizing actions. *EPPO Bulletin*, **44**, 187–194.

Bebber, D.P. (2015) Range-Expanding Pests and Pathogens in a Warming World. *Annual Review of Phytopathology*, **53**, 335–356.

Bebber, D.P., Castillo, Á.D. & Gurr, S.J. (2016) Modelling coffee leaf rust risk in Colombia with climate reanalysis data. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**, 20150458.

Bebber, D.P. & Gurr, S.J. (2015) Crop-destroying fungal and oomycete pathogens challenge food security. *Fungal Genetics and Biology*, **74**, 62–64.

Bebber, D.P., Holmes, T. & Gurr, S.J. (2014a) The global spread of crop pests and pathogens. *Global Ecology and Biogeography*, **23**, 1398–1407.

Bebber, D.P., Holmes, T., Smith, D. & Gurr, S.J. (2014b) Economic and physical determinants of the global distributions of crop pests and pathogens. *New Phytologist*, **202**, 901–910.

Bebber, D.P., Ramotowski, M.A.T. & Gurr, S.J. (2013) Crop pests and pathogens move polewards in a warming world. *Nature Climate Change*, **3**, 985–988.

Berzitis, E.A., Minigan, J.N., Hallett, R.H. & Newman, J.A. (2014) Climate and host plant availability impact the future distribution of the bean leaf beetle (Cerotoma trifurcata). *Global Change Biology*, **20**, 2778–2792.

Boakes, E.H., McGowan, P.J.K., Fuller, R.A., Chang-qing, D., Clark, N.E., O'Connor, K. & Mace, G.M. (2010) Distorted views of biodiversity: spatial and temporal bias in species occurrence data. *PLoS Biol*, **8**, e1000385.

Bregaglio, S., Cappelli, G. & Donatelli, M. (2012) Evaluating the suitability of a generic fungal infection model for pest risk assessment studies. *Ecological Modelling*, **247**, 58–63.

Chapman, D., Purse, B.V., Roy, H.E. & Bullock, J.M. (2017) Global trade networks determine the distribution of invasive non-native species. *Global Ecology and Biogeography*, **26**, 907–917.

Crooks, J.A. (2005) Lag times and exotic species: The ecology and management of biological invasions in slow-motion. *Ecoscience*, **12**, 316–329.

DEFRA (2018) UK Plant Health Risk Register.

Department of Agriculture, Fisheries and Forestry (2013) *Final pest risk analysis report for Drosophila suzukii*, Australian Government, Canberra, Australia.

Ding, J., Mack, R.N., Lu, P., Ren, M. & Huang, H. (2008) China's booming economy is sparking and accelerating biological invasions. *BioScience*, **58**, 317.

Dormann, C.F., Schymanski, S.J., Cabral, J., Chuine, I., Graham, C., Hartig, F., Kearney, M., Morin, X., Römermann, C., Schröder, B. & Singer, A. (2012) Correlation and process in species distribution models: bridging a dichotomy. *Journal of Biogeography*, **39**, 2119–2131.

Elith, J. & Leathwick, J.R. (2009) Species Distribution Models: Ecological Explanation and Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 677–697.

EPPO (2008) *Report of a Pest Risk Analysis for Xanthomonas axonopodis pv. allii*, European and Mediterranean Plant Protection Organization, Paris.

Eschen, R., Holmes, T., Smith, D., Roques, A., Santini, A. & Kenis, M. (2014) Likelihood of establishment of tree pests and diseases based on their worldwide occurrence as determined by hierarchical cluster analysis. *Forest Ecology and Management*, **315**, 103–111.

FAO (2004) *ISPM 21. Pest risk analysis for regulated non-quarantine pests*, IPPC, FAO, Rome.
Fears, R., Aro, E.-M., Pais, M.S. & ter Meulen, V. (2014) How should we tackle the global risks to
plant health? *Trends in Plant Science*, **19**, 206–208.
Fisher, M.C., Henk, D.A., Briggs, C.J., Brownstein, J.S., Madoff, L.C., McCraw, S.L. & Gurr, S.J.
(2012) Emerging fungal threats to animal, plant and ecosystem health. *Nature*, **484**, 186–194.
Flood, J. & Day, R. (2016) Managing risks from pests in global commodity networks – policy
perspectives. *Food Security*, **8**, 89–101.
Guo, L., Zhu, X.-Q., Hu, C.-H. & Ristaino, J.B. (2010) Genetic Structure of Phytophthora infestans
Populations in China Indicates Multiple Migration Events. *Phytopathology*, **100**, 997–1006.
Huang, D., Zhang, R., Kim, K.C. & Suarez, A.V. (2012) Spatial Pattern and Determinants of the First
Detection Locations of Invasive Alien Species in Mainland China. *Plos One*, **7**, e31734.
IMF (2013) IMF Data and Statistics.
Jeger, M., Bragard, C., Caffier, D., Candresse, T., Chatzivassiliou, E., Dehnen-Schmutz, K., Gilioli,
G., Grégoire, J.-C., Jaques Miret, J.A., MacLeod, A., Navarro, M.N., Niere, B., Parnell, S.,
Potting, R., Rafoss, T., Rossi, V., Urek, G., Van Der Werf, W., West, J., Winter, S., Gardi, C.,
Mosbach-Schulz, O., Koufakis, I. & Van Bruggen, A. (2017) Pest risk assessment of
Diaporthe vaccinii for the EU territory. *EFSA Journal*, **15**, e04924.
Jeger, M., Bragard, C., Caffier, D., Candresse, T., Chatzivassiliou, E., Dehnen-Schmutz, K., Grégoire,
J.-C., Jaques Miret, J.A., MacLeod, A., Navajas Navarro, M., Niere, B., Parnell, S., Potting,
R., Rafoss, T., Rossi, V., Urek, G., Van Bruggen, A., Van Der Werf, W., West, J., Winter, S.,
Hart, A., Schans, J., Schrader, G., Suffert, M., Kertész, V., Kozelska, S., Mannino, M.R.,
Mosbach-Schulz, O., Pautasso, M., Stancanelli, G., Tramontini, S., Vos, S. & Gilioli, G.
(2018) Guidance on quantitative pest risk assessment. *EFSA Journal*, **16**, e05350.
Jiménez-Valverde, A. (2012) Insights into the area under the receiver operating characteristic curve
(AUC) as a discrimination measure in species distribution modelling. *Global Ecology and
Biogeography*, **21**, 498–507.
Jones, K.E., Patel, N.G., Levy, M.A., Storeygard, A., Balk, D., Gittleman, J.L. & Daszak, P. (2008)
Global trends in emerging infectious diseases. *Nature*, **451**, 990–993.
Kent, S. (1994) Words of estimative probability. *Studies in Intelligence*, **8**, 49–65.
Klayman, D.L. (1985) Qinghaosu (Artemisinin): An Antimalarial Drug from China. *Science*, **228**,
1049–1055.
Kriticos, D.J. (2012) Regional climate-matching to estimate current and future sources of biosecurity
threats. *Biological Invasions*, **14**, 1533–1544.
Kriticos, D.J., Morin, L., Leriche, A., Anderson, R.C. & Caley, P. (2013) Combining a Climatic
Niche Model of an Invasive Fungus with Its Host Species Distributions to Identify Risks to
Natural Assets: Puccinia psidii Sensu Lato in Australia. *PLoS ONE*, **8**, e64479.
Liebhold, A.M., McCullough, D.G., Blackburn, L.M., Frankel, S.J., Von Holle, B. & Aukema, J.E.
(2013) A highly aggregated geographical distribution of forest pest invasions in the USA.
Diversity and Distributions, **19**, 1208–1216.
MacLeod, A., Jones, G.D., Anderson, H.M. & Mumford, R.A. (2016) Plant health and food security,
linking science, economics, policy and industry. *Food Security*, **8**, 17–25.
MacLeod, A., Pautasso, M., Jeger, M.J. & Haines-Young, R. (2010) Evolution of the international
regulation of plant pests and challenges for future plant health. *Food Security*, **2**, 49–70.
Monfreda, C., Ramankutty, N. & Foley, J.A. (2008) Farming the planet: 2. Geographic distribution of
crop areas, yields, physiological types, and net primary production in the year 2000. *Global
Biogeochemical Cycles*, **22**, GB1022.
Ordonez, N., Seidl, M.F., Waalwijk, C., Drenth, A., Kilian, A., Thomma, B.P.H.J., Ploetz, R.C. &
Kema, G.H.J. (2015) Worse Comes to Worst: Bananas and Panama Disease—When Plant
and Pathogen Clones Meet. *PLOS Pathogens*, **11**, e1005197.
Paini, D.R., Sheppard, A.W., Cook, D.C., Barro, P.J.D., Worner, S.P. & Thomas, M.B. (2016) Global
threat to agriculture from invasive species. *Proceedings of the National Academy of Sciences*,
113, 7575–7579.
Paini, D.R., Worner, S.P., Cook, D.C., De Barro, P.J. & Thomas, M.B. (2010) Threat of invasive
pests from within national borders. *Nature Communications*, **1**, 115.

Patpour, M., Hovmøller, M.S., Shahin, A.A., Newcomb, M., Olivera, P., Jin, Y., Luster, D., Hodson, D., Nazari, K. & Azab, M. (2015) First Report of the Ug99 Race Group of Wheat Stem Rust, Puccinia graminis f. sp. tritici, in Egypt in 2014. *Plant Disease*, **100**, 863–863.

Pyšek, P., Jarošík, V., Hulme, P.E., Kühn, I., Wild, J., Arianoutsou, M., Bacher, S., Chiron, F., Didžiulis, V., Essl, F., Genovesi, P., Gherardi, F., Hejda, M., Kark, S., Lambdon, P.W., Desprez-Loustau, M.-L., Nentwig, W., Pergl, J., Poboljšaj, K., Rabitsch, W., Roques, A., Roy, D.B., Shirley, S., Solarz, W., Vilà, M. & Winter, M. (2010) Disentangling the role of environmental and human pressures on biological invasions across Europe. *Proceedings of the National Academy of Sciences*, **107**, 12157–12162.

Pyšek, P., Richardson, D.M., Pergl, J., Jarošík, V., Sixtová, Z. & Weber, E. (2008) Geographical and taxonomic biases in invasion ecology. *Trends in Ecology & Evolution*, **23**, 237–244.

R Development Core Team (2017) *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna.

Robinet, C., Kehlenbeck, H., Kriticos, D.J., Baker, R.H.A., Battisti, A., Brunel, S., Dupin, M., Eyre, D., Faccoli, M., Ilieva, Z., Kenis, M., Knight, J., Reynaud, P., Yart, A. & van der Werf, W. (2012) A Suite of Models to Support the Quantitative Assessment of Spread in Pest Risk Analysis. *PLoS ONE*, **7**, e43366.

Santini, A., Ghelardini, L., De Pace, C., Desprez-Loustau, M.L., Capretti, P., Chandelier, A., Cech, T., Chira, D., Diamandis, S., Gaitniekis, T., Hantula, J., Holdenrieder, O., Jankovsky, L., Jung, T., Jurc, D., Kirisits, T., Kunca, A., Lygis, V., Malecka, M., Marcais, B., Schmitz, S., Schumacher, J., Solheim, H., Solla, A., Szabò, I., Tsopelas, P., Vannini, A., Vettraino, A.M., Webber, J., Woodward, S. & Stenlid, J. (2013) Biogeographical patterns and determinants of invasion by forest pathogens in Europe. *New Phytologist*, **197**, 238–250.

Savary, S., Bregaglio, S., Willocquet, L., Gustafson, D., D'Croz, D.M., Sparks, A., Castilla, N., Djurle, A., Allinne, C., Sharma, M., Rossi, V., Amorim, L., Bergamin, A., Yuen, J., Esker, P., McRoberts, N., Avelino, J., Duveiller, E., Koo, J. & Garrett, K. (2017) Crop health and its global impacts on the components of food security. *Food Security*, **9**, 311–327.

Scimago Lab (2017) Scimago Journal & Country Rank.

Shimwela, M.M., Narouei-Khandan, H.A., Halbert, S.E., Keremane, M.L., Minsavage, G.V., Timilsina, S., Massawe, D.P., Jones, J.B. & van Bruggen, A.H.C. (2016) First occurrence of Diaphorina citri in East Africa, characterization of the Ca. Liberibacter species causing huanglongbing (HLB) in Tanzania, and potential further spread of D. citri and HLB in Africa and Europe. *European Journal of Plant Pathology*, **146**, 349–368.

Skelsey, P., Cooke, D.E.L., Lynott, J.S. & Lees, A.K. (2016) Crop connectivity under climate change: future environmental and geographic risks of potato late blight in Scotland. *Global Change Biology*, n/a-n/a.

Soberón, J. (2007) Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters*, **10**, 1115–1123.

Soberón, J. & Nakamura, M. (2009) Niches and distributional areas: concepts, methods, and assumptions. *Proceedings of the National Academy of Sciences*, **106**, 19644–19650.

Soberón, J. & Peterson, A.T. (2005) Interpretation of models of fundamental ecological niches and species' distributional areas. *Biodiversity Informatics*, **2**, 1–10.

Sutherst, R.W. (2014) Pest species distribution modelling: origins and lessons from history. *Biological Invasions*, **16**, 239–256.

Venette, R.C., Kriticos, D.J., Magarey, R.D., Koch, F.H., Baker, R.H.A., Worner, S.P., Gómez Raboteaux, N.N., McKenney, D.W., Dobesberger, E.J., Yemshanov, D., De Barro, P.J., Hutchison, W.D., Fowler, G., Kalaris, T.M. & Pedlar, J. (2010) Pest Risk Maps for Invasive Alien Species: A Roadmap for Improvement. *BioScience*, **60**, 349–362.

Wangai, A.W., Redinbaugh, M.G., Kinyua, Z.M., Miano, D.W., Leley, P.K., Kasina, M., Mahuku, G., Scheets, K. & Jeffers, D. (2012) First report of maize chlorotic mottle virus and Maize Lethal Necrosis in Kenya. *Plant Disease*, **96**, 1582–1582.

Ward, M. (2016) Action against pest spread—the case for retrospective analysis with a focus on timing. *Food Security*, **8**, 77–81.

Westphal, M.I., Browne, M., MacKinnon, K. & Noble, I. (2008) The link between international trade and the global distribution of invasive alien species. *Biological Invasions*, **10**, 391–398.