**Chromatin modifiers alter recombination between divergent DNA sequences**

Ujani Chakraborty, Beata Mackenroth, and Eric Alani

Department of Molecular Biology and Genetics, Cornell University, Ithaca, New York, United States of America

Key words: heteroduplex rejection, homologous recombination, mismatch repair, histone chaperones, chromatin modifiers

Corresponding Author:

Eric Alani, Ph.D.

Cornell University Department of Molecular Biology and Genetics

459 Biotechnology Building

Ithaca, NY 14853-2703, United States of America

Email: eea3@cornell.edu

Tel: 607-254-4811

Running title: Chromatin modifiers alter anti-recombination

**Abstract**

Recombination between divergent DNA sequences is actively prevented by heteroduplex rejection mechanisms. In baker's yeast such anti-recombination mechanisms can be initiated by the recognition of DNA mismatches in heteroduplex DNA by MSH proteins, followed by recruitment of the Sgs1-Top3-Rmi1 helicase-topoisomerase complex to unwind the recombination intermediate. We previously showed that the repair/rejection decision during single-strand annealing recombination is temporally regulated by MSH protein levels and by factors that excise non-homologous single-stranded tails. These observations, coupled with recent studies indicating that mismatch repair factors interact with components of the histone chaperone machinery, encouraged us to explore roles for epigenetic factors and chromatin conformation in regulating the decision to reject vs. repair recombination between divergent DNA substrates. This work involved the use of an inverted repeat recombination assay thought to measure sister chromatid repair during DNA replication. Our observations are consistent with the histone chaperones CAF-1 and Rtt106 and the histone deacetylase Sir2 acting to suppress heteroduplex rejection and the Rpd3, Hst3 and Hst4 deacetylases acting to promote heteroduplex rejection. These observations and double mutant analysis have led to a model in which nucleosomes located at DNA lesions stabilize recombination intermediates and compete with mismatch repair factors that mediate heteroduplex rejection.

**Introduction**

Homologous recombination (HR) is a conservative DNA repair pathway that is critical for repairing DNA double strand breaks (DSBs).  This process is regulated to prevent recombination between divergent DNA sequences (reviewed in George and Alani 2012). Such regulation, which can prevent deleterious chromosomal rearrangements, is initiated through the recognition of mismatches in heteroduplex DNA that forms during strand invasion steps involving single-stranded DNA from a broken chromosome and a divergent duplex donor.  In the yeast *Saccharomyces cerevisiae*, the Msh2-Msh6 and Msh2-Msh3 mismatch repair (MMR) complexes recognize mismatches in heteroduplex DNA and recruit the RecQ family helicase-topoisomerase complex Sgs1-Top3-Rmi1 to unwind the recombination intermediate in a process known as heteroduplex rejection (Datta *et al*. 1996; Chen and Jinks-Robertson 1999; Nicholson *et al*. 2000; Myung *et al*. 2001; Spell and Jinks-Robertson 2004a; Sugawara *et al*. 2004; Goldfarb and Alani 2005; Chakraborty *et al*. 2016). If rejection does not occur, MSH complexes initiate repair of the mismatches in heteroduplex DNA after/while the break is repaired.  We refer to this regulation as the repair/rejection decision.

An important aspect of the repair/rejection decision is that tolerating multiple mismatches in heteroduplex DNA formed from divergent substrates can lead to chromosomal rearrangements, but a highly efficient rejection system can result in low repair efficiency of DNA breaks.  Various factors are likely to influence this decision (reviewed in Chakraborty and Alani 2016). For example, we found that MSH protein levels influence the repair/rejection decision during single strand annealing (SSA; Chakraborty *et al*. 2016).  During SSA, HR is initiated by a DSB located between two closely spaced repeat sequences.  Resection of the DSB promotes annealing of homologous sequences, followed by the clipping of 3' non-homologous tails that must be excised before repair steps is completed.  During SSA involving divergent repeat sequences, modest overexpression of Msh6 resulted in a significant increase in heteroduplex rejection due to a decrease in the availability of Msh2-Msh3 to clip the 3' tails. Thus 3' tail clipping during SSA serves as a regulatory step, with rejection favored prior to 3' tail removal.

Consistent with these findings, Anand *et al.* (2017) showed in a break induced repair recombination system that 3' non-homologous tails promote heteroduplex rejection, and the absence of such tails prevents it. These observations indicate a crosstalk between the rejection machinery and the local environment that likely occurs prior to steps in HR that involve repair DNA synthesis.

Recent studies have indicated that chromatin structure can influence HR outcomes. Several nucleosome remodeling complexes have been shown to be recruited to DSBs in steps thought to increase chromatin accessibility and allow strand resection and presynaptic filament formation (reviewed in Hauer and Gasser 2017). Such chromatin remodelers were shown to promote chromatin mobility during DSB formation, and the increased mobility correlated to more efficient repair by HR (Dion *et al*. 2012; Mine-Hattab and Rothstein 2012; Neumann *et al*. 2012; Hauer *et al.* 2017). Also, histone chaperones, which act in DNA replication-dependent nucleosome assembly, have been implicated in DNA repair. These chaperones include CAF-1, Asf1 and Rtt106, which are all involved in DNA replication-dependent nucleosome assembly (Tyler *et al.* 1999; Tagami *et al*. 2004; Huang *et al*. 2005). For example, nucleosome assembly mediated by CAF-1 during DNA repair is coupled with DNA synthesis and requires an interaction with PCNA (Gaillard *et al*. 1996; Tyler *et al*. 1999; Linger and Tyler 2005; Polo, Roche and Almouzni 2006; Pietrobon *et al*. 2014), and CAF-1 and Asf1 play roles in restoring chromatin after DNA repair in budding yeast at the repair sites by turning off the DNA damage checkpoint (Chen *et al*. 2008; Kim and Haber 2009; Diao *et al.* 2017).

Histone modifications have also been shown to affect genome stability. For example, deacetylation of an acetylated lysine residue at amino acid 56 in Histone 3 (H3K56) by Hst3 and Hst4 is required for the suppression of mutations and gross chromosomal rearrangements, and acetylation of H3K56 by Rtt109 is also required for suppression of mutations (Kadyrova *et al*. 2013). In contrast, histone deacetylases such as Rpd3L and Hda1 promote trinucleotide repeat expansions associated with various neurodegenerative diseases (Debacker *et al*. 2012). Thus, histone acetylation and deacetylation are likely to play important roles in various stages of HR.

In support of this, Tamburini and Tyler (2005) showed that histone acetyltransferases such as Gcn5 and Esa1 are recruited to an HO endonuclease induced DSB in *S. cerevisiae* followed at a later stage by recruitment of histone deacetylases such as Sir2, Hst1 and Rpd3. Furthermore, they showed that mutating acetylable lysine residues in Histone subunit 4, or deleting *GCN5* or *RPD3* caused inviability in response to HO endonuclease induced lesions repaired primarily by HR. These observations suggest that after DSB formation, histone acetylases participate in nucleosome removal in the vicinity of the break and "open up" the chromatin structure increasing the accessibility of the underlying DNA to repair factors. Once DNA repair is underway, histone deacetylases likely modify the chromatin to a "closed" conformation that serves as a commitment step to complete the repair process and restore chromatin to its original state.

The NAD-dependent histone deacetylase Sir2, known for its role in transcriptional silencing and heterochromatin formation, also plays direct roles in forming a repressive local chromatin environment around most euchromatic replication origins (Gartenberg and Smith 2016; Hoggard *et al.* 2018).  Sir2 is also involved in DNA repair pathways such as non-homologous end joining and nucleotide excision repair (Tsukamoto 1997; Boulton and Jackson 1998; Guintini *et al.* 2017).  *sir2* mutants are hypersensitive to DNA damaging agents, and a number of studies have also reported that in response to DNA DSBs, a significant fraction of the histone bound SIR complex was displaced from subtelomeric regions and relocated to sites of DSBs in a DNA checkpoint dependent manner, suggesting that the recruitment of SIR complexes reflects the assembly of a repressed chromatin state following DNA repair (Martin *et al.* 1999; McAinsh *et al.* 1999; Mills *et al.* 1999).

A number of studies have shown that nucleosome assembly on newly synthesized DNA during replication and MMR are mutually inhibitory processes. MMR during DNA replication is thought to be restricted to the short time window between the formation of the mismatch and the chaperone-assisted assembly of nucleosomes on the newly replicated DNA (Li *et al.* 2009; Kadyrova, Blanko and Kadyrov 2011; Schopf *et al.* 2012; Blanko, Kadyrova and Kadyrov 2016).

The human MSH2-MSH6 complex has been shown to interact with CAF-1 *in vitro* (Schopf *et al.* 2012). Additionally, human MSH2-MSH6 inhibits CAF-1 and ASF1A-dependent packaging of a DNA mismatch into a nucleosome, and deposition of the $(H3-H4)_2$ tetramers on DNA protects the discontinuous daughter strand from unnecessary degradation during MMR (Blanko, Kadyrova and Kadyrov 2016). Moreover, Pietrobon *et al.* (2014) showed in fission yeast that CAF-1 stabilizes D-loops in an HR pathway, by counteracting their disassembly mediated by the RecQ family helicase Rqh1 when cells replicate a damaged template.

How is the balance between heteroduplex rejection and homologous recombination repair maintained? We tested in baker's yeast roles for histone acetylases, deacetylases, and chaperones in regulating the repair/rejection decision. To our knowledge, this is the first study to look at possible roles of epigenetic factors and chromatin modifiers in modulating heteroduplex rejection. We show that histone chaperones CAF-1 and Rtt106 suppress heteroduplex rejection in steps dependent on mismatch recognition. Additionally, the histone deacetylase Sir2 appears to act in a common pathway with CAF-1 and/or Rtt106 to suppress rejection. However, other factors involved in nucleosome assembly during DNA replication such as Asf1 and Rtt109 do not affect rejection efficiency. Similarly, histone acetylases such as Gcn5 that assemble early at recombination sites, and other histone deacetylases such as Hst1 do not affect the efficiency of rejection. However, mutants lacking the Hst3, Hst4, or Rpd3 histone deacetylases show defects in rejection. Taken together, these results are consistent with the idea that nucleosomes at DNA lesions, which are likely to be localized independently of DNA synthesis, stabilize recombination intermediates and thus prevent access by anti-recombination factors.

**Materials and Methods**

**Yeast strains and plasmids.** Yeast strains used in this study are listed in Table 1 and were constructed and grown using standard techniques (Rose *et al.* 1990). Gene disruptions used to make the strains are described in Tables 1-3 (*geneXΔ::KANMX*) and were obtained by PCR amplification (details provided upon request) of chromosomal DNA derived from the yeast

knockout collection (Brachmann *et al.* 1998). These DNA fragments and the *pol30-8* integrating

*2μ* vector described below were introduced into yeast strains using standard transformation

procedures (Gietz and Schiestl 1991).  The presence of mutant alleles was confirmed by PCR

analysis of chromosomal DNA, and in some cases DNA sequencing of the amplified DNA

fragments. The mutations comprising the *pol30-8 allele* (R61A, D63A) were introduced into

pEAA578 (*POL30::KANMX*) by Q5 site-directed mutagenesis (New England Biolabs) to create

the single step integrating vector pEAI438 (*pol30-8::KANMX*). Plasmids pRS426 (*2μ, URA3*;

Christianson *et al.* 1992) and pEAM272 (*MSH2, MSH6, 2μ, URA3;* Chakraborty *et al.* 2018)

were used in the MSH overexpression experiments presented in Table 3.

**Inverted repeat recombination assay.** Strains used to measure homologous and

homeologous recombination are listed in Table 1. Strains lacking plasmids were initially struck

onto synthetic complete plates and those containing *2μ* plasmids were struck onto minimal

dropout media plates (Rose *et al.* 1990). A total of 10-116 single colonies per strain were then

inoculated into 5 ml of synthetic complete or minimal dropout medium containing 4% galactose

and 2% glycerol and grown to saturation for ~2 days at 30°C. Appropriate dilutions of cells were

plated onto minimal media (2% galactose, 2% glycerol) plates lacking histidine and the amino

acid required to maintain the *2μ* plasmid (selective) and onto minimal media (2% glucose) plates

lacking the amino acid required to maintain the *2μ* plasmid (permissive). Plates were incubated

for 4 days at 30°C (5 days for experiments involving *rpd3Δ*) and then scored for frequency of

His[+] colonies. Rates of homologous and homeologous recombination were calculated as

described (Nicholson *et al.* 2000; Spell and Robertson 2004b). Pairwise Mann–Whitney U tests

were performed between mutant and corresponding wild-type of each strain. Differences were

considered significant when $p < 0.05$.

**Repetition of experiments.**  The inverted repeat recombination assays were repeated on 2-4 days, with roughly an equal number of repetitions per day, and 2-4 independent transformants were analyzed for each genotype.

**Data availability.** Strains and plasmids are available upon request.

**Results**

**The absence of histone chaperones CAF-1 and Rtt106, but not Asf1 and Rtt109, improves anti-recombination.**  As summarized above, CAF-1 was found to stabilize D-loops in *S. pombe* that occur as the result of template switching at replication forks (Pietrobon *et al.* 2014). This observation encouraged us to test if the histone chaperones CAF-1 and Rtt106 play roles in stabilizing recombination intermediates during HR, and in turn suppressing rejection. We also tested roles for other histone chaperones and chromatin modifying enzymes associated with DNA replication.  These include Asf1, which binds to newly synthesized H3/H4 dimers that are then acetylated at H3K56 by Rtt109 (Tyler *et al.* 1999; Driscoll *et al.* 2007; Han *et al.* 2007), and Rtt109, which mediates histone modification that promotes binding of H3 to the histone chaperones CAF-1 and Rtt106, and subsequently, the binding of CAF-1 to PCNA, facilitating histone deposition near the replication fork (Li *et al.* 2008).  We were encouraged to study the above factors because much is known about the interactions and crosstalk between the replication apparatus and nucleosome assembly machinery. For example, CAF-1 interacts with the replication processivity clamp PCNA and Asf1 interacts with RFC, which loads PCNA, and the MCM helicase (Groth *et al.* 2007; Franco *et al.* 2005). In addition to their role in nucleosome assembly, Asf1 and Rtt109 indirectly promote nucleosome disassembly through H3K56 acetylation (Adkins *et al.* 2004; Adkins and Tyler 2004; Schwabish and Struhl 2006; Korber *et al.* 2006).

We used an inverted repeat recombination assay in baker's yeast to study roles for chromatin modifiers in regulating the repair/rejection decision (Nicholson *et al.* 2000; Figure 1A).

In this assay, spontaneous recombination events reorient *HIS3* and intron sequences to yield a functional *HIS3* gene. Such events, which are thought to be initiated by DNA lesions that occur during or shortly after the replication of the recombination substrates, are consistent with repair through sister chromatid gene conversion (Chen and Jinks-Robertson 1998). We calculated the ratio of recombination rates in strains containing identical DNA substrates to those seen in strains containing divergent DNA substrates (creating four single nucleotide mismatches or four 4-nucleotide loop mismatches in a 350 bp substrate; Table 1) to determine the efficiency of heteroduplex rejection.  In wild-type strains this ratio is 15 to 18 for single nucleotide mismatches; in MMR defective strains the ratio is ~1, indicating that homologous and divergent recombination rates are similar in MMR defective mutants (Table 2; Figure 1B Chakraborty et al., 2018).

The *cac1Δ* mutation (deletion of the large subunit of CAF-1) increased rejection efficiency by ~3-fold, compared to wild-type, in recombination assays involving base-base or 4-nt loop mismatches (Table 2; Figure 1B). The *rtt106Δ* mutation conferred a 2.2-fold increase in rejection efficiency as measured in the base-base mismatch recombination assay (Table 2; Figure 1B).  *msh2Δ cac1Δ* and *msh2Δ rtt106Δ* mutants showed rejection ratios comparable to *msh2Δ* strains (Table 2; Figure 1B), indicating that the increases in rejection in *cac1Δ* and *rtt106Δ* strains were dependent on the rejection machinery.  Interestingly, the *cac1Δ rtt106Δ* double mutant showed a rejection ratio significantly higher (5.9-fold increase compared to wild-type) than the single mutants, suggesting a redundant function for these two factors analogous to redundant roles for these proteins in DNA replication (Table 2; Figure 1B; Li *et al.* 2008). Together, these observations suggested that deposition of histones by CAF-1 and Rtt106 during double-strand break repair stabilized recombination intermediates, making them less accessible to heteroduplex rejection factors.

In contrast, deleting Asf1 or Rtt109, factors thought to act upstream of CAF-1 and Rtt106, did not confer an effect on heteroduplex rejection (Table 2; Figure 1B). These factors are required to acetylate Histone H3 at K56, and the resulting H3K56ac-H4 dimers are then

transferred to histone chaperones such as CAF-1 and Rtt106 (Schneider *et al.* 2006; Han *et al.* 2007; Driscoll *et al.* 2007; Tsubota *et al.* 2007). Furthermore, the *asf1Δ* mutation suppressed the increase in rejection seen in *cac1Δ* or *rtt106Δ* mutants (Table 2; Figure 1B). Additionally, we performed recombination assays in 4-nt loop mismatch strains and found that the *asf1Δ* mutation had minimal if any effect on rejection, and that it suppressed the increased rejection of *cac1Δ* strains, in agreement with observations from base-base mismatch strains (Table 2).

   Previous studies showed that mutants defective in the CAF-1 complex are sensitive to DNA damaging agents, and have suggested that during DNA repair CAF-1 is recruited by the replication processivity clamp PCNA to facilitate DNA synthesis repair steps (Moggs *et al.* 2000; Linger and Tyler 2005). The Stillman group identified in baker's yeast an allele of *POL30* (gene encoding PCNA; *pol30-8* (R61A, D63A)) that displayed reduced binding to CAF-1 and compromised recruitment of CAF-1 to replicating DNA (Shibahara and Stillman 1999; Zhang *et al.* 2000), and Linger and Tyler (2005) showed that *pol30-8* mutant yeasts were similarly sensitive to DNA damaging agents as *cac2Δ* (subunit of the heterotrimeric CAF-1 complex) and *cac2Δ pol30-8* mutants. As shown in Table 2 and Figure 1B, the *pol30-8* allele, which conferred sensitivity to MMS, increased heteroduplex rejection by 1.6-fold.

**Deletion of the Sir2 silencing factor increases anti-recombination.** The silencing factors Sir2 and Sir3 are recruited by the histone chaperones CAF-1 and Rtt106 to sites of heterochromatin formation. In the absence of CAF-1 and Rtt106, Sir proteins are mis-localized (Huang *et al.* 2007). These observations encouraged us to test if the histone deacetylases Hst1, Hst3, Hst4 or Sir2 are involved in the regulation of rejection efficiency. We were further encouraged to test these factors because Tamburini and Tyler (2005) showed in *S. cerevisiae* that the histone acetylases Gcn5 and Esa1 (essential for viability) and the histone deacetylases Rpd3, Sir2 and Hst1 were recruited to an HO endonuclease induced lesion during HR. We were also interested in Hst3 and Hst4, which act to remove H3K56ac marks from newly generated chromatin in G2/M, because studies from Munoz-Galvan *et al.* (2013) suggested that

acetylation and deacetylation of H3K56 were important for selecting the sister chromatid as a template for repair of DSBs that occur during DNA replication.  As shown in Figure 1B and Table 2, the *sir2Δ* mutation increased the efficiency of heteroduplex rejection by 1.9- and 1.7-fold in the nucleotide substitution and 4-nt loop strains, respectively. Deletion of *HST3* or *HST4* conferred mild effects on the rejection ratio (1.3- and 1.6-fold lower than wild-type respectively). However, the *hst3Δ hst4Δ* double mutant showed a 4.0-fold *reduction* in anti-recombination of base-base mismatches compared to wild-type strains.

The increased heteroduplex rejection phenotype observed in *sir2Δ* strains and the opposite phenotype seen in *hst3Δ hst4Δ* strains encouraged us to test if deletion mutations in the histone deacetylase Hst1, the acetylase Gcn5, and Rpd3, a modifier of Sir2 function, affected rejection.  As shown in Table 2 and Figure 1B, *hst1Δ* and *gcn5Δ* mutations did not confer an effect.  We then tested a strain deleted for Rpd3, which encodes a subunit of both the Rdp3S and Rpd3L histone deacetylase complexes that act in chromatin remodeling.  Several studies reported that Rpd3 and Sir2 have antagonistic effects on silent chromatin propagation and replication timing (Zhou *et al*. 2009; Thurtle-Schmidt *et al*. 2016; Yoshida *et al*. 2014; Ehrentraut *et al*. 2010). As shown in Table 2 and Figure 1B, the *rpd3Δ* mutation decreased rejection 2.3-fold in the nucleotide substitution strain. Together these observations provide evidence that chromatin modifiers can suppress and enhance heteroduplex rejection (see Discussion).

**SIR2, CAC1, and RTT106 appear to function in the same anti-recombination pathway.**
To determine if CAF-1 and Rtt106 function in the same pathway as Sir2 for preventing heteroduplex rejection, we examine the phenotype of double mutants*. cac1Δ sir2Δ*  and *rtt106Δ sir2Δ* mutant strains exhibited rejection ratios similar to those of single mutants, suggesting epistasis (Table 2; Figure 1B). However, the rejection ratio dropped in *sir2Δ cac1Δ rtt106Δ* mutants compared to any of the double mutant combinations, indicating a more complex genetic interaction (see below).

Lastly, we asked if the increased rejection seen in *sir2Δ* strains was epistatic to that seen in strains bearing pEAM272 (*2μ, MSH2, MSH6* plasmid).  In wild-type strains bearing pEAM272 the Msh2-Msh6 complex is overexpressed by ~8-fold (Chakraborty *et al*. 2018). Wild-type strains containing pEAM272 display a 3.7-fold increase, compared to those containing the dummy vector pRS426, in the rejection ratio, consistent with an increased concentration of Msh2-Msh6 resulting in improved rejection by increasing the likelihood of mismatch recognition in heteroduplex DNA.  As shown in Table 3, *sir2Δ* strains containing pEAM272 display an even higher rejection ratio (1.9-fold higher than *sir2Δ* with pRS426, 6.1-fold higher than wild-type with pRS426), suggesting that the *SIR2* effect on anti-recombination is not epistatic to that seen in the wild-type strain containing pEAM272. One interpretation of this observation is that the Sir2 effect on rejection occurs in steps that compete with mismatch recognition. Curiously, *rtt106Δ* strains containing pEAM272 showed a rejection ratio that was similar to the wild-type lacking pEAM272 (Table 3), indicating a more complex phenotype reminiscent of the decreased rejection ratio seen in *sir2Δ cac1Δ rtt106Δ* triple mutants.

**Altered recombination rates in chromatin remodeling mutants do not correlate with their heteroduplex rejection phenotypes.**  *msh2Δ* mutants, which are defective in heteroduplex rejection, display elevated levels of homologous recombination (Table 2; Datta *et al.* 1997; Nicholson *et al*. 2000). Previously the Jinks-Robertson group hypothesized that the increased homologous recombination seen in *msh2Δ* reflects the fact that the length of perfect homology required to avoid heteroduplex rejection (610 bp) is larger than the length of the 350 bp repeats present in the inverted repeat substrate (Datta *et al.* 1997).  As shown in Table 2, the rates of recombination between homologous sequences in the mutants analyzed in this study vary but do not appear to correlate to changes in their repair/rejection ratios. *asf1Δ* and *rtt109Δ* mutants show recombination levels higher than *msh2Δ*, but a homologous/divergence recombination ratio similar to wild-type.  In contrast, *cac1Δ* and *rtt106Δ*, which show homologous recombination levels between wild-type and *msh2Δ*, and *sir2Δ*, which shows recombination

levels similar to *msh2Δ*, displayed homologous/divergence recombination ratios higher than wild-type.  A similar lack of correlation was seen for the double mutant combinations presented in Table 2 and Figure 1B.  This information suggests that overall levels of homologous recombination do not impact the rejection ratio.  One possible exception is the *hst3Δ hst4Δ* double mutant, which shows very high rates of homologous recombination and is compromised for heteroduplex rejection (see Discussion). A challenge in this analysis is that the inverted repeat assay detects spontaneous recombination events, and thus it is difficult to precisely determine if the changes in homologous recombination seen in chromatin modification mutants is due to increased DNA damage or other mechanisms.

**Discussion**

We focused on understanding roles for chromatin structure and modifications in regulating the heteroduplex rejection/DNA repair decision.  Improving repair at the cost of reduced fidelity can lead to gene conversion, chromosomal rearrangement and loss of heterozygosity, whereas high fidelity can compromise repair efficiency (reviewed in Chakraborty and Alani 2016).  We found that the histone chaperones CAF-1 and Rtt106 act to suppress heteroduplex rejection in processes that involve the deacetylase Sir2. In contrast, a large set of factors involved in nucleosome assembly during DNA replication or in modifying histones (Asf1, Rtt109, Hst1, Gcn5*)* do not affect rejection efficiency. These results are consistent with a pathway in which the presence of nucleosomes at DNA lesions acts to stabilize recombination intermediates and inhibit anti-recombination (Figure 2).

The model presented in Figure 2 outlines a recombination event involving divergent sequences initiated by a DSB.  DNA mismatches in heteroduplex DNA that form during strand invasion are recognized by MMR proteins which in turn recruit the Sgs1-Top3-Rmi1 helicase-topoisomerase complex to unwind and reject the recombination intermediate.  In this model, heteroduplex rejection is repressed by the presence of nucleosomes in a process regulated by CAF-1 and Rtt106, representing a regulatory step in the repair/rejection decision.  In this model

maintenance of nucleosomes by the histone chaperones CAF-1 and Rtt106 acts to stabilize heteroduplex DNA. This is followed by Sir2 deacetylating the newly deposited histones and further creating a tightly packed "closed" chromatin structure that prevents rejection and promotes repair through DNA synthesis steps.

The model presented in Figure 2 is supported by the following observations: 1. Deleting *CAC1* or *RTT106* resulted in increased heteroduplex rejection in strains containing both base-base and 4-nt loop mismatches. Furthermore, *cac1Δ rtt106Δ* strains showed an even higher increase in heteroduplex rejection, consistent with CAF-1 and Rtt106 acting redundantly in nucleosome assembly (Table 2; Li *et al.* 2008).  2. A number of studies have shown that MMR and nucleosome assembly during replication are mutually inhibitory processes and that human MutSα interacts physically with CAF-1 (Li *et al.* 2009; Kadyrova, Blanko and Kadyrov 2011; Schopf *et al.* 2012; Blanko, Kadyrova and Kadyrov 2016). Thus, in a manner analogous to their interaction during post-replicative MMR, nucleosome deposition could suppress MMR mediated heteroduplex rejection, forcing the latter process to occur in the time window after strand invasion and before nucleosome deposition. 3. Various studies have shown in fission yeast and humans that RecQ helicases (Rqh1 in *S. pombe* and BLM and WRN in humans) physically interact with the large subunit of CAF-1 (Pietrobon *et al.* 2014; Jiao *et al.* 2004; Jiao *et al.* 2007). Additionally, Pietrobon *et al.* (2014) showed that CAF-1 suppresses D-loop disassembly by Rqh1 during template switching. Based on these observations, it is possible that CAF-1 physically interacts with Sgs1 in budding yeast and counteracts its unwinding activity during heteroduplex rejection. 4. The removal of 3' non-homologous tails during SSA acts as a temporal switch; rejection is favored before tail removal, prior to DNA synthesis steps, and repair, after (Chakraborty *et al.* 2016).  Thus, analogous to 3' non-homologous tail removal during SSA, nucleosome maintenance by CAF-1 and Rtt106, followed by nucleosome deacetylation by Sir2 during HR could provide another type of temporal commitment step that regulates the rejection vs repair decision. 5. The *sir2Δ* mutation improved rejection efficiency and *sir2Δ cac1Δ* and *sir2Δ rtt106Δ* strains showed rejection efficiencies similar to single

mutants, suggesting that these factors act in common pathways (Table 2).  In support of this idea, Tamburini and Tyler (2005) showed that Sir2 localizes to sites of DSBs after histone acetylases such as Gcn5 and Esa1, and Hoggard *et al*. (2018) show that Sir2 functions to repress replicative origins at euchromatic regions. These data suggest that Sir2 is likely to localize to sites of HR at later stages of repair, likely after the deposition of nucleosomes, in order to modify histones by deacetylating them and further compacting the repair substrates making it harder for the rejection machinery to act. Together, they suggest that CAF-1 and Rtt106 function redundantly to deposit nucleosomes on recombination intermediates that stabilize the DNA heteroduplex and suppress rejection.

**asf1Δ mutation suppresses the hyper-rejection phenotype seen in *cac1Δ* and *rtt106Δ* strains.**  Curiously, the *asf1Δ* and *rtt109Δ* mutations did not alter heteroduplex rejection ratios, and *asf1Δ* suppressed the increased ratio seen in *cac1Δ* and *rtt106Δ* strains (both base-base and 4-nt loop mismatch substrates). This was surprising, given that Asf1 and Rtt109 act upstream of CAF-1 and Rtt106 in the nucleosome deposition pathway. Asf1 and Rtt109 have been implicated in nucleosome removal during DNA replication and transcriptional activation (Groth *et al*. 2007; Ransom *et al*. 2010; Adkins *et al*. 2004; Adkins and Tyler 2004; Schwabish and Struhl 2006; Korber *et al.* 2006).  Such a nucleosome removal activity could also act during DNA recombination, and in its absence, lead to chromatin acting to stabilize strand invasion intermediates that are refractory to heteroduplex rejection.  In this model, a lack of, or delay in, nucleosome deposition in *cac1Δ* or *rtt106Δ* mutants resulting in increased rejection could be compensated for by the lack of nucleosome removal in *asf1Δ* mutants. Such a scenario is supported by recent *in vivo* and *in vitro* findings indicating that histones are present at ssDNA (Adkins *et al*. 2017; Huang *et al*. 2018). Adkins *et al*. (2017) also showed *in vitro* that histones remain bound to ssDNA as resection proceeds for longer distances.

**Does CAF-1 regulation of heteroduplex rejection depend on its interaction with PCNA?**

CAF-1 has been implicated in DNA synthesis coupled nucleosome deposition via its interaction with PCNA (Shibahara and Stillman 1999; Zhang *et al*. 2000; Krawitz *et al*. 2002).  We found that strains bearing the *pol30-8* allele, which significantly weakens the PCNA-CAF-1 interaction, showed heteroduplex rejection ratios higher than wild-type but lower than *cac1Δ* (Figure 1B). One explanation for this phenotype is that CAF-1 regulation of heteroduplex rejection partially depends on its interaction with PCNA.  Alternatively, CAF1-PCNA interactions are required but the *pol30-8* allele does not fully disrupt CAF1-PCNA interactions *in vivo*.  In support of the former possibility, Hoek *et al*. (2011) found that a mutation in human CAF-1 that causes a defect in PCNA interactions (N-terminal truncation of the p150 subunit of CAF-1) does not affect the recruitment of CAF-1 to sites of DNA damage or confer sensitivity to DNA damaging agents. They also showed direct interactions between CAF-1 and the KU complex and 14-3-3 proteins, both of which are involved in DNA damage responses.  Additionally, CAF-1 has been shown to interact with the RecQ family BLM helicase which inhibits its activity *in vitro*, and the WRN helicase, also a RecQ family helicase, interacts with CAF-1 in a DNA damage-dependent manner (Jiao *et al*. 2004; Jiao *et al*. 2007). Pietrobon *et al*. (2014) also showed in fission yeast that CAF-1 promotes template switching at the replication fork by counteracting D-loop disassembly by Rqh1, a RecQ family helicase. Finally, Huang *et al*. (2018) have suggested that CAF-1 along with ASF1 are involved in chromatin assembly on ssDNA prior to Rad51 nucleo-protein filament formation, implying that it occurs in a PCNA independent manner, as PCNA might be expected to localize much later during DNA synthesis steps.

**MMR factors are likely to associate with sites of recombination early in the process.** Our data suggest that the deposition of nucleosomes and their subsequent modification are likely to suppress rejection and limit the time window during which the rejection machinery can act to unwind recombination intermediates involving divergent substrates. Previously we showed that during SSA, removal of 3' non-homologous tails serves a similar role with respect to providing a

limited time frame for rejection to occur (Chakraborty *et al.* 2016).  Several studies also support

this idea.  For example, Li *et al.* (2013) showed in mammalian cells that MSH2-MSH6 is

recruited via a PWWP motif in MSH6 to H3K36me3 histones before or during early S-phase. In

the context of rejection, this would ensure that mismatch recognition proteins are localized to

chromatin before replication fork stalling related recombination events occur. Although yeast

Msh6 and human MSH3 proteins do not have PWWP motifs, one can imagine that these

proteins may be recruited by other histone marks to chromatin. Additionally, MSH proteins have

been shown to localize rapidly to DSBs, even in the absence of a donor template, and act in the

DNA damage response (Evans *et al.* 2000; Lyndaker *et al.* 2008; Hong *et al.* 2008; Burdova *et*

*al.* 2015). Interestingly, overexpression of Msh2 and Msh6 increased the rejection ratio further in

*sir2Δ* strains, suggesting that mismatch recognition effects on rejection are distinct and are likely

to precede changes in chromatin structure (Table 3).  Taken together, these data suggest that

mismatch recognition proteins localize to sites of recombination either before DSBs are formed

or soon afterwards, and that chromatin could alter this localization and thus impact heteroduplex

rejection.

**Why do *rpd3Δ* strains display defects in heteroduplex rejection?**  A large-scale screen for

mutants defective in trinucleotide repeat instability identified a mutation in Sin3, which is a

subunit of the histone deacetylases Rpd3L and Rpd3S (Debacker *et al.* 2012).  In their analysis,

*sin3* mutants displayed significant reductions (9 to 18-fold) in expansion rates for a trinucleotide

repeat reporter.  Trinucleotide repeat expansions are also suppressed by disruption of factors

involved in heteroduplex rejection such as Msh2 and Msh3; such MSH factors are hypothesized

to promote trinucleotide repeat expansions by stabilizing slipped strand intermediates or through

altered MMR (McMurray 2010).  Based on these observations, the defect in heteroduplex

rejection observed in *rpd3Δ* strains could reflect the down-regulation of MSH factors that act in

mismatch recognition. Alternatively, the *rpd3Δ* phenotype results from improved localization of

Sir2, which acts to suppress heteroduplex rejection, to an initiating DSB site, thus lowering

rejection. The latter explanation fits with observations obtained from Zhou *et al*. (2009), who showed that in the absence of Rpd3, increased localization was seen for Sir2 at telomeres and homolgous mating type loci (HM) leading to an extension of silent chromatin in these areas.

**Can the heteroduplex rejection machinery be saturated?**  As shown in Table 2 and Figure 1B, the *hst3Δ hst4Δ* double mutant appears compromised for anti-recombination and displays very high rates of homologous recombination.  Kadyrova *et al.* (2013) showed that *hst3Δ hst4Δ* mutants display very high mutation rates that result from base substitutions, 1-bp insertions/deletions, and spontaneous gross chromosomal rearrangements. The rate of mutations in *hst3Δ hst4Δ*, as measured in forward mutation and reversion assays, was similar to that seen in MMR defective strains, and *msh2Δ hst3Δ hst4Δ* triple mutants displayed greater than additive mutation rates compared to *msh2Δ* and *hst3Δ hst4Δ*.  Based on these and other observations they proposed that Hst3 and Hst4 participate in genetic stability mechanisms that work with mismatch repair and replicative polymerase proofreading mechanisms to suppress spontaneous mutagenesis.  In this framework, the decreased rejection seen in *hst3Δ hst4Δ* strains could result from high rates of mutagenesis saturating the MMR machinery and thus reducing the pool of MSH proteins available to participate in heteroduplex rejection. Another possibility is that the lack of Hst3 and Hst4 leads to higher levels of H3K56 acetylation, which in turn favors nucleosome deposition by CAF-1 and Rtt106 factors that selectively bind to H3K56 acetylated histones. Such a situation could stabilize the strand invasion intermediate and suppress anti-recombination. In contrast, the presence of Hst3 and Hst4 would promote rejection by deacetylating H3K56 and thus suppress nucleosome deposition by CAF-1 and Rtt106. In support of this idea, Celic *et al*. (2006) showed that H3K56 sites are hyperacetylated in yeast lacking Hst3 and Hst4.  Additional studies will be required to distinguish between these models.

**Do chromatin modification factors have indirect effects on heteroduplex rejection?**

Studies in yeast showed that the Fun30, RSC and INO80 chromatin remodelers promote resection of DNA DSB ends (Chen *et al*. 2012; Lademann *et al*. 2017; Daley *et al*. 2015), and studies in human cells showed that ASF1 protects such ends from excessive resection (Huang *et al*. 2018). Could changes in resection rates impact heteroduplex rejection? Resection of DNA DSB ends is a critical initiating step in homologous recombination because it generates 3' single strand tails that participate in the formation of heteroduplex DNA. Thus, changes in resection rates could impact the stability of recombination intermediates by altering heteroduplex tract lengths. Consistent with this idea, a recent study in yeast showed that defects in resection as well as increased amounts of the single-strand binding protein RPA enhanced the efficiency of repair through an ectopic donor locus (Lee *et al*. 2016).  We were unable to test effects on resection because His$^+$ recombinants result from the repair of spontaneous DNA lesions. However, Datta *et al.* (1997) estimated in the system we used, the length of perfect homology required to initiate stable heteroduplex formation (20 bp), and avoid heteroduplex rejection (610 bp, which is larger than the 350 bp repeat). These estimates, the work of Lee *et al.* (2016), and our finding that homologous recombination is not decreased in the chromatin remodeling mutants analyzed, suggest to us that the *increased* rejection seen in *cac1Δ, rtt106Δ*, and *sir2Δ* mutants is not likely due to alterations in resection rates, though we cannot exclude the possibility that greater resection could lead to longer heteroduplex tract intermediates with more mismatches that are substrates for rejection. Also, we cannot exclude the possibility that the *decreased* rejection seen in *hst3Δ hst4Δ* and *rpd3Δ* mutants resulted from severely limited resection if only short heteroduplex tracts form that contain perfect homology tracts, without branch migration steps that would lead to the formation of mismatches.

Lastly, we recognize challenges using the ratio of homologous/divergent recombination as a surrogate for heteroduplex rejection efficiency. Such a measure could be misleading if chromatin modifying mutants alter recombination rates through pathways that are insensitive to heteroduplex rejection. In such a scenario the homologous/divergent recombination ratio would

appear to change without impacting the heteroduplex rejection machinery as a result of similarly increasing or decreasing homologous and divergent recombination. This scenario could explain the decreased rejection seen in *hst3Δ hst4Δ* mutants where homologous recombination events are dramatically elevated.  However, arguments against this possibility include: 1. The altered recombination rates seen in chromatin remodeling mutants did not correlate to changes in the homologous/divergent recombination ratio (Table 2).  2. The homologous/divergent recombination ratio was similarly disrupted in *msh2Δ*, *cac1Δ msh2Δ* and *rtt106Δ msh2Δ* mutants (Table 2), suggesting that the heteroduplex rejection machinery was responding in a similar fashion in *cac1Δ* and *rtt106Δ* mutants. The above concerns (resection rates, efficiency of rejection) will need to be further explored in a system where heteroduplex rejection can be monitored through DSB events induced at specific sites (e.g. HO, I-SceI).

**Acknowledgments**

We thank members of the Alani Lab, Jessica Tyler, Mike Fasullo, Michael Lichten and Scott Keeney for helpful comments and advice, and Sue Jinks-Robertson for providing us with the inverted repeat recombination strains SJR769, GCY615 and GCY559.  U.C., B.M. and E.A. were supported by National Institutes of Health (NIH) grant GM53085. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of General Medical Sciences or the National Institutes of Health.

**References**

Adkins MW, Howar SR, Tyler JK (2004) Chromatin disassembly mediated by the histone chaperone Asf1 is essential for transcriptional activation of the yeast *PHO5* and *PHO8* genes Mol Cell 14: 657-666.

Adkins MW, Tyler JK (2004) The histone chaperone Asf1p mediates global chromatin disassembly in vivo. J Biol Chem 279: 52069-52074.

Adkins NL, Swygert SG, Kaur P, Niu H, Grigoryev SA, Sung P, Wang H, Peterson CL (2017) Nucleosome-like, single-stranded DNA (ssDNA)-histone octamer complexes and the implication for DNA double-strand break repair. J Biol Chem 292: 5271-5281.

Anand R, Beach A, Li K, Haber J (2017) Rad51-mediated double-strand break repair and mismatch correction of divergent substrates. Nature 544: 377-380.

Blanko ER, Kadyrova LY, Kadyrov FA (2016) DNA mismatch repair interacts with CAF-1- and ASF1A-H3-H4-dependent histone (H3-H4)2 tetramer deposition. J Biol Chem 291: 9203–9217.

Boulton SJ, Jackson SP (1998) Components of the Ku-dependent non-homologous end-joining pathway are involved in telomeric length maintenance and telomeric silencing. EMBO J 17: 1819-1828.

Brachmann, CB, Davies A, Cost GJ, Caputo E, Li J, *et al.* (1998) Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR mediated gene disruption and other applications. Yeast 14: 115–132.

Burdova K, Mihaljevic B, Sturzenegger A, Chappidi N, Janscak P (2015) The Mismatch-Binding Factor MutSbeta Can Mediate ATR Activation in Response to DNA Double-Strand Breaks. Mol Cell 59: 603–614.

Celic I, Masumoto H, Griffith WP, Meluh P, Cotter RJ, Boeke JD, Verreault A (2006) The sirtuins hst3 and Hst4p preserve genome integrity by controlling histone h3 lysine 56 deacetylation. Curr Biol 16: 1280-1289.

Chakraborty, U., and E. Alani, (2016) Understanding how mismatch repair proteins participate in the repair/anti-recombination decision. FEMS Yeast Res 16**:** fow071.

Chakraborty U, Dinh TA, Alani E. (2018) Genomic Instability Promoted by Overexpression of Mismatch Repair Factors in Yeast: A Model for Understanding Cancer Progression. Genetics 209: 439-456.

Chakraborty U, George CM, Lyndaker AM, Alani E (2016) A delicate balance between repair and replication factors regulates recombination between divergent DNA sequences in Saccharomyces cerevisiae.  Genetics 202: 525-540.

Chen CC, Carson JJ, Feser J, Tamburini B, Zabaronick S, Linger J, Tyler JK (2008) Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. Cell 134: 231-243.

Chen W, Jinks-Robertson S (1998) Mismatch repair proteins regulate heteroduplex formation during mitotic recombination in yeast. Mol Cell Biol 18: 6525-6537.

Chen W, Jinks-Robertson S (1999) The role of the mismatch repair machinery in regulating mitotic and meiotic recombination between diverged sequences in yeast. Genetics 151: 1299–1313.

Chen X, Cui D, Papusha A, Zhang X, Chu CD, Tang J, Chen K, Pan X, Ira G (2012) The Fun30 nucleosome remodeller promotes resection of DNA double-strand break ends. Nature 489: 576-580.

Christianson TW, Sikorski RS, Dante M, Shero JH, Hieter P (1992) Multifunctional yeast high-copy-number shuttle vectors. Gene 110**:** 119-122.

Datta A, Adjiri A, New L, Crouse GF, Jinks-Robertson S (1996) Mitotic crossovers between diverged sequences are regulated by mismatch repair proteins in Saccharomyces cerevisiae. Mol Cell Biol 16: 1085–1093.

Datta A, Hendrix M, Lipsitch, M, Jinks-Robertson S (1997) Dual roles for DNA sequence identity and the mismatch repair system in the regulation of mitotic crossing-over in yeast. Proc Natl Acad Sci USA 94: 9757-9762.

Daley JM, Niu H, Miller AS, Sung P (2015) Biochemical mechanism of DSB end resection and its regulation.  DNA Repair 32: 66–74.

Debacker K, Frizzell A, Gleeson O, Kirkham-McCarthy L, Mertz T, Lahue RS (2012) Histone deacetylase complexes promote trinucleotide repeat expansions. PLoS Biol 10:e1001257. doi: 10.1371/journal.pbio.1001257.

Diao LT, Chen CC, Dennehey B, Pal S, Wang P, Shen ZJ, Deem A, Tyler JK (2017) Delineation of the role of chromatin assembly and the Rtt101Mms1 E3 ubiquitin ligase in DNA damage checkpoint recovery in budding yeast. PLoS ONE 12: e0180556.

Dion V, Kalck V, Horigome C, Towbin BD, Gasser SM (2012) Increased mobility of double-strand breaks requires Mec1, Rad9 and the homologous recombination machinery Nat Cell Biol 14: 502-509.

Driscoll R, Hudson A, Jackson SP (2007) Yeast Rtt109 promotes genome stability by acetylating histone H3 K56 on lysine 56.  Science 315: 649-652.

Ehrentraut S, Weber JM, Dybowski JN, Hoffmann D, Ehrenhofer-Murray AE (2010) Rpd3-dependent boundary formation at telomeres by removal of Sir2 substrate. Proc Natl Acad Sci USA 107: 5522-5527.

Evans E, Sugawara N, Haber JE, Alani E (2000) The Saccharomyces cerevisiae Msh2 mismatch repair protein localizes to recombination intermediates in vivo. Mol Cell 5: 789–799.

Franco AA, Lam WM, Burgers PM, Kaufman PD (2005) Histone deposition protein Asf1 maintains DNA replisome integrity and interacts with replication factor C. Genes Dev 19: 1365-1375.

Gaillard PH, Martini EM, Kaufman PD *et al*. (1996) Chromatin assembly coupled to DNA repair: a new role for chromatin assembly factor I. Cell 86: 887–896.

Gartenberg MR, Smith JS (2016) The nuts and bolts of transcriptionally silent chromatin in Saccharomyces cerevisiae. Genetics 203: 1563-1599.

George CM, Alani E (2012) Multiple cellular mechanisms prevent chromosomal rearrangements involving repetitive DNA. Crit Rev Biochem Mol Biol 47: 297-313.

Gietz RD, Schiestl RH (1991) Applications of high efficiency lithium acetate transformation of intact yeast cells using single-stranded nucleic acids as carrier. Yeast 7**:** 253-263.

Goldfarb T, Alani E (2005) Distinct roles for the Saccharomyces cerevisiae mismatch repair proteins in heteroduplex rejection, mismatch repair and nonhomologous tail removal. Genetics 169: 563-574.

Groth A, Rocha W, Verreault A, Almouzni G (2007) Chromatin challenges during DNA replication and repair. Cell 128: 721–733.

Guintini L, Tremblay M, Toussaint M, D'Amours A, Wellinger RE, Wellinger RJ, Conconi A (2017)  Repair of UV-induced DNA lesions in natural *Saccharomyces cerevisiae* telomeres is moderated by Sir2 and Sir3, and inhibited by yKu-Sir4 interaction. Nucleic Acids Res 45: 4577–4589.

Han J, Zhou H, Horazdovsky B, Zhang K, Xu RM, Zhang Z (2007) Rtt109 acetylates histone H3 lysine 56 and functions in DNA replication.  Science 315: 653-655.

Hauer MH, Gasser SM (2017) Chromatin and nucleosome dynamics in DNA damage and repair. Genes Dev 31: 2204–2221.

Hauer MH, Seeber A, Singh V, Thierry R, Sack R, Amitai A, Kryzhanovska M, Eglinger J, Holcman D, Owen-Hughes T, Gasser SM (2017) Histone degradation in response to DNA damage enhances chromatin dynamics and recombination rates. Nat Struct Mol Biol 24: 99–107.

Hoek M, Myers MP, Stillman B (2011) An analysis of CAF-1-interacting proteins reveals dynamic and direct interactions with the KU complex and 14-3-3 proteins. J Biol Chem 286: 10876-10887.

Hoggard TA, Chang F, Perry KR, Subramanian S, Kenworthy J, Chueng J, Shor E, Hyland EM, Boeke JD, Weinreich M, Fox CA (2018) Yeast heterochromatin regulators Sir2 and Sir3 act directly at euchromatic DNA replication origins. PLoS Genet. 2018 May 24;14(5):e1007418. doi: 10.1371/journal.pgen.1007418

Hong Z, Jiang J, Hashiguchi K, Hoshi M, Lan L, Yasui A (2008) Recruitment of mismatch repair proteins to the site of DNA damage in human cells.  J Cell Sci 121: 3146–3154.

Huang TH, Fowler F, Chen CC, Shen ZJ, Sleckman B, Tyler JK (2018) The histone chaperones ASF1 and CAF-1 promote MMS22L-TONSL-mediated Rad51 loading onto ssDNA during homologous recombination in human cells. Mol. Cell 69: 879–892.

Huang S, Zhou H, Katzmann D, Hochstrasser M, Atanasova E, Zhang Z (2005) Rtt106p is a histone chaperone involved in heterochromatin-mediated silencing. Proc Natl Acad Sci USA 102: 13410-13415.

Huang S, Zhou H, Tarara J, Zhang Z (2007) A novel role for histone chaperones CAF-1 and Rtt106p in heterochromatin silencing. EMBO J 26: 2274-2283.

Jiao R, Bachrati CZ, Pedrazzi G, Kuster P, Petkovic M, Li JL, Egli D, Hickson ID, Stagljar I (2004) Physical and functional interaction between the Bloom's syndrome gene product and the largest subunit of chromatin assembly factor 1. Mol Cell Biol 24: 4710-4719.

Jiao R, Harrigan JA, Shevelev I, Dietschy T, Selak N, Indig FE, Piotrowski J, Janscak P, Bohr VA, Stagljar I (2007) The Werner syndrome protein is required for recruitment of chromatin assembly factor 1 following DNA damage.  Oncogene 26: 3811-3822.

Kadyrova LY, Mertz TM, Zhang Y *et al.* (2013) A Reversible Histone H3 Acetylation Cooperates with Mismatch Repair and Replicative Polymerases in Maintaining Genome Stability. PLoS Genet 9: e1003899.

Kadyrova LY, Blanko ER, Kadyrov FA (2011) CAF-I-dependent control of degradation of the discontinuous strands during mismatch repair. Proc Natl Acad Sci USA 108: 2753-2758.

Kim JA, Haber JE (2009) Chromatin assembly factors Asf1 and CAF-1 have overlapping roles in deactivating the DNA damage checkpoint when DNA repair is complete. Proc Natl Acad Sci USA 106: 1151-1156.

Korber P, Barbaric S, Luckenbach T, Schmid A, Schermer UJ, Blaschke D, Hörz W (2006) The histone chaperone Asf1 increases the rate of histone eviction at the yeast *PH05* and *PH08* promoters. J Biol. Chem 281: 5539-5545.

Krawitz DC, Kama T, Kaufman PD (2002) Chromatin assembly factor I mutants defective for PCNA binding require Asf1/Hir proteins for silencing. Mol Cell Biol 22: 614-625.

Lademann CA, Renkawitz J, Pfander B, Jentsch S (2017) The INO80 Complex Removes H2A.Z to Promote Presynaptic Filament Formation during Homologous Recombination. Cell Rep 19: 1294-1303.

Lee, C.-S., Wang, R.W., Chang, H.-H., Capurso, D., Segal, M.R., and Haber, J.E. (2016) Chromosome position determines the success of double-strand break repair. Proc Natl Acad Sci USA 113: E146–E154.

Li F, Mao G, Tong D, Huang J, Gu L, Yang W, Li GM (2013) The histone mark H3K36me3 regulates human DNA mismatch repair through its interaction with MutSα. Cell 153: 590–600.

Li F, Tian L, Gu L, Li GM (2009) Evidence that nucleosomes inhibit mismatch repair in eukaryotic cells. J Biol Chem 284: 33056–33061.

Li Q, Zhou H, Wurtele H, Davies B, Horazdovsky B, *et al.* (2008) Acetylation of histone H3 lysine 56 regulates replication-coupled nucleosome assembly. Cell 134: 244–255.

Linger J, Tyler JK (2005) The yeast histone chaperone chromatin assembly factor 1 protects against double-strand DNA-damaging agents. Genetics 171: 1513–1522.

Lyndaker AM, Goldfarb T, Alani E (2008) Mutants defective in Rad1-Rad10-Slx4 exhibit a unique pattern of viability during mating-type switching in Saccharomyces cerevisiae. Genetics 179: 1807-1821.

Martin SG, Laroche T, Suka N, Grunstein M, Gasser SM (1999). Relocalization of telomeric Ku and SIR proteins in response to DNA strand breaks in yeast. Cell 97: 621-633.

McAinsh AD, Scott-Drew S, Murray JA, Jackson SP (1999) DNA damage triggers disruption of telomeric silencing and Mec1p-dependent relocation of Sir3p. Curr Biol 9: 963-966.

McMurray CT (2010) Mechanisms of trinucleotide repeat instability during human development. Nat Rev Genet 11: 786-799.

Mills KD, Sinclair DA, Guarente L (1999) MEC1-dependent redistribution of the Sir3 silencing protein from telomeres to DNA double-strand breaks. Cell 97: 609-620.

Miné-Hattab J, Rothstein R (2012) Increased chromosome mobility facilitates homology search during recombination Nat Cell Biol 14: 510-517.

Moggs JG, Grandi P, Quivy JP, Jónsson ZO, Hübscher U, Becker PB, Almouzni G (2000) A CAF-1-PCNA-mediated chromatin assembly pathway triggered by sensing DNA damage. Mol Cell Biol 20: 1206-1218.

Muñoz-Galván S, Jimeno S, Rothstein R, Aguilera A (2013) Histone H3K56 acetylation, Rad52, and non-DNA repair factors control double-strand break repair choice with the sister chromatid. PLoS Genet 9: e1003237.

Myung K, Datta A, Chen C, Kolodner RD (2001) SGS1, the Saccharomyces cerevisiae homologue of BLM and WRN, suppresses genome instability and homeologous recombination. Nat Genet 27: 113-116.

Neumann FR, Dion V, Gehlen LR, Tsai-Pflugfelder M, Schmid R, Taddei A, Gasser SM (2012) Targeted INO80 enhances subnuclear chromatin movement and ectopic homologous recombination Genes Dev 26: 369-383.

Nicholson A, Hendrix M, Jinks-Robertson S, Crouse GF (2000) Regulation of mitotic homeologous recombination in yeast: functions of mismatch repair and nucleotide excision repair genes. Genetics 154: 133–146.

Pietrobon V, Fréon K, Hardy J *et al.* (2014) The Chromatin Assembly Factor 1 Promotes Rad51-Dependent Template Switches at Replication Forks by Counteracting D-Loop Disassembly by the RecQ-Type Helicase Rqh1. PLoS Biol 12: e1001968.

Polo SE, Roche D, Almouzni G (2006) New histone incorporation marks sites of UV repair in human cells. Cell 127: 481–493.

Ransom M, Dennehey BK, Tyler JK (2010) Chaperoning histones during DNA replication and repair. Cell 140: 183-195.

Rose MD, Winston F, Hieter P (1990) Methods in Yeast Genetics. Cold Spring Harbor Laboratory Press.

Schöpf B, Bregenhorn S, Quivy J-P, Kadyrov FA, Almouzni G, Jiricny J (2012) Interplay between mismatch repair and chromatin assembly. Proc Natl Acad Sci USA 109: 1895-1900.

Schneider J, Bajwa P, Johnson FC, Bhaumik SR, Shilatifard A (2006) Rtt109 is required for proper H3K56 acetylation: a chromatin mark associated with the elongating RNA polymerase II. J Biol Chem 281: 37270–37274.

Schwabish MA Struhl K (2006) Asf1 mediates histone eviction and deposition during elongation by RNA polymerase II. Mol Cell 22: 415-422.

Shibahara K, Stillman B (1999) Replication-dependent marking of DNA by PCNA facilitates CAF-1-coupled inheritance of chromatin. Cell 96: 575–585.

Spell RM, Jinks-Robertson S (2004a) Examination of the roles of Sgs1 and Srs2 helicases in the enforcement of recombination fidelity in Saccharomyces cerevisiae. Genetics 168: 1855-1865.

Spell RM, Jinks-Robertson S (2004b) Determination of mitotic recombination rates by fluctuation analysis in Saccharomyces cerevisiae. Methods Mol Biol 262: 3-12.

Sugawara N Goldfarb T, Studamire B, Alani E, Haber JE (2004) Heteroduplex rejection during single-strand annealing requires Sgs1 helicase and mismatch repair proteins Msh2 and Msh6 but not Pms1. Proc Natl Acad Sci USA 101: 9315-9320.

Tagami H, Ray-Gallet D, Almouzni G, Nakatani Y (2004) Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis Cell, 116: 51-61.

Tamburini BA, Tyler JK (2005). Localized histone acetylation and deacetylation triggered by the homologous recombination pathway of double-strand DNA repair. Mol Cell Biol 25: 4903-4913.

Thurtle-Schmidt DM, Dodson AE, Rine J (2016) Histone deacetylases with antagonistic roles in Saccharomyces cerevisiae heterochromatin formation. Genetics 204: 177–190.

Tsubota T, Berndsen CE, Erkmann JA, Smith CL, Yang L, *et al*. (2007) Histone H3-K56 acetylation is catalyzed by histone chaperone-dependent complexes. Mol Cell 25: 703–712.

Tsukamoto Y, Kato J, Ikeda H. (1997) Silencing factors participate in DNA repair and recombination in Saccharomyces cerevisiae. Nature 388: 900-903.

Tyler JK, Adams CR, Chen SR, Kobayashi R, Kamakaka RT, Kadonaga JT (1999) The RCAF complex mediates chromatin assembly during DNA replication and repair. Nature 402: 555-560.

Yoshida K, Bacal J, Desmarais D, Padioleau I, Tsaponina O, *et al.* (2014) The histone deacetylases Sir2 and Rpd3 act on ribosomal DNA to control the replication program in budding yeast.  Mol Cell 54: 691–697.

Zhang Z, Shibahara K, Stillman B (2000) PCNA connects DNA replication to epigenetic inheritance in yeast. Nature 408: 221–225.

Zhou J, Zhou BO, Lenzmeier BA, Zhou JQ (2009) Histone deacetylase Rpd3 antagonizes Sir2-dependent silent chromatin propagation. Nucl Acids Res 37: 3699-3713.

**Table 1.  Strains and plasmids used in this study.**

| | |
|---|---|
| SJR328 | *MATα, ade2-101, his3Δ200, ura3-Nhe, lys2ΔRV::hisG, leu2-R* |
| SJR769 | SJR328, cβ2/cβ2:*LEU2*, homologous 350 bp substrates |
| GCY615 | SJR328, cβ2/cβ2-ns:*LEU2*, predicted to form four, base-base mismatches in the 350 bp cβ2 substrates |
| GCY559 | SJR328, cβ2/cβ2-4L:*LEU2*, predicted to form four 4-nt loops in the 350 bp cβ2 substrates |
| EAY1605,1606 | SJR769, *msh2Δ::hisG-URA3-hisG* |
| EAY1609,1610 | GCY615, *msh2Δ::hisG-URA3-hisG* |
| EAY3853-3856 | SJR769, *hst3Δ::KANMX* |
| EAY3857-3860 | GCY615, *hst3Δ::KANMX* |
| EAY3865-3868 | SJR769, *hst4Δ::KANMX* |
| EAY3869-3872 | GCY615, *hst4Δ::KANMX* |
| EAY3877-3880 | SJR769, *hst3Δ::KANMX hst4Δ::NATMX* |
| EAY3881-3884 | GCY615, *hst3Δ::KANMX hst4Δ::NATMX* |
| EAY3964-3967 | SJR769, *hst1Δ::KANMX* |
| EAY3968-3971 | GCY615, *hst1Δ::KANMX* |
| EAY3841-3844 | SJR769, *sir2Δ::KANMX* |
| EAY3845-3848 | GCY615, *sir2Δ::KANMX* |
| EAY3849-3852 | GCY559, *sir2Δ::KANMX* |
| EAY4002-4005 | SJR769, *rpd3Δ::KANMX* |
| EAY4006-4009 | GCY615, *rpd3Δ::KANMX* |
| EAY4047-4050 | SJR769, *gcn5Δ::KANMX* |
| EAY4051-4054 | GCY615, *gcn5Δ::KANMX* |
| EAY3889-3892 | SJR769, *cac1Δ::KANMX* |
| EAY3893-3896 | GCY615, *cac1Δ::KANMX* |
| EAY3897-3900 | GCY559, *cac1Δ::KANMX* |
| EAY3972,3974,3975 | SJR769, *rtt106Δ::KANMX* |
| EAY3976-3978 | GCY615, *rtt106Δ::KANMX* |
| EAY3901-3904 | SJR769, *asf1Δ::KANMX* |
| EAY3905-3908 | GCY615, *asf1Δ::KANMX* |
| EAY3909-3912 | GCY559, *asf1Δ::KANMX* |
| EAY4127-4129 | SJR769, *pol30-8::KANMX* |
| EAY4130-4132 | GCY615, *pol30-8::KANMX* |
| EAY3925, 3926 | SJR769, *cac1Δ::KANMX asf1Δ::NATMX* |
| EAY3929, 3930 | GCY615, *cac1Δ::KANMX asf1Δ::NATMX* |
| EAY3933, 3934 | GCY559, *cac1Δ::KANMX asf1Δ::NATMX* |
| EAY4018-4021 | SJR769, *rtt106Δ::KANMX asf1Δ::NATMX* |
| EAY4022-4025 | GCY615, *rtt106Δ::KANMX asf1Δ::NATMX* |
| EAY3913-3916 | SJR769, *rtt109Δ::KANMX* |
| EAY3917-3920 | GCY615, *rtt109Δ::KANMX* |
| EAY4010-4013 | SJR769, *rtt106Δ::KANMX cac1Δ::NATMX* |
| EAY4014-4017 | GCY615, *rtt106Δ::KANMX cac1Δ::NATMX* |
| EAY4027-4029 | SJR769, *sir2Δ::KANMX cac1Δ::NATMX* |
| EAY4030-4033 | GCY615, *sir2Δ::KANMX cac1Δ::NATMX* |
| EAY4034-4037 | SJR769, *rtt106Δ::KANMX sir2Δ::NATMX* |
| EAY4038-4041 | GCY615, *rtt106Δ::KANMX sir2Δ::NATMX* |
| EAY4042-4044 | SJR769, *rtt106Δ::KANMX cac1Δ::NATMX sir2Δ::HYGMX* |

| EAY4045,4046 | GCY615, *rtt106Δ::KANMX cac1Δ::NATMX sir2Δ::HYGMX* |
| EAY4069-4072 | SJR769, *cac1Δ::KANMX msh2Δ::hisG-URA3-hisG* |
| EAY4073-4076 | GCY615, *cac1Δ::KANMX msh2Δ::hisG-URA3-hisG* |
| EAY4077-4080 | SJR769, *rtt106Δ::KANMX msh2Δ::hisG-URA3-hisG* |
| EAY4081-4084 | GCY615, *rtt106Δ::KANMX msh2Δ::hisG-URA3-hisG* |

SJR328, SJR769, GCY615, and GCY559 were obtained from Sue Jinks-Robertson and are

described in Nicholson *et al.* (2000).

**Table 2. Recombination rates as measured in the inverted repeat reporter assay.**

| Strain | Relevant genotype | Rate of His$^+$ recombination (x10$^{-6}$) | Homologous rate/ Homeologous rate[a] |
|--------|-------------------|-------------------------------------------|--------------------------------------|
| Cβ2-Cβ2 | wild-type | 0.52 (0.46-0.57) [b] | |
| | *msh2Δ* | 2.08 (1.35-2.87) ** | |
| | *cac1Δ* | 1.76 (0.98-2.43) ** | |
| | *rtt106Δ* | 1.74 (1.33-2.43) ** | |
| | *asf1Δ* | 4.20 (3.04-4.41) ** | |
| | *rtt109Δ* | 4.04 (1.75-5.95) ** | |
| | *cac1Δ asf1Δ* | 1.01 (0.52-1.95) ** | |
| | *rtt106Δ asf1Δ* | 1.97 (1.13-2.80) ** | |
| | *cac1Δ rtt106Δ* | 2.67 (1.83-3.48) ** | |
| | *pol30-8* | 0.80 (0.59-1.85) ** | |
| | *hst3Δ* | 1.44 (1.28-1.76) ** | |
| | *hst4Δ* | 3.46 (3.09-4.19) ** | |
| | *hst3Δ hst4Δ* | 13.2 (5.18-32.9) ** | |
| | *hst1Δ* | 1.24 (1.01-1.52) ** | |
| | *sir2Δ* | 2.22 (2.07-2.66) ** | |
| | *rpd3Δ* | 0.26 (0.17-0.29) ** | |
| | *gcn5Δ* | 0.90 (0.77-1.13) ** | |
| | *sir2Δ cac1Δ* | 3.61 (3.29-7.07) ** | |
| | *sir2Δ rtt106Δ* | 6.59 (2.84-11.9) ** | |
| | *sir2Δ cac1Δ rtt106Δ* | 6.11 (3.99-8.34) ** | |

|  |  |  |  |
|---|---|---|---|
|  | *msh2Δ cac1Δ* | 5.17 (4.26-5.65) ** |  |
|  | *msh2Δ rtt106Δ* | 5.66 (3.50-7.35) ** |  |
| Cβ2/Cβ2-ns | wild-type | 0.029 (0.025-0.032) | 18 |
|  | *msh2Δ* | 2.36 (1.89-2.67) ** | 0.88 |
|  | *cac1Δ* | 0.032 (0.022-0.048) | 55 |
|  | *rtt106Δ* | 0.045 (0.020-0.095) | 39 |
|  | *asf1Δ* | 0.24 (0.20-0.29) ** | 18 |
|  | *rtt109Δ* | 0.24 (0.16-0.30) ** | 17 |
|  | *cac1Δ asf1Δ* | 0.054 (0.039-0.23) ** | 19 |
|  | *rtt106Δ asf1Δ* | 0.13 (0.080-0.42) ** | 15 |
|  | *cac1Δ rtt106Δ* | 0.025 (0.012-0.068) | 107 |
|  | *pol30-8* | 0.028 (0.020-0.034) | 28 |
|  | *hst3Δ* | 0.11 (0.079-0.13) ** | 14 |
|  | *hst4Δ* | 0.32 (0.26-0.42) ** | 11 |
|  | *hst3Δ hst4Δ* | 2.91 (1.57-6.05) ** | 4.5 |
|  | *hst1Δ* | 0.055 (0.029-0.082) ** | 22 |
|  | *sir2Δ* | 0.063 (0.041-0.079) ** | 35 |
|  | *rpd3Δ* | 0.033 (0.032-0.050) * | 7.8 |
|  | *gcn5Δ* | 0.062 (0.039-0.12) ** | 15 |
|  | *sir2Δ cac1Δ* | 0.062 (0.014-0.14) | 58 |
|  | *sir2Δ rtt106Δ* | 0.12 (0.078-0.18) ** | 57 |
|  | *sir2Δ cac1Δ rtt106Δ* | 0.24 (0.17-0.41) ** | 26 |
|  | *msh2Δ cac1Δ* | 5.32 (3.86-6.19) ** | 0.97 |
|  | *msh2Δ rtt106Δ* | 3.83 (3.36-4.48) ** | 1.5 |

| | | | |
|---|---|---|---|
| Cβ2/Cβ2-4L | wild-type | 0.062 (0.040-0.11) | 8.3 |
| | *cac1Δ* | 0.082 (0.058-0.17) | 21 |
| | *asf1Δ* | 0.57 (0.42-0.71) ** | 7.3 |
| | *cac1Δ asf1Δ* | 0.27 (0.20-0.38) ** | 3.7 |
| | *sir2Δ* | 0.16 (0.11-0.19) ** | 14 |

Recombination rates and 95% confidence intervals were calculated as described in Materials and Methods. The genotypes of the parental strains are shown in Table 1. Cβ2/Cβ2 = homologous substrate; Cβ2/Cβ2-ns = base-base mismatch substrate; Cβ2/Cβ2-4L = 4-nt loop mismatch substrate. [a]Homologous rate (Cβ2-Cβ2)/homeologous rate for strains with the same genotype or overexpression plasmid. [b]Numbers in parentheses indicate 95% confidence intervals. *significantly different from wild-type of the same strain (p<0.05, Mann-Whitney test). **significantly different from wild-type of the same strain (p<0.01, Mann-Whitney test).

**Table 3.  Effect of MSH2 and MSH6 overexpression on recombination and heteroduplex rejection.**

| strain | genotype, plasmid | Rate of His$^+$ recombination (x10$^{-6}$) | Homologous rate/ Homeologous rate[a] |
|---|---|---|---|
| SJR769; | wild-type, *2µ-dummy*[#] | 0.64 (0.47-0.90) | |
| Cβ2-Cβ2 | wild-type, *2µ-MSH2-MSH6* [#] | 6.00 (4.09-9.40) ** | |
| | *sir2Δ, 2µ-dummy* | 3.57 (3.09-5.03) ** | |
| | *sir2Δ, 2µ-MSH2-MSH6* | 6.72 (5.04-8.43) ** | |
| | *rtt106Δ, 2µ-dummy* | 1.66 (1.43-4.04) ** | |
| | *rtt106Δ, 2µ-MSH2-MSH6* | 2.69 (2.27-4.19) ** | |
| GCY615; | wild-type, *2µ-dummy*[#] | 0.042 (0.021-0.076) | 15 |
| Cβ2-Cβ2 | wild-type, *2µ-MSH2-MSH6* [#] | 0.11 (0.05-0.23) * | 56 |
| -ns | *sir2Δ, 2µ-dummy* | 0.074 (0.040-0.14) * | 48 |
| | *sir2Δ, 2µ-MSH2-MSH6* | 0.074 (0.051-0.13) * | 91 |
| | *rtt106Δ, 2µ-dummy* | 0.041 (0.029-0.093) | 40 |
| | *rtt106Δ, 2µ-MSH2-MSH6* | 0.21 (0.14-0.65) ** | 13 |

Recombination rates and 95% confidence intervals were calculated as described in Materials and Methods. The genotypes of the parental strains are shown in Table 1. Cβ2/Cβ2 = homologous substrate; Cβ2/Cβ2-ns = base-base mismatch substrate. [a]Homologous rate (Cβ2-Cβ2)/homeologous rate for strains with the same genotype or overexpression plasmid. [b]Numbers in parentheses indicate 95% confidence intervals. *significantly different from wild-type of the same strain (p<0.05, Mann-Whitney test). **significantly different from wild-type of the same strain (p<0.01, Mann-Whitney test). [#]Published data from Chakraborty *et al*. (2018).

**Figure legends**

**Figure 1.  Roles for histone chaperones, acetylases and deacetylases in regulating heteroduplex rejection efficiency**.  A. Schematic of an intron-based recombination assay involving inverted repeat sequences (white and gray boxes) that form a functional *HIS3* reporter following HR (adapted from Nicholson *et al*. 2000). His$^+$ recombinants are thought to result from gene conversion between inverted repeat sequences on sister chromatids. Identical or divergent repeats predicted to form base-base (cβ2/cβ2-ns) and 4-nt loop (cβ2/cβ2-4L) mismatches in heteroduplex DNA were analyzed. B. Ratios of homologous to divergent recombination rates (calculated as described in the Materials and Methods) are shown for base-base mismatches as a measure of heteroduplex rejection efficiency for wild-type or the indicated mutant strains. See Table 2 for the presentation of quantitative measurements.  The dashed lines in panel B indicate the ratio of homologous to homeologous recombination rates for wild-type.

**Figure 2. A model outlining possible steps during homologous recombination where chromatin modifiers regulate the repair/rejection decision.** We hypothesize that factors promoting nucleosome assembly/maintenance such as CAF-1 and Rtt106 deposit nucleosomes at DNA lesions independent of DNA synthesis on ssDNA prior to strand invasion (1). Such deposition/maintenance, coupled with the formation of Sir2-dependent chromatin marks that promote a closed chromatin state, stabilizes D-loops to suppress rejection. Additionally, we hypothesize that CAF-1 physically interacts with Msh6 and possibly Sgs1 to inhibit the rejection factors from unwinding heteroduplex substrates (2). Finally, DNA synthesis coupled nucleosome deposition by CAF-1 and Rtt106 combined with Sir2 dependent modifications of these nucleosomes is likely to stabilize heteroduplex D-loop intermediates, making it more difficult for rejection factors to unwind the intermediates. In this model MSH proteins are recruited to a DSB either directly or through specific histone marks, and nucleosome maintenance limits the time frame in which anti-recombination can be performed and promotes repair of the broken

chromosome and ultimately mismatch repair steps (see text for additional details). Factors/steps that promote rejection are highlighted in pink boxes and those that promote repair are highlighted in yellow boxes.

A.

His⁻ → sister chromatid gene conversion → His⁺

B.

Figure 1

**DSB**

Histones deposited
by CAF-I and Rtt106
on ssDNA

Mismatch
recognition

CAF-I inhibits
MutSα and Sgs1

Nucleosomes
deposited by
CAF-I and Rtt106

Repair: Deposited
histones deacetylated by
Sir2 stabilizes D-loop

Rejection: Msh2-Msh6 recruits STR
complex and unwinds heteroduplex
prior to nucleosome deposition

Msh2-Msh6

Nucleosome

Mismatch

Sgs1-Top3-Rmi1

CAF-1

Rtt106

Sir2

Rpd3

Promotes
repair

Promotes
rejection

Figure 2