A Data-Driven Transcriptional Taxonomy of Adipogenic Chemicals to Identify White and

Brite Adipogens

Stephanie Kim,*[1,2,] Eric Reed,*[1,3,4] Stefano Monti,[#1,3,4] Jennifer Schlezinger[#1,2]

*Co-first authors

[#]Co-senior authors

[1]Boston University Superfund Research Program, Boston University, MA 02118 USA

[2]Boston University School of Public Health, Department of Environmental Health, MA 02118

USA

[3]Boston University School of Medicine, Department of Computational Medicine, MA 02118

USA

[4]Boston University Bioinformatics Program, MA 02115, USA

Corresponding author:
Jennifer J. Schlezinger, Ph.D.
Boston University School of Public Health
Dept. of Environmental Health
715 Albany Street, R-405
Boston, MA 02118
Phone: 617-358-1708
Email: jschlezi@bu.edu
22
Stefano Monti, Ph.D.
Boston University School of Medicine
Division of Computational BioMedicine
72 East Concord Street, E-611
Boston, MA – 02118
Phone: 617-358-7087
Email : smonti@bu.edu
30

31   **Running Title:**
32   Classification Taxonomy for Adipogenic Chemicals
33

34   **Acknowledgements:**
35
36   **Declaration of competing financial interests (CFI):**
37   The authors declare they have no actual or potential competing financial interests.
38
39   **Grant Information:**
40   Superfund Research Program [P42 ES007381]
41   BU-Joslin Pilot & Feasibility Program Award (2016)

Abstract:

Background: Chemicals in disparate structural classes activate specific subsets of PPARγ's transcriptional programs to generate adipocytes with distinct phenotypes.

Objectives: Our objectives were to 1) establish a novel classification method to predict PPARγ ligands and modifying chemicals, and 2) create a taxonomy to group chemicals based on their effects on PPARγ's transcriptome and downstream metabolic functions. We tested the hypothesis that environmental adipogens highly ranked by the taxonomy, but segregated from therapeutic PPARγ ligands, would induce white but not brite adipogenesis.

Methods: 3T3-L1 cells were differentiated in the presence of 76 chemicals (negative controls, nuclear receptor ligands known to influence adipocyte biology, potential environmental PPARγ ligands). Differentiation was assessed by measuring lipid accumulation. mRNA expression was determined by RNA-Seq and validated by RT-qPCR. A novel classification model was developed using an amended random forest procedure. A subset of environmental contaminants identified as strong PPARγ agonists were analyzed by their effects on lipid handling, mitochondrial biogenesis and cellular respiration in 3T3-L1 cells and human preadipocytes.

Results: We used lipid accumulation and RNA sequencing data to develop a classification system that 1) identified PPARγ agonists, and 2) sorted chemicals into likely white or brite adipogens. Expression of *Cidec* was the most efficacious indicator of strong PPARγ activation. Two known environmental PPARγ ligands, tetrabromobisphenol A and triphenyl phosphate, which sorted distinctly from therapeutic ligands, induced white adipocyte genes but failed to induce *Pgc1a* and *Ucp1*, and induced fatty acid uptake but not mitochondrial biogenesis in 3T3-L1 cells. Moreover, two chemicals identified as highly ranked PPARγ agonists, tonalide and

quinoxyfen, induced white adipogenesis without the concomitant health-promoting

characteristics of brite adipocytes in mouse and human preadipocytes.

Discussion: A novel classification procedure accurately identified environmental chemicals as

PPARγ ligands distinct from known PPARγ-activating therapeutics. The computational and

experimental framework has general applicability to the classification of as-yet uncharacterized

chemicals.

Introduction:

Since 1980, the prevalence of obesity has been increasing in the United States (Flegal et al. 2016). Further, in 2015, it was estimated that a total of 108 million children and 604 million adults were obese worldwide (GBD 2017). This poses a major public health threat since overweight and obesity increase the risk of metabolic syndrome, which, in turn, sets the stage for metabolic diseases, such as type 2 diabetes, cardiovascular disease, nonalcoholic fatty liver disease and stroke (Park et al. 2003). The Endocrine Society's latest scientific statement on the obesity pathogenesis states that obesity is a disorder of the energy homeostasis system, rather than just a passive accumulation of adipose, and that environmental factors, including chemicals, confer obesity risk (Schwartz et al. 2017). The rapid increases in obesity and metabolic diseases correlate with substantial increases in environmental chemical production and exposures over the last few decades, and experimental evidence in animal models demonstrates the ability of a broad spectrum of various environmental metabolism-disrupting chemicals to induce adiposity and metabolic disruption (Heindel et al. 2017).

Adipocytes are essential for maintaining metabolic homeostasis as they are the storage depot of free fatty acids and release hormones that can modulate body fat mass (Rosen and Spiegelman 2006). Adipogenesis is a highly regulated process that involves a network of transcription factors acting at different time points during differentiation (Farmer 2006). Peroxisome proliferator activated receptor γ (PPARγ) is a ligand activated, nuclear receptor, which is required for adipocyte formation and function (Tontonoz et al. 1994), as well as metabolic homeostasis. In both PPARγ haploinsufficient (Gumbilai et al. 2016) and knockout (He et al. 2003; Jiang et al. 2014; O'Donnell et al. 2016; Zhang et al. 2004) rodent models, there was a lack of adipocyte formation and metabolic disruption.

PPARγ regulates energy homeostasis by activating expression of genes involved in both

storage of excess energy as lipids in white adipocytes and energy utilization by triggering

mitochondrial biogenesis, fatty acid oxidation and thermogenesis in brite (brown-in-white) and

brown adipocytes. The white adipogenic, brite/brown adipogenic and insulin sensitizing

activities of PPARγ are regulated distinctly through ligand-specific post-translational

modifications (Banks et al. 2015; Choi et al. 2010; Choi et al. 2011; Qiang et al. 2012) and co-

regulator recruitment (Burgermeister et al. 2006; Feige et al. 2007; Ohno et al. 2012; Villanueva

et al. 2013). While adult humans have long been thought to not have brown adipose tissue,

humans with minimal brite adipocyte populations are at higher risk for obesity and type 2

diabetes (Claussnitzer et al. 2015; Sidossis and Kajimura 2015; Timmons and Pedersen 2009).

Growing evidence supports the hypothesis that environmental PPARγ ligands induce

phenotypically distinct adipocytes. Tributyltin (TBT) induces the formation of an adipocyte with

lower adiponectin expression and altered glucose homeostasis (Regnier et al. 2015).

Furthermore, TBT failed to induce expression of genes associated with browning of adipocytes

(e.g. *Ppara*, *Pgc1a*, *Cidea*, *Elovl3*, *Ucp1*) in differentiating 3T3-L1 adipocytes (Kim et al. 2018;

Shoucri et al. 2018). As a result, TBT-induced adipocytes failed to up-regulate mitochondrial

biogenesis and had low levels of cellular respiration (Kim et al. 2018; Shoucri et al. 2018). The

structurally similar environmental PPARγ ligand, triphenyl phosphate, also fails to induce brite

adipogenesis, and this correlates with an inability to prevent PPARγ from being phosphorylated

at Serine 273 (S273) (Kim et al. 2020).

The EPA developed the Toxicity Forecaster (ToxCast™) program to use high-throughput

screening assays to prioritize chemicals and inform regulatory decisions regarding thousands of

environmental chemicals (Kavlock et al. 2012). Several ToxCast™ assays can measure the

ability of chemicals to bind to or activate PPARγ, and these assays have been used to generate a

toxicological priority index (ToxPi) that were expected to predict the adipogenic potential of

chemicals in cell culture models (Auerbach et al. 2016). Yet, it was shown that the results of

ToxCast™ PPARγ assays do not always correlate well with activity measured in a laboratory

setting and that the ToxPi designed for adipogenesis was prone to predicting false positives

(Janesick et al. 2016). Furthermore, the ToxCast/ToxPi approach cannot distinguish between

white and brite adipogens (Pereira-Fernandes et al. 2014).

In this study, we investigate differences in cellular response between adipogenic and non-

adipogenic compounds, as well as the heterogeneity of response across adipogenic compounds.

Our ultimate goal is to create a method for identification of novel adipogenic compounds using

the taxonomic organization of known and predicted adipogenic compounds based on their

divergent transcriptional response. To this end, we generated phenotypic and transcriptomic data

from adipocytes differentiated in the presence of 76 different chemicals. We combined the cost-

effective generation of agonistic transcriptomic data by 3'Digital Gene Expression – a highly

multiplexed RNA-Seq technology – with a new classification method to predict PPARγ-

activating and modifying chemicals. Further, we investigated metabolic-related outcome

pathways as effects of the chemical exposures. We created a data-driven taxonomy to

specifically classify chemicals into distinct categories based on their various interactions with

and effects on adipogenesis, presumably through their interaction with PPARγ. Based on the

taxonomy-based predictions, we tested the phenotype (white *vs.* brite adipocyte functions) of

environmental adipogens predicted to fail to induce brite adipogenesis in 3T3-L1 cells and

primary human adipocytes.

Methods:

Chemicals

149   DMSO was purchased from American Bioanalytical (Natick, MA). CAS numbers,

150   sources and catalog numbers of experimental chemicals are provided in **Table S1**. Human

151   insulin, dexamethasone, 3-isobutyl-1-methylxanthine (IBMX), and all other chemicals were from

152   Sigma-Aldrich (St. Louis, MO) unless noted.

Cell Culture

3T3 L1 (RRID:CVCL_0123, Lot # 63343749, ATCC, Manassas, VA) cells were

originally derived from a Swiss mouse embryonic fibroblast line (Green and Kehinde, 1975).

Cells were maintained in Growth Medium (high-glucose DMEM (Corning, 10-013-CV) with

10% calf serum (Sigma), 100 U/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin

B). All experiments were conducted with cells between passages 3 and 9. Experimental

conditions are outlined in **Table 1** and **Figure S1A**. For experiments, cells were plated in

Growth Medium and incubated for 4 days, at which time the cultures are confluent for 2 days.

"Naïve" pre-adipocytes were cultured in every experiment and grown in Growth Medium for the

duration of an experiment. On day 0, differentiation was induced by replacing the medium with

Differentiation Medium (DMEM,10% fetal bovine serum (FBS, Sigma-Aldrich), 100 U/ml

penicillin, 100 µg/ml streptomycin, 250 nM dexamethasone (Figures 1-3), 1 nM dexamethasone

(Figures 5-10) or no dexamethasone (Figure S3), 167 nM human insulin, 0.5 mM IBMX). Also

on day 0, single experimental wells were treated with vehicle (DMSO, 0.2% final concentration),

rosiglitazone (positive control, 200 nM) or test chemicals. On days 3 and 5 of differentiation,

medium was replaced with Maintenance Medium (DMEM, 10% FBS, 167 nM human insulin,

100 U/ml penicillin, 100 µg/ml streptomycin), and the cultures were re-dosed. On Day 7 of

differentiation, medium was replaced with Adipocyte Medium (DMEM, 10% FBS, 100 U/ml

penicillin, 100 µg/ml streptomycin), and the cultures were re-dosed. On day 10, cytotoxicity was

assessed by microscopic inspection, with cultures containing more than 10% rounded cells

excluded from consideration. For most chemicals only a single concentration was tested;

however, for those chemicals inducing toxicity the concentration was reduced until no toxicity

was observed (See **Table S1** for information on maximum tested and maximum non-toxic

concentrations tested). Wells with healthy cells were harvested for analysis of gene expression,

lipid accumulation, fatty acid uptake, mitochondrial biogenesis, mitochondrial membrane

potential, and cellular respiration.

OP9 cells (RRID:CVCL_4398, Lot # 63544739, ATCC) are a bone marrow stromal cell

line derived from newborn calvaria of the (C57BL/6xC3H)F2-op/op mouse (Nakano *et al.*, 1994).

Cells were maintained in αMEM (Gibco, 12-561-056) with 20% FBS, 26 mM sodium bicarbonate,

100 U/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B. Cells were plated in 24

well plates at 50,000 cells were well in 500 µl medium and incubated for 4 days. Induction and

maintenance of adipogenesis and treatment were as described for 3T3 L1 cells, except that the

dexamethasone concentration was 125 nM.

Primary human subcutaneous pre-adipocytes from five individual female patients were

obtained from the Boston Nutrition Obesity Research Center (Boston, MA). The patients were

32-51 years of age and had BMIs ranging from 26.0-30.7. One patient was pre-diabetic, and four

were non-diabetic. Adipocytes were differentiated as previously described (Lee and Fried 2014).

Experimental conditions are outlined in **Table 1** and **Figure S1B**. The pre-adipocytes were

maintained in Growth Medium (αMEM with 10% FBS, 100 U/ml penicillin, 100 µg/ml

streptomycin, 0.25 µg/ml amphotericin B). For experiments, human pre-adipocytes were plated

in Growth Medium and grown to confluence (3-5) days. "Naïve" pre-adipocytes were cultured in

every experiment and grown in Growth Medium for the duration of an experiment. On day 0,

differentiation was induced by replacing the Growth Medium with Differentiation Medium

(DMEM/F12, 25 mM NaHCO$_3$, 100 U/ml penicillin, 100 µg/ml streptomycin, 33 µM d-Biotin,

17 µM pantothenate, 100 nM dexamethasone, 100 nM human insulin, 0.5 mM IBMX, 2 nM T$_3$,

10 µg/ml transferrin). Also on day 0, single experimental wells also were treated with vehicle

(DMSO, 0.1% final concentration), rosiglitazone (positive control, 4 µM) or test chemicals. On

day 3 of differentiation, medium was replaced with fresh Differentiation Medium, and the

cultures were re-dosed. On days 5, 7, 10, and 12 of differentiation, the medium was replaced

with Maintenance Medium (DMEM/F12, 25 mM NaHCO$_3$, 100 U/ml penicillin, 100 µg/ml

streptomycin, 3% FBS, 33 µM d-Biotin, 17 µM pantothenate, 10 nM dexamethasone, 10 nM

insulin), and the cultures were re-dosed. Following 14 days of differentiation and dosing, cells

were harvested for analysis of gene expression, lipid accumulation, fatty acid uptake,

mitochondrial biogenesis, and cellular respiration.

Lipid Accumulation

3T3-L1 cells or human preadipocytes were plated in 24 well plates at 50,000-100,000

cells per well in 0.5 ml maintenance medium at initiation of the experiment. Dosing is outlined in

Table 1. Medium was removed from the differentiated cells, and they were rinsed with PBS. The

cells were then incubated with Nile Red (1 µg/ml in PBS) for 15 min in the dark. Fluorescence

(λex= 485 nm, λem= 530 nm) was measured using a Synergy2 plate reader (BioTek Inc.,

Winooski, VT). The fluorescence in all experimental wells was normalized by subtracting the

fluorescence measured in naïve pre-adipocyte cultures within each experiment and reported as

"Naïve Corrected RFU."

Transcriptome Profiling

3T3-L1 cells were plated in 24 well plates at 50,000 cells per well in 0.5 ml maintenance

medium at initiation of the experiment. Dosing is outlined in **Table 1**. Total RNA was extracted

and genomic DNA was removed using the Direct-zol MagBead RNA Kit and following

manufacturer's protocol (Zymo Research, Orange, CA). RNA concentrations and contamination

were determined spectrophotometrically using a Nanodrop (ND-1000, ThermoFisher). A final

concentration of 5 ng RNA/ul was used for each sample. For each chemical, 3-5 biological

replicates were profiled and carefully randomized across six 96-well plates, including 26 DMSO

vehicle controls, and 16 naïve pre-adipocyte cultures. Sequencing and gene expression

quantification was carried out by the MIT Technology Lab at Broad Institute (Cambridge, MA).

RNA libraries were prepared using a highly multiplexed 3' Digital Gene Expression (3' DGE)

protocol developed by (Xiong et al. 2017) and sequenced on an Illumina NextSeq 500,

generating between 2.13E8 and 3.87E8 reads and a mean of 3.02E8 reads per lane across 96

samples. All reads containing bases with Phred quality scores < Q10 were removed. The

remaining reads were aligned to mouse reference genome, GRCm38, and counted in 21,511

possible transcripts annotations. Only instances of uniquely aligned reads were quantified (i.e.,

reads that aligned to only one transcript). Furthermore, multiple reads with the same unique

molecular identifier (UMI), aligning to the same gene were quantified as a single count.

Reverse Transcription (RT)-qPCR

3T3-L1 cells or human preadipocytes were plated in 24-well plates at 100,000 cells per

well in 0.5 ml maintenance medium at initiation of the experiment. Dosing is outlined in **Table**

1. Total RNA was extracted and genomic DNA was removed using the 96-well Direct-zol

MagBead RNA Kit (Zymo Research). RNA concentrations and contamination were determined

spectrophotometrically using a Nanodrop. cDNA was synthesized from total RNA using the

iScript™ Reverse Transcription System (BioRad, Hercules, CA). All qPCR reactions were

performed using the PowerUp™ SYBR Green Master Mix (Thermo Fisher Scientific, Waltham,

MA). The qPCR reactions were performed using a 7500 Fast Real-Time PCR System (Applied

Biosystems, Carlsbad, CA): UDG activation (50°C for 2 min), polymerase activation (95°C for 2

min), 40 cycles of denaturation (95°C for 15 sec) and annealing (various temperatures for 15

sec), extension (72°C for 60 sec). The primer sequences and annealing temperatures are provided

in **Table S2**. All primers were obtained from Integrated DNA Technologies (Coralville, IA).

Relative gene expression was determined using the Pfaffl method to account for differential

primer efficiencies (Pfaffl 2001), using the geometric mean of the Cq values for beta-2-

microglobulin (*B2m*) and 18s ribosomal RNA (*Rn18s*) for mouse gene normalization and of

ribosomal protein L27 (*RPL27*) and *B2M* for human gene normalization. The Cq value from

naïve, pre-adipocyte cultures within each experiment was used as the reference point for the

experiment. Data are reported as "Relative Expression."

Cell Number Analysis

3T3-L1 cells or human preadipocytes were plated in 96-well, black-sided plates at 12,500

cells per well in 0.2 ml maintenance medium at initiation of the experiment. Dosing is outlined in

Table 1. Cellular protein was measured by Janus Green staining and measuring absorbance (595

nM) using a Synergy2 plate reader. The absorbance in experimental wells was normalized by

dividing by the absorbance measured in naïve pre-adipocyte cultures within the experiment and

reported as "Relative Cell Density."

Fatty Acid Uptake

3T3-L1 cells or human preadipocytes were plated in 96-well, black-sided plates at 12,500

cells per well in 0.2 ml maintenance medium at initiation of the experiment. Dosing is outlined in

Table 1. Fatty acid uptake was measured by treating differentiated cells with 100 µL of Fatty

Acid Dye Loading Solution (Sigma-Aldrich, MAK156). Fluorescence intensity (λex= 485nm,

λem= 530nm) measured at time zero and after a 10-minute incubation using a Synergy2 plate

reader. The fluorescence at time zero was subtracted from the fluorescence at the end of the

incubation and as "RFU."

Mitochondrial Biogenesis

3T3-L1 cells or human preadipocytes were plated in 24-well plates at 100,000 cells per

well in 0.5 ml maintenance medium at initiation of the experiment. Dosing is outlined in **Table**

1. Mitochondrial biogenesis was measured in differentiated cells using the MitoBiogenesis In-

Cell Elisa Colorimetric Kit, following the manufacturer's protocol (Abcam). The expression of

the mitochondrial protein SDH was measured and normalized to total protein content via Janus

Green staining. Absorbance (OD 405nm for SDH, and OD 595nm for Janus) was measured

using a Synergy2 plate reader. The absorbance ratios of SDH/Janus are reported as "Relative

Mitochondrial Protein Expression."

Oxygen Consumption

3T3-L1 cells were plated in 24-well plates at 100,000 cells per well in 0.5 ml

maintenance medium at initiation of the experiment. Dosing is outlined in **Table 1**. After 10 days

of differentiation, cells were gently trypsinized, diluted in 400 ul Adipocyte Medium, and 80 ul

per well was transferred to duplicate wells of a 96-well Agilent Seahorse plate. After 24 hrs

incubation, the medium was changed to Seahorse XF Assay Medium without glucose (1mM

sodium pyruvate, 1mM GlutaMax, pH 7.4) and cultures incubated at 37°C in a non-CO_2

incubator for 1 hr. To measure mitochondrial respiration, the Agilent Seahorse XF96 Cell Mito

Stress Test Analyzer (available at BUMC Analytical Instrumentation Core) was used, following

the manufacturer's standard protocol. The compounds and their concentrations used to determine

oxygen consumption rate (OCR) included 1) 0.5 μM oligomycin, 1.0 μM carbonyl cyanide-p-

trifluoromethoxyphenylhydrazone (FCCP) and 2 μM rotenone/2 μM antimycin. After the

Seahorse analysis, cells were fixed in 4% paraformaldehyde and stained with Janus Green.

Respiration rates were normalized by dividing by the Janus Green absorbance and reported as

"Relative OCR." Basal respiration was determined by subtracting non-mitochondrial respiration

from the last rate measurement before the injection of oligomycin. Maximum respiration was

determined by subtracting non-mitochondrial respiration from the maximum rate measurement

after the injection of FCCP. Spare capacity was determined by subtracting the basal respiration

from the maximum respiration.

Adiponectin Secretion

3T3-L1 cells were plated in 24-well plates at 100,000 cells per well in 0.5 ml

maintenance medium at initiation of the experiment. Dosing is outlined in **Table 1**. After 9 days

of differentiation, the medium was replaced with washout medium (DMEM+0.1% BSA), and

cultures were incubated for 24 hrs. The washout medium was replaced, the cells redosed and the

incubation continued for 24 hrs. Samples of medium were collected, diluted 1:200 and analyzed

for adiponectin by ELISA following the manufacturer's instructions (Mouse

Adiponectin/Acrp30 Quantikine ELISA Kit, R&D Systems, MRP300). Absorbance (450 nM)

was measured using a Synergy2 plate reader and concentrations calculated relative to a standard

curve.

Statistical Analyses

All statistical analyses were performed in R (v 3.4.3) and Prism 7 (GraphPad Software,

Inc., La Jolla, CA). All R code used for processing and analysis of transcriptome profiles is

publicly available on GitHub (https://github.com/montilab/Adipogen2020) and were carried out

using several R packages. Normalization and differential gene expression analysis was

performed using *limma* (v 3.34.9) (Ritchie et al. 2015). Batch correction was performed using

ComBat (v 3.26.0) (Leek et al. 2012). Gene set projection was performed using *GSVA* (v 1.30.0)

(Hanzelmann et al. 2013). Statistical testing of partial correlation estimates was performed using

ppcor (v 1.1) (Kim 2015).

For 3T3-L1 experiments the biological replicates correspond to independently plated

experiments. For human primary preadipocyte experiments the biological replicates correspond

to distinct individuals' preadipocytes (5 individuals in all). For each experiment, naïve,

undifferentiated 3T3-L1 cells were included for within-experiment normalization. The Nile Red

data and the qPCR data were not normally distributed; therefore, the data were log transformed

before statistical analyses. One-factor ANOVAs (Dunnett's) were performed to analyze the

qPCR and phenotypic data and determine differences from vehicle-treated cells. Data are

presented as means $\pm$ standard error (SE).

Transcriptome Data Processing

The number of counted reads per sample transcriptome profile varied widely with a range

of 7.90E1 to 2.27E6 (Mean = 2.25E5, SD = 2.94E5). To remove technical noise introduced by

low overall expression quantification of individual samples, we performed an iterative

clustering-based approach to determine sets of samples which segregate as a result of low total

read counts. Each iteration included four steps: removal of low count genes, normalization, plate-

level batch correction, and hierarchical clustering. Low count genes, defined as having mean

counts <1 across all samples, were removed to reduce statistical noise introduced by inaccurate

quantification of consistently lowly expressed transcripts. Normalization was performed using

Trimmed Mean of M-values (TMM), the default method employed by *limma* (v 3.34.9) (Ritchie

et al. 2015). Batch correction was performed by *ComBat* (v 3.26.0) (Leek et al. 2012).

Hierarchical clustering was performed on the 3000 genes with the largest median absolute

deviation (MAD) score, using Euclidean distance and 1-Pearson correlation as the distance

metric for samples and genes, respectively, and Ward's agglomerative method (Ward 1963).

Clusters of samples clearly representative of low expression quantification were removed. This

process was repeated until no such low expression outlier sample cluster was present (four

iterations). For the remaining samples, low count genes were removed, and samples were

normalized and batch corrected by the same procedure.

Following sample- and gene-level quality control filtering, the final processed data set

included expression levels of 9,616 genes for each of 234 samples. These 234 samples included

2-4 remaining replicates of each compound, 25 DMSO vehicle controls, and 15 naïve pre-

adipocyte cultures. Sequencing data from 3'DGE have been deposited to GEO (Accession:

GSE124564).

PPARγ Activating/Modifier Classification

A classification model was inferred from the *training set* consisting of 37 known PPARγ-

modifying compounds and 23 known non-PPARγ modifying compounds, including vehicle, to

predict the label of the *test set* of 17 potential PPARγ-modifying compounds (**Table S1**). Known

compounds were selected based on a literature search for experimental evidence of modification

of PPARγ activity, including PPARγ binding assays, coactivator recruitment or computational

modeling (definitive ligands), PPARγ-driven reporter assays (at least 25% of the rosiglitazone-

induced maximum), expression of PPARγ target genes and/or differentiation of 3T3 L1 or

multipotent stromal cells into adipocytes in the absence of a known PPARγ ligand (**Table S1**).

Known non-PPARγ modifying compounds were selected based on a literature search for

chemicals that influence adipogenesis but that are ligands for other nuclear receptors (e.g., aryl

hydrocarbon receptor, glucocorticoid receptor) or selected based on their structural dissimilarity

from known PPARγ ligands and lack of evidence of PPARγ activation (**Table S1**). The model

inference was based on an amended random forest procedure developed to better account for the

presence of biological replicates in the data. Specifically, for each classification tree, samples

and genes were bagged using sampling techniques consistent with (Breiman 2001). In particular,

samples were bootstrapped (i.e., sampled with replacement), and genes were subsampled by the

square root of the number of represented genes. To account for chemical-level variability and to

prevent replicates of the same chemical exposure from being separated, we implemented an extra

step, which we denote as "bag-merging", to summarize values from replicate samples of the

same chemical exposure after bootstrap sampling: within each "bag" of samples, replicates of the

same chemical exposure were merged to their mean expression. For prediction of test data,

replicates of the same chemical exposure were merged to their mean expression and run through

the trained random forest, such that each tree generates a vote of either 0 or 1, PPARγ-modifying

negative or positive, respectively. The mean of these votes across all trees is a value between 0

and 1, which can be interpreted as the pseudo-probability of the chemical exposure being

PPARγ-modifying.

Prior to generating the final predictive model, the expected performance of this

classification approach for predicting PPARγ-modification status with this data set was assessed

using 10-fold cross validation on the training set of known PPARγ-modification status. For each

fold, samples were stratified at the chemical exposure level, such that each fold included 6

distinct compounds, and all replicates of the same compound were only included in either the

training or the test folds. Next, prior to training each random forest model, gene filtering based

on between *vs.* within exposure variance using ANOVA was performed. Genes with an F-

statistics associated with an FDR corrected p-value > 0.05 were filtered out. Thresholds for

determining class membership based on voting was determined by running the training folds

through the random forest and selecting the threshold producing the highest F1-score, i.e., the

harmonic mean of precision and sensitivity. Performance was assessed in terms of area under the

curve (AUC), as well as precision, sensitivity, specificity, F1-score, and balanced accuracy, i.e.,

the mean of specificity and sensitivity. All random forests were generated using 2000 decision

trees.

The performance of this procedure was compared to three alternative random forest

strategies. In the first, which we denote as "pre-merging", the mean gene expression across

replicates was computed, and a classic random forest was applied to the classification of each

merged chemical profile. In the second, which we denote as "classic", replicate samples were

treated as independent perturbations and classified based on a classic random forest. Finally, for

the third, which we denote as "pooled", the mean of votes across replicates from the "classic"

strategy were used to estimate class membership per compound. To compare the classifiers'

performance, we repeated the 10-fold CV procedure 10-times to generate a distribution of

performance statistics for each strategy.

The final classification model used to predict the unlabeled chemicals was built using the

full training set of 59 labelled chemicals, including vehicle, and 1,199 genes after performing the

same ANOVA gene filtering approach used for cross validation as described above. The full list

of genes used in this model, including chemical-specific log fold-changes, F-statistics, nominal

P-values and FDR corrected P-values are provided in **Excel File 1**. The relative importance of

each gene in each random forest model was measured using the Gini importance, which

evaluates the mean decrease of label impurity when a particular gene is used to separate

chemicals in a given tree (Breiman 2001). Maximum purity is achieved (Gini=0) when all

chemicals within a branch have the same label.

Adipogen Clustering

Known and potential PPARγ activators/modifiers were clustered based on their test

statistics from univariate analysis comparing each chemical exposure or naïve pre-adipocyte

culture to vehicle using *limma* (v 3.34.9) (Ritchie et al. 2015). In order to assess taxonomic

differences between different exposure outcomes, a recursive unsupervised procedure, which we

denote as "*K2Taxonomer*" (Reed et al. 2020), was developed, whereby the set of PPARγ

modifying compounds underwent recursive partitioning into subgroups. At each iteration of the

procedure, the top 10% of genes was selected based on estimation of non-random changes in

gene expression across the current set or subset of compounds via the sum of squared test

statistics. Chemicals were then separated into two clusters based on their Euclidean distance,

using Ward's agglomerative method (Ward 1963) and the *cutree* R function. The procedure was

recursively applied to each of the two identified subgroups of chemicals and repeated until the

two-cluster split would result in a single chemical in the terminal subgroup. To obtain and

measure the most stable clusters, each iteration was bootstrapped 200 times by resampling gene-

level statistics with replacement. The most common clusters were used, and the proportion of

total bootstrapping iterations that included these identical clustering assignments was reported.

In order to derive gene-signatures of each split, differential analysis was performed to

compare chemicals from the two clusters at that split. In these models, biological replicate status

was accounted for using the duplicate correlation procedure using *limma* (v 3.34.9) (Ritchie et al.

2015). From these models, differential signatures were defined whereby genes were assigned to

one of four genesets based on two criteria: their differential expression between the two chemical

clusters at the split, and – within each chemical cluster – the differential expression between

chemicals and vehicle. In particular, for a particular gene, the difference between mean

expression between the two chemical clusters must have $|\log2(\text{Fold-Change})| > 1$ (i.e., Fold-

Change>2) and an FDR Q-value < 0.1. Each gene was then assigned to one-of-four genesets: 1)

genes with higher expression in the "left" chemical group vs. the "right" chemical group, and

with higher expression in the left chemical group vs. vehicle; 2) genes with higher expression in

the left chemical group vs. the right chemical group, and lower expression in the left chemical

group vs. vehicle; 3) genes with higher expression in the right chemical group vs. the left

chemical group, and with higher expression in the right chemical group vs. vehicle; 2) genes

with higher expression in the right chemical group vs. the left chemical group, and with lower

expression in the right chemical group vs. vehicle. Since the results of direct differential analysis

between the two chemical clusters are not indicative of overall up- or down-regulation, these

designations were determined based on the aggregate of the comparisons of each chemical or

naïve pre-adipocyte culture to vehicle. Specifically, a gene was assigned to a cluster based on

maximum absolute value of the mean of the chemical or naïve pre-adipocyte culture versus

vehicle derived test statistics used for clustering. Direction of regulation was then determined

based on the sign of the mean of these test statistics. Functional enrichment, comparing these

gene sets to independently annotated gene sets was carried out via Fisher's exact test. These

gene sets include those of the full set of Gene Ontology (GO) Biological Processes gene set

compendia downloaded from MSigDB, c5.bp.v6.2.symbols.gmt (Subramanian et al. 2005;

Liberzon et al. 2011), as well two gene sets derived from publicly available microarray

expression data from an experiment using mouse embryonic fibroblasts to compare wild-type

samples with mutant samples that do not undergo phosphorylation of PPARγ at Ser273, GEO

accession number GSE22033 (Choi et al. 2010). These additional gene sets were comprised of

genes, measured to be significantly higher or lower in expression (FDR Q-Value < 0.05) in

mutant samples, based on differential analysis of RMA normalized expression with *limma* (v

3.34.9). The full set of differential gene expression analysis and functional enrichment analysis

results, including the chemicals belonging to each subgroup, are provided in **Excel File 2**.

Human Transcriptome Analysis

To assess the human relevance of the gene signatures derived from the adipogen

clustering in chemical-exposed 3T3-L1 cells, we analyzed projections of these signatures onto

the transcriptional profiles from 770 subjects who were part of the Metabolic Syndrome in Men

(METSIM) study. Although comprised of only male subjects, the METSIM data set was chosen

for this analysis because it is the largest publicly available human subject data set, GEO

accession number GSE70353 (Civelek et al. 2017), which includes gene expression profiles from

subcutaneous adipose tissue, as well as a comprehensive set of cardio-metabolic measurements.

Affymetrix Human Genome U219 Array Microarray CEL files were annotated to unique Entrez

gene IDs, using a custom CDF file from BrainArray (HGU219_Hs_ENSG_22.0.0). Each of the

four possible gene sets derived from adipogen clustering were then projected on each of the 770

human transcriptome profiles using gene set variation analysis (GSVA), using *GSVA* (v 1.30.0)

(Hanzelmann et al. 2013), resulting in an enrichment score for each gene set and sample.

For each projected gene set, we tested for relationships between single-sample

enrichment scores and clinical measurements. Of notice, many of the clinical measurements are

correlated with each other, such that confounding is likely to generate many spurious results. To

overcome this problem, for each gene set projection, we tested the significance of the partial

correlation between single-sample enrichment scores and each of the clinical variables while

controlling for the remaining ones, including age, using *ppcor* (v 1.1) (Kim 2015). Given that

the relationships between single-sample enrichment scores and any one clinical variable are not

assumed to be linear, these partial correlations were calculated from Spearman correlation

estimates. P-values were adjusted across all combinations of gene set and clinical measurement.

Measurements with at least one comparison with a gene set projection yielding an FDR Q-value

< 0.1 are reported. To reduce expected redundancies in the measurements, of the 23 initial

quantitative measurements included in these data, we ran this analysis on a subset of 12

measurements (**Table S3**). Of note, fat free mass % was chosen in this subset of 12

measurements over body mass index because it has been shown to be better associated with risk

and presence of metabolic syndrome (Liu et al. 2013).

Results

Classification of novel taxonomic subgroups of adipogens

Potential adipogens (chemicals that change the differentiation and/or function of

adipocytes) were identified by review of the literature for reports of PPARγ agonism or

modulation of adipocyte differentiation or by the ToxPi designed to identify chemicals in the

ToxCast dataset that have potential to be PPARγ ligands/modifiers (**Table S1**). Our

classification groups were labeled as "Yes", "No", or "Potential", based on the chemical's

potential ability to act as a ligand of or modify PPARγ (i.e., to alter its post translational

modifications) as noted in the "PPARγ Ligand or Modifier" column in **Table S1**.

The classic mouse pre-adipocyte model, 3T3-L1 cells, was differentiated and treated with

vehicle (DMSO) or with each of the 76 test chemicals 4 times over the ten-day course of

differentiation (concentrations are reported in **Table S1**). In order to maximize the number of

chemicals that could be characterized, each chemical was tested at a single, maximal, non-toxic

dose. We also limited the maximum concentrations to 20 μM because concentrations above this

would not be reached in humans and because most (although not all) chemicals are not toxic at

or below 20 μM. Lipid accumulation was determined after 10 days. Lipid accumulation spanned

from significantly lower to significantly higher than vehicle-treated cells (**Figure 1**). **Excel File 3**

contains numerical data from all cell culture experiments. Lipid accumulation was highly

correlated with expression of adipocyte specific genes (e.g., *Cidec* and *Fabp4* (Danesch et al.

1992), **Figure S2**); therefore, we considered lipid accumulation a biomarker of adipocyte

differentiation in this system. Of the 27 chemicals that significantly induced lipid accumulation,

18 were known PPARγ ligands/modifiers and 9 were potential PPARγ ligands/modifiers (**Figure**

1, **Table S1**). Nine of the 17 potential adipogens generated significantly higher differentiation

than Vh-treated cells, including Alleth, Cyazo, Fenth, Fludiox, Quinox, TDCPP, Tebu, Tona, and

TTBP.

We turned to two other models to investigate why these differences from other studies

could be occurring. First, we differentiated the 3T3 L1 cells in the presence of the same suite of

chemicals, but in the absence of dexamethasone in the differentiation cocktail. The adipogenic

effects of steroids (e.g., melengesterol, cotisosterone, dexamethasone) were significant when

dexamethasone was not included in the cocktail (**Figure S3a**). Although, adipogenesis in the

presence and absence of 250 nM dexamethasone was moderately correlated overall (**Figure**

S3b). Second, we differentiated OP9 cells, a late preadipocyte model (Wolins et al. 2006), in the

presence of the same suite of chemicals. OP9 cells were more sensitive to differentiation

stimulated by RXR ligands (**Figure S4a**). However, adipocyte differentiation stimulated in OP9

cells was still moderately correlated with differentiation stimulated in 3T3 L1 cells differentiated

with 250 nM dexamethasone (**Figure S4b**).

Following the analysis of lipid accumulation, RNA was isolated from the cells, and the

transcriptome was characterized by RNA-Seq. The transcriptome data were then used to develop

a classification model to identify adipogens likely to act as PPARγ activators/modifiers. More

specifically, the *training set* of 59 chemicals with "Yes"/"No" labels was used to build the

classifier, which was then applied to the prediction of the *test set* of 17 "Potential" ligands. When

predicting PPARγ activation/modifier status ("Yes" *vs.* "No"), the mean area-under-the-curve

(AUC), precision, sensitivity, specificity, F1-score, and balanced accuracy from repeated 10-fold

cross validation (over the training set) of the random forest with bag merging procedure was

0.87, 0.85, 0.79, 0.76, 0.82, and 0.77, respectively (**Figure 2A**). We observed the most drastic

improvement of measured balanced accuracy, precision, and specificity by the bag merging

procedure compared to other assessed strategies (**Figure S5**). The full set of cross-validation

results for each performance metrics and all 10 repetitions can be found in **Excel File 1**. The first

two metrics in particular (AUC and precision) reflect expectation of relatively few false positive

results compared to the other strategies. In the final model fitted on the entire training set, the

voting threshold that produced the highest F1-score was 0.53. When we applied the classifier to

the test set of the 17 chemicals of unknown interaction with PPARγ, 13 had random forest votes

greater than this value (**Table 3**). Of these 13 compounds, four had votes > 0.88. These

chemicals included quinoxyfen, tonalide, allethrin, and fenthion. These compounds were

predicted as PPARγ activators/modifiers with high confidence and were selected for further

functional analyses. Of the 1,215 genes that passed ANOVA filtering and were included in the

random forest models, ribosomal protein L13 (*Rpl13*) and cell death Inducing DFFA Like

Effector C (*Cidec*) were measured to have the highest Gini importance (**Figure 2B**); *Rpl13* was

primarily expressed at a lower level and *Cidec* was primarily expressed at a higher level in cells

treated with known PPARγ activators/modifiers (**Figure S6**). Gene specific ANOVA filtering

results and importance estimates, as well as out-of-bag voting estimates for each compound in

the training set can be found in **Excel File 1**.

Adipogen Taxonomy Discovery

The taxonomy derived by the *K2Taxonomer* (Reed et al. 2020) procedure recapitulated

many known characteristics shared by adipogens included in this study (**Figure 3**). For example,

three terminal subgroups were labelled in **Figure 3** based on their shared characteristics. These

include: flame retardants (tetrabromobisphenol A (TBBPA) and triphenyl phosphate (TPhP)),

phthalates (MBUP, MEHP, MBZP, and BBZP), and RXR agonists (TBT and LG268).

Interestingly, we observed two subgroups containing all of the four thiazolidinediones, with

rosiglitazone (Rosig) segregating with the non-thiazolidinedione S26948 and pioglitazone, MCC

555, and troglitazone segregating together. Comprehensive functional annotation of the gene

signatures and enriched pathways for comparisons between sub-groups at each split can be found

in **Excel File 2**.

All of these terminal subgroups fell within a larger module containing 25 chemicals,

highlighted by expression patterns consistent with higher adipogenic activity including up-

regulation of genes significantly enriched in pathways involved in adipogenesis and lipid

metabolism (Soukas et al. 2001). In addition, these chemicals also demonstrated consistent

down-regulation of extracellular component genes. This effect was strongest in cells exposed to

thiazolidinediones and flame retardants, two classes of chemicals well-described to be strong

PPARγ agonists (Berger et al. 1996; Fang et al. 2015a; Riu et al. 2011). The subgroup of

thiazolidinediones, including S26948, was characterized by up-regulation of genes involved in

beta-oxidation, the process by which fatty acids are metabolized.

The gene expression profiles of the remaining 17 chemicals, including naïve pre-

adipocytes, demonstrated markedly less up-regulation of genes regulated by PPARγ. Of these 17

perturbations, a subgroup of 8 chemicals (BADGE, PrPar, 15dPGJ2, SR1664, METBP, DINP,

BuPA, and Fenth) included the reference vehicle signature. Compared to the next closest

subgroup, expression profiles of these compounds were characterized by up-regulation of

adipogenesis-related pathways indicative of modest PPARγ agonism. Additionally, a subgroup

comprised of 9CRA, DBT, LG754, and ATRA exposures and naïve pre-adipocyte signatures

was characterized by down-regulation of genes involved in adipogenesis and lipid metabolism,

indicating repression of PPARγ activity. Interestingly, both protectin D1 (Prote) and resolvin E1

(Resol) clustered closely in a subgroup with the cyclin-dependent kinase (CDK) inhibitor,

roscovitine (Rosco), which is known to induce insulin sensitivity and brite adipogenesis (Wang

et al. 2016).

In summary, our top-down taxonomy discovery approach identified subgroups of

adipogens characterized by differential transcriptomic activity at each split. Annotation of these

transcriptomic signatures revealed clear differences in the set and magnitude of perturbations to

known adipocyte biological processes by subgroups of chemicals. Membership of these

subgroups confirmed many expectations, such as subgroups comprised solely of phthalates,

thiazolidinediones, or flame retardants

Relationship between taxonomic subgroups and the human adipose transcriptome

Given that our taxonomic clustering was based on adipogen exposures in a mouse model,

we sought to establish its relevance to the relationship between human adipose tissue function

and markers of cardio-metabolic health. To this end, we projected the gene signatures derived as

either significantly increased or decreased gene sets in specific taxonomic subgroups onto a

publicly available clinical data set of gene expression profiles from subcutaneous adipose tissue

of 770 male subjects (Civelek et al. 2017) and assessed the relationship of these projections to a

set of 12 clinical measurements. **Figure 4A-B** shows the partial correlation between plasma

adiponectin and projection of gene sets, either increased or decreased in specific subgroups

(further details are provided in **Excel File 4**). **Figure 4C** shows these relationships for the

remaining measurements which demonstrated a statistically significant partial correlation for at

least one projection, FDR Q-value < 0.10. For positive associations, i.e. in the same direction,

the higher expressed gene sets had positive partial correlation measurements, while the lower

expressed gene sets had negative partial correlation measurements. The opposite was true for

negative associations.

We observed concordant results for putative markers of metabolic health, plasma

adiponectin and fat free mass %, which were positively associated with projection of gene

signatures from two terminal subgroups which include all TZD and non-TZD type 2 diabetic

drugs, Trogl, MCC555, Piogl, Rosig, and S26948, as well as a terminal subgroup which includes

Rosco, Resol, and Protec. Similarly, we observed concordant results for putative markers of

metabolic dysfunction, interleukin-1 receptor antagonist and fasting plasma insulin, which

included a primary subgroup comprised of terminal subgroups characterized by weak PPARγ

agonism, PPARγ modification, or PPARγ activity repression.

Taken together, these results confirm the ability of our mouse-based, in vitro-derived

signatures of capturing salient functional aspects of healthy and unhealthy metabolic functions in

human subjects.

Investigation of the white and brite adipocyte taxonomy

We aimed to better assess how the distinction between gene expression patterns

translated into functional differences in the induced adipocytes. Therefore, we selected chemicals

from representative groups of related adipogens for genotypic and phenotypic characterization.

We compared a strong PPARγ therapeutic agonist that also modifies PPARγ phosphorylation

(Rosig) (Choi et al., 2010), a chemical that modifies only PPARγ phosphorylation (Rosco)

(Wang et al., 2016), a weak PPARγ agonist and endogenous molecule (15dPGJ2) (Foreman et

al., 1995) and two known environmental PPARγ ligands (TBBPA and TPhP) (Chappell et al.

2018; Pillai et al. 2014). Protocols for differentiating 3T3 L1 cells are quite variable (reviewed

and tested in Zhao et al. 2019), and there was some concern that the use of 250 nM

dexamethasone would influence the differentiation process; therefore, we tested the effect of the

ligands on gene expression in media with two dexamethasone concentrations (1 and 250 nM).

3T3-L1 cells were differentiated and treated with vehicle (DMSO), Rosig (positive control),

Rosco, 15dPGJ2, TBBPA and TPhP. Cell density and gene expression were determined after 10

days. Rosig significantly increased cell number, while the other chemicals had no effect, positive

or negative (**Figure S7**).

We determined if changes in gene expression correlated with expression of genes

previously shown to be associated with white and brite/brown adipocytes (Qiang et al. 2012). As

expected, Rosig, TBBPA, and TPhP significantly induced *Pparg2, Plin1, Fabp4* and *Cidec*

expression in both dexamethasone conditions, while 15dPGJ2 only induced expression of these

genes with 250 nM dexamethasone (**Figure 5A, Figure S8A**). *Pparg2* expression was

moderately correlated between the two dexamethasone conditions (Spearman's r = 0.06786, p =

0.0469), while *Plin1*, *Fabp4* and *Cidec* were highly correlated between conditions (Spearman's r

= 0.9286, p = 0.013; r = 0.8929, p = 0.0034; r = 0.9524, r < 0.0001, respectively). For genes

expressed in brite adipocytes, a different pattern emerged, with little correlation between the two

dexamethasone conditions. *Adipoq* is expressed by both white and brown adipocyte depots; it

was induced by Rosig, TBBPA, and TPhP in both dexamethasone conditions. 15dPGJ2 and

Rosco only induced *Adipoq* expression with 250 nM dexamethasone (**Figure 5A, Figure S8A**).

For brite adipocyte genes, Rosig was the only ligand that induced expression of *Ppara*, *Pgc1a*,

Elovl3, *Cidea*, *Ucp1*, *Acaa2* in both dexamethasone conditions (**Figure 5B**, **Figure S8B**). Rosco

also induced expression of the brite adipocyte genes, but only in the 250 nM dexamethasone

condition (**Figure S8B**). TBBPA and TPhP induced expression of *Ppara*, *Elovl3*, *Cidea*, *Acaa2*

in low dexamethasone conditions only and did not induce expression of *Pgc1a* or *Ucp1* in either

dexamethasone condition (**Figure 5B**, **Figure S8B**).

Next, we determined if changes in gene expression correlated with changes in adipocyte

function. 3T3-L1 cells were differentiated in the presence of 1 nM dexamethansone and treated

as described for the mRNA expression analyses. Compared to vehicle-treated cells, Rosig and

TPhP significantly induced the rate of fatty acid uptake (**Figure 6A**) and Rosig, TPhP and

TBBPA significantly increased total lipid accumulation (**Figure 6B**). Lipid accumulation was

highly correlated with adiponectin secretion (**Figure 7**, Spearman's r = 0.9565, p < 0.0001).

Overall, lipid accumulation was highly correlated with expression of all genes examined, with

the exception of *Pgc1a* and *Ucp1* (**Figure S9**). Only Rosig significantly induced mitochondrial

biogenesis (**Figure 8**). Rosig also significantly increased cellular respiration (**Figure 9**). Rosco,

15dPGJ2, TBBPA and TPhP did not change cellular respiration.

*** Identification of adipogens that favor white adipogenesis***

653   Quinoxyfen (Quino) and tonalide (Tonal) were two of the environmental chemicals that

received the highest PPARγ ligand/modifier vote and segregated distinctly from the therapeutic

ligands (**Table 2**). Thus, we tested the hypothesis that Quino and Tonal are adipogens that do not

induce gene expression or metabolic phenotypes indicative of high energy expenditure or brite

adipogenesis. 3T3-L1 cells were differentiated in the presence of 1 nM dexamethasone and

treated with vehicle (DMSO), rosiglitazone (positive control), Quino, or Tonal. These

concentrations were determined to be non-toxic with only Rosig increasing cell number (**Figure**

S10A). In 3T3-L1 cells, Rosig and Quino significantly increased the rate of fatty acid uptake, but

Rosig, Quino and Tonal all significantly increased total lipid accumulation (**Figure 10A**). Rosig

and Quino, but not Tonal, significantly increased adiponectin secretion (**Figure 10B**).

Interestingly, only Quino significantly increased the expression of *Pparg2*. Both Rosig and

Quino significantly increased expression of *Plin1*, *Cidec*, and *Cidea*, while Tonal had little

efficacy in inducing expression of adipocyte-specific genes (**Figure 10C**). While both Rosig and

Quino significantly upregulated expression of *Cidea*, only Rosig significantly induced *Pgc1a* and

Ucp1 (**Figure 10D**). In spite of both Rosig and Quino both increasing mitochondrial biogenesis

(**Figure 10E**), only Rosig significantly increased cellular respiration (**Figure 10F**).

669   Last, we investigated whether results in our mouse model, 3T3-L1 cells, could be

recapitulated in a human model. Primary, human subcutaneous preadipocytes were differentiated

and treated with vehicle (DMSO), rosiglitazone (positive control), Quino, or Tonal. Quino and

Tonal significantly induced lipid accumulation (**Figure 11A**). Tonal, but not Quino, increased

cell number (**Figure S10B**). Both Quino and Tonal induced *CIDEC* expression but failed to

induce *CIDEA* expression (**Figure 11B**). In contrast to 3T3-L1 cells, Quino and Tonal did not

increase fatty acid uptake over that induced by the hormonal cocktail in the differentiated

primary human adipocytes (**Figure 11C**). Quino and Tonal did not induce mitochondrial

biogenesis (**Figure 11D**).

Discussion

The chemical environment has changed dramatically in the past 40 years, and an

epidemic increase in the prevalence of obesity has occurred over the same time period (Baillie-

Hamilton 2002). Yet, it is still unclear how chemical exposures may be contributing to adverse

metabolic health effects. New tools are needed not just to identify potential adipogens, but to

provide information on the type of adipocyte that is formed. Here, we have both developed a new

analytical framework for adipogen identification and characterization and tested its utility in

hypothesis generation. We showed that adipogens segregate based on distinct patterns of gene

expression. Furthermore, subgroups within the taxonomy containing therapeutics shared gene

expression patterns in human adipose tissue in accordance with positive markers of metabolic

health, specifically plasma adiponectin and fat free mass %, suggesting that these gene

expression are related to metabolic health in humans. We identified two environmental

contaminants segregating outside the therapeutic groups for hypothesis testing. Our results

support the conclusion that quinoxyfen and tonalide have a limited capacity to induce the health-

promoting effects of mitochondrial biogenesis and brite adipocyte differentiation.

Adipogen taxonomy identifies environmental chemicals that favor white adipogenesis

Potential adipogens (chemicals that change the differentiation and/or function of

adipocytes) were identified by review of the literature and based on reports of PPARγ agonism

or modulation of adipocyte differentiation, as well as through querying ToxCast data. Not all of

the chemicals previously identified as adipogens induced significant lipid accumulation.

Mono(2-ethylhexyl) phthalate (MEHP)(Feige et al. 2007), SR1664 (Choi et al. 2011), and 15-

deoxy-$\Delta^{12,14}$-prostaglandin J2 (15dPGJ2)(Forman et al. 1995) are PPARγ agonists that were

expected to increase adipocyte differentiation but did not. LG268 (Boehm et al. 1994) and TBT

are RXR agonists that were also expected to significantly increase adipocyte differentiation but

did not. T007 is a PPARγ antagonist (Lee et al. 2002) that was expected to decrease adipocyte

differentiation but did not. We hypothesize that this likely resulted from the fact that we did not

apply any chemical above 20 μM (with the exception of fenthion). Concentrations in the 100 μM

range have been used in previous studies (e.g. DOSS (Temkin et al. 2016); parabens (Hu et al.

2013); phthalates (Hurst and Waxman 2003; Feige et al. 2007)). The RXR agonists LG268 and

TBT also were expected to significantly increase adipocyte differentiation but did not.

We tested the chemicals in multiple differentiation scenarios (3T3-L1s with 0, 1 or 250

nM dexamethasone and in OP9 cells). Each differentiation scenario generated highly correlated

results, but with some deviations. 3T3-L1 cells are the most used model in the study of

adipogens (Green and Kehinde 1975; Ruiz-Ojeda et al. 2016), with a broad range of "standard"

differentiation cocktails. Not surprisingly, in 3T3 L1 cells differentiated in the absence of

dexamethasone, glucocorticoid receptor agonists significantly induced lipid accumulation. A

significant drawback of this model, however, is the variability of the efficacy of differentiation,

which has been shown to depend on the commercial source, passage number, culture dishes used

and medium volume used (Sheng et al. 2014; Kassotis et al. 2017). On the other hand, an

important feature of 3T3-L1 cells is their ability to differentiate into both white and brite

adipocytes (Vernochet et al. 2009). Thus, we suggest that the minimum amount of

dexamethasone to support efficient differentiation, without inducing significant background

differentiation, should be identified for a given lot of cells before proceeding with testing.

We have previously shown that TBT induced adipogenesis with greater efficacy in OP9

cells than in 3T3-L1 cells (Kassotis et al. 2017; Andrews et al., 2020), and also showed here that

OP9 cells more efficaciously responded to RXR ligands, in general, likely because they are more

committed to adipogenesis as late pre-adipocytes (Wolins et al., 2006) than 3T3 L1 cells, which

are early pre-adipocytes (Lane and Tang, 2005). OP9 are significantly less finicky than 3T3-L1

cells in terms of the experimental conditions required for efficient adipocyte differentiation,

likely because they are further along in commitment to adipogenesis than 3T3-L1 cells (Wolins

et al., 2006; Kassotis et al., 2017). However, they have not been tested for their ability to

differentiate into brite adipocytes.

We identified four compounds as high-confidence PPARγ ligands/modifiers: quinoxyfen,

tonalide, allethrin, and fenthion. In the final model used for these predictions, biologically

informative genes emerged as important for predicting PPARγ ligand/modification status,

specifically the down-regulation of *Rpl13* and the up-regulation of *Cidec. Rpl13* is involved in

ribosomal machinery and is down-regulated during human adipogensis (Marcon et al. 2017).

Cidec is a lipid droplet structural gene, the expression of which was positively correlated with

adipocyte lipid droplet size, insulin levels, and glycerol release (Ito et al. 2010). Of these four

compounds, quinoxyfen and tonalide are of particular public health concern. Quinoxyfen is

among a panel of pesticides with different chemical structures and modes of action (i.e.,

zoxamide, spirodiclofen, fludioxonil, tebupirimfos, forchlorfenuron, flusilazole, acetamaprid,

and pymetrozine) that induced adipogenesis and adipogenic gene expression in 3T3-L1 cells

(Janesick et al. 2016). Quinoxyfen is a fungicide widely used to prevent the growth of powdery

mildew on grapes (Duncan et al. 2018). We chose to test tonalide because it was reported to

strongly increase adipogenesis in 3T3-L1 cells, although it was concluded that this response was

not due to direct PPARγ activation (Pereira-Fernandes et al. 2013). Our results differed in this

regard. Tonalide bioaccumulates in adipose tissue of many organisms including humans, and

exposure is widespread because of its common use in cosmetics and cleaning agents (Kannan et

al. 2005). Combined, tonalide and galaxolide constitute 95% of the polycyclic musks used in the

EU market and 90% of that of the US market (HERA 2004).

Our results support the conclusion that quinoxyfen and tonalide are adipogenic

chemicals, likely to be acting through PPARγ. In clustering analysis, quinoxyfen and tonalide

were among the largest subgroup of eight potential strong PPARγ agonists. Notably, this cluster

included both synthetic/therapeutic (nTZDpa, tesaglitazar, telmisartan) and environmental

compounds (tributyl phosphate and triphenyltin) and was characterized by general up-regulation

of pathways of adipogenic activity. However, quinoxyfen and tonalide generated adipocytes that

were phenotypically distinct from adipocytes induced by therapeutics such as rosiglitazone. That

environmental PPARγ ligands can induce a distinct adipocyte phenotype has been shown

previously for TBT (Kim et al. 2018; Regnier et al. 2015; Shoucri et al. 2018) and TPhP (Kim et

al. 2020). Ultimately, binding assays or computational ligand binding domain modeling will be

needed to confirm that quinoxyfen and tonalide are PPARγ agonists.

We tested the effect of quinoxyfen and tonalide on human subcutaneous preadipocyte

differentiation. There was a strong association between abdominal adiposity and metabolic

syndrome (Cornier et al. 2008). Classically, visceral adipose tissue has been thought to be the

driver of metabolic dysfunction; however, there is an alternative explanation that visceral

adiposity results secondarily from the dysfunction of subcutaneous adipose tissue in the upper

body (Jensen 2008; Lee et al. 2017). Subcutaneous adipose tissue represents 85% of all body fat

(Frayn and Karpe 2014) and thus has a large overall capacity for generating brite adipocytes.

Additionally, lack of browning capacity of human subcutaneous adipocytes was associated with

insulin resistance in a group of male Swedish individuals (Yang et al. 2003). As hypothesized

based on the taxonomical analysis, quinoxyfen and tonalide induced white adipocyte functions

such as increased lipid accumulation, but in contrast to rosiglitazone, did not induce

mitochondrial biogenesis.

We hypothesize that the differences in adipocyte phenotype that are induced by

environmental adipogens and likely PPARγ ligands (e.g., TBBPA, TPhP, quinoxyfen, tonalide)

could result from the conformation that PPARγ assumes when liganded with these chemicals

rather than with therapeutic agents. Differences in conformation not only determine the efficacy

to which PPARγ is activated but also the transcriptional repertoire (Chrisman et al. 2018). We

have shown recently, for instance, that TPhP did not protect PPARγ from phosphorylation at

s273 and did not induce brite adipogenesis; however, when PPARγ cannot be phosphorylated at

s273, TPhP could induce brite adipogens (Kim et al. 2020). Structural analyses would need to be

conducted with each environmental ligand specifically to determine the contribution of

conformational changes to the biological activity.

Access to post-translational modification sites and coregulator binding surfaces depends

upon the structure that PPARγ assumes. The white adipogenic, brite/brown adipogenic and

insulin sensitizing activities of PPARγ are regulated separately through differential co-regulator

recruitment (Villanueva et al. 2013) and post-translational modifications (Choi et al. 2010; Choi

787 et al. 2011), with ligands having distinct abilities to activate each of PPARγ's functions. Suites of

788 genes have been shown to be specifically regulated by the acetylation status of PPARγ (SirT1-

789 mediated) (Qiang et al. 2012), by the phosphorylation status of PPARγ (ERK/MEK/CDK5-

790 mediated) (Choi et al. 2010; Wang et al. 2016) and/or by the recruitment of Prdm16 to PPARγ

791 (Seale et al. 2007). Future work will investigate the connections between the phosphorylation

792 status of PPARγ liganded with environmental PPARγ ligands, the recruitment and release of

793 coregulators, and the ability of PPARγ to recruit transcriptional machinery to specific DNA-

794 binding sites. It will be important to determine the metabolic effects of chemicals like TPhP,

795 quinoxyfen and tonalide *in vivo*.

796 ***Analytical approaches for adipogen characterization***

797  In this study, we performed a high-throughput, cost-effective transcriptomic screening to

798 profile adipocytes formed from 3T3-L1 preadipocytes exposed to a panel of compounds of

799 known and unknown adipogenic impact. Common to toxicogenomic projects, this panel-based

800 study design allowed for characterization of the extent to which each chemical modified

801 differentiation (in this case, adipogenesis as related to the change in lipid accumulation). It also

802 supported the exploration of how subsets of chemicals influence multiple biological processes

803 that determine the functional status of a cell (in this case, processes that determine white *vs*. brite

804 adipogenesis). Exploration of these biological processes allowed for the prediction of the

805 phenotypic impact of previously unclassified compounds, as well as for the characterization of

806 the heterogeneity of the cellular activity of compounds with similar known phenotypic impact.

807 Here, we performed both types of analyses: first through the implementation and application of

808 random forest classification models to identify potential adipogens, and second via the recursive

clustering of the data to identify and characterize taxonomic subgroups of known and potential

PPARγ ligands/modifiers.

For both analyses, we introduced amendments to commonly used machine learning

procedures, to improve accuracy and resolution of the acquired result. For the classification task,

we amended the random forest algorithm to tailor it to study designs typically adopted in

toxicogenomic projects (see Methods). With the addition of an extra step to average the

expression across replicates of the bootstrapped samples, we observed consistently higher

performance across conventional metrics than with the standard algorithm. For the clustering

task, we employed a procedure where we recursively divide sets of chemicals into two subgroups

and assessed the robustness of each division, as well as annotated transcriptional drivers of each

division. As a result, we were not limited to interpreting the clustering results as mutually

exclusive groups, but rather as a taxonomy of subgroups where sets of compounds shared some

transcriptional impact and differ in others, as was expected given the dynamic nature of the

modifications by which compounds directly and indirectly affect PPARγ activity.

Future work will generalize random forest method to incorporate more complex study

designs. To this end, the classification approach adopted in this project is being developed as a

random forest software tool soon to be made available as an R package, allowing for the

interchanging independent functions at different steps of the algorithm. The strength and utility

of this approach extends beyond toxicogenomic studies, and can be used in a variety of

applications of high-throughput screening, including drug discovery, such as the Connectivity

Map (CMAP) (Subramanian et al. 2017), and longitudinal molecular epidemiology studies, such

as the Framingham Heart Study (Mahmood et al. 2014).

Adipogen portal

Given the breadth of results generated by this analysis, our description here is far from

exhaustive.   As such, we have created an interactive website

(https://montilab.bu.edu/adipogenome/) to support the interactive exploration of these results at

both the gene and pathway-level. A help page illustrating how to navigate this web portal is

provided on the website as well. The portal is built around a point-and-click dendrogram of the

clustering results as in **Figure 3**. Selecting a node of this dendrogram will populate the rest of

the portal with the chemical lists, differential analysis, and pathway level hyper-enrichment

results for each subgroup defined by a split. For instance, selecting node "H" will show the

chemicals in each subgroup to the right (Group 1 = Honokiol, T007907; Group 2 = Prote, Resol,

and Rosco), as well as the differential gene signature for each group below. Selecting *Cidec*, the

top gene in the Group 2 signature, displays hyper-enrichment results for gene sets which include

Cidec and have a nominal p-value < 0.50. The hyper-enrichment results for all genes can be

found below this table. Finally, selecting a gene set name will display the gene set members at

the bottom frame of the portal, with gene hits in bold. All tables are query-able and

downloadable.

<u>Conclusions</u>

Emerging data implicate contributions of environmental metabolism-disrupting

chemicals to perturbations of pathways related to metabolic disease pathogenesis, such as

disruptions in insulin signaling and mitochondrial activity. There is still a gap in identifying and

examining how environmental chemicals can act as obesity-inducing and metabolism-disrupting

chemicals. Our implementation of novel strategies for classification and taxonomy development

can help identify environmental chemicals that are acting on PPARγ. Further, our approach

provides a basis from which to investigate effects of adipogens on not just the generation of

adipocytes, but potentially pathological changes in their function. To this end, we have shown

how two environmental contaminants, quinoxyfen and tonalide, are inducers of white

adipogenesis. Finally, we emphasize that these approaches are generally suitable for analyses of

transcriptomic data generated from toxicogenomic screening studies. For example, provided such

data, *K2Taxonomer* (Reed et al. 2020) can be applied to characterize molecular patterns driving

segregation of experimental groups and infer dysregulated biological activity inherent to

subgroups of experimental groups, thereby providing insight to inform Adverse Outcome

Pathway (AOP) models of response to chemical exposures (Brockmeier et al. 2017).

References:

Andrews, F. V., Kim, S. M., Edwards, L., & Schlezinger, J. J. 2020. Identifying adipogenic
chemicals: Disparate effects in 3T3-L1, OP9 and primary mesenchymal multipotent cell models.
Toxicology in Vitro, 67: 104904.

Auerbach S, Filer D, Reif D, Walker V, Holloway AC, Schlezinger J, et al. 2016. Prioritizing
environmental chemicals for obesity and diabetes outcomes research: A screening approach
using toxcast high-throughput data. Environ Health Perspect 124:1141-1154.

Baillie-Hamilton PF. 2002. Chemical toxins: a hypothesis to explain the global obesity epidemic.
J Altern Complement Med 8(2):185-92.

Banks AS, McAllister FE, Camporez JP, Zushin PJ, Jurczak MJ, Laznik-Bogoslavski D, et al.
2015. An erk/cdk5 axis controls the diabetogenic actions of ppargamma. Nature 517:391-395.

Berger J, Bailey P, Biswas C, Cullinan CA, Doebber TW, Hayes NS, et al. 1996.
Thiazolidinediones produce a conformational change in peroxisomal proliferator-activated
receptor-gamma: Binding and activation correlate with antidiabetic actions in db/db mice.
Endocrinology 137:4189-4195.

Boehm MF, Zhang L, Badea BA, White SK, Mais DE, Berger E, Suto CM, Goldman ME,
Heyman RA. 1994. Synthesis and structure-activity relationships of novel retinoid X receptor-
selective retinoids. J Med Chem 37:2930-41.

Breiman L. 2001. Random forests. Machine Learning
45:https://doi.org/10.1023/A:1010933404324.

Brockmeier EK, Hodges G, Hutchinson TH, Butler E, Hecker M, Tollefsen KE, Garcia-Reyero
N, Kille P, Becker D, Chipman K, Colbourne J, Collette TW, Cossins A, Cronin M, Graystock P,
Gutsell S, Knapen D, Katsiadaki I, Lange A, Marshall S, Owen SF, Perkins EJ, Plaistow S,
Schroeder A, Taylor D, Viant M, Ankley G, Falciani F. 2017. The Role of Omics in the
Application of Adverse Outcome Pathways for Chemical Risk Assessment. Toxicol Sci.
158:252-262.

Burgermeister E, Schnoebelen A, Flament A, Benz J, Stihle M, Gsell B, et al. 2006. A novel
partial agonist of peroxisome proliferator-activated receptor-gamma (ppargamma) recruits
ppargamma-coactivator-1alpha, prevents triglyceride accumulation, and potentiates insulin
signaling in vitro. Mol Endocrinol 20:809-830.

Cano-Sancho G, Smith A, La Merrill MA. 2017. Triphenyl phosphate enhances adipogenic
differentiation, glucose uptake and lipolysis via endocrine and noradrenergic mechanisms.
Toxicol In Vitro 40:280-288.

Chappell, V. A., Janesick, A., Blumberg, B., & Fenton, S. E. 2018. Tetrabromobisphenol-A
Promotes Early Adipogenesis and Lipogenesis in 3T3-L1 Cells. Toxicological Sciences, 166(2),
332–344.

Choi JH, Banks AS, Estall JL, Kajimura S, Bostrom P, Laznik D, et al. 2010. Anti-diabetic drugs
inhibit obesity-linked phosphorylation of ppargamma by cdk5. Nature 466:451-456.

Choi JH, Banks AS, Kamenecka TM, Busby SA, Chalmers MJ, Kumar N, et al. 2011.
Antidiabetic actions of a non-agonist ppargamma ligand blocking cdk5-mediated
phosphorylation. Nature 477:477-481.

Chrisman IM, Nemetchek MD, de Vera IMS, Shang J, Heidari Z, Long Y, et al. 2018. Defining a
conformational ensemble that directs activation of ppargamma. Nat Commun 9:1794.

Civelek M, Wu Y, Pan C, Raulerson CK, Ko A, He A, et al. 2017. Genetic regulation of adipose
gene expression and cardio-metabolic traits. Am J Hum Genet 100:428-443.

Claussnitzer M, Dankel SN, Kim KH, Quon G, Meuleman W, Haugen C, et al. 2015. Fto obesity
variant circuitry and adipocyte browning in humans. The New England journal of medicine
373:895-907.

Cornier MA, Dabelea D, Hernandez TL, Lindstrom RC, Steig AJ, Stob NR, et al. 2008. The
metabolic syndrome. Endocr Rev 29:777-822.

Danesch U, Hoeck W, Ringold GM. 1992. Cloning and transcriptional regulation of a novel
adipocyte-specific gene, fsp27. Caat-enhancer-binding protein (c/ebp) and c/ebp-like proteins
interact with sequences required for differentiation-dependent expression. J Biol Chem
267:7185-7193.

Duncan H, Abad-Somovilla A, Abad-Fuentes A, Agullo C, Mercader JV. 2018.
Immunochemical rapid determination of quinoxyfen, a priority hazardous pollutant.
Chemosphere 211:302-307.

Fang M, Webster TF, Ferguson PL, Stapleton HM. 2015a. Characterizing the peroxisome
proliferator-activated receptor (ppargamma) ligand binding potential of several major flame
retardants, their metabolites, and chemical mixtures in house dust. Environ Health Perspect
123:166-172.

Fang M, Webster TF, Stapleton HM. 2015b. Effect-directed analysis of human peroxisome
proliferator-activated nuclear receptors (ppargamma1) ligands in indoor dust. Environ Sci
Technol 49:10065-10073.

Farmer SR. 2006. Transcriptional control of adipocyte formation. Cell metabolism 4:263-273.

Feige JN, Gelman L, Rossi D, Zoete V, Metivier R, Tudor C, et al. 2007. The endocrine disruptor
monoethyl-hexyl-phthalate is a selective peroxisome proliferator-activated receptor gamma
modulator that promotes adipogenesis. J Biol Chem 282:19152-19166.

Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. 2016. Trends in obesity
amoung adults in the United States 2005-2014. JAMA 315:2284-91.

Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, Evans RM. 1995. 15-Deoxy-delta
12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. Cell
83:803-12.

Frayn KN, Karpe F. 2014. Regulation of human subcutaneous adipose tissue blood flow.
International journal of obesity (2005) 38:1019-1026.

GBD. 2017. Health effects of overweight and obesity in 195 countries over 25 years. The New
England journal of medicine 377:13-27.

Green H, Kehinde O. (1975). An established preadipose cell line and its differentiation in culture.
II. Factors affecting the adipose conversion. Cell, 5(1), 19–27.

Gumbilai V, Ebihara K, Aizawa-Abe M, Ebihara C, Zhao M, Yamamoto Y, et al. 2016. Fat mass
reduction with adipocyte hypertrophy and insulin resistance in heterozygous ppargamma mutant
rats. Diabetes 65:2954-2965.

Hanzelmann S, Castelo R, Guinney J. 2013. Gsva: Gene set variation analysis for microarray and
rna-seq data. BMC Bioinformatics 14:7.

He W, Barak Y, Hevener A, Olson P, Liao D, Le J, et al. 2003. Adipose-specific peroxisome
proliferator-activated receptor gamma knockout causes insulin resistance in fat and liver but not
in muscle. Proc Natl Acad Sci U S A 100:15712-15717.

Heindel JJ, Blumberg B, Cave M, Machtinger R, Mantovani A, Mendez MA, et al. 2017.
Metabolism disrupting chemicals and metabolic disorders. Reproductive toxicology (Elmsford,
NY 68:3-33.

HERA. 2004. Human & environmental risk assessment on ingredients of household cleaning
products: Polycyclic musks. Available: https://www.heraproject.com/files/29-E-
04_pcm_HHCB_AHTN_HERA_Environmenta_DISCLed26.pdf [accessed Dec. 27 2018].

Hiromori Y, Ido A, Aoki A, Kimura T, Nagase H, Nakanishi T. 2016. Ligand activity of group
15 compounds possessing triphenyl substituent for the rxr and ppargamma nuclear receptors.
Biol Pharm Bull 39:1596-1603.

Hu P, Chen X, Whitener RJ, Boder ET, Jones JO, Porollo A, et al. 2013. Effects of parabens on
adipocyte differentiation. Toxicol Sci 131:56-70.

Hurst CH, Waxman DJ. 2003. Activation of ppara and pparg by environmental phthalate
monoesters. Toxicol Appl Pharmacol.

Ito M, Nagasawa M, Hara T, Ide T, Murakami K. 2010. Differential roles of cidea and cidec in
insulin-induced anti-apoptosis and lipid droplet formation in human adipocytes. Journal of lipid
research 51:1676-1684.

Janesick AS, Dimastrogiovanni G, Vanek L, Boulos C, Chamorro-Garcia R, Tang W, et al. 2016.
On the utility of toxcast and toxpi as methods for identifying new obesogens. Environ Health
Perspect 124:1214-1226.

Jensen MD. 2008. Role of body fat distribution and the metabolic complications of obesity. J
Clin Endocrinol Metab 93:S57-63.

Jiang Y, Berry DC, Tang W, Graff JM. 2014. Independent stem cell lineages regulate adipose
organogenesis and adipose homeostasis. Cell Rep 9:1007-1022.

Kannan K, Reiner JL, Yun SH, Perrotta EE, Tao L, Johnson-Restrepo B, et al. 2005. Polycyclic
musk compounds in higher trophic level aquatic organisms and humans from the united states.
Chemosphere 61:693-700.

Kassotis CD, Masse L, Kim S, Schlezinger JJ, Webster TF, Stapleton HM. 2017.
Characterization of adipogenic chemicals in three different cell culture systems: Implications for
reproducibility based on cell source and handling. Sci Rep 7:42104.

Kavlock R, Chandler K, Houck K, Hunter S, Judson R, Kleinstreuer N, et al. 2012. Update on
epa's toxcast program: Providing high throughput decision support tools for chemical risk
management. Chem Res Toxicol 25:1287-1302.

Kim S. 2015. ppcor: An R Package for a Fast Calculation to Semi-partial Correlation
Coefficients. Commun Stat Appl Methods 22:665–674; doi:10.5351/CSAM.2015.22.6.665.

Kim S, Li A, Monti S, Schlezinger JJ. 2018. Tributyltin induces a transcriptional response
without a brite adipocyte signature in adipocyte models. Arch Toxicol 92:2859-2874.

Kim S, Rabhi N, Blum B, Hekman R, Wynne K, Emili A, et al. 2020. Triphenyl phosphate is a
selective pparγ modulator that does not induce brite adipogenesis in vitro and in vivo. Arch
Toxicol. 94:3087-3103.

Lane, M. D., & Tang, Q. Q. 2005. From multipotent stem cell to adipocyte. *Birth Defects Res A*
Clin Mol Teratol, *73*(7), 476–477.

Lee G, Elwood F, McNally J, Weiszmann J, Lindstrom M, Amaral K, Nakamura M, Miao S,
Cao P, Learned RM, Chen JL, Li Y. 2002. T0070907, a selective ligand for peroxisome
proliferator-activated receptor gamma, functions as an antagonist of biochemical and cellular
activities. J Biol Chem 277:19649-57.

Lee JJ, Pedley A, Therkelsen KE, Hoffmann U, Massaro JM, Levy D, et al. 2017. Upper body
subcutaneous fat is associated with cardiometabolic risk factors. Am J Med 130:958-966 e951.

Lee MJ, Fried SK. 2014. Optimal protocol for the differentiation and metabolic analysis of
human adipose stromal cells. Methods Enzymol 538:49-65.

Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. 2012. The sva package for removing
batch effects and other unwanted variation in high-throughput experiments. Bioinformatics
(Oxford, England) 28:882-883.

le Maire A, Grimaldi M, Roecklin D, Dagnino S, Vivat-Hannah V, Balaguer P, Bourguet W.
2009. Activation of RXR-PPAR heterodimers by organotin environmental endocrine disruptors.
EMBO Rep 10:367-73.

Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP. 2011.
Molecular signatures database (MSigDB) 3.0. Bioinformatics 27:1739–1740.

Liu, P., Ma, F., Lou, H., & Liu, Y. 2013. The utility of fat mass index vs. body mass index and
percentage of body fat in the screening of metabolic syndrome. BMC Public Health, 13, 629.

Mahmood SS, Levy D, Vasan RS, Wang TJ. 2014. The framingham heart study and the
epidemiology of cardiovascular disease: A historical perspective. Lancet 383:999-1008.

Marcon BH, Holetz FB, Eastman G, Origa-Alves AC, Amoros MA, de Aguiar AM, et al. 2017.
Downregulation of the protein synthesis machinery is a major regulatory event during early
adipogenic differentiation of human adipose-derived stromal cells. Stem Cell Res 25:191-201.

O'Donnell PE, Ye XZ, DeChellis MA, Davis VM, Duan SZ, Mortensen RM, et al. 2016.
Lipodystrophy, diabetes and normal serum insulin in ppargamma-deficient neonatal mice. PloS
one 11:e0160636.

Ohno H, Shinoda K, Spiegelman BM, Kajimura S. 2012. Ppargamma agonists induce a white-to-
brown fat conversion through stabilization of prdm16 protein. Cell metabolism 15:395-404.

Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. 2003. The metabolic syndrome: Prevalence and associated risk factor findings in the us population from the third national health and nutrition examination survey, 1988-1994. Arch Intern Med 163:427-436.

Pereira-Fernandes A, Demaegdt H, Vandermeiren K, Hectors TL, Jorens PG, Blust R, et al. 2013. Evaluation of a screening system for obesogenic compounds: Screening of endocrine disrupting compounds and evaluation of the ppar dependency of the effect. PloS one 8:e77481.

Pereira-Fernandes A, Vanparys C, Vergauwen L, Knapen D, Jorens PG, Blust R. 2014. Toxicogenomics in the 3t3-l1 cell line, a new approach for screening of obesogenic compounds. Toxicol Sci 140:352-363.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time rt-pcr. Nucleic acids research 29:e45.

Qiang L, Wang L, Kon N, Zhao W, Lee S, Zhang Y, et al. 2012. Brown remodeling of white adipose tissue by sirt1-dependent deacetylation of ppargamma. Cell 150:620-632.

Reed ER, Monti S. 2020. Multi-resolution characterization of molecular taxonomies in bulk and single-cell transcriptomics data. bioRxiv; doi:10.1101/2020.11.05.370197.

Regnier SM, El-Hashani E, Kamau W, Zhang X, Massad SL, Sargis RM. 2015. Tributyltin differentially promotes development of a phenotypically distinct adipocyte. Obesity (Silver Spring, Md In press.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. 2015. Limma powers differential expression analyses for rna-sequencing and microarray studies. Nucleic acids research 43:e47.

Riu A, Grimaldi M, le Maire A, Bey G, Phillips K, Boulahtouf A, et al. 2011. Peroxisome proliferator-activated receptor gamma is a target for halogenated analogs of bisphenol a. Environ Health Perspect 119:1227-1232.

Rosen ED, Spiegelman BM. 2006. Adipocytes as regulators of energy balance and glucose homeostasis. Nature 444:847-853.

Ruiz-Ojeda, F. J., Ruperez, A. I., Gomez-Llorente, C., Gil, A., & Aguilera, C. M. 2016. Cell Models and Their Application for Studying Adipogenic Differentiation in Relation to Obesity: A Review. Int J Mol Sci, 17(7): 1040.

Schwartz MW, Seeley RJ, Zeltser LM, Drewnowski A, Ravussin E, Redman LM, et al. 2017. Obesity pathogenesis: An endocrine society scientific statement. Endocr Rev 38:267-296.

Seale P, Kajimura S, Yang W, Chin S, Rohas LM, Uldry M, et al. 2007. Transcriptional control of brown fat determination by prdm16. Cell metabolism 6:38-54.

Sheng X, Tucci J, Malvar J, Mittelman SD. 2014. Adipocyte differentiation is affected by media height above the cell layer. Int J Obesity (Lond) 38(2):315-20.

Shoucri BM, Hung VT, Chamorro-Garcia R, Shioda T, Blumberg B. 2018. Retinoid x receptor activation during adipogenesis of female mesenchymal stem cells programs a dysfunctional adipocyte. Endocrinology 159:2863-2883.

Sidossis L, Kajimura S. 2015. Brown and beige fat in humans: Thermogenic adipocytes that control energy and glucose homeostasis. J Clin Invest 125:478-486.

Soukas A, Socci ND, Saatkamp BD, Novelli S, Friedman JM. 2001. Distinct transcriptional
profiles of adipogenesis in vivo and in vitro. J Biol Chem 276:34167-34174.

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. 2005. Gene
set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression
profiles. Proc Natl Acad Sci U S A 102:15545-15550.

Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, et al. 2017. A next
generation connectivity map: L1000 platform and the first 1,000,000 profiles. Cell 171:1437-
1452 e1417.

Takacs ML, Abbott BD. 2007. Activation of mouse and human peroxisome proliferator-activated
receptors (alpha, beta/delta, gamma) by perfluorooctanoic acid and perfluorooctane sulfonate.
Toxicol Sci 95:108-117.

Temkin AM, Bowers RR, Magaletta ME, Holshouser S, Maggi A, Ciana P, et al. 2016. Effects of
crude oil/dispersant mixture and dispersant components on ppargamma activity in vitro and in
vivo: Identification of dioctyl sodium sulfosuccinate (doss; cas #577-11-7) as a probable
obesogen. Environ Health Perspect 124:112-119.

Timmons JA, Pedersen BK. 2009. The importance of brown adipose tissue. The New England
journal of medicine 361:415-416; author reply 418-421.

Tontonoz P, Hu. E, Spiegelman BM. 1994. Stimulation of adipogenesis in fibroblasts by pparg2,
a lipid-activated transcription factor. Cell 79:1147-1156.

Vernochet, C., Peres, S. B., Davis, K. E., McDonald, M. E., Qiang, L., Wang, H., Scherer, P. E.,
& Farmer, S. R. 2009. C/EBPalpha and the corepressors CtBP1 and CtBP2 regulate repression of
select visceral white adipose genes during induction of the brown phenotype in white adipocytes
by peroxisome proliferator-activated receptor gamma agonists. Mol Cell Biol, 29(17), 4714–
4728.

Villanueva CJ, Vergnes L, Wang J, Drew BG, Hong C, Tu Y, et al. 2013. Adipose subtype-
selective recruitment of tle3 or prdm16 by ppargamma specifies lipid storage versus thermogenic
gene programs. Cell metabolism 17:423-435.

Wang H, Liu L, Lin JZ, Aprahamian TR, Farmer SR. 2016. Browning of white adipose tissue
with roscovitine induces a distinct population of ucp1+ adipocytes. Cell metabolism 24:835-847.

Ward JH. 1963. Hierarchical Grouping to Optimize an Objective Function. Journal of the
American Statistical Association 58:236–244.

Wolins, N. E., Quaynor, B. K., Skinner, J. R., Tzekov, A., Park, C., Choi, K., & Bickel, P. E.
(2006). OP9 mouse stromal cells rapidly differentiate into adipocytes: characterization of a
useful new model of adipogenesis. J Lipid Res, 47(2), 450–460.

Xiong Y, Soumillon M, Wu J, Hansen J, Hu B, van Hasselt JGC, et al. 2017. A comparison of
mrna sequencing with random primed and 3'-directed libraries. Sci Rep 7:14626.

Yang X, Enerback S, Smith U. 2003. Reduced expression of foxc2 and brown adipogenic genes
in human subjects with insulin resistance. Obesity research 11:1182-1191.

Zhang J, Fu M, Cui T, Xiong C, Xu K, Zhong W, et al. 2004. Selective disruption of ppargamma
2 impairs the development of adipose tissue and insulin sensitivity. Proc Natl Acad Sci U S A
101:10703-10708.

Zhao X, Hu H, Wang C, Bai L, Wang Y, Wang W, Wang J. 2019. A comparison of methods for
effective differentiation of frozen-thawed 3T3-L1 cell. Anal Biochem 568:57-64.

1106

1107

1108

1109   **Tables:**

1110   **Table 1.** Summary of experimental conditions

	3T3-L1	HUMAN PREADIPOCYTES
Exposure Period (days)	10	14
Number of times dosed	4	6
Positive Control	Rosiglitazone	Rosiglitazone
Negative Control (Vehicle)	DMSO	DMSO

1111

1112

1113 **Table 2.** Amended random forest classification results for 17 compounds that are potential

1114 PPARγ Ligands/Modifiers.

CHEMICAL NAME	KNOWN SOURCE/USE	PPARγ LIGAND/ MODIFIER SCORE (VOTE ± 95% CI)
d-cis,trans-Allethrin	Insecticide	0.91 ± 0.01
Tonalide	Musk (fragrance)	0.90 ± 0.01
Quinoxyfen	Fungicide	0.90 ± 0.01
Fenthion	Insecticide	0.88 ± 0.01
2,4,6-Tris(tert-butyl)phenol	Antioxidant (industrial)	0.80 ± 0.02
Prallethrin	Insecticide	0.78 ± 0.02
Tebuconazole	Fungicide	0.78 ± 0.02
Fludioxonil	Fungicide	0.77 ± 0.02
Tris(1,3-dichloro-2-propyl) phosphate	Flame retardant	0.76 ± 0.02
Cyazofamid	Pesticide	0.72 ± 0.02
Perfluorooctanoic acid	Fluorosurfactant	0.59 ± 0.02
Triphenyl phosphite	Pesticide	0.57 ± 0.02
Tris(1-chloro-2-propyl) phosphate	Flame retardant	0.54 ± 0.02
Triphenylphosphine oxide [a]	Crystallizing aid, byproduct	0.49 ± 0.02
Diphenyl phosphate [a]	Metabolite of TPhP	0.47 ± 0.02
Dioctyl sulfosuccinate sodium [a]	Surfactant	0.41 ± 0.02
Perfluorooctanesulfonic acid [a]	Fluorosurfactant	0.40 ± 0.02

1115 [a] Below the highest F1-score threshold.
1116

1117

1118

1119

1120

1121   **Figures:**

1122

Figure 1. Lipid accumulation in differentiated and treated 3T3-L1 pre-adipocytes.

Confluent 3T3 L1 cells were differentiated using a standard hormone cocktail for 10 days.

During differentiation, cells were treated with vehicle (Vh, 0.2% DMSO, final concentration),

rosiglitazone (100 nM) or test chemicals (**Table S1**). On days 3, 5, and 7 of differentiation, the

medium was replaced and the cultures re-dosed. Following 10 days of differentiation and dosing,

cells were analyzed for lipid accumulation by Nile Red staining. Nile Red fluorescence was

normalized by subtracting the fluorescence measured in naïve pre-adipocyte cultures within each

experiment and reported as "Naïve Corrected RFU." Numerical data are provided in **Excel File**

3. Data are presented as mean ± SE (n=4). Statistically different from Vh-treated (highlighted in

green) (*p<0.05, ANOVA, Dunnett's).

1133

1134

1135

Figure 2. Amended random forest classification performance and gene importance of final

classification model.

(A) Performance of random forest classification procedure based 10-fold cross validation.

The plot shows the receiver operating characteristic curves associated with the AUC for 10

repetitions of performance estimation. The values represent the mean estimate across repetitions

for each evaluation metric. The standard deviations these distributions are shown in parenthesis.

The full set of performance estimates for each repetition, performance metric, and classification

procedure considered are shown in **Excel File 1**. **(B)** Gini importance versus ranking of genes

used in the final random forest model. The names of the top 2 genes are highlighted.

Compound-specific gene expression of *Rpl13* and *Cidec* are shown in **Figure S4**.

1146

1147

1148

1149 **Figure 3.** Chemical taxonomy of adipogens based on K2 clustering of the 3'DGE data.

1150 The dendrogram shows the taxonomy-driven hierarchical grouping of test chemical exposures of

1151 3T3-L1 cells or naïve pre-adipocytes. Each split is labeled with a letter, and the proportion of

1152 gene-level bootstraps which produced the resulting split is shown. Highlights of hyper-

1153 enrichment of gene ontology (GO) biological processes are shown.

1154

1155

Figure 4. Associations between plasma adiponectin levels and projections of the 3T3-L1 derived chemical taxonomy gene signatures onto human adipose tissue gene expression.

(A) Each adjoined triangle in the dendrogram denotes a set of genes derived from the 3T3-L1 data, either higher (left) or lower (right) expression at a given node. Triangle texture and color pertain to the direction of the partial correlation of the projection of these gene sets with plasma adiponectin levels, such that positive and negative partial correlations are indicated by solid red

and checkered blue triangles, respectively. Interpretation of these associations is in the key on the

upper right. All colored triangles reach a significance threshold of FDR Q-value < 0.10. **(B)**

Plots of the partial correlation measurements for gene set projections and plasma adiponectin

levels. The top and bottom plots are indicative of gene sets that are either increased in expression

or decreased in expression in a particular sub-group, respectively. All colored points reach a

significance threshold of FDR Q-value < 0.10, as in (A). **(C)** Results of all partial correlation

analyses of projection of taxonomy gene sets with clinical measurements with at least one

comparison reaching significance, FDR Q-value < 0.10. The interpretation of these associations

is the same as in (A).

1172

1173

1174

Figure 5. White and brite gene expression in differentiated and treated 3T3-L1 adipocytes.

Confluent 3T3 L1 cells were differentiated using a standard hormone cocktail for 10 days.

During differentiation, cells were treated with vehicle (Vh, 0.2% DMSO, final concentration),

rosiglitazone (Rosig, 200 nM), roscovitine (Rosco, 4 μM), 15dPGJ2 (500 nM), TBBPA (20 μM)

and TPhP (10 μM). On days 3, 5, and 7 of differentiation, the adipocyte maintenance medium

was replaced and the cultures re-dosed. Following 10 days of differentiation and dosing, cells

were analyzed for gene expression by RT-qPCR. Gene expression levels were normalized to the

geometric mean of the expression levels of B2m and Rn18s and expressed as "Relative

Expression" in comparison to naïve, pre-adipocyte cultures using the Pfaffl method. **(A)** Genes

related to white adipogenesis. **(B)** Genes related to brite adipogenesis. Numerical data are

provided in **Excel File 3**. Data are presented as mean $\pm$ SE of n=4 independent experiments.

Statistically different from Vh-treated (highlighted in green) (*p<0.05, **p<0.01, ***p<0.001,

****p<0.0001, ANOVA, Dunnett's).

1188

1189

Figure 6. Fatty acid uptake and lipid accumulation in differentiated and treated 3T3-L1 adipocytes.

Differentiation and dosing were carried out as described in Figure 5. Cells were analyzed after 10 days of differentiation. **(A)** Fatty acid uptake was analyzed using a dodecanoic acid fluorescent fatty acid substrate. The fluorescence at time zero was subtracted from the fluorescence at the end of the 10 min incubation and reported as "RFU." **(B)** Lipid accumulation was analyzed by Nile Red staining. Nile Red fluorescence was normalized by subtracting the fluorescence measured in naïve pre-adipocyte cultures within each experiment and reported as "Naïve Corrected RFU." Numerical data are provided in **Excel File 3**. Data are presented as means ± SE (n=4). Statistically different from Vh-treated (highlighted in green) (*p<0.05, ***p<0.001, ANOVA, Dunnett's).

Figure 7. Adiponectin secretion in differentiated and treated 3T3-L1 adipocytes.

Differentiation and dosing were carried out as described in Figure 5. On day 9 of differentiation,

cells were switch to washout medium (DMEM+0.1% BSA) and incubated 24hrs. The washout

medium was replaced and the cells were redosed. Medium was collected after a further 24hrs to

adiponectin analysis by ELISA. Concentrations were calculated from absorbance values relative

to a standard curve and reported as ng/ml. Numerical data are provided in **Excel File 3**. Data are

presented as means ± SE (n=4). Statistically different from Vh-treated (highlighted in green)

(*$p<0.05$, **$p<0.01$, ANOVA, Dunnett's).

1215

Figure 8. Mitochondrial biogenesis in differentiated and treated 3T3-L1 adipocytes.

Differentiation and dosing were carried out as described in Figure 5. Following 10 days of

differentiation, mitochondrial biogenesis was analyzed by measuring mitochondria-specific

proteins by ELISA. The absorbance ratios of SDH/Janus are reported as "Relative Mitochondrial

Protein Expression." Numerical data are provided in **Excel File 3**. Data are presented as means ±

SE (n=4). Statistically different from Vh-treated (highlighted in green) (*p<0.05, ANOVA,

Dunnett's).

1223

1224

1225

Figure 9. Cellular respiration in differentiated and treated 3T3-L1 adipocytes.

Differentiation and dosing were carried out as described in Figure 5. Following 10 days of

differentiation, mitochondrial respiration was analyzed by Seahorse Assay. Oxygen consumption

rate was normalized for cell number by dividing by the Janus Green absorbance in each well and

reported as "Relative OCR." Numerical data are provided in **Excel File 3**. Data are presented as

means ± SE (n=4). Statistically different from Vh-treated (highlighted in green) (*$p < 0.05$,

ANOVA, Dunnett's).

1233

1234

1235

1236

Figure 10. Quinoxyfen- and tonalide-induced adipogenesis in 3T3-L1 pre-adipocytes.

Confluent 3T3 L1 cells were differentiated using a standard hormone cocktail for 10 days.

During differentiation, cells were treated with vehicle (Vh, 0.2% DMSO, final concentration),

rosiglitazone (Rosig, 200 nM), quinoxyfen (Quino, 10 µM) or tonalide (Tonal, 4 µM). On days 3,

5, and 7 of differentiation, the adipocyte maintenance medium was replaced and the cultures re-

dosed. Following 10 days of differentiation and dosing, cultures were analyzed. (**A**) Lipid

handling. For fatty acid uptake, the fluorescence at time zero was subtracted from the

fluorescence at the end of the 10 min incubation and reported as "RFU." Lipid accumulation was

determined from Nile Red fluorescence, which was normalized by subtracting the fluorescence

measured in naïve pre-adipocyte cultures within each experiment and reported as "Naïve

Corrected RFU." (**B**) Adiponectin concentrations in the medium measured by ELISA and

calculated from absorbance values relative to a standard curve and reported as ng/ml. Gene

expression levels determined by RT-qPCR and were normalized to the geometric mean of the

expression levels of *B2m* and *Rn18s* and expressed as "Relative Expression" in comparison to

naïve, pre-adipocyte cultures using the Pfaffl method. (**C**) White adipocyte gene expression. (**D**)

Brite adipocyte gene expression. (**E**) Mitochondrial biogenesis was determined by measuring

protein concentrations by ELISA. Absorbance ratios of SDH/Janus are reported as "Relative

Mitochondrial Protein Expression." (**F**) Cellular respiration was measured by Seahorse Assay.

Oxygen consumption rate was normalized for cell number by dividing by the Janus Green

absorbance in each well and reported as "Relative OCR." Numerical data are provided in **Excel**

File 3. Data are presented as means ± SE (n=4-6). Statistically different from Vh-treated

(highlighted in green) (*$p<0.05$, **$p<0.01$, ***$p<0.001$, ANOVA, Dunnett's).

1259

Figure 11. Tonalide- and quinoxyfen-induced adipogenesis in primary human adipocytes.

Confluent primary human preadipocytes were differentiated using a standard human adipocyte

hormone cocktail for 14 days. During differentiation, cells were treated with vehicle (Vh, 0.1%

DMSO, final concentration), rosiglitazone (Rosig, 4 μM), quinoxyfen (Quino, 4 μM) or tonalide

(4 μM). On days 3, 5, 7, 10, and 12 of differentiation, the medium was replaced and the cultures

re-dosed. Following 14 days of differentiation and dosing, cultures were analyzed for **(A)** lipid

handling, **(B)** white (*Cidec*) and brite (*Cidea*) gene expression, and **(C)** mitochondrial biogenesis.

Individual data are presented with the mean indicated by a line. Gene expression levels were

normalized to the geometric mean of the expression levels of *B2M* and *RPL27* and expressed as

relative expression in comparison to naïve, pre-adipocyte cultures. Numerical data are provided

1270   in **Excel File 3**. Statistically different from Vh-treated (highlighted in green) (*p<0.05,

1271   **p<0.01, ***p<0.001, ANOVA, Dunnett's).

1272