Alveolar macrophage chromatin is modified to orchestrate host response to *Mycobacterium bovis* infection

Thomas Jonathan Hall [1], Douglas Vernimmen [2, *], John Andrew Browne [1], Michael P. Mullen [3], Stephen Vincent Gordon [4, 5], David Evan MacHugh [1, 4, *] and Alan Mark O'Doherty [1]

[1] Animal Genomics Laboratory, UCD School of Agriculture and Food Science, College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

[2] The Roslin Institute, University of Edinburgh, Easter Bush Campus, Midlothian, EH25 9RG, UK.

[3] Bioscience Research Institute, Athlone Institute of Technology, Dublin Road, Athlone, Co. Westmeath, N37 HD68, Ireland.

[4] UCD School of Veterinary Medicine, University College Dublin, Belfield, Dublin, D04 W6F6, Ireland.

[5] UCD Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin, D04 V1W8, Ireland.

* Corresponding authors:

Douglas Vernimmen   douglas.vernimmen@roslin.ed.ac.uk

David E. MacHugh   david.machugh@ucd.ie

Keywords

ChIP-seq, chromatin, macrophage, mycobacteria, *Mycobacterium bovis*, tuberculosis

Abstract

Background: Bovine tuberculosis is caused by infection with *Mycobacterium bovis*, which can also cause disease in a range of other mammals, including humans. Alveolar macrophages are the key immune effector cells that first encounter *M. bovis* and how the macrophage epigenome responds to mycobacterial pathogens is currently not well understood.

Results: Here, we have used chromatin immunoprecipitation sequencing (ChIP-seq), RNA-seq and miRNA-seq to examine the effect of *M. bovis* infection on the bovine alveolar macrophage (bAM) epigenome. We show that H3K4me3 is more prevalent, at a genome-wide level, in chromatin from *M. bovis*-infected bAM compared to control non-infected bAM; this was particularly evident at the transcriptional start sites of genes that determine programmed macrophage responses to mycobacterial infection (e.g. M1/M2 macrophage polarisation). This pattern was also supported by the distribution of RNA Polymerase II (PolII) ChIP-seq results, which highlighted significantly increased transcriptional activity at genes demarcated by permissive chromatin. Identification of these genes enabled integration of high-density GWAS data, which revealed genomic regions associated with resilience to infection with *M. bovis* in cattle.

Conclusions: Through integration of these data, we show that bAM transcriptional reprogramming occurs through differential distribution of H3K4me3 and PolII at key immune genes. Furthermore, this subset of genes can be used to prioritise genomic variants from a relevant GWAS data set.

Background

Bovine tuberculosis (bTB) is a chronic infectious disease of livestock, particularly domestic cattle (*Bos taurus* and *Bos indicus*), which causes more than $3 billion in losses to global agriculture annually [1, 2]. The disease can also significantly impact wildlife including, for example, several deer species, American bison (*Bison bison*), African buffalo (*Syncerus caffer*), the brushtail possum (*Trichosurus vulpecula*) and the European badger (*Meles meles*) [3-6]. The etiological agent of bTB is *Mycobacterium bovis*, a pathogen with a genome sequence that is 99.95% identical to *M. tuberculosis*, the primary cause of human tuberculosis (TB) [7]. In certain agroecological milieus *M. bovis* can also cause zoonotic TB with serious implications for human health [8-10].

Previous studies have shown that the pathogenesis of bTB disease in animals is similar to TB disease in humans and many of the features of *M. tuberculosis* infection are also characteristic of *M. bovis* infection in cattle [11-13]. Transmission is via inhalation of contaminated aerosol droplets and the primary site of infection is the lungs where the bacilli are phagocytosed by alveolar macrophages, which normally can contain or destroy intracellular bacilli [14, 15]. Disease-causing mycobacteria, however, can persist and replicate within alveolar macrophages via a bewildering range of evolved mechanisms that subvert and interfere with host immune responses [16-19]. These mechanisms include: recruitment of cell surface receptors on the host macrophage; blocking of macrophage phagosome-lysosome fusion; detoxification of reactive oxygen and nitrogen intermediates (ROI and RNI); harnessing of intracellular nutrient supply and metabolism; inhibition of apoptosis and autophagy; suppression of antigen presentation; modulation of macrophage signalling pathways; cytosolic escape from the phagosome; and induction of necrosis, which leads to immunopathology and shedding of the pathogen from the host [20-25].

Considering the dramatic perturbation of the macrophage by intracellular mycobacteria,

we and others have demonstrated that bovine and human alveolar macrophage transcriptomes

are extensively reprogrammed in response to infection with *M. bovis* and *M. tuberculosis* [26-

31]. These studies have also revealed that differentially expressed gene sets and dysregulated

cellular networks and pathways are functionally associated with many of the macrophage

processes described above that can control or eliminate intracellular microbes.

For many intracellular pathogens it is now also evident that the infection process

involves alteration of epigenetic marks and chromatin remodelling that may profoundly alter

host cell gene expression [32-35]. For example, distinct DNA methylation changes are

detectable in macrophages infected with the intracellular protozoan *Leishmania donovani*,

which causes visceral leishmaniasis [36]. Recent studies using cells with a macrophage

phenotype generated from the THP-1 human monocyte cell line have provided evidence that

infection with *M. tuberculosis* induces alterations to DNA methylation patterns at specific

inflammatory genes [37] and across the genome in a non-canonical fashion [38].

With regards to host cell histones, the intracellular pathogen *Chlamydia trachomatis*

secretes NEU, a SET domain-containing effector protein that functions as a histone

methyltransferase and induces chromatin modifications favourable to the pathogen [39]. It

has also been shown that *Legionella pneumophila*—the causative agent of Legionnaires'

disease—secretes a SET domain-containing histone methyltransferase, RomA, which targets

histone H3 to downregulate host genes and promote intracellular replication [40]. In the

context of mycobacterial infections, Yaseen et al. have shown that the Rv1988 protein,

secreted by virulent mycobacteria, localises to the chromatin upon infection and mediates

repression of host cell genes through methylation of histone H3 at a non-canonical arginine

residue [41]. In addition, chromatin immunoprecipitation sequencing (ChIP-seq) analysis of

H3K4 monomethylation (a marker of poised or active enhancers), showed that regulatory

sequence motifs embedded in subtypes of Alu SINE transposable elements are key components of the epigenetic machinery modulating human macrophage gene expression during *M. tuberculosis* infection [42].

In light of the profound macrophage reprogramming induced by mycobacterial infection, and previous work demonstrating a role for host cell chromatin modifications, we have used ChIP-seq and RNA sequencing (RNA-seq) to examine gene expression changes that reflect host-pathogen interaction in bovine alveolar macrophages (bAM) infected with *M. bovis*. The results obtained support an important role for dynamic chromatin remodelling in the macrophage response to mycobacterial infection, particularly with respect to M1/M2 polarisation. Genes identified from ChIP-seq and RNA-seq results were also integrated with GWAS data to prioritise genomic regions and SNPs associated with bTB resilience. Finally, the suitability of bAM for ChIP-seq assays and the results obtained demonstrate that these cells represent an excellent model system for unravelling the epigenetic and transcriptional circuitry perturbed during mycobacterial infection of vertebrate macrophages.

Results

M. bovis infection induces trimethylation of H3K4 at key immune function related loci in bovine alveolar macrophages

Previous studies have shown that bAM undergo extensive gene expression reprogramming following infection of *M. bovis* [29], with approximately one third of the genome significantly differentially expressed within bovine macrophages 24 hours after infection [28]. Changes of this magnitude are comparable to those observed in previous experiments that have examined the chromatin remodelling that accompanies mycobacterial infection of macrophages, where trimethylation of lysine 4 of Histone H3 (H3K4me3) was shown to correlate with active transcription [42, 43].

We used chromatin immunoprecipitation sequencing (ChIP-seq) to examine histone modification changes that occur after *M. bovis* infection of bAM from sex- and aged-matched Holstein-Friesian cattle. The aim was to determine genome-wide changes in the distribution of H3K4me3 and H3K27me3, and PolII occupancy at the response genes [44]. After data quality control and filtering, ~760 million paired end reads were aligned to the UMD 3.1 bovine genome build at an average alignment rate of 96.23%. Correlation plots of genome-wide H3K4me3, H3K27me3 and PolII sequencing reads from infected and non-infected bAMs showed high correlation between samples (Pearson's correlation coefficient: 0.93–0.97) for all three ChIP-seq targets (supplementary figure 1); indicating that the observed differences in histone modifications between samples are limited and localised to specific regions of the genome.

Differential peaks between conditions were called, compared and visualised with IGV to determine where differences in H3K4me3, H3K27me3 and PolII occupancy occur between control and infected bAM (Figure 1). ChIP seq peaks are defined as areas of the genome enriched by read counts after alignment to the reference genome.

Peak differences for H3K4me3 occurred at multiple locations across the genome and

were estimated by the fold enrichment of a peak normalised against input control DNA that

had not undergone antibody enrichment. Differential peaks in each condition were defined by

several criteria: 1) the fold enrichment of each peak had to be larger than 10 in at least one

condition [45]; 2) the identified peaks had a *P*-value cut off of 0.05; 3) the peaks being

compared in each condition were no more than 500 bp up- and downstream of each other; 4)

the peaks were classified as different using log-likelihood ratios and affinity scores, with

using MACS2 and diffBind, respectively; and 5) visual inspection of the tracks of the peaks

confirmed the computationally determined differences in each condition.

Peaks that occurred in a particular sample indicate that H3K4me3 and PolII are highly

correlated with condition (Figure 2A); this demonstrates that the differences in H3

modifications are a result of infection rather than genomic differences between animals.

Analysis of genome-wide H3K4me3 revealed significant peak differences between control

and infected samples at multiple sites in the genome under these criteria, with some of these

differences occurring at the transcriptional start site of 233 genes. (Figure 2 A-D and

supplementary figure 4). Supplementary figure 1 demonstrates that the differences in

H3K4me3 and Pol II peaks are minor, with cells from both conditions sharing most peaks and

differing by only 1.8–2.95% in peaks across the genome. Principal component analysis

(PCA) of the H3K4me3 mark and PolII data indicated that these H3K4me3 and PolII peak

differences are strongly associated with *M. bovis* infection of bAM (supplementary figure 3).

Changes in H3K4me3 are accompanied with immune related transcriptional reprogramming

Previous studies have shown that increased H3K4me3 is frequently accompanied by an increase in PolII occupancy and elevated expression of proximal genes [46, 47]. In the present study, we observed that H3K4me3 is accompanied by an increase in PolII occupancy (Figure 1 and supplementary figure 4). For a small number of genes (24 out of 233) where the H3K4me3 peak was larger in the control than the infected samples, PolII occupancy was greater in control bAM for 20 genes (83.3%) and greater in infected bAM for three genes (12.5%). Conversely, where the H3K4me3 peak was larger in the infected bAM, PolII occupancy was greater in the infected samples for 127 genes (60.4%) and greater in the control bAM for 14 genes (6.6%). The remaining 60 genes (25%) did not exhibit H3K4me-associated PolII occupancy in either control or infected samples. Figure 3A illustrates this trend, showing that PolII occupancy normally accompanies H3K4me3.

To establish if H3K4me3 mark patterns were correlated with changes in gene expression, control non-infected bAM and bAM infected *M. bovis* AF2122/97 from four animals 24 hpi (including the two animals used for ChIP-seq) were used to generate eight RNA-seq libraries. After quality control and filtering, ~250 million reads were mapped to the bovine genome, with 72% total read mapping, overall. RNA-seq analysis revealed 7,757 differentially expressed genes ($\log_2$FC > 0: 3,723 genes; $\log_2$FC < 0: 4,034 genes; FDR < 0.1). Of the 233 genes identified in the ChIP-seq analysis, 232 (99.6%) were differentially expressed with these criteria (see supplementary file 2). Of the genes that exhibited H3K4me3 peaks that were larger in the infected bAMs, 21 (10%) were downregulated and 189 (90%) were upregulated. Of the genes that exhibited larger H3K4me3 peaks in the control group, 22 (91.6%) were downregulated and 2 (8.4%) were upregulated (Figure 3A).

This pattern of directional gene expression correlating with H3K4me3 for the control and infected samples is consistent with the literature [46, 47].

Existing published RNA-seq data generated by our group using *M. bovis*-infected ($n = $ 10) and control non-infected bAM ($n = $ 10) at 24 hpi [29], was also examined in light of the results from the present study. For the 232 genes identified here, a Pearson correlation coefficient of 0.85 was observed for two data sets (Figure 3D), thus demonstrating that gene expression differences between *M. bovis*-infected and control non-infected bAM are consistent across experiments, even where samples sizes are markedly different.

Transcriptional reprogramming is coupled with differential microRNA expression

We have previously demonstrated that differential expression of immunoregulatory microRNAs (miRNAs) is evident in bAM infected with *M. bovis* compared to non-infected control bAM [31, 48]. To investigate the expression of miRNA in bAM used for the ChIP-seq analyses, miRNA was extracted and sequenced from the samples used for the RNA-seq analysis. After quality control and filtering, ~100 million reads were mapped to the bovine genome, with 79% total reads mapping, overall. Twenty-three differentially expressed miRNAs were detected at 24 hpi ($\log_2$FC > 0: 13; $\log_2$FC < 0: 10; FDR < 0.10). Of the 232 genes identified in the ChIP-seq/RNA-seq analysis, 93 are potential targets for the 23 differentially expressed miRNAs (supplementary data 3). Further examination revealed that multiple immune genes, such as *BCL2A1* (bta-mir-874), *ARG2* (bta-mir-101)*, TMEM173* (aka *STING*) (bta-mir-296-3p) and *STAT1* (bta-mir-2346), are potential regulatory targets for these miRNAs (Figure 3B). This observation therefore supports the hypothesis that miRNAs function in parallel with chromatin modifications to modulate gene expression in response to infection by *M. bovis*.

The H3K4me3, PolII, K27me3 and RNA-seq data were subsequently integrated to

evaluate the relationship between histone modifications and gene expression changes. Three

dimensional plots were generated to visualize the global differences between H3K4me3,

PolII and gene expression in infected and non-infected bAM (Figure 3D). These plots show

that reduction of H3K4me3 in infected cells is associated with a decrease in gene expression

and an absence of PolII occupancy. Genome-wide H3K27me3 was also investigated to

determine whether methylation of this residue was altered in response to *M. bovis* infection

and if it was related to gene expression. No significant differences for H3K27me3 between

control and infected bAM were detected, indicating that repression of gene expression

through H3K27me3 does not play a role in the bAM response to *M. bovis* at 24hpi. However,

supplementary figure 2 indicates that the presence of a H3K27me3 peak in both control and

infected cells at the TSS of a H3k4me3 enriched gene correlated well with a lower or

complete lack of Pol II occupancy.

Pathway analysis reveals H3K4me3 marks are enriched for key immunological genes

To identify biological pathways associated with genes identified through the ChIP-seq

analyses, we integrated the ChIP-seq, RNA-seq and miRNA-seq data and created a list of 232

genes that were present in each data set. Pathway analyses were carried out using three

pathway tools: Ingenuity Pathway Analysis (IPA), Panther and DAVID [49-51]. IPA

revealed an association with *respiratory illness* and the *innate immune response*

(supplementary file 2). Panther was used to examine the gene ontology categories of the 232

genes (Figure 4A); this revealed enrichment for metabolic processes, response to stimuli and

cellular processes, indicating that increased H3K4me3 in response to *M. bovis* infection

occurs at TSS of genes associated with the immune response.

The final part of the pathway analysis was performed using DAVID [51]. DAVID uses

a list of background genes and query genes (in this case the 232 common genes across data

sets) and identifies enriched groups of genes with shared biological functions. The DAVID

analysis demonstrated that the 232 genes are involved in several signalling pathways,

including the PI3K/AKT/mTOR, JAK-STAT and RIG-I-like signalling pathways (Figure 4B

and the top 10 pathways are detailed in supplementary file 3).

GWAS integration prioritises bovine SNPs associated with resilience to *M. bovis* infection

Previous work used high-density SNP (597,144 SNPs) data from 841 Holstein-

Friesian bulls for a genome-wide association study (GWAS) to detect SNPs associated with

susceptibility/resistance to *M. bovis* infection [52]. Using a permutation-based approach to

generate null SNP distributions, we leveraged these data to show that genomic regions within

100 kb up- and downstream of each of the 232 genes exhibiting differential H3K4me3 ChIP-

seq peaks are significantly enriched for additional SNPs associated with resilience to *M. bovis*

infection.

In total, 12,056 SNPs within the GWAS data set were located within 100 kb of the

232 H3K4me3 genes. Of these SNPs, up to 26 were found to be significantly associated with

bTB susceptibility, depending on the distance interval of each gene. Interestingly, 22 SNPs

found within 25 kb of 11 genes were found to be most significant at P and q values < 0.05,

with declining significance of association as the region extended beyond 25 kb (Figure 4C

and supplementary file 3). Significant SNPs were detected in proximity to the following

genes: *SAMSN1, CTSL, TNFAIP3, CLMP, ABTB2, RNFT1, MIC1, MIC2, EDN1, ARID5B*,

all of which had significant differential enrichment of H3K4me3.

Discussion

H3K4me3 mark occurs at key immune genes

Our study has generated new information regarding host-pathogen interaction during the initial stages of *M. bovis* infection. We demonstrate that chromatin is remodelled through differential H3K4me3 and that PolII occupancy is altered at key immune genes in *M. bovis*-infected bAM. This chromatin remodelling correlates with changes in the expression of genes that are pivotal for the innate immune response to mycobacteria [28, 29, 53]. Our work supports the hypothesis that chromatin modifications of the host macrophage genome play an essential role during intracellular infections by mycobacterial pathogens [54, 55].

The top pathways identified were the JAK-STAT signalling pathway, the PI3K/AKT/mTOR signalling pathway and the RIG-I-like receptor signalling pathway. In mammals, the JAK-STAT pathway is the principal signalling pathway that modulates expression of a wide array of cytokines and growth factors, involved in cell proliferation and apoptosis [56]. The JAK-STAT signalling pathway and its regulators are also associated with coordinating an effective host response to mycobacterial infection [57, 58]. Two JAK-STAT associated stimulating factors and a ligand receptor that exhibited increased H3K4me3 marks in infected samples were encoded by the *OSM*, *CSF3* and *CNTFR* genes, respectively [59, 60]. *OSM* has previously been shown to be upregulated in cells infected with either *M. bovis* or *M. tuberculosis* [29, 61, 62]. Our work shows that this increased expression in response to mycobacteria is facilitated by H3K4me3-mediated chromatin accessibility. The protein encoded by *CSF3* has also been implicated as an immunostimulator in the response to mycobacterial infection due to its role in granulocyte and myeloid haematopoiesis [63]. *CNTFR* encodes a ligand receptor that stimulates the JAK-STAT pathway and shows increased expression in other studies of mycobacterial infection [28, 29]. Following stimulation of *JAK* through ligand receptor binding, *STAT1* expression is increased. STAT1,

a signal transducer and transcription activator that mediates cellular responses to interferons

(IFNs), cytokines and growth factors, is a pivotal JAK-STAT component and a core

component in the response to mycobacterial infection [64]. Here, the TSS of *STAT1* was

associated with an increased deposition of H3K4me3. Interestingly, upregulation of *STAT1*

was associated with a downregulation of bta-miR-2346, predicted to be a negative regulator

of STAT1 (see supplementary file 3). Overall, these results show that major components of

the JAK-STAT pathway undergo chromatin remodelling mediated via H3K4me3, thereby

facilitating activation and propagation of the JAK-STAT pathway through chromatin

accessibility.

Key genes encoding components of the PI3K/AKT/mTOR pathway, such as *IRF7*,

RAC1 and *PIK3AP1*, were also identified as having increased H3K4me3 in *M. bovis* infected

macrophages. PI3K/AKT/mTOR signalling contributes to a variety of processes that are

critical in mediating aspects cell growth and survival [65]. Phosphatidylinositol-3 kinases

(PI3Ks) and the mammalian target of rapamycin (mTOR) are integral to coordinating innate

immune defences [66]. The PI3K/AKT/mTOR pathway is an important regulator of type I

interferon production via activation of the interferon-regulatory factor 7, IRF7. RAC1 is a

key activator of the PI3K/AKT/mTOR pathway and, in its active state, binds to a range of

effector proteins to regulate cellular responses such as secretory processes, phagocytosis of

apoptotic cells, and epithelial cell polarization [67]. In addition, *in silico* analysis of our

differentially expressed miRNAs predicted that several miRNAs, such as bta-miR-1343-3p,

bta-miR-2411-3p and bta-miR-1296, regulate *RAC1*. *PIK3AP1* expression was also

increased, in line with previous mycobacterial infection studies [28, 29]. Hence as observed

with the JAK-STAT pathway, H3K4me3 at these key PI3K/AKT/mTOR pathway genes acts

to regulate the innate response to mycobacterial infection.

Initiation of the RIG-I-like receptor signalling pathway generally occurs following viral

infection through sensing of viral RNAs by cytoplasmic RIG-I-like receptors (RLRs) and

activation of transcription factors that drive production of type I IFNs [68]. A number of

bacterial species induce type I IFN independently of TLRs, potentially through the RIG-I-like

pathway [69, 70]. While type I IFNs have a well characterised role in the inhibition of viral

replication, their role during bacterial infection is less well defined [71-73]. In humans and

mice, *M. tuberculosis*-induced type I IFN is associated with TB disease progression and

impairment of host resistance [74-76]. In our study, genes encoding multiple components of

the RIG-I-like receptor signalling pathway, such as *TRIM25, ISG15, IRF7* and *IKBKE,* were

enriched for K4me3 and PolII occupancy in *M. bovis*-infected bAM. These results

demonstrate that the RIG-I-like pathway activation in *M. bovis*-infected bAM is driven, to a

large extent, by reconfiguration of the host chromatin.

H3K4me3 enriched loci are also flanked by genomic polymorphisms associated with

resilience to *M. bovis* infection. Integration of our data with GWAS data from 841 bulls that

have robust phenotypes for bTB susceptibility/resistance revealed 22 statistically significant

SNPs within 25 kb of 11 H3K4me3 enriched genes. Most of these genes are involved in host

immunity, with *CTSL, TNFAIP3,* and *RNFT1* directly implicated in the human response to

M. tuberculosis infection [77-79]. The reprioritisation of genomic regions and array-based

SNPs using integrative genomics approaches will be relevant for genomic prediction and

genome-enabled breeding and may facilitate fine mapping efforts and the identification of

targets for genome editing of cattle resilient to bTB.

H3K4me3 deposition at host macrophage genes and immunological evasion by *M. bovis*

The present study has revealed elevated H3K4me3 deposition and PolII occupancy at key immune genes that are involved in the innate response to mycobacterial infection. In addition, we also identified several immune genes that had differential H3K4me3 and expression, where the expression change may be detrimental to the ability the host macrophage to clear infection. An example of this is *ARG2*, which exhibited increased H3K4me3 deposition, PolII occupancy and expression (Log$_2$FC = 3.415, P_{adj} = 7.52 ×10^{-16}) in infected cells. However, it is also interesting to note that the integrated expression output of *ARG2* may also be determined by the bta-miR-101 miRNA, a potential silencer of *ARG2* expression, which was observed to be upregulated in infected cells. Elevated levels of arginase 2, the protein product of the *ARG2* gene, have previously been shown to shift macrophages to an M2 phenotype [80, 81], which is anti-inflammatory and exhibits decreased responsiveness to IFN-gamma and decreased bactericidal activity [82]. Hence, it may be hypothesised that *M. bovis* infection triggers H3K4me3 deposition at the TSS of *ARG2* to drive an M2 phenotype and generate a more favourable niche for the establishment of infection. Similarly to *ARG2,* increased expression of *BCL2A1* in *M. bovis*-infected bAM may also facilitate development of a replicative milieu for intracellular mycobacteria. Increased expression of *BCL2A1* is associated with decreased macrophage apoptosis [83], which would otherwise restrict replication of intracellular pathogens.

In comparison to control non-infected bAM, the *TMEM173* (aka *STING*) gene exhibited substantially decreased expression in *M. bovis*-infected bAM (Log$_2$FC = -3.225, P_{adj} = 8.64 ×10^{-11}). *TMEM173* encodes transmembrane protein 173, which drives IFN production and as such is a major regulator of the innate immune response to viral and bacterial infections, including *M. bovis* and *M. tuberculosis* [28, 84, 85]. Downregulation of

TMEM173 indicates that *M. bovis* can actively reduce or block methylation of H3K4 at this

gene in infected macrophages, thereby enhancing the pathogen's intracellular survival. In this

regard, we have recently shown that infection of bAM with *M. tuberculosis*, which is

attenuated in cattle, causes increased *TMEM173* expression compared to infection with *M.*

bovis [28].

The molecular mechanisms that pathogens employ to manipulate the host genome to

subvert or evade the immune response are yet to be fully elucidated. Hijacking the host's own

mechanisms for chromatin modulation is one potential explanation that has garnered attention

in recent years [33, 35]. These modulations of the host chromatin in bAMs may be mediated

through *M. bovis*-derived signals transmitted through bacterial metabolites, RNA-signalling

or secreted peptides [41, 86-88].

Conclusions

Elucidation of the mechanisms used by pathogens to establish infection, and ultimately

cause disease, requires an intimate knowledge of host-pathogen interactions. Using

transcriptomics and epigenomics, we have identified altered expression of major host

immune genes following infection of primary bovine macrophages with *M. bovis*. We have

shown that reprogramming of the alveolar macrophage transcriptome occurs mainly through

increased deposition of H3K4me3 at key immune function genes, with additional gene

expression modulation via miRNA differential expression. This modulation of gene

expression drives a shift of the macrophage phenotype towards the more replicate-permissive

M2 macrophage phenotype. We have also identified that alveolar macrophages infected with

M. bovis exhibit differentially expressed genes (in regions with modified chromatin) that are

enriched for significant SNPs from GWAS data for bTB resilience; this shows the power of

integrative approaches to identify key loci to be targeted for selective breeding programmes.

Our data confirm the emerging concept that pathogens can hijack host chromatin, through

manipulation of H3K4me3, to subvert host immunity and to establish infection.

Methods

Preparation and infection of bAMs

Alveolar macrophages and *M. bovis* 2122 were prepared as described previously [89] with minor adjustments, 2×10^6 macrophages were seeded in 60 mm tissue culture plates and challenged with *M. bovis* at an MOI of 10:1 (2×10^7 bacteria per plate) for 24 h; parallel non-infected controls were prepared simultaneously.

Preparation of nucleic acids for sequencing

Sheared fixed chromatin was prepared exactly as described in the truChIP™ Chromatin Shearing Kit (Covaris) using 2×10^6 macrophage cells per AFA tube. Briefly, cells were washed in cold PBS and 2.0 ml of fixing Buffer A was added to each plate, to which 200 µl of freshly prepared 11.1% formaldehyde solution was added. After 10 min on a gentle rocker the crosslinking was halted by the addition of 120 µl of quenching solution E, cells were washed with cold PBS, released from the plate using a cell scraper and re-suspended in 300 µl Lysis Buffer B for 10 min with gentle agitation at 4°C to release the nuclei. The nuclei were pelleted and washed once in Wash Buffer C and three times in Shearing Buffer D3 (X3) prior to been resuspended in a final volume of 130 µl of Shearing Buffer D3. The nuclei were transferred to a micro AFA tube and sonicated for 8 min each using the Covaris E220e as per the manufacturer's instructions. Chromatin immunoprecipitation of sonicated DNA samples was carried out using the Chromatin Immunoprecipitation (ChIP) Assay Kit (Merck KGaD) and anti-H3K4me3 (05-745R) (Merck KGaD), Pol II (H-224) (Santa Cruz Biotechnology, Inc.) or anti-H3K27me3 (07-449) (Merck KGaD) as previously described [90]. RNA was extracted from infected ($n = 4$) and control ($n = 4$) bAM samples using the RNeasy Plus Mini

400 Kit (Qiagen) as previously described [91]. All 8 samples exhibited excellent RNA quality

401 metrics (RIN >9).

Sequencing

Illumina TruSeq Stranded mRNA and TruSeq Small RNA kits were used for mRNA-

seq and small RNA-seq library preparations and the NEB Next Ultra ChIPseq Library Prep

kit (New England Biolabs) was used for ChIP-seq library preparations. Pooled libraries were

sequenced by Edinburgh Genomics (http://genomics.ed.ac.uk) as follows: paired-end reads (2

× 75 bp) were obtained for mRNA and ChIP DNA libraries using the HiSeq 4000 sequencing

platform and single-end read (50 bp) were obtained for small RNA libraries using the HiSeq

2500 high output version 4 platform.

ChIP-seq bioinformatics analysis

Computational analyses for all bioinformatic processes were performed on a 72-CPU

compute server with Linux Ubuntu (version 16.04.4 LTS). An average of 54 M paired end

75bp reads were obtained for each histone mark. At each step of data processing, read quality

was assessed via FastQC (version 0.11.5) [92]. Any samples that indicated adapter

contamination were trimmed via Cutadapt (version 1.15) [93]. Raw read correlation plots

were generated via EaSeq (version 1.05) [94]. The raw reads were aligned to UMD 3.1.1

bovine genome assembly using Bowtie2 (version 2.3.0) [95]. The mean alignment rate for the

histone marks was 96.23%. The resulting SAM files were converted and indexed into BAM

files via Samtools (version 1.3.1) [96]. After alignment, samples were combined and sorted

into 14 files, based on the animal (A1 or A2), the histone mark (K4/K27/PolII) and treatment

(control or infected) i.e. A1-CTRL-K4. Peaks were called by using alignment files to

determine where the reads have aligned to specific regions of the genome, and then

comparing that alignment to the input samples as a normalization step.

The peak calling was carried out via MACS (version 2.1.1.20160309) [97]. The K4me3

mark was called in sharp peak mode and K27me3 and Pol II were called in broad peak mode,

as per the user guide. Peak tracks were generated in macs and visualized with the Integrative

Genome Viewer (version 2.3) [98]. Union peaks were generated by combining and merging

overlapping peaks in all samples for each histone mark. Differential peak calling was called

via macs using the bdgdiff function. Peaks images were generated by visually assessing all

three marks in tandem across the entire bovine genome with IGV. The significance of peaks

was determined by sorting peaks for each mark in each treatment by P value and then fold

enrichment with a cut-off of 2 and a P value threshold of 0.05 [99]. Peaks from each animal

in each condition for each mark were cross referenced with the IGV images and differential

peak caller to determine a difference in fold enrichment for each observed peak difference

between conditions. This required comparing peak start and end sites, chromosomes, P and q

values for each summit, summit locations and normalised fold enrichment of a peak against

the input sample (see supplementary file 1 for peak sets). Any peaks that exhibited a

difference of 4 or greater fold enrichment, a P value of less than 0.05, an FDR (q value) less

than 0.05 and that were also identified by the differential peak caller were selected for further

analysis (see supplementary file 1 for peaks at TSS that met some but not all of the above

criteria). Peaks that were then classified to be different between conditions in all three data

sets were examined to determine their proximity to TSS. Differential peaks were also called

using the R package DiffBind (version 2.80) [100]. DiffBind includes functions to support

the processing of peak sets, including overlapping and merging peak sets, counting

sequencing reads overlapping intervals in peak sets, and identifying statistically significantly

differentially bound sites based on evidence of binding affinity (measured by differences in

read densities, see supplementary info 1). For H3K27me3 DiffBind differential peak calling,

the initial MACS2 peak list, consisting of 64,264 total peaks (see supplementary info 1), was

merged and reduced to a smaller group of larger, broader peaks to reduce noise and false

positive discovery (Figure 2B).

RNA-seq bioinformatics analysis

An average of 44 M paired end 75 bp reads were obtained for each of the eight samples

(four control, four infected). Adapter sequence contamination and paired-end reads of poor

quality were removed from the raw data. At each step, read quality was assessed with FastQC

(version 0.11.5). Any samples that indicated adapter contamination were trimmed via

Cutadapt (version 1.15). The raw reads were aligned to the UMD 3.1.1 bovine transcriptome

using Salmon (version 0.8.1) [101]. Aligned reads were also counted in Salmon and the

resulting quantification files were annotated at gene level via tximport (version 3.7) [102].

The annotated gene counts were then normalised and differential expression analysis

performed with DESeq2 (version 1.20.0) [103], correcting for multiple testing using the

Benjamini-Hochberg method [104]. Genes identified from ChIP-seq as exhibiting differential

histone modifications were cross referenced with the RNA-seq data set to determine

significant $\log_2$FC between *M. bovis*-infected and control non-infected. Additionally, this

RNA-seq data was cross referenced with RNA-seq data from a previous study that

investigated bAM infected with *M. bovis* [29].

MicroRNA-seq bioinformatics analysis

A mean of 26 M paired-end 50 bp reads were obtained for each of the eight samples

(four control, four infected). At each step of data processing, read quality was assessed via

fastqc (version 0.11.5). Any samples that exhibited adapter contamination were trimmed via

Cutadapt (version 1.15) and all reads smaller than 17 bp were removed from the analysis.

Raw reads were mapped to UMD3.1 using Bowtie (version 1.2.2). miRNA detection,

identification and quantification was carried out with mirdeep2 (version 0.0.91). Isoform

analysis was also performed using mirdeep2. Differential expression analysis was performed

using DESeq2, correcting for multiple testing with the Benjamini-Hochberg method. Any

miRNAs that were significantly differentially expressed (FDR < 0.10) were selected for

further analysis. To determine if significantly differentially expressed miRNAs target genes

selected in the ChIP seq analysis, miRmap [105] was used to predict the likelihood a specific

miRNA targets one or more of the genes based on three criteria: delta G binding, probability

exact and phylogenetic conservation of seed site, which is then combined into a single

scoring metric (miRmap score). Any predicted gene targets with miRmap score $\geq$ 0.70 were

included in the analysis (see supplementary info 3).

Pathway analysis

488   Pathway analysis was carried out on any gene that had a differential peak between

489   control and infected samples. Pathway analysis and gene ontology was carried out by DAVID

490   (version 6.8), Ingenuity pathway analysis (01-13) and PANTHER (version 13.1) [49, 106].

491   KEGG pathways were selected by choosing pathways that had the highest number of genes

492   identified in the ChIP seq data and had a FDR < 0.05.

Integration of GWAS data.

495   GWAS data for genetic susceptibility to *M. bovis* infection previously generated by

496   Richardson and colleagues [52] were analysed to determine if subsets of SNPs selected

according to their distance to H3K4me3 and PolII active loci were enriched for significant GWAS hits. The nominal P values used in this study were generated using single SNP regression analysis in a mixed animal model as described previously [52]. In summary, high-density genotypes ($n = 597,144$) of dairy bulls ($n = 841$) used for artificial insemination were associated with deregressed estimated breeding values for bTB susceptibility that had been calculated from epidemiological information on 105,914 daughters and provided by the Irish Cattle Breeding Federation (ICBF). In this study, the significance of the distribution of SNP nominal P values (from Richardson et al 2016) within and up to 100 kb up and downstream to genes identified as having differential H3K4me3 and PolII activity on bTB susceptibility were estimated in R using q value (FDRTOOL) and permutation analysis (custom scripts). A total of 1000 samplings (with replacement) from the HD GWAS P value data set ($n = 597,144$) representing the size of each of selected SNP subsets were generated. The q values for each SNP P value subset and all its permuted equivalents were calculated using the FDRTOOL library in R. The subsequent significance level (P_{perm}) assigned to each of the SNP subsets was equivalent to the proportion of permutations in which at least the same number of q values < 0.05 as the SNP subset were obtained, i.e. by chance.

Figure legends

Figure 1. Track visualization of *M. bovis* induced H3K4me3 and PolII occupancy with relative change in expression at three immune response associated genes.

Examples of signal tracks illustrating peaks of H3K27me3 (top two tracks), H3K4me3 (middle two tracks) and PolII (bottom two tracks) in infected (red) and non-infected (blue) bovine alveolar macrophages, with the bovine reference genome on the bottom of each panel reading left to right. Accompanying each track image is the expression of the corresponding gene, with normalised counts of infected cells in red and control in blue. The *ARG2* gene exhibited an increase in H3K4me3 at 24 hpi as evidenced by the larger red H3K4me3 and red PolII peaks. The *IFITM2* gene also exhibited larger H3K4me3 and PolII peaks in infected samples; however, in contrast to this, *SIRT3*, which is located ~20kb upstream from *IFITM2* gene, had no significant change in either peak. *TMEM173* (aka *STING*) exhibits an opposite pattern to most genes identified as having differential H3K4me3, where a larger peak is observed in control samples rather than infected.

Figure 2. *M.bovis* induced histone modifications occur genome wide at key immune loci.

(**A**) Correlation heatmaps of differential peaks for H3K4me3, H3K27me3 and PolII. Every peak location that is not consistent between each animal in each condition (i.e. a peak only occurs in the control group) is compared to determine if these inconsistent peaks are correlated with the animal or the condition. The differential peaks in H3K4me3 and PolII correlate highly with condition, whereas there was no significant global differences in the distribution of H3K27me3. (**B**) Venn diagrams of differential peaks for H3K4me3, H3K27me3 and PolII. Each condition shares the majority of peaks. Where differences occur at TSS of genes, these genes are frequently associated with immune function. (**C**) Volcano

plots of differential peaks for H3K4me3 and PolII. The y-axis shows significance as FDR and

the x-axis indicates increase in affinity for control (left) and infected (right). Significant sites

are denoted in blue. (**E**) Boxplots of differential peaks for H3K4me3 and PolII. Infected bAM

are shown in red and control bAM are shown in blue. The left two boxes of each plot show

distribution of reads over all differentially bound sites in the infected and control groups. The

middle two boxes of each plot show the distribution of reads in differentially bound sites that

increase in affinity in the control group. The far right boxes in each plot show the distribution

of reads in differentially bound sites that increase in affinity in the infected group.

Figure 3. H3K4me3 is accompanied by functional changes in PolII occupancy, gene

expression and gene regulation.

(**A**) Scatter plots of H3K4me3 against PolII occupancy and gene expression. The first plot is

the difference of peaks for H3K4me3 between conditions, ranging from negative to positive

values, with negative being a larger peak in the control samples (blue dots) and the positive

values being a larger peak in the infected samples (red dots), on the y-axis. The x-axis

represents the $\log_2$FC for each of the 232 genes, with each gene as a single data point. The

second plot also has H3K4me3 on the y-axis but with peak differences in PolII on the x-axis,

with negative and positive values corresponding to greater occupancy in the control and

infected samples, respectively. The final plot shows $\log_2$FC relative to PolII occupancy. (**B**)

Plots of normalised miRNA-seq counts. Each plot represents the normalised counts of a

miRNA that was detected as exhibiting differential expression. Bta-miR-101 interacts with

ARG2, bta-miR-296-3p with *TMEM173* (aka *STING*), bta-miR-874 with *BCL2A1* and bta-

miR-2346 with *STAT1*. Red bars indicate infected and blue represent control samples.

(**C**) Correlation and Venn diagram for both RNA-seq studies. The x-axis of the scatter plot

represents the $\log_2$FC for each of the 232 genes from this study and the y-axis represents the

$\log_2$FC for each of the 232 genes from the previous study [29]. The Venn diagram shows the

global overlap of differentially expressed genes from both studies with an FDR cut off of

<0.1. (**D**) 3-D plots for all three data sets. A combination of all three scatter plots from Figure

2A. Data points are genes. Blue genes are those that exhibited greater H3K4me3 in control

bAM, red exhibited greater H3K4me3 in infected bAM.

Figure 4. Gene ontology enrichment and pathway analysis.

(**A**) Gene ontology pie charts generated through PANTHER pathway analysis. 232 genes

cluster by gene ontology under three main categories: *Biological process*, *Cellular*

component and *Molecular function*. (**B**) KEGG pathway images containing genes identified

from the ChIP-seq and RNA-seq analysis. Gene symbols coloured in yellow were identified

in the ChIP-seq and RNA-seq analysis. Gene symbols coloured in red were also targeted by

one or more differentially expressed miRNAs. Up or down red arrows indicate greater

H3K4me3 in infected or control, respectively. Up or down yellow arrows indicate log2FC

increase or decrease of the associated gene, respectively. (**C**) Line graph showing different

genomic ranges from genes that are enriched for significant SNPs from GWAS data for bTB

resilience. The bars represent the number of SNPs that occupy each range from each ChIP-

seq enriched gene, with more SNPs correlating with a greater distance. The blue plotted line

represents the negative $\log_{10}$ probability that the significant SNPs found at each distance at

0.05 FDR q value are significant by chance, with SNPs at 25 kb exhibiting the lowest

probability. The null SNP P value distribution for each data point was generated from 1000

permutations of random SNPs corresponding to the number of SNPs observed in a particular

genomic range. (D) Genes enriched for SNPs significantly associated with resilience to *M.*

bovis infection. SNP IDs and functional information obtained from the GeneCards® database

[107] are also shown.

Abbreviations

bAM: bovine alveolar macrophage/s; bTB: bovine tuberculosis; ChIP: chromatin immunoprecipitation; FDR: false discovery rate; GWAS: genome-wide association study; H3K27me3: histone H3 lysine 27 tri-methylation; H3K4me3: histone H3 lysine 4 tri-methylation; hpi: hours post infection; log$_2$FC: log$_2$ fold change; PolII: RNA Polymerase II; SNP: single nucleotide polymorphism; TB: tuberculosis.

Acknowledgments

The authors would also like to thank FAANG–Europe for awarding A.M.O.D. a short-term scientific mission (STSM) grant. We would also like to acknowledge Edinburgh Genomics for generation of sequencing data.

Author Contributions

Conceptualization, A.M.O.D., D.E.M. and T.J.H.; Software and Formal Analysis, T.J.H. and M.P.M.; Investigation, A.M.O.D., D.V. and J.A.B.; Resources, D.E.M., S.V.G. and D.V.; Data Curation, T.J.H.; Writing – original draft, T.J.H, A.M.O.D. and D.E.M.; Writing – review and editing, D.V., S.V.G and M.P.M.; Supervision and Project Administration, D.E.M. and A.M.O.D.; Funding Acquisition; D.E.M, S.V.G, D.V. and A.M.O.D.

Competing Interests

The authors declare no competing interests.

Ethics Statement and Consent to Participate

All animal procedures were performed according to the provisions of Statutory Instrument No. 543/2012 (under Directive 2010/63/EU on the Protection of Animals used for Scientific Purposes). Ethical approval was obtained from the University College Dublin

611   Animal Ethics Committee (protocol number AREC-13-14-Gordon). No human subjects were

612   used for this study.

Consent for Publication

614   Not applicable.

Funding

This study was supported by Science Foundation Ireland (SFI) Investigator Programme Awards to D.E.M. and S.V.G. (grant nos. SFI/08/IN.1/B2038 and SFI/15/IA/3154); a European Union Framework 7 Project Grant to D.E.M. (no: KBBE-211602-MACROSYS); an EU H2020 COST Action short-term scientific mission (STSM) grant to A.M.O.D. (reference code: COST-STSM-ECOST-STSM-CA15112-050317-081648); a University of Edinburgh Chancellor's Fellowship to D.V.; and Institute Strategic Grant funding to the Roslin Institute (grant nos. BBS/E/D/10002070 and BBS/E/D/20002172). The funding agencies had no role in the study design, collection, analysis, and interpretation of data, and no role in writing the manuscript.

References

References

1.   Steele JH: **Introduction (Part 2 Regional and Country Status Reports).** In
Mycobacterium bovis infection in animals and humans. Edited by Thoen CO, Steele JH.
Ames, IA, USA: Iowa State University Press; 1995: 169-172

2.   Waters WR, Palmer MV, Buddle BM, Vordermeier HM: **Bovine tuberculosis**
vaccine research: historical perspectives and recent advances. *Vaccine* 2012, **30:**2611-
2622.

3.   Fitzgerald SD, Kaneene JB: **Wildlife reservoirs of bovine tuberculosis worldwide:**
hosts, pathology, surveillance, and control. *Vet Pathol* 2013, **50:**488-499.

4.   Gormley E, Corner LAL: **Pathogenesis of *Mycobacterium bovis* infection: the**
badger model as a paradigm for understanding tuberculosis in animals. *Front Vet Sci*
2017, **4:**247.

5.   Malone KM, Gordon SV: ***Mycobacterium tuberculosis* complex members adapted**
to wild and domestic animals. *Adv Exp Med Biol* 2017, **1019:**135-154.

6.   Palmer MV: ***Mycobacterium bovis*: characteristics of wildlife reservoir hosts.**
Transbound Emerg Dis 2013, **60 Suppl 1:**1-13.

7.   Garnier T, Eiglmeier K, Camus JC, Medina N, Mansoor H, Pryor M, Duthoy S,
Grondin S, Lacroix C, Monsempe C, et al: **The complete genome sequence of**
***Mycobacterium bovis*.** *Proc Natl Acad Sci U S A* 2003, **100:**7877-7882.

8.   Olea-Popelka F, Muwonge A, Perera A, Dean AS, Mumford E, Erlacher-Vindel E,
Forcella S, Silk BJ, Ditiu L, El Idrissi A, et al: **Zoonotic tuberculosis in human beings**
caused by *Mycobacterium bovis* – a call for action. *Lancet Infect Dis* 2017, **17:**e21-e25.

9.   Thoen CO, Kaplan B, Thoen TC, Gilsdorf MJ, Shere JA: **Zoonotic tuberculosis. A**
comprehensive ONE HEALTH approach. *Medicina (B Aires)* 2016, **76:**159-165.

10.   Vayr F, Martin-Blondel G, Savall F, Soulat JM, Deffontaines G, Herin F:
Occupational exposure to human *Mycobacterium bovis* infection: A systematic review.
PLoS Negl Trop Dis 2018, **12:**e0006208.

11.   Buddle BM, Vordermeier HM, Hewinson RG: **Experimental infection models of**
tuberculosis in domestic livestock. *Microbiol Spectr* 2016, **4**.

12.   Waters WR, Maggioli MF, McGill JL, Lyashchenko KP, Palmer MV: **Relevance of**
bovine tuberculosis research to the understanding of human disease: historical
perspectives, approaches, and immunologic mechanisms. *Vet Immunol Immunopathol*
2014, **159:**113-132.

13.   Williams A, Orme IM: **Animal models of tuberculosis: an overview.** *Microbiol*
Spectr 2016, **4**.

14.   Kaufmann SHE, Dorhoi A: **Molecular determinants in phagocyte-bacteria**
interactions. *Immunity* 2016, **44:**476-491.

15. Weiss G, Schaible UE: **Macrophage defense mechanisms against intracellular bacteria.** *Immunol Rev* 2015, **264:**182-203.

16. Awuh JA, Flo TH: **Molecular basis of mycobacterial survival in macrophages.** *Cell Mol Life Sci* 2017, **74:**1625-1648.

17. Schorey JS, Schlesinger LS: **Innate immune responses to tuberculosis.** *Microbiol Spectr* 2016, **4**.

18. Cambier CJ, Falkow S, Ramakrishnan L: **Host evasion and exploitation schemes of** *Mycobacterium tuberculosis*. *Cell* 2014, **159:**1497-1509.

19. de Chastellier C: **The many niches and strategies used by pathogenic mycobacteria for survival within host macrophages.** *Immunobiology* 2009, **214:**526-542.

20. Stutz MD, Clark MP, Doerflinger M, Pellegrini M: *Mycobacterium tuberculosis***: Rewiring host cell signaling to promote infection.** *J Leukoc Biol* 2018, **103:**259-268.

21. Chaurasiya SK: **Tuberculosis: Smart manipulation of a lethal host.** *Microbiol Immunol* 2018, **62:**361-379.

22. BoseDasgupta S, Pieters J: **Macrophage-microbe interaction: lessons learned from the pathogen** *Mycobacterium tuberculosis*. *Semin Immunopathol* 2018, **40:**577-591.

23. Queval CJ, Brosch R, Simeone R: **The macrophage: A disputed fortress in the battle against** *Mycobacterium tuberculosis*. *Front Microbiol* 2017, **8:**2284.

24. Hussain Bhat K, Mukhopadhyay S: **Macrophage takeover and the host-bacilli interplay during tuberculosis.** *Future Microbiol* 2015, **10:**853-872.

25. Ehrt S, Schnappinger D: **Mycobacterial survival strategies in the phagosome: defence against host stresses.** *Cell Microbiol* 2009, **11:**1170-1178.

26. Papp AC, Azad AK, Pietrzak M, Williams A, Handelman SK, Igo RP, Jr., Stein CM, Hartmann K, Schlesinger LS, Sadee W: **AmpliSeq transcriptome analysis of human alveolar and monocyte-derived macrophages over time in response to** *Mycobacterium tuberculosis* **infection.** *PLoS ONE* 2018, **13:**e0198221.

27. Jensen K, Gallagher IJ, Johnston N, Welsh M, Skuce R, Williams JL, Glass EJ: **Variation in the early host-pathogen interaction of bovine macrophages with divergent** *Mycobacterium bovis* **strains in the United Kingdom.** *Infect Immun* 2018, **86**.

28. Malone KM, Rue-Albrecht K, Magee DA, Conlon K, Schubert OT, Nalpas NC, Browne JA, Smyth A, Gormley E, Aebersold R, et al: **Comparative 'omics analyses differentiate** *Mycobacterium tuberculosis* **and** *Mycobacterium bovis* **and reveal distinct macrophage responses to infection with the human and bovine tubercle bacilli.** *Microb Genom* 2018**:**[Epub ahead of print].

29. Nalpas NC, Magee DA, Conlon KM, Browne JA, Healy C, McLoughlin KE, Rue-Albrecht K, McGettigan PA, Killick KE, Gormley E, et al: **RNA sequencing provides exquisite insight into the manipulation of the alveolar macrophage by tubercle bacilli.** *Sci Rep* 2015, **5:**13629.

30. Lavalett L, Rodriguez H, Ortega H, Sadee W, Schlesinger LS, Barrera LF: **Alveolar**
macrophages from tuberculosis patients display an altered inflammatory gene
expression profile. *Tuberculosis* 2017, **107:**156-167.

31. Vegh P, Magee DA, Nalpas NC, Bryan K, McCabe MS, Browne JA, Conlon KM,
Gordon SV, Bradley DG, MacHugh DE, Lynn DJ: **MicroRNA profiling of the bovine**
alveolar macrophage response to *Mycobacterium bovis* **infection suggests pathogen**
survival is enhanced by microRNA regulation of endocytosis and lysosome trafficking.
Tuberculosis 2015, **95:**60-67.

32. Bierne H, Hamon M, Cossart P: **Epigenetics and bacterial infections.** *Cold Spring*
Harb Perspect Med 2012, **2:**a010272.

33. Hamon MA, Cossart P: **Histone modifications and chromatin remodeling during**
bacterial infections. *Cell Host Microbe* 2008, **4:**100-109.

34. Niller HH, Minarovits J: **Patho-epigenetics of infectious diseases caused by**
intracellular bacteria. *Adv Exp Med Biol* 2016, **879:**107-130.

35. Rolando M, Gomez-Valero L, Buchrieser C: **Bacterial remodelling of the host**
epigenome: functional role and evolution of effectors methylating host histones. *Cell*
Microbiol 2015, **17:**1098-1107.

36. Marr AK, MacIsaac JL, Jiang R, Airo AM, Kobor MS, McMaster WR: *Leishmania*
donovani **infection causes distinct epigenetic DNA methylation changes in host**
macrophages. *PLoS Pathog* 2014, **10:**e1004419.

37. Zheng L, Leung ETY, Wong HK, Lui G, Lee N, To K-F, Choy KW, Chan RCY, Ip
M: **Unraveling methylation changes of host macrophages in** *Mycobacterium tuberculosis*
infection. *Tuberculosis* 2016, **98:**139-148.

38. Sharma G, Sowpati DT, Singh P, Khan MZ, Ganji R, Upadhyay S, Banerjee S,
Nandicoori VK, Khosla S: **Genome-wide non-CpG methylation of the host genome**
during *M. tuberculosis* **infection.** *Sci Rep* 2016, **6:**25006.

39. Pennini ME, Perrinet S, Dautry-Varsat A, Subtil A: **Histone methylation by NUE, a**
novel nuclear effector of the intracellular pathogen *Chlamydia trachomatis***.** *PLoS Pathog*
2010, **6:**e1000995.

40. Rolando M, Sanulli S, Rusniok C, Gomez-Valero L, Bertholet C, Sahr T, Margueron
R, Buchrieser C: *Legionella pneumophila* **effector RomA uniquely modifies host**
chromatin to repress gene expression and promote intracellular bacterial replication.
Cell Host Microbe 2013, **13:**395-405.

41. Yaseen I, Kaur P, Nandicoori VK, Khosla S: **Mycobacteria modulate host**
epigenetic machinery by Rv1988 methylation of a non-tail arginine of histone H3. *Nat*
Commun 2015, **6:**8922.

42. Bouttier M, Laperriere D, Memari B, Mangiapane J, Fiore A, Mitchell E, Verway M,
Behr MA, Sladek R, Barreiro LB, et al: **Alu repeats as transcriptional regulatory**
platforms in macrophage responses to *M. tuberculosis* **infection.** *Nucleic Acids Res* 2016,
44:10571-10587.

43.   Arts RJW, Moorlag S, Novakovic B, Li Y, Wang SY, Oosting M, Kumar V, Xavier
RJ, Wijmenga C, Joosten LAB, et al: **BCG vaccination protects against experimental viral**
infection in humans through the induction of cytokines associated with trained
immunity. *Cell Host Microbe* 2018, **23:**89-100 e105.

44.   Sims RJ, 3rd, Nishioka K, Reinberg D: **Histone lysine methylation: a signature for**
chromatin function. *Trends Genet* 2003, **19:**629-639.

45.   Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein
BE, Bickel P, Brown JB, Cayting P, et al: **ChIP-seq guidelines and practices of the**
ENCODE and modENCODE consortia. *Genome Res* 2012, **22:**1813-1831.

46.   Clouaire T, Webb S, Skene P, Illingworth R, Kerr A, Andrews R, Lee JH, Skalnik D,
Bird A: **Cfp1 integrates both CpG content and gene activity for accurate H3K4me3**
deposition in embryonic stem cells. *Genes Dev* 2012, **26:**1714-1728.

47.   Barski A, Cuddapah S, Kartashov AV, Liu C, Imamichi H, Yang W, Peng W, Lane
HC, Zhao K: **Rapid recall ability of memory T cells is encoded in their epigenome.** *Sci*
Rep 2017, **7:**39785.

48.   Vegh P, Foroushani AB, Magee DA, McCabe MS, Browne JA, Nalpas NC, Conlon
KM, Gordon SV, Bradley DG, MacHugh DE, Lynn DJ: **Profiling microRNA expression in**
bovine alveolar macrophages using RNA-seq. *Vet Immunol Immunopathol* 2013, **155:**238-
244.

49.   Kramer A, Green J, Pollard J, Jr., Tugendreich S: **Causal analysis approaches in**
Ingenuity Pathway Analysis. *Bioinformatics* 2014, **30:**523-530.

50.   Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K,
Muruganujan A, Narechania A: **PANTHER: a library of protein families and subfamilies**
indexed by function. *Genome Res* 2003, **13:**2129-2141.

51.   Huang da W, Sherman BT, Lempicki RA: **Systematic and integrative analysis of**
large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009, **4:**44-57.

52.   Richardson IW, Berry DP, Wiencko HL, Higgins IM, More SJ, McClure J, Lynn DJ,
Bradley DG: **A genome-wide association study for genetic susceptibility to**
Mycobacterium bovis **infection in dairy cattle identifies a susceptibility QTL on**
chromosome 23. *Genet Sel Evol* 2016, **48:**19.

53.   Alcaraz-Lopez OA, Garcia-Gil C, Morales-Martinez C, Lopez-Rincon G, Estrada-
Chavez C, Gutierrez-Pabello JA, Esquivel-Solis H: **Divergent macrophage responses to**
Mycobacterium bovis **among naturally exposed uninfected and infected cattle.** *Immunol*
Cell Biol 2017, **95:**436-442.

54.   Cheng J, Blum R, Bowman C, Hu D, Shilatifard A, Shen S, Dynlacht BD: **A role for**
H3K4 monomethylation in gene repression and partitioning of chromatin readers. *Mol*
Cell 2014, **53:**979-992.

55.   LaMere SA, Thompson RC, Komori HK, Mark A, Salomon DR: **Promoter H3K4**
methylation dynamically reinforces activation-induced pathways in human CD4 T cells.
Genes Immun 2016, **17:**283-297.

56.   Rawlings JS, Rosler KM, Harrison DA: **The JAK/STAT signaling pathway.** *J Cell*
Sci 2004, **117:**1281-1283.

57.   Manca C, Tsenova L, Freeman S, Barczak AK, Tovey M, Murray PJ, Barry C,
Kaplan G: **Hypervirulent *M. tuberculosis* W/Beijing strains upregulate type I IFNs and**
increase expression of negative regulators of the Jak-Stat pathway. *J Interferon Cytokine*
Res 2005, **25:**694-701.

58.   Cliff JM, Kaufmann SH, McShane H, van Helden P, O'Garra A: **The human**
immune response to tuberculosis and its treatment: a view from the blood. *Immunol Rev*
2015, **264:**88-102.

59.   Pastuschek J, Poetzsch J, Morales-Prieto DM, Schleussner E, Markert UR, Georgiev
G: **Stimulation of the JAK/STAT pathway by LIF and OSM in the human granulosa**
cell line COV434. *J Reprod Immunol* 2015, **108:**48-55.

60.   Marino VJ, Roguin LP: **The granulocyte colony stimulating factor (G-CSF)**
activates Jak/STAT and MAPK pathways in a trophoblastic cell line. *J Cell Biochem*
2008, **103:**1512-1523.

61.   Polena H, Boudou F, Tilleul S, Dubois-Colas N, Lecointe C, Rakotosamimanana N,
Pelizzola M, Andriamandimby SF, Raharimanga V, Charles P, et al: *Mycobacterium*
tuberculosis **exploits the formation of new blood vessels for its dissemination.** *Sci Rep*
2016, **6:**33162.

62.   O'Kane CM, Elkington PT, Friedland JS: **Monocyte-dependent oncostatin M and**
TNF-alpha synergize to stimulate unopposed matrix metalloproteinase-1/3 secretion
from human lung fibroblasts in tuberculosis. *Eur J Immunol* 2008, **38:**1321-1330.

63.   Martins A, Han J, Kim SO: **The multifaceted effects of granulocyte colony-**
stimulating factor in immunomodulation and potential roles in intestinal immune
homeostasis. *IUBMB Life* 2010, **62:**611-617.

64.   Tsumura M, Okada S, Sakai H, Yasunaga S, Ohtsubo M, Murata T, Obata H, Yasumi
T, Kong XF, Abhyankar A, et al: **Dominant-negative STAT1 SH2 domain mutations in**
unrelated patients with Mendelian susceptibility to mycobacterial disease. *Hum Mutat*
2012, **33:**1377-1387.

65.   Yu JS, Cui W: **Proliferation, survival and metabolism: the role of**
PI3K/AKT/mTOR signalling in pluripotency and cell fate determination. *Development*
2016, **143:**3050-3060.

66.   Weichhart T, Saemann MD: **The PI3K/Akt/mTOR pathway in innate immune**
cells: emerging therapeutic applications. *Ann Rheum Dis* 2008, **67 Suppl 3:**iii70-74.

67.   Yip SC, El-Sibai M, Coniglio SJ, Mouneimne G, Eddy RJ, Drees BE, Neilsen PO,
Goswami S, Symons M, Condeelis JS, Backer JM: **The distinct roles of Ras and Rac in PI**
3-kinase-dependent protrusion during EGF-stimulated cell migration. *J Cell Sci* 2007,
120:3138-3146.

68.   Loo YM, Gale M, Jr.: **Immune signaling by RIG-I-like receptors.** *Immunity* 2011,
34:680-692.

69.	Charrel-Dennis M, Latz E, Halmen KA, Trieu-Cuot P, Fitzgerald KA, Kasper DL,
Golenbock DT: **TLR-independent type I interferon induction in response to an**
extracellular bacterial pathogen via intracellular recognition of its DNA. *Cell Host*
Microbe 2008, **4:**543-554.

70.	Dixit E, Kagan JC: **Intracellular pathogen detection by RIG-I-like receptors.** *Adv*
Immunol 2013, **117:**99-125.

71.	Boxx GM, Cheng G: **The roles of type I interferon in bacterial infection.** *Cell Host*
Microbe 2016, **19:**760-769.

72.	Mancuso G, Midiri A, Biondo C, Beninati C, Zummo S, Galbo R, Tomasello F,
Gambuzza M, Macri G, Ruggeri A, et al: **Type I IFN signaling is crucial for host**
resistance against different species of pathogenic bacteria. *J Immunol* 2007, **178:**3126-
3133.

73.	O'Connell RM, Saha SK, Vaidya SA, Bruhn KW, Miranda GA, Zarnegar B, Perry
AK, Nguyen BO, Lane TF, Taniguchi T, et al: **Type I interferon production enhances**
susceptibility to *Listeria monocytogenes* infection. *J Exp Med* 2004, **200:**437-445.

74.	Berry MP, Graham CM, McNab FW, Xu Z, Bloch SA, Oni T, Wilkinson KA,
Banchereau R, Skinner J, Wilkinson RJ, et al: **An interferon-inducible neutrophil-driven**
blood transcriptional signature in human tuberculosis. *Nature* 2010, **466:**973-977.

75.	Mayer-Barber KD, Andrade BB, Oland SD, Amaral EP, Barber DL, Gonzales J,
Derrick SC, Shi R, Kumar NP, Wei W, et al: **Host-directed therapy of tuberculosis based**
on interleukin-1 and type I interferon crosstalk. *Nature* 2014, **511:**99-103.

76.	Manca C, Tsenova L, Bergtold A, Freeman S, Tovey M, Musser JM, Barry CE, 3rd,
Freedman VH, Kaplan G: **Virulence of a *Mycobacterium tuberculosis* clinical isolate in**
mice is determined by failure to induce Th1 type immunity and is associated with
induction of IFN-alpha /beta. *Proc Natl Acad Sci U S A* 2001, **98:**5752-5757.

77.	Nepal RM, Mampe S, Shaffer B, Erickson AH, Bryant P: **Cathepsin L maturation**
and activity is impaired in macrophages harboring *M. avium* and *M. tuberculosis*. *Int*
Immunol 2006, **18:**931-939.

78.	Meenu S, Thiagarajan S, Ramalingam S, Michael A, Ramalingam S: **Modulation of**
host ubiquitin system genes in human endometrial cell line infected with *Mycobacterium*
***tuberculosis*.** *Med Microbiol Immunol (Berl)* 2016, **205:**163-171.

79.	Silver RF, Walrath J, Lee H, Jacobson BA, Horton H, Bowman MR, Nocka K, Sypek
JP: **Human alveolar macrophage gene responses to *Mycobacterium tuberculosis* strains**
H37Ra and H37Rv. *Am J Respir Cell Mol Biol* 2009, **40:**491-504.

80.	Hardbower DM, Asim M, Murray-Stewart T, Casero RA, Jr., Verriere T, Lewis ND,
Chaturvedi R, Piazuelo MB, Wilson KT: **Arginase 2 deletion leads to enhanced M1**
macrophage activation and upregulated polyamine metabolism in response to
***Helicobacter pylori* infection.** *Amino Acids* 2016, **48:**2375-2388.

81.   Lewis ND, Asim M, Barry DP, de Sablet T, Singh K, Piazuelo MB, Gobert AP,
Chaturvedi R, Wilson KT: **Immune evasion by *Helicobacter pylori* is mediated by**
induction of macrophage arginase II. *J Immunol* 2011, **186:**3632-3641.

82.   Huang Z, Luo Q, Guo Y, Chen J, Xiong G, Peng Y, Ye J, Li J: ***Mycobacterium***
***tuberculosis*-induced polarization of human macrophage orchestrates the formation and**
development of tuberculous granulomas in vitro. *PLoS ONE* 2015, **10:**e0129744.

83.   Vogler M: **BCL2A1: the underdog in the BCL2 family.** *Cell Death Differ* 2012,
19:67-74.

84.   Manzanillo PS, Shiloh MU, Portnoy DA, Cox JS: ***Mycobacterium tuberculosis***
activates the DNA-dependent cytosolic surveillance pathway within macrophages. *Cell*
Host Microbe 2012, **11:**469-480.

85.   McNab F, Mayer-Barber K, Sher A, Wack A, O'Garra A: **Type I interferons in**
infectious disease. *Nat Rev Immunol* 2015, **15:**87-103.

86.   Woo V, Alenghat T: **Host-microbiota interactions: epigenomic regulation.** *Curr*
Opin Immunol 2017, **44:**52-60.

87.   Silmon de Monerri NC, Kim K: **Pathogens hijack the epigenome: a new twist on**
host-pathogen interactions. *Am J Pathol* 2014, **184:**897-911.

88.   Sharma G, Upadhyay S, Srilalitha M, Nandicoori VK, Khosla S: **The interaction of**
mycobacterial protein Rv2966c with host chromatin is mediated through non-CpG
methylation and histone H3/H4 binding. *Nucleic Acids Res* 2015, **43:**3922-3937.

89.   Magee DA, Conlon KM, Nalpas NC, Browne JA, Pirson C, Healy C, McLoughlin
KE, Chen J, Vordermeier HM, Gormley E, et al: **Innate cytokine profiling of bovine**
alveolar macrophages reveals commonalities and divergence in the response to
***Mycobacterium bovis* and *Mycobacterium tuberculosis* infection.** *Tuberculosis (Edinb)*
2014, **94:**441-450.

90.   Vernimmen D, Lynch MD, De Gobbi M, Garrick D, Sharpe JA, Sloane-Stanley JA,
Smith AJ, Higgs DR: **Polycomb eviction as a new distant enhancer function.** *Genes Dev*
2011, **25:**1583-1588.

91.   O'Doherty AM, O'Shea LC, Fair T: **Bovine DNA methylation imprints are**
established in an oocyte size-specific manner, which are coordinated with the expression
of the DNMT3 family proteins. *Biol Reprod* 2012, **86:**67.

92.   Andrews S: **FastQC: a quality control tool for high throughput sequence data.**
Babraham Research Campus, Cambridge, UK: Bioinformatics Group, Babraham Institute;
2016.

93.   Martin M: **Cutadapt removes adapter sequences from high-throughput**
sequencing reads. *2011* 2011, **17:**3.

94.   Lerdrup M, Johansen JV, Agrawal-Singh S, Hansen K: **An interactive environment**
for agile analysis and visualization of ChIP-sequencing data. *Nat Struct Mol Biol* 2016,
23:349-357.

95.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2.** *Nat*
Methods 2012, **9:**357-359.

96.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, Genome Project Data Processing S: **The Sequence Alignment/Map format and**
SAMtools. *Bioinformatics* 2009, **25:**2078-2079.

97.   Feng J, Liu T, Zhang Y: **Using MACS to identify peaks from ChIP-Seq data.** *Curr*
Protoc Bioinformatics 2011, **Chapter 2:**Unit 2 14.

98.   Thorvaldsdottir H, Robinson JT, Mesirov JP: **Integrative Genomics Viewer (IGV):**
high-performance genomics data visualization and exploration. *Brief Bioinform* 2013,
14:178-192.

99.   Wilbanks EG, Facciotti MT: **Evaluation of algorithm performance in ChIP-seq**
peak detection. *PLoS ONE* 2010, **5:**e11471.

100.   Stark R, Brown G: **DiffBind: differential binding analysis of ChIP-Seq peak data.**
http://bioconductor.org/packages/release/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf.
2011.

101.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C: **Salmon provides fast and**
bias-aware quantification of transcript expression. *Nat Methods* 2017, **14:**417-419.

102.   Soneson C, Love MI, Robinson MD: **Differential analyses for RNA-seq:**
transcript-level estimates improve gene-level inferences. *F1000Res* 2015, **4:**1521.

103.   Love MI, Huber W, Anders S: **Moderated estimation of fold change and**
dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014, **15:**550.

104.   Benjamini Y, Hochberg Y: **Controlling the false discovery rate - a practical and**
powerful approach to multiple testing. *J R Stat Soc Ser B Method* 1995, **57:**289-300.

105.   Vejnar CE, Zdobnov EM: **MiRmap: comprehensive prediction of microRNA**
target repression strength. *Nucleic Acids Res* 2012, **40:**11673-11683.

106.   Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, Thomas PD: **PANTHER**
version 11: expanded annotation data from Gene Ontology and Reactome pathways,
and data analysis tool enhancements. *Nucleic Acids Res* 2017, **45:**D183-D189.

107.   Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI,
Nudel R, Lieder I, Mazor Y, et al: **The GeneCards Suite: from gene data mining to**
disease genome sequence analyses. *Curr Protoc Bioinformatics* 2016, **54:**1.30.31-31.30.33.

930
931

Figure 1

Figure 2

Figure 3

Figure 4

A

Molecular function

Biological process

Cellular component

B

JAK-STAT signalling pathway

PI3-AKT signalling pathway

RIG-I-like signalling pathway

C

D

SNP	Gene	Function
rs109467646	*ABTB2*	Protein heterodimerization activity
rs42340336	*ARID5B*	RNA polymerase II regulatory region sequence-specific DNA binding
rs43291594	*CLMP*	Cell-cell adhesion
rs43291608	*CLMP*	Cell-cell adhesion
rs43291613	*CLMP*	Cell-cell adhesion
rs135643197	*CLMP*	Cell-cell adhesion
rs43291628	*CLMP*	Cell-cell adhesion
rs43292253	*CLMP*	Cell-cell adhesion
rs43292266	*CLMP*	Cell-cell adhesion
rs43292281	*CLMP*	Cell-cell adhesion
rs43292303	*CLMP*	Cell-cell adhesion
rs43292326	*CLMP*	Cell-cell adhesion
rs43293441	*CLMP*	Cell-cell adhesion
rs29014510	*CLMP*	Cell-cell adhesion
rs134561028	*CTSL*	Histone binding
rs42247932	*CTSL*	Histone binding
rs110265685	*EDN1*	Regulation of endothelin secretion
rs133489745	*MIC1*	MICOS complex
rs135121106	*RNFT1*	Integral component of membrane
rs133286573	*SAMSN1*	Phosphotyrosine residue binding
rs134127062	*TNFAIP3*	Ligase activity
rs137535115	*TNFAIP3*	Ligase activity