Full Title:

Pilot study to evaluate the effect of a calcium and vitamin D-containing oral

bolus on serum calcium levels in early lactation Holstein dairy cows

Short title:

Calcium supplementation and serum calcium in early lactation dairy cows

D.A. Shock,[1][*][¶] D.L. Renaud,[1,2,¶] S.M. Roche, [1,¶], R. Genore [1,3&], R. Reynen[3,&], M. E. Olson[4,&]

[1]Agricultural Communications and Epidemiological Research (ACER) Consulting, Guelph, Ontario,

Canada.

[2]Department of Population Medicine, University of Guelph, Guelph, Ontario, Canada.

[3]Heartland Veterinary Services, Listowel, Ontario, Canada

[4]Solvet Animal Health, Calgary, Alberta, Canada.

*Corresponding author:

Email: dshock@acerconsult.ca (DS)

[¶]These authors contributed equally to this work.

[&]These authors also contributed equally to this work.

Abstract

The initiation of lactation challenges the ability of the modern lactating cow to maintain calcium homeostasis, and typically results in a drop in blood calcium levels; leading to mobilization of calcium reserves from skeletal stores. As such, the recommendation to provide supplemental calcium at parturition to older cows has become an industry-standard practice. Mature cows were treated at calving and 12 hours later with either the novel calcium bolus (Calboost (CB) (Solvet, Calgary, Alberta, Canada)), or a commercially available calcium bolus (Bovikalc (BK) (Boehringer Ingelheim Vetmedica Inc., St. Joseph, MO)). Blood was collected from animals at 0, 1, 6, 12, 13, and 24 hours following calving, and the resulting serum samples were analyzed. Overall, there was no statistical difference between the two calcium boluses for blood calcium levels within the first 24 hours following parturition (P = 0.50). Cows treated with both oral calcium boluses experienced a significant increase in serum calcium by 1 hour after parturition, however, this increase was not sustained through subsequent sampling times. This pilot study demonstrates that both boluses have a similar effect in the elevation of blood calcium.

Introduction

The initiation of milk production challenges the ability of the lactating dairy cow to maintain normal calcium concentrations. Once lactation begins, blood calcium levels drop leading to mobilization of calcium reserves from skeletal stores [1,2]. Clinical and subclinical hypocalcaemia are common issues in the dairy industry [3–5], and are both associated with a variety of negative sequelae including an increased risk of displaced abomasum, mastitis, retained placenta, dystocia, and ketosis [6,7]. Given these negative connotations, it is imperative to improve the calcium status of a newly freshened dairy cow.

Oral supplementation with calcium chloride and calcium sulfate in a fat coated bolus has shown to have

significant effects on improving calcium status in the period following calving [8]. Clinically, Oetzel and

Miller [9] found that the administration of the bolus to lame cows and high producing cows lead to

fewer health events and higher milk production in these groups, respectively leading to an economic

benefit of $3 to $8 per cow [10]. There has been very strong adoption of the oral calcium supplement by

dairy producers throughout North America; however, many of the alternative products to Bovikalc

(Boehringer Ingelheim Vetmedica Inc., St. Joseph, MO) have not been tested to demonstrate similar

efficacy in influencing blood calcium levels.

Therefore, the primary objective of this pilot study was to assess the ability of a novel oral calcium bolus

containing calcium chloride, calcium sulfate, and vitamin D3 (Calboost [Solvet, Calgary, Alberta, Canada])

to influence the serum calcium level in recently fresh, mature Holsteins over the first 24 hours following

parturition, relative to a positive control (Bovikalc). We hypothesized that there would be no difference

between the two oral boluses with respect to serum calcium levels over the first 24 hours following

administration.

Materials and methods

Ethics statement

The present field-based study was conducted in compliance with the research guidelines set forth by the

Canadian Council on Animal Care after appropriate Animal Use Protocol review by the Institutional

Animal Care and Use Committee (Alberta Agriculture, Airdrie Alberta, Canada).

Sample size calculation

Based on previous research, oral calcium supplementation should increase the serum calcium

concentration by 0.25 mmol/L (0.125 mmol/L standard deviation) relative to levels at calving [11]. A

total sample size of 20 mature cows (10 each in Calboost and Bovikalc groups) was calculated, assuming

a 95% confidence level and 80% power.

74   Study herd and treatment protocols

The study was conducted at a commercial dairy farm milking 500 cows in southwestern Ontario,

Canada, between March and September 2018. Average milk production per cow was 38 kg, at 3.9% fat

and 3.1% protein, at an average days in milk of 155. The dry cow ration contained straw, corn silage, and

protein supplement, and is considered a low calcium, low phosphorous, low potassium, high magnesium

diet that does not contain anionic salts. Historically, the farm averages less than 5% incident risk of

clinical hypocalcemia. This farm was chosen due to proximity with the veterinary clinic that assisted with

sample handling and analysis, as well as technical capability to collect and record relevant samples and

data.

Cows were randomly allocated to receive either the Calboost bolus (CB) or Bovikalc bolus (BK) according

to a randomization sheet that was generated using the RAND command in Microsoft Excel. In this

experiment, a positive control was used, as the practice of oral calcium supplementation at calving is an

industry-standard recommendation, and few producers capable of conducting field-based research are

willing to accept the risk of not prophylactically treating all mature cows with oral calcium. First lactation

cows, and those animals whose calvings were unobserved were excluded from enrollment. The farm

staff was responsible for bolus administration, blood sample collection and storage, and reporting any

disease events. They were blinded to the treatment groups throughout the trial.

The bolus was given orally at calving and again at 12 hours post-calving. Blood samples were collected

via venipuncture of the coccygeal vein. Blood was taken at 0, 1, 6, 12, 13, and 24 hours relative to

calving. Due to practicalities associated with sampling enrolled cows, the sampling window was allowed

to vary by ± 1 hour of the protocol designation. Samples were allowed to clot, centrifuged for 5 minutes

at 800 g, and serum was transferred into sterile storage vials. The vials were stored at -20°C until

submitted for analysis. Samples were analyzed to determine total serum calcium via a photometric

method (Abaxis Vetscan vs2, Abaxis Inc., Union City, CA).

Following treatments, participating farm personnel were instructed to record treatment assignment and

all pertinent health and disease events (including death and culling) for all cows that had calved after

study commencement.

Data collection

102   Variables of interest for analysis included date and time of calving, calcium bolus administration, and

103   blood collection, lactation number, dystocia, non-lactating days prior to calving, disease dates,

104   culling/death dates, and occurrence of twin pregnancies.

Statistical analysis

106   All datasets were imported as comma-separated files into the STATA 14 statistical software program

107   (StataCorp LP, College Station, TX) for statistical analysis. The cow was considered the unit of analysis for

108   this study.

Descriptive statistics and univariable analysis

Descriptive statistics were generated for the final dataset. Serum calcium, phosphorous, and magnesium

levels were explored through graphical plots. Specific comparisons between the serum calcium, days

dry, lactation, season of sampling, and the presence of twins at calving in CB and BK cows were made,

with univariable statistical comparisons made using Fisher exact tests or Kruskall-Wallis one-way

analysis of variance.

Statistical comparisons were made between potential predictor variables and outcomes of interest. For

this study, the outcome of interest was serum calcium level (in mmol/L). Univariable linear regression

modelling was employed to study variables of interest. Variables explored for each model included:

treatment group (CB vs BK), lactation group (2 and over 3), non-lactating days prior to parturition,

season of calving, sampling time (0, 1, 6, 12, 13, and 24 hours relative to calving), dystocia (0 = eutotic

calving, 1 = dystotic calving), and presence of twins at calving. Variables that had moderate statistical

associations with the outcome of interest (defined at a liberal P-value < 0.2) were included in

subsequent multivariable models.

Multivariable regression analysis

A repeated measures mixed linear regression model was constructed, using random effects to model

sample number within cow, using a first-order autoregressive correlation structure between time points.

A step-wise backwards elimination process was employed, where variables identified as potentially

associated with the outcome of interest in the initial univariable screening were included in a full

multivariable model. Those variables having P-values for partial F-tests or type III tests of fixed effects

greater than 0.05 were eliminated from the model after assessing whether they were part of biologically

plausible interaction terms or had a confounding effect on the outcome of interest (e.g. a > 20% change

in coefficient values when the term is removed from the model). Continuous variables were assessed for

linearity with predicted model outcomes through visually inspecting a LOWESS curve (local weight

scatterplot smoothing) for linear relationship, as well as the significance ($P < 0.05$) of a quadratic term in

the model. If a continuous variable did not have a linear relationship with the outcome of interest, it was

subsequently categorized based on biologically relevant cut-points, or if appropriate, a quadratic term

was retained in the model. Variables retained in the final model were assessed for collinearity through

the examination of Pearson or Spearman rank-order correlation coefficients. When high correlation was

found between variables (> 0.6), the most biologically appropriate variable was chosen for inclusion in

the final model. Standardized residuals were generated and visually assessed for normality and

homoscedasticity for the linear mixed regression model. If the residuals were heteroskedastic or not

normally distributed, appropriate transformations were performed on the outcome of interest.

Results and Discussion

Table 1 outlines the descriptive characteristics of trial animals. The BK group had a lower absolute

number of animals in their second and third lactation, although this difference was not statistically

significant. The twinning rate in each group was relatively low, with 2 twin calvings in the BK group and 1

in the CB group. Mean non-lactating days was numerically greater for CB cows, but this difference was

not statistically significant.

Table 1. Descriptive demographic comparisons between CalBoost (CB) and Bovikalc (BK) of cows enrolled on a commercial dairy farm in southwestern Ontario, Canada.

Variable	CalBoost	BoviKalc	P-Value
Lactation (frequency)			
2nd and 3rd lactation	5	7	0.65
> 3rd lactation	5	3	
Calving Season (frequency)			
March-April	3	1	0.41
May-July	1	3	
August-September	6	6	
Twins (frequency)			
Yes	1	2	1.00
No	9	8	
Mean Days Dry (SE)	73.2 (8.1)	61.3 (6.3)	0.23

Figure 1. a. outlines a descriptive comparison in serum calcium levels between CB and BK-treated

animals. Of note, CB cows had lower serum calcium at parturition relative to BK animals, but the CB

group had numerically higher serum calcium levels in subsequent sampling times (Figure 1.a.). Figures

1.b. and 1.c. outline raw mean (SD) serum phosphorus and magnesium levels measured in study cows,

respectively. There were no significant differences between CB and BK groups for all blood metabolites

(P > 0.05) measured on univariable analysis.

Figure 1. a. Raw mean (SD) serum calcium (mmol/L) plot of Calboost (solid line) and Bovikalc (dashed

line) groups over the first 24 hours following calving.

Figure 1. b. Raw mean (SD) serum phosphorus (mmol/L) plot of Calboost (solid line) and Bovikalc

(dashed line) groups over the first 24 hours following calving.

Figure 1. c. Raw mean (SD) serum magnesium (mmol/L) plot of Calboost (solid line) and Bovikalc

(dashed line) groups over the first 24 hours following calving.

Of note, there were 2 animals of 20 that were sampled outside of the appropriate window (time ± 1

hour) for the 12-hour samples, both of which were in the CB treatment group. Analyses were conducted

both including and excluding these animals, with no bearing on the outcome of the experiment.

Therefore, the data for these animals remained in the analysis (N = 20) to avoid excessive model

overfitting.

Residual analysis from raw serum calcium data showed considerable heteroskedasticity, therefore

natural logarithms were calculated for each calcium value. Upon subsequent iterations of the model,

residuals were homoscedastic and normally distributed with this transformation. Multivariable,

repeated measures linear regression model output is given in Table 2.

Table 2. Multivariable, repeated measures linear regression model output evaluating the effect that

Calboost (CB) relative to Bovikalc (BK) has on log-transformed serum calcium values within the first 24

hours following parturition.

	Coefficient	SE	P-Value	95% Confidence Interval	
Treatment					
BK[a]	Referent				
CB[b]	0.03	0.05	0.50	-0.06	0.12
Sample No.					
1	Referent				
2	0.18	0.05	0.001	0.08	0.29
3	0.07	0.06	0.23	-0.04	0.18
4	0.04	0.06	0.52	-0.08	0.15
5	0.01	0.06	0.80	-0.10	0.13
6	-0.01	0.06	0.82	-0.13	0.10
Calving Month					
March-April	Referent				
May-July	-0.04	0.07	0.54	-0.17	0.09
August-September	0.16	0.05	0.002	0.06	0.27
Days Dry	0.01	0.00	<0.001	0.01	0.02

Days Dry (quadratic)	0.00007	0.00002	0.001	0.00	0.00
Cow Lactation:					
2nd and 3rd	Referent				
3rd+	-0.08	0.05	0.11	-0.17	0.02
Constant	-0.07	0.15	0.65	-0.36	0.23

[a]BK = Bovikalc; [b]CB = Calboost

Serum calcium levels did not differ at any point between BK- and CB-treated cows (P = 0.50). Serum

calcium levels rose dramatically at the 1-hour mark following parturition; however, these levels

subsequently dropped and were not significantly different from baseline (t = 0) levels for the remainder

of the 24-hour sampling period. Cows that calved in the months of August and September had

significantly higher serum calcium levels (P = 0.002) relative to cows calving in March and April. Older

cows tended (P = 0.11) to have lower serum calcium levels. This variable was included in the model, as it

had a significant confounding effect (> 20% change in coefficient value) on the treatment variable of

interest (CB versus BK-treatment). Previous research has noted older cows to be at an increased risk of

clinical milk fever [1]. Overall, Figure 2 outlines serum calcium levels for BK- and CB-treated cows over

the experimental period.

Figure 2. Least square mean (SE) serum calcium values for Calboost (solid line) and Bovikalc (dashed

line) treated cows over the first 24 hours following parturition.

The manufacturing and marketing of novel supplements (i.e. mineral boluses, vitamins, botanicals) are

not subjected to as rigorous a regulatory process relative to veterinary pharmaceutical products. As

such, there is a paucity of peer-reviewed literature evaluating such novel products in the field. The

present study, though not designed to be a noninferiority or superiority study, does display that the

novel product is able to cause an elevation in serum calcium of recently calved cows of a magnitude

similar to another commercially available, efficacious product of similar composition.

Subclinical hypocalcemia (serum calcium levels < 2.0 mmol/l) is a pervasive issue in the modern dairy

industry. All cows in the current study experienced subclinical hypocalcemia. Previous research has

found that the incidence to be approximately 50% in mature (2^{nd} or greater lactation) Holstein cows

[4,12]. Indeed, the risk of displaced abomasum, ketosis, retained placenta, and metritis was 3.7, 5.5,

3.4, and 4.3 times more likely, respectively, in cows that experience subclinical hypocalcemia relative to

normocalcemic cohorts [13]. Overall, the costs associated with subclinical hypocalcemia are estimated

to be approximately $125 per case [2]. For the average 500 cow herd, assuming 65% of the herd is in

their 2^{nd} lactation or greater, and a 50% incidence of subclinical mastitis, the cost would be

approximately $20,312.50. Clearly, significant efforts should be made to prevent such losses.

Cows that received both calcium supplements experienced significant increases in serum calcium levels;

however, these levels were not sustained for the remainder of the 24 hours following parturition. These

results are consistent with recent literature that found oral calcium supplementation effectively

increased serum calcium levels at the 1 and 24 hours postpartum relative to untreated controls [8,11].

One important difference between the current experiment and the cited studies was the use of negative

control animals. As the current study was a pilot field-based trial, and calcium supplementation is a

pervasive practice throughout the industry, it was difficult (potentially unethical) to recruit a suitable

herd that would allow cows at high risk for hypocalcemia to remain untreated. On the other hand,

numerous studies evaluating postpartum, serial serum calcium levels in cows not receiving

supplemental calcium either remain constant or significantly decrease relative to baseline levels at

calving [8,11,14–16]. Therefore, although no definitive conclusions can be made, it would be a fair

assumption that had a negative control group been included in the experiment, their serum calcium

would likely not have increased at the 1-hour sample post-calving, and that calcium-supplemented cows

would likely have had higher serum calcium levels at this timepoint.

225　Conclusion

226　Subclinical hypocalcemia is a prevalent disorder of the modern commercial dairy cow. The current pilot

227　study found oral supplementation with two commercially available calcium boluses was associated with

228　a significant increase in serum calcium levels at 1-hour post-calving.

229　References

1.　Goff JP. The monitoring, prevention, and treatment of milk fever and subclinical hypocalcemia in

dairy cows. Vet J [Internet]. 2008;176(1):50–7. Available from:

http://dx.doi.org/10.1016/j.tvjl.2007.12.020

2.　Oetzel GR. Oral calcium supplementation in peripartum dairy cows. Vet Clin North Am - Food

Anim Pract [Internet]. 2013;29(2):447–55. Available from:

http://dx.doi.org/10.1016/j.cvfa.2013.03.006

3.　DeGaris PJ, Lean IJ. Milk fever in dairy cows: A review of pathophysiology and control principles.

Vet J [Internet]. 2008;176(1):58–69. Available from: http://dx.doi.org/10.1016/j.tvjl.2007.12.029

4.　Reinhardt TA, Lippolis JD, McCluskey BJ, Goff JP, Horst RL. Prevalence of subclinical hypocalcemia

in dairy herds. Vet J [Internet]. 2011;188(1):122–4. Available from:

http://dx.doi.org/10.1016/j.tvjl.2010.03.025

5.　McLaren CJ, Lissemore KD, Duffield TF, Leslie KE, Kelton DF, Grexton B. The relationship between

herd level disease incidence and a return over feed index in Ontario dairy herds. Can Vet J.

2006;47(8):767–73.

6.   Curtis CR, Erb HN, Sniffen CJ, Smith RD, Powers PA, Smith MC, et al. Association of parturient

hypocalcemia with eight periparturient disorders in Holstein cows. J Am Vet Med Assoc.

1983;183(5):559–61.

7.   Seifi HA, LeBlanc SJ, Leslie KE, Duffield TF. Metabolic predictors of post-partum disease and

culling risk in dairy cattle. Vet J [Internet]. 2011;188(2):216–20. Available from:

http://dx.doi.org/10.1016/j.tvjl.2010.04.007

8.   Sampson JD, Spain JN, Jones C, Carstensen L. Effects of calcium chloride and calcium sulfate in an

oral bolus given as a supplement to postpartum dairy cows. Vet Ther. 2009;10(3):131–9.

9.   Oetzel GR, Miller BE. Effect of oral calcium bolus supplementation on early-lactation health and

milk yield in commercial dairy herds. J Dairy Sci [Internet]. 2012;95(12):7051–65. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0022030212007229

10.   McArt JAA, Oetzel GR. A stochastic estimate of the economic impact of oral calcium

supplementation in postparturient dairy cows. J Dairy Sci [Internet]. 2015;98(10):7408–18.

Available from: http://linkinghub.elsevier.com/retrieve/pii/S0022030215005354

11.   Domino AR, Korzec HC, McArt JAA. Field trial of 2 calcium supplements on early lactation health

and production in multiparous Holstein cows. J Dairy Sci [Internet]. 2017;100(12):9681–90.

Available from: http://linkinghub.elsevier.com/retrieve/pii/S0022030217308998

12.   Horst RL, Goff JP, Reinhardt TA. Role of vitamin D in calcium homeostasis and its use in

prevention of bovine periparturient paresis. Acta Vet Scand. 2003;Suppl.(97):35–50.

13.   Rodríguez EM, Arís A, Bach A. Associations between subclinical hypocalcemia and postparturient

diseases in dairy cows. J Dairy Sci [Internet]. 2017;100(9):7427–34. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0022030217306331

14.   Martinez N, Sinedino LDP, Bisinotto RS, Daetz R, Risco CA, Galvão KN, et al. Effects of oral calcium

267   supplementation on productive and reproductive performance in Holstein cows. J Dairy Sci

268   [Internet]. 2016;99(10):8417–30. Available from:

269   http://linkinghub.elsevier.com/retrieve/pii/S0022030216304519

270   15.   Miltenburg CL, Duffield TF, Bienzle D, Scholtz EL, LeBlanc SJ. Randomized clinical trial of a calcium

271   supplement for improvement of health in dairy cows in early lactation. J Dairy Sci [Internet].

272   2016;99(8):6550–62. Available from:

273   http://linkinghub.elsevier.com/retrieve/pii/S0022030216303137

274   16.   Roberts K, Bennison J, McDougall S. Effect of treatment with oral Ca boluses following calving on

275   concentrations of Ca in serum in pasture-based dairy cows. N Z Vet J [Internet]. 2018;0(0):1–7.

276   Available from: https://www.tandfonline.com/doi/full/10.1080/00480169.2018.1520654

Figure

Figure

Figure

Figure