Connecting micro and macroevolution using genetic incompatibilities and natural selection on

additive genetic variance

Greg M. Walter[1]*, J. David Aguirre[2], Melanie J Wilkinson[1], Mark W. Blows[1], Thomas J. Richards[3] and

Daniel Ortiz-Barrientos[1]

[1]University of Queensland, School of Biological Sciences, St. Lucia QLD 4072, Australia

[2]Massey University, Institute of Natural and Mathematical Sciences, Auckland 0745, New Zealand

[3]Swedish University of Agricultural Science, Department of plant Biology, Uppsala, 75007, Sweden

* Corresponding Author: Greg M. Walter

9   Email: g.walter@bristol.ac.uk

10   Address: School of Biological Sciences, Life Sciences Building

11   University of Bristol, Bristol, UK BS8 1TQ

12   Phone: +44 7377 074 175

13 **Running head:** Divergent natural selection on additive genetic variance

14 **Data archiving:** Data will be deposited in Dryad on acceptance.

15 **Keywords:** natural selection, trade-offs, additive genetic variance, selection gradient, response to

16 selection, adaptive divergence, adaptive radiation, genetic incompatibilities, reproductive isolation,

17 macroevolution

Abstract

Evolutionary biologists have long sought to identify the links between micro and macroevolution to better understand how biodiversity is created. Despite the pursuit, it remains a challenge to understand how allele frequency changes correlate with the evolution of morphological diversity, and the build-up of reproductive isolation. To connect micro and macroevolution, we tested the adaptive importance of alleles underlying genetic incompatibilities, and the consequences for predicting evolutionary trajectories. Using a quantitative genetics crossing design, we produced an F4 Advanced Recombinant Form (ARF) between four contrasting ecotypes, which we phenotyped in the glasshouse (N=770) and transplanted into the four natural habitats (N=14,265 seeds), alongside the parental ecotypes. F2 hybrid breakdown was associated with the loss of extreme phenotypes and environment-specific genetic variation in field performance. We found evidence of genetic trade-offs among environments, but only in axes describing smaller amounts of genetic variance for fitness. Habitats that showed stronger patterns of adaptive divergence for native versus foreign ecotypes, also showed lower genetic variance in fitness of the ARF. Integrating data from the field and glasshouse predicted patterns of selection on morphological traits in a similar direction to the parental ecotypes. Overall, our results provide strong empirical evidence linking ecotype specific alleles with fitness trade-offs, phenotypic divergence and the rise in genetic incompatibilities among recently derived ecotypes. Our data connects microevolutionary change with macroevolution through adaptive radiation, where increases in environment specific alleles create changes in the distribution of genetic variance, ameliorating genetic constraints to adaptation as adaptive divergence proceeds.

Introduction

Despite the fundamental importance of divergent natural selection for adaptive evolution, research lacks a

clear empirical relationship between micro and macroevolution, largely due to the difficulty of estimating

natural selection in the wild (Pujol et al. 2018), and connecting it to the accumulation of reproductive

isolation (Baack et al. 2015). There are some examples where divergent natural selection acting on genes

of major effect (e.g., *Eda* gene in sticklebacks) describe patterns of adaptive divergence (Barrett et al.

2008), but few studies have connected divergence in genetic variation underlying quantitative traits with

observed evolutionary trajectories and the accumulation of reproductive isolation. Consequently, a more

detailed understanding of natural selection is required to understand if alleles underlying adaptation are

also those contributing to reproductive isolation, and how this creates adaptive radiation.

Alleles can confer an adaptive advantage in one environment, but with neutral or deleterious effects in

other environments, leading to fitness tradeoffs (Anderson et al. 2011; Anderson et al. 2013). If such

environment specific alleles evolve with reduced or restricted gene flow, they will be novel in relation to

genotypes from alternative environments and fail when tested in alternative genetic backgrounds, creating

reproductive isolation (Coyne and Orr 2004). Under this scenario, environment specific alleles will lead

to Bateson-Dobzhansky-Muller genetic incompatibilities when incompatible with alternative genetic

backgrounds (Bateson 1909; Dobzhansky 1937; Muller 1942), meaning alleles underlying adaptive traits

in one ecotype may lead to hybrid breakdown when they are introgressed into an alternative ecotype

(Gavrilets 2003; Kondrashov 2003; Navarro and Barton 2003). If alleles underlying adaptation to

contrasting environments concomitantly create fitness trade-offs and reproductive isolation, we can use

changes in environment specific allele frequencies to connect micro and macroevolution.

Natural selection is unlikely to affect single traits in isolation, favoring beneficial combinations of traits

and the evolution of multivariate phenotypes (Lande 1979; Cheverud 1982). Adaptation will be

constrained when traits share genetic variance, and genetic architecture rather than natural selection

determines evolutionary trajectories (Lande and Arnold 1983; Arnold 1992; Arnold et al. 2008). In this

way, adaptive alleles will likely increase in frequency, but only if selection on genetically correlated traits

allow it. Genetic correlations are expected to remain stable, at least in the short term, which would make

rapid adaptive divergence leading to adaptive radiation difficult (Walsh and Blows 2009). However,

recent studies have shown that G-matrices can potentially evolve rapidly, questioning the role of

constraints in adaptive radiation (Doroszuk et al. 2008; Eroukhmanoff and Svensson 2011; Walter et al.

2018a). If genetic correlations can evolve in response to natural selection, then environment-specific

allele frequency changes can overcome genetic constraints to promote rapid adaptive divergence (Sgrò

and Hoffmann 2004), and the path to adaptive radiation will be more straightforward.

We explore these predictions within the adaptive radiation of an Australian native wildflower, *Senecio*

pinnatifolius. We focus on four ecotypes within this species complex including two coastal types found

on sandy dunes (Dune ecotype, *Senecio pinnatifolius* var. *pinnatifolius*) and rocky headlands (Headland

ecotype, *S. pinnatifolius* var. *maritimus*), and two inland ecotypes that occur in moist sub-tropical

rainforest (Tableland ecotype, *S. pinnatifolius* var. *serratus*) and dry sclerophyll woodland (Woodland

ecotype, *S. pinnatifolius* var. *dissectifolius*) (Ornduff 1964; Ali 1969; Radford et al. 2004). Ecotypes are

generally semi-perennial, with the exception of the annual Woodland ecotype. They are obligate

outcrossers and share insect pollinators. Previous work has shown that these ecotypes arose as a result of

adaptation to divergent natural selection (Roda et al. 2013a; Melo et al. 2014; Walter et al. 2016),

resulting in distinct environment-specific plant morphologies (Walter et al. 2018a). It is likely this

association is the result of the accumulation of adaptive, environment specific alleles, evidenced by the

existence of fitness trade-offs (Walter et al. 2018b) and immigrant inviability among contrasting habitats

(Melo et al. 2014; Richards and Ortiz-Barrientos 2016; Richards et al. 2016; Walter et al. 2016). Artificial

hybridization among ecotypes produced vigorous F1 offspring, with hybrid breakdown observed at the F2

generation as a strong reduction in reproductive capacity and suggesting incompatible alleles have arisen

among ecotypes. Fitness was recovered in the subsequent F3 generation, suggesting genetic

incompatibilities arose as a breakup of coadapted gene complexes (Walter et al. 2016). Overall, these

ecotypes display strong evidence of having undergone a recent adaptive radiation.

Here, we employ a combination of glasshouse and field experiments to explore the implications of

simulated gene flow during adaptive radiation. We use a quantitative genetic crossing design to create an

F4 generation advanced recombinant form (ARF) between the four ecotypes. Alongside the parental

ecotypes, we phenotyped the ARF in the glasshouse, and performed a large-scale field transplant across

the four natural habitats to quantify divergent natural selection. We tested three predictions: 1) If

environment specific adaptive alleles underlie both phenotypic divergence and genetic incompatibilities,

F2 breakdown should result in a reduction in phenotypic variance in the ARF population. 2) A purge of

environment specific alleles following F2 breakdown should reduce genetic trade offs in the ARF

exposed to contrasting environments in the field, and 3) If trait differences between ecotypes are the

result of divergent natural selection, selection gradients measured on the ARF under field conditions

should align with the direction of trait divergence in the parental ecotypes.

Methods

Crossing design

To create the ARF we first sampled seeds from one natural population from each of the four ecotypes,

which we germinated and grew at the University of Queensland glasshouses. We sampled seeds for the

Dune and Headland ecotypes at Lennox Head, NSW (*-28.783005, 153.594018* and *-28.813117,*

153.605319, respectively), from the Tableland ecotype at O'Reilley's Rainforest Retreat, Qld (-

28.230508, 153.135078) and the Woodland ecotype at Upper Brookfield, Qld (*-27.479946, 152.824709*).

At each location, we collected seeds from 24-49 plants separated from each other by at least 10 m to

minimize the likelihood of sampling close relatives. Two seeds from each individual sampled were

germinated and grown in the University of Queensland glasshouses, which then formed the base

population for our crossing design, outlined below. To grow plants, we first scarified each seed and

placed them in glass petri dishes containing moist filter paper. After leaving them in the dark for two days

we transferred the germinated seeds to a 25°C constant temperature growth room with 12h:12h light:dark

photoperiod. After one week, we transferred the seedlings to the glasshouse and transplanted them into

85mm pots containing a mixture of 70% pine bark and 30% coco peat with slow release osmocote

fertilizer and 830g/m^3 of Suscon Maxi insecticide. We conducted controlled crosses on mature plants by

rubbing two mature flower heads together, labeling the flower heads and collecting the seeds as they

emerged.

We created the ARF ensuring each ecotype contributed equally and ensuring that at each generation (see

Figure 1C), all full-sibling families (hereafter, 'families') contributed equally to the next generation. First,

we grew plants for the base population from seeds sampled from the natural populations and performed

crosses among the ecotypes ($n = 41$-60 individuals/ecotype) to create all combinations of F1 hybrids ($n =$

12 crossing combinations; $n = 20$-25 families/cross type). We then mated among all combinations of

crosses in the F1 generation such that all F2 families ($n = 24$ crossing combinations; $n = 17$-22 families

/cross type) possessed a grandparent from each of the original parental ecotypes (e.g., $F1_{Dune,Headland} \times$

$F1_{Tableland,Woodland}$). Given strong reductions in intrinsic fitness was observed in a previous Dune x

Headland F2 hybrid (Walter et al. 2016), we maximized the number of F1 crosses to produce 458 F2

families in total. We grew one individual from each family. Reductions in fitness were observed as F2

hybrid sterility (42% of F2 individuals were successfully mated compared to >90% in F1 hybrids) and

reduced fertility (49% reduction in seed set compared to F1 hybrids) (Walter et al. 2016). Consequently,

we divided the F2 individuals that produced flowers into three replicate crossing lines to maintain

replicates of the construction of the ARF. We then randomly mated among all F2 individuals within each

line ($n = 4$-12 families/F2 cross type; total F2 families crossed N = 202) to produce the F3 generation (N

= 259 families), ensuring that each family contributed equally. We then produced the F4 generation by

first growing one individual from each F3 family and randomly designating each individual as a sire or

dam. We then mated 115 sires to 114 dams in a full-sibling, half-sibling crossing design to produce 198

families for the F4 generation. The numbers of families and individuals used to create each generation of

the ARF are listed in Table S1.

In the following analyses we examine results from two experiments using the ARF. In experiment 1, we

grew the ARF in the glasshouse to estimate genetic variance underlying morphological traits. In

experiment 2, we transplanted seeds of the ARF into the four habitats to compare the fitness of the ARF

with the parental ecotypes. We then used the field fitness of the ARF (experiment 2) to quantify the

genetic covariance in performance among transplant habitats and identify genotype-by-environment

interactions that would indicate genetic trade-offs among habitats. Finally, to quantify differences in

natural selection among habitats, we combined the data from both experiments and estimated the genetic

covariance between morphological traits in the glasshouse (experiment 1) and field fitness in each habitat

(experiment 2).

Experiment 1: Glasshouse phenotypes

To estimate genetic variance underlying morphological traits we grew four individuals from each full

sibling family of the ARF (n = 198 full-sibling families, total N = 770 individuals) in 30 cell growth trays

containing the same potting media described above. Alongside the ARF we grew four individuals from

~25 full sibling families for each of the parental ecotypes (N = 366 individuals). Plants were grown in a

25°C controlled temperature room with a 12h:12h day:night photoperiod. After eight weeks of growth we

measured plant height and sampled one fully mature leaf for each plant. We used the software 'Lamina'

to analyse the scanned leaf and quantify six variables relating to leaf size and leaf shape (Bylesjo et al.

2008). Using the outputs of Lamina, we quantified leaf morphology using leaf area, leaf area^2 / leaf

perimeter^2 as a measure of leaf complexity, leaf circularity, number of indents standardized by leaf

perimeter, leaf indent width and leaf indent depth.

Experiment 2: Field transplant

Seeds from the F4 generation of the ARF were transplanted into each of the habitats. At each transplant

habitat, we planted 18 seeds from each full-sibling family (n = 198) divided equally amongst six

experimental blocks (habitat n ≈ 3,500 seeds, total N = 14,265 seeds). Alongside the ARF we

transplanted seeds from the populations of parental ecotypes used to create the ARF (for each population

n = 180 seeds/habitat) (analysed previously in Walter et al. 2016). See Walter et al. (2016); Walter et al.

(2018b) for a detailed description of the field experiment. Briefly, we glued each seed to a toothpick

using non-drip glue and planted them in 25mm x 25mm plastic grids in March 2014. Field observations

suggested that seeds in the natural populations can germinate year-round given sufficient rain. Given we

wanted to standardise germination time to estimate post-germination development and survival, to

replicate natural germination conditions we suspended shadecloth (50%) 15cm above each experimental

block and watered them daily for three weeks. During the initial three-week period we measured

emergence and mortality daily. Following the initial three weeks we measured survival and development

at weeks 4, 5, 7 and 9, and then monthly until 20 months at which time there were fewer than 20% of

germinated plants remained, and we ceased the experiment. The measures of fitness we recorded were:

whether each seedling emerged, whether each seedling reached 10 leaves (as a measure of seedling

establishment) and produced a bud (reached maturity). All measures of fitness were collected as binary

data.

Implementation of Bayesian models

In the subsequent analyses we implemented Bayesian models to 1) compare field performance

(experiment 2) of the ARF with the parental ecotypes, 2) identify whether genotype-by-environment

interactions create genetic trade-offs among transplant habitats (experiment 2), 3) estimate genetic

(co)variance of morphological traits for the ARF (experiment 1), and 4) estimate the genetic covariance

between morphological traits (experiment 1) and field performance (experiment 2), to identify differences

in natural selection on morphological traits, among habitats.

All Bayesian models were implemented using R (R Core Team 2016) within the package 'MCMCglmm'

(Hadfield 2010). From each model we extracted 1,000 Markov chain Monte Carlo (MCMC) samples,

which provided the posterior distribution for the parameters we were estimating. For each analysis, we

implemented Markov chains of different lengths (listed in Table S2), while ensuring that we included a

sufficient burn-in period and thinning interval to sample the parameters with autocorrelation values of

less than 0.05 and effective sample sizes exceeding 85% of the total number of samples, for all

parameters. We used uninformative parameter expanded priors and checked their sensitivity by re-

implementing all models while adjusting the parameters and ensuring the posterior distribution did not

change.

For the analyses estimating genetic variance, comparing estimates of genetic variance with zero provides

an uninformative test of significance because estimates are restricted to be greater than zero (positive-

definite). To create an informative significance test, we re-implemented each model with randomized

data, created by shuffling the parental information. For each model implemented on the observed data, we

re-implemented the same model on 1,000 randomizations of the data, and extracted the posterior mean for

each randomization. We then compared the distribution of means from models conducted on the

randomizations, to the mean of the observed posterior distribution. If the mean of the observed

distribution occurred outside the 95% Highest Posterior Density (HPD) interval for the random

distribution, we took this as evidence that we captured biologically important information for the

comparison of interest. As we were only interested in estimating the posterior mean of models

implemented on each randomization of the data, we could reduce computing time by reducing the total

number of sampling iterations. To do so, we maintained the same burn-in period and sampling interval to

ensure an identical mixing of MCMC chains, reducing only the total number of sampling iterations to the

number required to obtain a stable estimate of the mean. We calculated the number of sampling iterations

required using the models implemented on the observed data, which was different for each of the analyses

outlined below (Table S2).

Comparing ARF and ecotype morphology

To compare differences in multivariate phenotype between the ARF and parental ecotypes, we

implemented a multivariate analysis of variance (MANOVA) on the seven morphological traits measured

in experiment 1. We first standardized all seven morphological traits to a mean of zero, and standard

deviation of one before including them as a multivariate response variable. To test whether the ARF was

phenotypically different to each ecotype we conducted a separate MANOVA for each pairwise

comparison between the parental ecotypes, and the ARF. We used a bonferroni corrected α-value of

0.0125 (α = 0.05 / *n*, where *n* represents the number of tests). To visualize differences among all ecotypes

and the ARF we estimate **D**, the variance-covariance matrix representing multivariate phenotypic

divergence. To do so, we first conducted another MANOVA that included all ecotypes (but not the ARF).

From this, we extracted the sums of squares and cross-product matrices for the ecotypes ($SSCP_H$) and

error terms ($SSCP_E$) to calculate their mean-square matrices by dividing by the appropriate degrees of

freedom ($MS_H = SSCP_H / 3$; $MSE = SSCP_E / 365$). Using the mean-square matrices we calculated **D** =

($MS_H - MS_E$) / *nf*, where *nf* represents the number of measured individuals per genotype in an unbalanced

design, calculated using equation 9 in Martin et al. (2008). Our D-matrix then represents divergence in

multivariate mean phenotype, among the parental ecotypes, after removing the residual phenotypic

variation. To visualize the phenotypic space occupied by the ARF relative to the parental ecotypes, we

decomposed **D** into orthogonal axes (eigenvectors) and calculated the phenotype scores for the first two

eigenvectors for all ecotypes, and the ARF.

Comparing ARF and ecotype field performance

30   We estimated fitness at early life history stages for the ARF and parental ecotypes transplanted into all

31   four habitats. To do so, we created a dummy variable that represented the ARF and native versus foreign

32   ecotypes in each habitat. We then used MCMCglmm to implement the model,

$$y_{ijklm} = H_i + P_j + H_i{\times}P_j + B_{k(i)} + L_{l(j)} + e_{m(ijkl)} \,, \qquad (1)$$

where transplant habitat (H_i), ARF/ecotype (P_j) and their interaction ($H_i{\times}P_j$) were included as fixed

effects. Blocks within transplant habitat ($B_{k(i)}$) and replicate genetic lines within the ARF ($L_{l(j)}$) were

included as random effects, and $e_{m(ijkl)}$ represented the model error. We implemented equation 1 with

emergence, seedling establishment and maturity as a multivariate response variable (y_{ijklm}). As such, for

all ecotypes and the ARF, equation 1 calculated the probability of reaching maturity, conditional on the

previous life history stages.

Quantifying divergent natural selection

We used the ARF to investigate differences in natural selection among contrasting natural habitats. To do

so, we conducted two further analyses. First, to identify whether natural selection created genetic trade-

offs among habitats, indicated by a negative genetic covariance among habitats, we analysed field

performance using a genotype-by-environment covariance framework described below. Next, we

examined whether genetic selection on traits occurred in the direction of the native ecotypes. To do so,

we combined the morphology data from experiment 1, with field performance data in experiment 2 and

used the Robertson-Price Identity to estimate the genetic covariance between morphological traits and

field performance for each transplant habitat. We predicted that if natural selection on morphology

occurred in the direction of the original ecotypes, differences in selection gradients (among habitats)

would align with divergence in phenotype mean of the parental ecotypes.

 Genetic trade-offs among contrasting habitats

We investigated genotype-by-environment (G×E) interactions in the ARF using a character state

approach, where different environments represent different traits (Robinson and Beckerman 2013). To do

so, we used the field performance of the ARF and implemented

$$y_{ijklmn} = L_i + H_j + S_{k(il)} + D_{l(ik)} + B_{m(j)} + e_{n(ijklm)} \, , \qquad (2)$$

where replicate genetic line of the ARF (L_i) and transplant habitat (H_j) were included as fixed effects.

We included sire ($S_{k(il)}$), dam ($D_{l(ik)}$) and block within habitat ($B_{m(j)}$) as random effects, with

$e_{n(ijklm)}$ representing the residual error variance. For each term in the random component, we estimated

random intercepts for each habitat and the covariance among habitats. As such, for the sire and dam

components we estimated a 4×4 covariance matrix representing variance in each habitat, and covariance

among habitats. Information for estimating covariance among habitats is taken from individuals of the

same full-sibling families transplanted in each habitat. Consequently, we implemented equation 2 with a

heterogeneous residual covariance matrix. This allowed for different variances in each habitat, but fixed

residual covariances at zero because individuals (seeds) could not be planted in two habitats

simultaneously. We used three separate implementations of equation 2 for emergence, seedling

establishment and maturity included as binary univariate response variables ($\mathbf{y}_{ijklmn}$).

From equation 2, $\mathbf{S}_{l(im)}$ represents one quarter of the additive genetic variance in each habitat, and one

quarter of the additive genetic covariance between habitats. We multiplied the sire variance component

($\mathbf{S}_{l(im)}$) by four and used the posterior mean as our observed estimate of additive genetic (co)variance for

field performance among the four habitats. The diagonal of the resulting G-matrices represents additive

genetic variance within a transplant habitat, with the off diagonal representing the genetic covariance

between habitats.

Divergent natural selection on morphological traits

By linking genetic variance underlying morphological traits in the laboratory, with genetic variance

underlying field performance, we sought to quantify differences in natural selection among the transplant

habitats using the Robertson-Price Identity. A requirement for natural selection is genetic variance in both

morphological traits and field performance. The analysis of genetic variance underlying field

performance (described in the previous section) identified significant genetic variance for the ability to

reach maturity, in all four transplant habitats (see results). To identify the morphological traits with

genetic variation we used the morphology data from experiment 1 and implemented

$$y_{ijklmn} = L_i + S_{j(ik)} + D_{k(ij)} + S_{j(ik)} \times D_{k(ij)} + e_{n(ijk)} , \qquad (3)$$

where replicate genetic line of the ARF (L_i) was included as a fixed effect and sire ($S_{j(ik)}$), dam ($D_{k(ij)}$)

and their interaction ($S_{j(ik)} \times D_{k(ij)}$) were random effects, with $e_{n(ijk)}$ as the residual variance. We

implemented equation 3 with plant height and six leaf morphology traits as a multivariate response

variable (y_{ijkl}). To prevent traits on different scales affecting the analysis, we centered all traits to a mean

of zero and standardized to a standard deviation of one prior to analysis. We then calculated the additive

genetic (co)variance matrix as four times the sire variance component. As traits were standardized prior to

analysis, genetic variances represent heritabilities. We found only four traits with heritabilities greater

than 0.1 (plant height, leaf area, leaf perimeter^2 / area^2 and leaf indent width; see Table S3), which we

then combined with field performance to study natural selection in the subsequent analyses, described

below.

We estimated the genetic covariance between the morphological traits and field performance by

implementing the Robertson-Price Identity

$$R = s_g = cov(w, z) , \qquad (4)$$

where the response to selection (R) is analogous to the selection differential (s_g), calculated as the genetic

covariance between a trait (z) and fitness (w). Equation 4 then generalizes to multivariate form by

including more phenotypic traits and estimating a genetic variance-covariance matrix (**G**), with fitness as

the final trait. In this framework, s_g generalizes to the vector of selection gradients (s_g) representing the

multivariate response to selection. Estimating the response to selection in this way includes both direct

00 and indirect selection. To isolate the effect of direct selection on phenotypic responses, we can calculate

01 the genetic selection gradient by combining **G** and s_g with

$$\boldsymbol{\beta}_g = \boldsymbol{G}^{-1}\boldsymbol{s}_g , \qquad (5)$$

03 where $\boldsymbol{\beta}_g$ now represents a vector of genetic selection gradients (Lande and Arnold 1983; Rausher 1992),

04 after removing the effect of genetic correlations among traits.

05 To estimate the predicted response to selection (s_g) in the ARF we estimated the (co)variance between the

06 four morphology traits and field performance by implementing

$$\boldsymbol{y}_{ijklm} = \boldsymbol{L}_i + \boldsymbol{S}_{j(ik)} + \boldsymbol{D}_{k(ij)} + \boldsymbol{S}_{j(ik)} \times \boldsymbol{D}_{k(ij)} + \boldsymbol{B}_l + \boldsymbol{e}_{m(ijkl)} , \qquad (6)$$

where replicate genetic line ($\boldsymbol{L}_i$) was the only fixed effect. Sire ($\boldsymbol{S}_{j(ik)}$), dam ($\boldsymbol{D}_{k(ij)}$) and their interaction

($\boldsymbol{S}_{j(ik)} \times \boldsymbol{D}_{k(ij)}$) were included as random effects along with block within habitat ($\boldsymbol{B}_l$). The multivariate

response variable ($\boldsymbol{y}_{ijklm}$) included four phenotypic traits as well as ability to reach maturity in each

habitat. Fitness and morphology was measured on separate individuals (field versus glasshouse

experiments), and so similar to equation 2, we estimated a heterogeneous residual covariance matrix.

Multiplying the sire variance component ($\boldsymbol{S}_{j(ik)}$) by four (from equation 6) gave the additive genetic

variance-covariance matrix (**G**). Elements in the first four rows and columns represented **G** among

morphological traits. Covariance elements in the fifth column (and row) denote the genetic covariance

between each trait and fitness (s_g), with genetic variance in fitness in the final element (fifth row, fifth

column); see Stinchcombe et al. (2014) for details. We used four separate implementations of equation 6

for field performance as the ability to reach maturity in each of the four transplant habitats. We calculated

the additive genetic (co)variance matrix as four times the sire variance component and extracted s_g as the

vector of covariances between morphological traits and field performance (rows one to four of the fifth

column).

To identify whether we captured biologically meaningful differences in selection among habitats, we

conducted two analyses. First, s_g and $\boldsymbol{\beta}_g$ being vectors, we calculated the dot product (representing vector

length) of the observed and random matrices. If the observed length was greater than the length calculated

from the random distribution, we took this as evidence we detected biologically meaningful estimates of

selection for a given habitat (Stinchcombe et al. 2014; Walsh and Lynch 2018). Second, to quantify the

differences in s_g among the four transplant habitats we estimated variance is selection gradients using

$$\mathbf{Z} = \begin{pmatrix} \sigma^2(s_{g1}) & \sigma(s_{g1}, s_{g2}) & \dots & \sigma(s_{g1}, s_{gn}) \\ & \sigma^2(s_{g2}) & \ddots & \vdots \\ & & & \sigma^2(s_{gn}) \end{pmatrix}, \qquad (7)$$

where $\mathbf{Z}$ then represents the among-habitat variance in s_g for the nth trait along the diagonal. The off-

diagonal then contains the covariance in s_g among habitats, for each bivariate trait combination

(Chenoweth et al. 2010). In the same way, we used equation 7 to calculate $\mathbf{B}$, the among-habitat

(co)variance in $\boldsymbol{\beta}_g$. Comparing the eigenvalues of observed and random (for both $\mathbf{B}$ and $\mathbf{Z}$) provided tests

of significance. Observed eigenvalues with values greater than the random distribution of eigenvalues

suggested we captured greater among-habitat differences in selection than expected by random sampling.

Results

Comparing ARF and ecotype morphology

Ecotypes showed strong differences in leaf morphology (Figure 1A), with the ARF exhibiting large

variation, intermediate to the parental ecotypes (Figure 1B). Pairwise MANOVAs showed the ARF was

significantly different to all ecotypes (Dune: Wilks' $\lambda = 0.71$, $F_{1,857} = 50.782$, $P = <0.001$; Headland:

Wilks' $\lambda = 0.64$, $F_{1,863} = 69.747$, $P = <0.001$; Tableland: Wilks' $\lambda = 0.38$, $F_{1,866} = 192.36$, $P = <0.001$;

Woodland: Wilks' $\lambda = 0.22$, $F_{1,864} = 445.1$, $P = <0.001$). The MANOVA conducted on only the parental

ecotypes described a significant difference among ecotypes in multivariate mean phenotype (Wilks' $\lambda =$

0.03, $F_{3,362} = 117.86$, $P = <0.001$), where differences among ecotypes captured 64% of the total variance.

The first eigenvector of $\mathbf{D}$ (d_{max}) described 84% of phenotypic divergence mostly created by phenotypic

differences between the Tableland and Headland ecotypes (Figure 1D). The second eigenvector (d_2)

described 14% of variation in multivariate phenotype, describing differences between the Woodland and

Tableland ecotypes (Figure 1D). The ARF occupied an area in phenotypic space close to the Dune

48 ecotype, and intermediate between the Headland, Tableland and Woodland ecotypes. However, the ARF

49 mean was not similar to that of the overall mean of all ecotypes, but exhibited high phenotypic variance

50 that appeared to be missing some of the extreme phenotypes, especially from the Tableland and

51 Woodland ecotypes (Figure 1D).

52
Figure 1: A) Ecotypes vary dramatically in leaf morphology. **B)** The ARF exhibited large variation in leaf morphology,
visually intermediate among the original ecotypes. **C)** The ARF was created by equally mating among all ecotypes. **D)** The
distribution of ecotype and ARF scores for the first two axes of **D** showing the ARF (grey) occupying an area in phenotypic
space similar to the mean of all ecotypes (black), but lacking extreme phenotypes, especially of the Tableland and Woodland
ecotypes.

Comparing ARF and ecotype field performance

Given ecotypes have shown adaptation to their contrasting habitats (Walter et al. 2016; Walter et al.

2018b), we expected that as an intermediate form, the ARF would show intermediate performance

between native and foreign populations. We found the performance of the ARF was similar to the native

ecotypes for seedling establishment and maturity (Figure 2), suggesting hybrid vigor despite several

generations of recombination.

Figure 2: Field performance of the ARF compared to foreign (F) and native (N) ecotypes in each transplant habitat. Fitness measured as the probability of reaching **A)** seedling establishment, and **B)** maturity. Credible intervals represent 95% HPD intervals.

Genetic trade-offs among contrasting habitats

If habitat-specific natural selection creates genetic trade-offs between contrasting habitats, we expected

the ARF to show genetic variance for field fitness, and negative genetic covariance between contrasting

habitats. However, given the ARF lacked the extreme phenotypes of the Headland, Tableland and

Woodland ecotypes (Figure 1), and exhibited relatively high field performance (Figure 2), we might

expect low genetic variance associated with either zero genetic covariance or a positive genetic

covariance between habitats. We found that across the four habitats, additive genetic variance increased

as life history stages progressed (Figure 3). Observed estimates of genetic variance in field performance

were within the random distribution at emergence, but were greater than the random distribution for

maturity (Figure 3). In the headland and tableland habitats we detected lower genetic variance than

expected by chance for seedling establishment.

Decomposing the genetic covariance matrix described orthogonal axes of genetic variation underlying field fitness. Decomposing the matrix for each life history stage, we found the first three eigenvectors for maturity described more genetic variance than expected under random sampling (Figure 4). Interpreting the loadings of each eigenvector reveal how each habitat contributes to describing the genetic variance in fitness quantified by that eigenvector. Habitats with loadings of the same sign describe shared genetic variance for fitness, whereas loadings of different signs describe differences in genetic variance and provide evidence of fitness trade-offs. We found all habitats contributed equally to describing genetic variance underlying the first eigenvector, suggesting it described heterosis or shared genetic variation needed to function in stressful environments (Table 1). However, eigenvectors two and three provided evidence of genetic tradeoffs, describing genetic variance in fitness that differed between the woodland and dune ecotypes (e_2), and between the tableland, and the dune and woodland transplant habitats (e_3; Table 1). Eigenvector 4 did not describe biologically meaningful genetic variance (Figure 4), but described differences in genetic variance between the headland, and dune and tableland habitats. The

97 posterior mean G-matrices and genetic correlations for field performance are located in supplementary

98 Table S4.

99
Figure 4: Comparing the amount of genetic variance described by eigenvectors representing the observed (filled circles)
versus random matrices (unfilled circles and dashed lines), for each life history stage. Gray bars represent the amount of
genetic variance in the randomized matrices described by the observed eigenvectors. Only the first three eigenvectors for
maturity described more genetic variance than expected by random sampling. Credible intervals represent 95% HPD intervals.

Table 1: Eigenanalysis of the additive genetic (co)variance matrix for field performance at maturity. Loadings in bold are
greater than 0.25 to aid interpretation. HPD represents the observed 95% HPD credible intervals.

Eigenvectors	e_1	e_2	e_3	e_4
Eigenvalue	2.492	0.782	0.248	0.116
HPD	0.837-4.179	0.036-1.984	0.037-0.569	0.011-0.267
Proportion	0.685	0.215	0.068	0.032
Dune	**-0.56**	**0.61**	**0.45**	**-0.34**
Headland	**-0.34**	0.22	-0.04	**0.91**
Tableland	**-0.54**	-0.02	**-0.81**	**-0.23**
Woodland	**-0.53**	**-0.76**	**0.37**	0.00

Overall, our results showed strong patterns of adaptive divergence (Figure 2), and although there appears

to be a common genetic basis to fitness in all environments (e_1; Table 1) we also detected genetic trade-

offs for fitness among certain habitats (Table 1). Despite strong adaptive divergence in Figure 2, the

headland and tableland habitats exhibited weaker additive genetic variance for fitness (Figure 5), and

weaker genetic trade-offs with other environments (Table 1), when compared to the dune and woodland.

This suggested alleles lost during F2 hybrid breakdown contributed to both genetic incompatibilities and

adaptive genetic variation that was lost in the ARF, reducing genetic variance for field performance in

certain environments and producing weaker genetic trade-offs than expected. To test this, for each habitat

we compared the strength of adaptive divergence (Figure 2; native ecotype performance – foreign ecotype

performance) against the level of genetic variance exhibited by the ARF. As predicted, we found a strong

negative association for seedling establishment and a weaker negative association for maturity (Figure 5),

suggesting alleles associated with strong adaptive divergence were also responsible for genetic

incompatibilities.

20
Figure 5: Stronger local adaptation was negatively associated with the level of genetic variance. A comparison between the
strength of adaptive divergence (difference in fitness between the native ecotype and foreign ecotypes) versus the level of
genetic variance exhibited by the ARF, in the same habitat. Solid circles and lines represent seedling establishment, triangles
and dashed lines represent the ability to reach maturity. Credible intervals represent 95% HPD intervals. Estimating a
regression slope for each MCMC iteration showed a significant negative association at 88% HPD for seedling establishment,
but a non-significant relationship for maturity.

Natural selection on morphological traits

To quantify selection in each habitat we calculated s_g as the genetic covariance between morphological

traits measured in the glasshouse, and field performance measured in each of the four transplant habitats.

We then isolated direct selection by calculating $\boldsymbol{\beta}_g$, the genetic selection gradient for each habitat.

Comparing the length of observed and random s_g and $\boldsymbol{\beta}_g$ suggested we captured biologically meaningful

selection within each habitat (Figure S5A). To quantify differences in selection among habitats we

estimated **B** and **Z** as the among-habitat (co)variance in selection vectors. Comparison of observed and

random eigenvalues showed that both selection vectors exhibited greater differences among habitats than

expected by random sampling (Figure S5B), suggesting differences in our observed selection vectors

described biologically meaningful differences in natural selection among transplant habitats.

If differences in natural selection among the four habitats occurred in the direction of adaptive evolution,

we would expect differences in s_g, but not $\boldsymbol{\beta}_g$, to align with divergence in mean phenotype of the parental

ecotypes. Eigenanalyis of **B** and **Z** quantifies the axes that describe differences among the original

selection vectors, with the first axis for each matrix representing 83% (HPD 56-98%) and 81% (HPD 55-

98%) of the total variance, respectively. We tested whether the first axis from each selection vector

aligned with d_{max}, the axis describing the greatest difference in multivariate phenotype mean. To do so,

we calculated the angle between the first eigenvector of **B** and **Z**, and d_{max}. We found the alignment

between **Z** and **D**, but not **B,** was closer than expected with random sampling (Figure 6A). To

complement this analysis, we conducted a more extensive analysis using a covariance tensor approach,

which is provided in supplementary material. Results obtained from both analyses matched closely,

suggesting the response to selection, but not the direction of selection, aligned with divergence in parental

ecotype morphology.

49
Figure 6: Differences in s_g, but not $\boldsymbol{\beta}_g$, aligned with d_{max}, but differences in $\boldsymbol{\beta}_g$ aligned with d_2. **A)** The angle between the first
eigenvector of **Z**, and d_{max} was closer than expected by random sampling, but the first eigenvector of **B** did not show a close
alignment with d_{max}. Credible intervals represent 90% HPD intervals. **B)** Two-dimensional schematic approximately
representing the orientation of **B** and **Z** in relation to **D**, and d_{max}.

Discussion

The ARF exhibited a multivariate phenotype intermediate to the four parental ecotypes, and lacking in

some of the more extreme phenotypes. Field performance of the ARF was similar, or higher, than the

native ecotypes, suggesting heterosis has persisted. Despite this, we found trade-offs in field performance

among habitats, but only in axes describing smaller amounts of genetic variance underlying field fitness.

The ARF also showed reduced genetic variance for fitness when transplanted into habitats associated with

stronger adaptive divergence, providing evidence that divergent natural selection shaped genetic variation

in traits associated with fitness. Among-habitat differences in natural selection on morphological traits

aligned with the direction of morphological divergence of the original ecotypes, providing evidence for

divergent natural selection towards the original ecotypes, despite only one generation of selection.

Together, our results provide empirical evidence suggesting interactions between genetic

incompatibilities and divergent natural selection created adaptive radiation of *S. pinnatifolius* into four

contrasting habitats.

While there is abundant evidence implicating divergent natural selection in the accumulation of extrinsic

reproductive barriers, the contribution of natural selection to the evolution of genetic incompatibilities

during population divergence remains unresolved (Baack et al. 2015). Our results showed that genetic

incompatibilities were associated with the loss of extreme phenotypic variation, and the underlying genes

were likely adaptive. We suggest that environment-specific dominant alleles link extreme phenotypes

with natural selection and reproductive isolation to create adaptive radiation in these contrasting ecotypes.

This is because heterozygotes (with alleles from different ecotypes) at one or more loci underlie F2

hybrid breakdown, creating negative additive × dominant or dominant × dominant interactions (Demuth

and Wade 2005; Willett 2006), suggesting genetic incompatibilities at the F2 generation are largely

produced by dominant alleles (e.g., Sweigart et al. 2006; Latta et al. 2007). Our F2 hybrid was

constructed by mating between two completely unrelated F1 crosses (Figure 1C; e.g., F1$_{Dune,Headland}$ ×

F1$_{Tableland,Woodland}$), increasing heterozgosity compared to traditional F2 crosses between two populations,

and reducing the likelihood of homozygous recessive loci. Dominant alleles will be more visible to

selection, allowing them to increasing in frequency rapidly and create rapid adaptive divergence. Whether

these alleles then contribute to the evolution of stronger genetic incompatibilities (e.g., F1 hybrid

breakdown) remains unexplored.

F2 hybrid breakdown indicates population divergence as a build-up of coadapted gene complexes, created

when selection assembles beneficial combinations of alleles (Cutter 2012; Corbett-Detig et al. 2013). In

this system, it is likely the evolution of coadapted gene complexes, rather than speciation genes (Nosil

and Schluter 2011), were likely responsible for the rise of intrinsic reproductive isolation during the early

stages of speciation (Corbett-Detig et al. 2013). We can then view the evolution of these ecotypes from a

perspective that combines Fisher's geometric model (Fisher 1930) and Wright's focus on epistasis

(Wright 1931), where selection acts upon additive genetic variation by increasing allele frequencies at

independent loci, but limited recombination due to small population size or strong selection creates

coadapted gene complexes, making them functionally epistatic. The strength of coadaptation within a

population will then determine how genetic incompatibilities arise among populations and lead to

speciation.

The strength of divergence (and hence, reproductive isolation) among coadapted gene complexes will be

population and environment specific, and depend on the interaction between mutation, migration, drift

and selection. Previous studies of Dune-Headland parapatric pairs along the Australian coastline have

shown convergent evolution, suggesting multiple independent origins of these ecotypes (Roda et al.

2013a; Roda et al. 2013b; Roda et al. 2017). If the same dominant alleles important for adaptation to

these contrasting environments are repeatedly selected in the same environment, they may form

coadapted gene complexes within each population, with drift or local adaptation causing differences

among localities (Goodnight 2000). Whether locally adapted coadapted gene complexes between

locations of the same species will give rise to reproductive isolation remains unexplored, but could

provide important insights into the relationship between adaptation and divergence across a

heterogeneous landscape.

Genetic variance for life history and fitness traits is often low (e.g., McFarlane et al. 2014), and often

decreases with ontogeny (e.g., Aguirre et al. 2014). In contrast, our results suggested increases in genetic

variance with development, which has been found previously in laboratory conditions (Styga et al. 2018).

Changes in genetic variance underlying fitness have profound implications for understanding adaptation

and responses to environmental change (Sgrò and Hoffmann 2004). If genetic correlations among traits

under selection change during ontogeny, the effects of selection will not be linear over time and will

depend on changes in the combination of genetic variation and selection pressures over time. As different

trait combinations will be available to selection at different developmental points, patterns of adaptation

will be determined by the combination of traits visible when selection is strong (Bourret et al. 2017;

Styga et al. 2018). Consequently, it will be important to consider the relationship between changes in

genetic correlations and changes in natural selection, as development proceeds.

The alignment between differences in the response to selection (s_g), but not the genetic selection gradients

($\boldsymbol{\beta}_g$), and phenotypic divergence (**D**), suggests that constraints to adaptation would exist if the ARF was

left to evolve in the natural environments. This is because after one generation of selection, the mean

phenotype was expected to follow divergence towards the parental ecotypes, but selection in the absence

of genetic correlations among traits was in a direction different to phenotypic divergence. We must be

circumspect in this interpretation because estimation of $\boldsymbol{\beta}_g$ assumes we have included all traits under

selection, whereas s_g does not suffer from the same limitations (Morrissey et al. 2012; Stinchcombe et al.

2014).

In Walter et al. (2018a), we showed that genetic variance has evolved, and diverged among ecotypes,

which aligned with the direction of morphological evolution. This suggested that genetic constraints have

limited capacity to constrain adaptation during adaptive radiation, or genetic variance can evolve to

reduce genetic constraints as evolution proceeds. Given we observed divergence in ecotype **G**, but also

genetic constraints in the ARF after ecotype-specific adaptive alleles were lost, we believe the loss of

ecotype-specific adaptive alleles has re-created the constraints present during the very early stages of

adaptive divergence. Thus, adaptive radiation can occur when environment-specific alleles increase in

frequency, causing changes in the distribution of genetic variance and ameliorating genetic constraints as

adaptive divergence proceeds. Therefore, during adaptive radiation, the early stages of adaptive

divergence may be constrained (Lande and Arnold 1983; Arnold 1992; Arnold et al. 2008; Chenoweth et

al. 2010), but longer-term evolution is determined by the long-term correlated response to selection (Zeng

1988).

In this work, we identified patterns of phenotypic divergence and adaptive divergence among recently

derived ecotypes as a result of the accumulation of environment-specific alleles in response to natural

selection. We show that these alleles likely created fitness trade-offs between habitats, lead to outlier

phenotypes absent in alternative ecotypes and were likely the cause of genetic incompatibities. Through

these experiments we identify the connection between microevolutionary genetic changes and

macroevolutionary diversification in the context of an adaptive radiation.

References

Aguirre, J. D., M. W. Blows, and D. J. Marshall. 2014. The genetic covariance between life cycle stages
separated by metamorphosis. Proceedings of the Royal Society B-Biological Sciences 281.
Ali, S. I. 1969. *Senecio lautus* Complex in Australia .5. Taxonomic interpretations. Australian Journal of
Botany 17:161-176.
Anderson, J. T., C. R. Lee, C. A. Rushworth, R. I. Colautti, and T. Mitchell-Olds. 2013. Genetic trade-
offs and conditional neutrality contribute to local adaptation. Molecular Ecology 22:699-708.
Anderson, J. T., J. H. Willis, and T. Mitchell-Olds. 2011. Evolutionary genetics of plant adaptation.
Trends in Genetics 27:258-266.
Arnold, S. J. 1992. Constraints on Phenotypic Evolution. The American Naturalist 140:S85-S107.
Arnold, S. J., R. Burger, P. A. Hohenlohe, B. C. Ajie, and A. G. Jones. 2008. Understanding the
Evolution and Stability of the G-Matrix. Evolution 62:2451-2461.
Baack, E., M. C. Melo, L. H. Rieseberg, and D. Ortiz-Barrientos. 2015. The origins of reproductive
isolation in plants. New Phytologist.
Barrett, R. D. H., S. M. Rogers, and D. Schluter. 2008. Natural selection on a major armor gene in
threespine stickleback. Science 322:255-257.
Bateson, W. 1909, Heredity and Variation in Modern Lights. Oxford, UK, Cambridge University Press.
Bourret, A., M. Bélisle, F. Pelletier, and D. Garant. 2017. Evolutionary potential of morphological traits
across different life-history stages. Journal of Evolutionary Biology 30:616-626.
Bylesjo, M., V. Segura, R. Y. Soolanayakanahally, A. M. Rae, J. Trygg, P. Gustafsson, S. Jansson et al.
2008. LAMINA: a tool for rapid quantification of leaf size and shape parameters. BMC Plant
Biology 8.
Chenoweth, S. F., H. D. Rundle, and M. W. Blows. 2010. The contribution of selection and genetic
constraints to phenotypic divergence. The American Naturalist 175:186-196.
Cheverud, J. M. 1982. Phenotypic, Genetic, and Environmental Morphological Integration in the
Cranium. Evolution 36:499-516.
Corbett-Detig, R. B., J. Zhou, A. G. Clark, D. L. Hartl, and J. F. Ayroles. 2013. Genetic incompatibities
are widespread within species. Nature 504:135-137.
Coyne, J. A., and H. A. Orr. 2004, Speciation. Sunderland, MA, Sinauer Associates.
Cutter, A. D. 2012. The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities. Trends in
Ecology & Evolution 27:209-218.
Demuth, J. P., and M. J. Wade. 2005. On the theoretical and empirical framework for studying genetic
interactions within and among species. The American Naturalist 165:524-536.
Dobzhansky, T. G. 1937, Genetics and the origin of species. New York, Columbia University Press.
Doroszuk, A., M. W. Wojewodzic, G. Gort, and J. E. Kammenga. 2008. Rapid divergence of genetic
variance-covariance matrix within a natural population. The American Naturalist 171:291-304.
Eroukhmanoff, F., and E. I. Svensson. 2011. Evolution and stability of the G-matrix during the
colonization of a novel environment. Journal of Evolutionary Biology 24:1363-1373.
Fisher, R. A. 1930, The Genetical Theory of Natural Selection. Oxford, Oxford University Press.

Gavrilets, S. 2003. Perspective: models of speciation: what have we learned in 40 years? Evolution 57:2197-2215.

Goodnight, C. J. 2000. Quantitative trait loci and gene interaction: the quantitative genetics of metapopulations. Heredity (Edinb) 84:587-598.

Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. Journal of Statistical Software 33:1-22.

Kondrashov, A. S. 2003. Accumulation of Dobzhansky-Muller incompatibilities within a spatially structured population. Evolution 57:151-153.

Lande, R. 1979. Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:Body Size Allometry. Evolution 33:402-416.

Lande, R., and S. J. Arnold. 1983. The Measurement of Selection on Correlated Characters. Evolution 37:1210-1226.

Latta, R. G., K. M. Gardner, and A. D. Johansen-Morris. 2007. Hybridization, recombination, and the genetic basis of fitness variation across environments in *Avena barbata*. Genetica 129:167-177.

Martin, G., E. Chapuis, and J. Goudet. 2008. Multivariate Q_{st}–F_{st} Comparisons: A Neutrality Test for the Evolution of the G Matrix in Structured Populations. Genetics 180:2135-2149.

McFarlane, S. E., J. C. Gorrell, D. W. Coltman, M. M. Humphries, S. Boutin, and A. G. McAdam. 2014. Very low levels of direct additive genetic variance in fitness and fitness components in a red squirrel population. Ecology and Evolution 4:1729-1738.

Melo, M. C., A. Grealy, B. Brittain, G. M. Walter, and D. Ortiz-Barrientos. 2014. Strong extrinsic reproductive isolation between parapatric populations of an Australian groundsel. New Phytologist 203:323-334.

Morrissey, M. B., D. J. Parker, P. Korsten, J. M. Pemberton, L. E. Kruuk, and A. J. Wilson. 2012. The prediction of adaptive evolution: empirical application of the secondary theorem of selection and comparison to the breeder's equation. Evolution 66:2399-2410.

Muller, H. J. 1942. Isolating mechanisms, evolution and temperature. Biology Symposium 6.

Navarro, A., and N. H. Barton. 2003. Accumulating postzygotic isolation genes in parapatry: a new twist on chromosomal speciation. Evolution 57:447-459.

Nosil, P., and D. Schluter. 2011. The genes underlying the process of speciation. Trends in Ecology & Evolution 26:160-167.

Ornduff, R. 1964. Evolutionary Pathways of *Senecio lautus* Alliance in New Zealand and Australia. Evolution 18:349-360.

Pujol, B., S. Blanchet, A. Charmantier, E. Danchin, B. Facon, P. Marrot, F. Roux et al. 2018. The Missing Response to Selection in the Wild. Trends in Ecology & Evolution 33:337-346.

R Core Team. 2016.R: A language and environment for statistical computing, version v. 3.3.2.R Foundation for Statistical Computing, Vienna, Austria.

Radford, I. J., R. D. Cousens, and P. W. Michael. 2004. Morphological and genetic variation in the *Senecio pinnatifolius* complex: are variants worthy of taxonomic recognition? Australian Systematic Botany 17:29-48.

Rausher, M. D. 1992. The Measurement of Selection on Quantitative Traits: Biases Due to Environmental Covariances between Traits and Fitness. Evolution 46:616-626.

Richards, T. J., and D. Ortiz-Barrientos. 2016. Immigrant inviability produces a strong barrier to gene flow between parapatric ecotypes of *Senecio lautus*. Evolution 70:1239-1248.

Richards, T. J., G. M. Walter, K. McGuigan, and D. Ortiz-Barrientos. 2016. Divergent natural selection drives the evolution of reproductive isolation in an Australian wildflower. Evolution 70:1993-2003.

Robinson, M. R., and A. P. Beckerman. 2013. Quantifying multivariate plasticity: genetic variation in resource acquisition drives plasticity in resource allocation to components of life history. Ecology Letters 16:281-290.

Roda, F., L. Ambrose, G. M. Walter, H. L. L. Liu, A. Schaul, A. Lowe, P. B. Pelser et al. 2013a. Genomic evidence for the parallel evolution of coastal forms in the *Senecio lautus* complex. Molecular Ecology 22:2941-2952.

Roda, F., H. L. Liu, M. J. Wilkinson, G. M. Walter, M. E. James, D. M. Bernal, M. C. Melo et al. 2013b. Convergence and Divergence during the Adaptation to Similar Environments by an Australian Groundsel. Evolution 67:2515-2529.

Roda, F., G. M. Walter, R. Nipper, and D. Ortiz-Barrientos. 2017. Genomic clustering of adaptive loci during parallel evolution of an Australian wildflower. Molecular Ecology 26:3687-3699.

Sgrò, C. M., and A. A. Hoffmann. 2004. Genetic correlations, tradeoffs and environmental variation. Heredity 93:241-248.

Stinchcombe, J. R., A. K. Simonsen, and M. W. Blows. 2014. Estimating Uncertainty in Multivariate Responses to Selection. Evolution 68:1188-1196.

Styga, J. M., T. M. Houslay, A. J. Wilson, and R. L. Earley. 2018. Development of G: a test in an amphibious fish. Heredity.

Sweigart, A. L., L. Fishman, and J. H. Willis. 2006. A simple genetic incompatibility causes hybrid male sterility in *Mimulus*. Genetics 172:2465-2479.

Walsh, B., and M. W. Blows. 2009. Abundant Genetic Variation plus Strong Selection = Multivariate Genetic Constraints: A Geometric View of Adaptation. Annual Review of Ecology Evolution and Systematics 40:41-59.

Walsh, B., and M. Lynch. 2018, Evolution and selection of quantitative traits. Oxford, Oxford University Press.

Walter, G. M., J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018a. Evolution of genetic variance during adaptive radiation. The American Naturalist 191:E108–E128.

Walter, G. M., M. J. Wilkinson, J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018b. Environmentally induced development costs underlie fitness tradeoffs. Ecology 99:1391–1401.

Walter, G. M., M. J. Wilkinson, M. E. James, T. J. Richards, J. D. Aguirre, and D. Ortiz-Barrientos. 2016. Diversification across a heterogeneous landscape. Evolution 70:1979-1992.

Willett, C. S. 2006. Deleterious epistatic interactions between electron transport system protein-coding loci in the copepod *Tigriopus californicus*. Genetics 173:1465-1477.

Wright, S. 1931. Evolution in Mendelian Populations. Genetics 16:97-159.

Zeng, Z. B. 1988. Long-Term Correlated Response, Interpopulation Covariation, and Interspecific Allometry. Evolution 42:363-374.