Quantitatively Partitioning Microbial Genomic Traits among Taxonomic Ranks

across the Microbial Tree of Life

Taylor M. Royalty[1]

Andrew D. Steen[1,2,*]

[1]Department of Earth and Planetary Sciences, University of Tennessee, Knoxville

[2]Department of Microbiology, University of Tennessee, Knoxville

*asteen1@utk.edu

Abstract

Widely used microbial taxonomies, such as the NCBI taxonomy, are based on a combination of

sequence homology among conserved genes and historically accepted taxonomies, which were

developed based on observable traits such as morphology and physiology. A recently-proposed

alternative taxonomy, the Genome Taxonomy Database (GTDB), incorporates only sequence homology

of conserved genes and attempts to partition taxonomic ranks such that each rank implies the same

amount of evolutionary distance, regardless of its position on the phylogenetic tree. This provides the

first opportunity to completely separate taxonomy from traits, and therefore to quantify how taxonomic

rank corresponds to traits across the microbial tree of life. We quantified the enrichment of clusters of

orthologous gene functional categories (COG-FCs) as a proxy for traits within the lineages of 13,735

cultured and uncultured microbial lineages from a custom-curated genome database. On average, 41.4%

of the variation in COG-FC enrichment is explained by taxonomic rank, with domain, phylum, class,

order, family, and genus explaining, on average, 3.2%, 14.6%, 4.1%, 9.2%, 4.8%, and 5.5% of the

variance, respectively ($p<0.001$ for all). To our knowledge, this is the first work to quantify the variance

in metabolic potential contributed by individual taxonomic ranks. A qualitative comparison between the

COG-FC enrichments and genus-level phylogenies, generated from published concatenated protein

sequence alignments, further supports the idea that metabolic potential is taxonomically coherent at

higher taxonomic ranks. The quantitative analyses presented here constrain the integral relationship

between diversification of microbial lineages and the metabolisms which they host.

Importance

Recently there has been great progress in defining a complete taxonomy of bacteria and archaea,

which has been enabled by improvements in DNA sequencing technology and new bioinformatic

techniques. A new, algorithmically-defined microbial tree of life describes those linkages relying solely

on genetic data, which raises the question of how microbial traits relate to taxonomy. Here, we adopted

cluster of orthologous group functional categories as a scheme to describe the genomic contents of

microbes, which can be applied to any microbial lineage for which genomes are available. This simple

approach allows quantitative comparisons between microbial genomes with different gene composition

from across the microbial tree of life. Our observations demonstrate statistically significant patterns in

cluster of orthologous group functional categories at the taxonomic levels spanning from domain to

genus.

Introduction

The relationship between microbial taxonomy and function is a longstanding problem in

microbiology (1–3). Prior to the identification of the 16S rRNA gene as a taxonomic marker, microbial

phylogenetic relationships were defined by traits such as morphology, behavior, and metabolic capacity.

Cheap DNA sequencing has provided the ability to fortify those phenotype-based taxonomies with

quantitative determinations of differences between marker genes, but canonical taxonomies such as the

NCBI taxonomy continue to "reflect the current consensus in the systematic literature," which ultimately

derives from trait-based taxonomies (4). Recently, Parks et al. (5) formalized the genome taxonomy

database (GTDB), a phylogeny in which taxonomic ranks are defined by "relative evolutionary

divergence" in order to create taxonomic ranks that have uniform evolutionary meaning across the

microbial tree of life (5). This approach removes phenotype or traits entirely from taxonomic

assignment. Thus, an investigation of the relationship between traits and phylogeny has not been

possible until the recent publication of a microbial tree of life that is based solely on evolutionary

distance.

Comparing phenotypic characteristics of microorganisms across the tree of life is not currently

possible, because most organisms and lineages currently lack cultured representatives (6, 7). We

therefore used the abundance of different Clusters of Orthologous Groups (COGs) in microbial

genomes, a proxy for phenotype which is available for all microorganisms for which genomes are

available. Clusters of orthologous groups (COGs) are a classification scheme that defines protein

domains based on groups of proteins sharing high sequence homology (8). More than ~5,700 COGs

have been identified to date. COGs are placed into one of 25 metabolic functional categories (COG-

FCs), which represents a generalized metabolic function. Our analyses quantify the degree to which

taxonomic rank (genus through domain) predicts the COG-FC content of genomes, and illustrate which

lineages are relatively enriched or depleted in specific COG-FCs. These analyses constitute a step

towards being able to probabilistically predict the metabolic or functional similarity of microbes given

their taxonomic classification.

Results

The genomes analyzed in this work were compiled from a variety of different sources, including

RefSeq v92, JGI IMG/M, and Genbank, in order to include genomes created using diverse sequencing

and assembly techniques. The integration of RefSeq v92, JGI IMG/M, and Genbank databases resulted

in a total of 119,852 genomes within the custom-curated database. Raw data, GTDB taxonomy, and

associated accessions are provided in Dataset S1, which is explained in more detail in Supplement 3,

available here:

https://www.dropbox.com/sh/rnm6ount2aqkmvn/AACGgZhdrA0fSYnD0mNtIpaxa?dl=0]. Of these

genomes, we included only those that satisfied a set of criteria designed to ensure that each genus

contained enough genomes to allow statistically robust analysis (see Methods). This resulted in a set of

13,735 lineages, representing 22 bacterial phyla and 4 archaeal phyla, of which 67% have been grown in

culture (Table 1).

Most predicted open reading frames for most lineages could be assigned to a COG-FC. Across

all phyla, an average of 84.3% $\pm$ 7.8% of open reading frames were assigned to a COG-FC (Fig S1).

Genomes of the same phylum tended to group together in an initial principal component analysis (PCA)

of raw COG-FC abundance (Fig. 1A). Since this analysis was based on absolute abundance of COG-FCs

in genomes, rather than relative abundance, we hypothesized that the relationship between COG-FC

abundance and phylum was largely a consequence of genome size, which is phylogenetically conserved

(9). Consistent with this possibility, position on PC 1 correlated closely with genome size (R^2=0.88; Fig

1B). We therefore normalized each COG-FC abundance, for each genome, to a prediction of COG-FC

abundance as a function of genome size derived from a generalized additive model (GAM; Fig S2;

summary statistics in table S1). Each GAM model was statistically significant ($p < 0.001$), and all but

five COG-FCs had deviance explained (analogous to adjusted R^2) of more than 50%. We interpret

analyses of these genome size-normalized data sets as reflecting the enrichment or depletion of COG-FC

abundance, relative to that expected for a given genome size. PCA of these COG-FC enrichments

showed that species-level lineages still tended to group by phylum, even though the inter-phyla

gradients in genome size were no longer apparent (Fig 1C, D).

To quantify the degree to which taxonomic rank explains the distribution of COG-FC

enrichments among individual genomes, we performed a permutation multivariate ANOVA

(PERMANOVA) using the following taxonomic ranks: domain, phylum, class, order, family, and genus,

as well as culture-status (cultured versus uncultured lineage). The rank of species was excluded from the

analysis as every lineage was unique, and thus, species would explain 100% of the data. Every rank

significantly influenced the distribution of COG-FC enrichments ($p < 0.001$), but the fraction of variance

that each rank explained differed substantially: phylum explained the most variance (14.6%), followed

by order (9.2%), genus (5.5%), family (4.8%), and class (4.1%). Domain explained only 3.1% of

variance in COG-FC enrichment, the least of any taxonomic rank. Culture-status was a significant

correlate of COG-FC abundance ($p < 0.001$) but had virtually no explanatory power, with variance

explained <0.001%. This observation is consistent with no particular COG-FC being systematically

enriched or depleted in uncultured microbes relative to cultured microbes.

The variability in COG-FC enrichment across different phyla was explored in addition to mean

COG-FC composition for individual phyla (Fig 3). Here, the distance in PCA space between each

lineage and the centroid of all lineages in its phyla is a measure of how different the genome is, in terms

of COG-FC distribution, from the typical lineage in that phylum. Among all phyla, COG-FC distribution

of *Crenarchaeota* was the most variable, followed by *Patescibacteria* and *Cyanobacterota*. The least

variable phyla were the *Synergistota*, *Marinisomatota*, and *Fibrobacterorta*, respectively (Fig 3A). We

explored the possibility that variance in COG-FC distribution was a function of the number of lineages

in the phylum. In other words, did COG-FC content of some genomes simply seem less variable because

they had been under-sampled? A plot of the average distance of lineages from their phylum's centroid

versus the number of lineages in the phylum reveals an apparent increase in variability between 10 and

roughly 100 genomes, at which point average distance being approaching an asymptote (Fig 3B).

Comparison of Akaike Information Criterion (AIC) of a saturating model of the data (Eq. 1), fit using

nonlinear least squares, to a linear regression indicated that the saturating model described the

relationship substantially better (ΔAIC = 10.5). Coefficient *A* of the saturating model, which represents

the value of the asymptote, was estimated to be 0.75 +/- 0.15 ($p < 0.001$). Coefficient B, which

represents how quickly the function approaches the asymptote, was 0.43 +/- 0.30 (p = 0.17). Coefficient

C, an offset to handle the fact that all the log-transformed distances have negative values, was -1.63 +/-

0.14 (p < 0.001).

Finally, we quantified the average enrichment or depletion of each COG-FC in each genus These

values were then visualized on a genus-level phylogenetic tree (Fig 4) built from concatenated ribosomal

protein sequences published by Parks et al. (5). Data underlying Fig. 4 are presented in Supplemental

Data Set 2. Several notable features appear in COG-FC enrichments at the phylum level. For example,

among the four Archaeal phyla represented here, *Thermoplasmatota* appears unique, with substantial

enrichments in cell motility and depletion in every other category. In general, the bacteria appeared

more variable at the phylum level than the archaea. The clade consisting of *Bacteroidota*, *Spirochaetota*,

and *Verrucomicrobiota* were notably depleted in the less-variable COG-FCs, including energy

production and conversion, amino acid transport and metabolism, and carbohydrate transport and

metabolism, among others. Another prominent feature is the near-ubiquitous enrichment of cell motility,

secondary metabolites biosynthesis, transport, and catabolism, lipid transport and metabolism, and

intracellular trafficking, secretion, and vesicular transport COGs in Proteobacteria. A notable dichotomy

in the enrichment of RNA processing and modification within the Proteobacteria mirrors the division of

the two largest clades within the proteobacteria. Overall, enrichment data qualitatively appears

consistent with phylogenetic relationships, albeit, occurring on different taxonomic levels.

To gain a sense of "notable" COG-FCs associated with different phyla, we calculated the mean

COG-FC across all lineages in a given phyla and compared these values against the 85th and 15th

percentiles for all lineages in our custom-curated database. All COG-FCs which were significantly ($p <$

0.05; based on a 10^5-iteration bootstrap analysis) greater or less than the 85th and 15th percentiles,

respectively, are shown in Table 2. Each archaeal phylum was enriched or depleted three-to-nine COG-

FCs, whereas most bdacteria phyla were enriched or depleted in in 3-4 COG-FCs. A few exceptions

arose, such as *Fibrobacterota* was deplete in 8 COG-FCs, *Nitrospirota* A was enriched in 4 and depleted

in 5, and *Proteobacteria* was the only phylum not heavily enriched or depleted in any COG-FCs.

Enrichment data, along with associated GTDB taxonomic assignments, used for generating Fig 4 is

available in Dataset S2.

Discussion

We find that the abundance of each COG-FC in genomes scales with the genome size, so PCA

analysis of COG-FC enrichments or depletions successfully distinguished phyla only once COG-FC

abundance was normalized to the abundance that was predicted based on genome size (Fig 1C, D). The

relationship between COG-FC abundance and genome size was clearly nonlinear, however, with higher

slopes at lower genome sizes, indicating that a simple linear model would not accurately capture that

relationship (Fig S2).

The PERMANOVA (Fig 2) and analysis of diversity of genomic composition within phyla (Fig

3) showed that microbial lineages exhibit characteristic enrichments of COG-FC, and that the extent of

variation varies among taxonomic ranks. Of all the taxonomic ranks, phylum was the most powerful

predictor of COG-FC enrichment, which is consistent with observations that phylum can be informative

of microbial function (e.g., 10–12). Surprisingly, lower taxonomic ranks such as genus and family had

little explanatory power. Many studies focus on metabolic coherence of individual traits and regularly

find traits conserved on the family level (2, 3, 13). The discrepancy between previous observations and

our observation likely relates to how we characterize patterns in metabolic potential. The tradeoff of the

approach used here is that, by analyzing COG-FCs, we lose information about specific genes or

potential metabolic functions but gain the ability to apply a consistent analysis across an entire genome

and across the entire microbial tree of life. Thus, the extent that observed patterns (Fig 1) reflect

phenotypically-expressed differences among lineages is unknown. Nonetheless, the statistical robustness

of the relationship between all taxonomic ranks and COG-FC patterns suggests that evolutionary

processes (e.g., horizontal gene transfer, vertical gene transfer, duplications, deletions, etc.) control the

preponderance of different COG-FCs across lineages.

The role that individual evolutionary processes play in influencing COG-FC enrichments at a

given taxonomic rank is likely variable. For instance, horizontal gene transfer is more common among

more closely related lineages (14) and thus, likely promotes increased levels of similarity at lower

taxonomic ranks. At higher taxonomic ranks, vertical processes may be more important. The asymptote

in the mean $\log_{10}$-distance from the centroid as function of lineages in a phylum suggests that identifying

more lineages for more poorly represented lineages should expand the diversity of COG-FCs that are

found, whereas phyla that were adequately sampled (at least ~1000 lineages) exhibited comparable

variability in COG-FC distributions (Fig 3B). Since many more than ~1000 distinct lineages of each

phylum are likely to exist (15), we propose that the taxonomic rank of phylum implies a fairly consistent

degree of diversity in COG-FC distribution. To the extent that phenotype matches genotype at the level

of COG-FC distributions, therefore, we expect that typical phyla exhibit similar phenotypic diversity. A

notable exception is the phylum *Crenarchaeota*, which were far more diverse than would be expected

based on the number of lineages sampled. The *Crenarchaeota*, as defined in the GTDB, collapsed

members of several phyla that had been designated separately under previous taxonomies, including

lineages that had previously been assigned as *Crenarchaeota*, *Thaumarchaeota*, *Euryarchaeota*,

Verstraetearchaeota, *Korarchaeota*, and *Bathyarchaeota* (5). It is possible that the relationship between

marker genes used in the GTDB and the rest of the genome is unusual for this clade, compared to other

phyla, or that the GTDB classification of *Crenarchaeota* is lacking in some other way.

Although the ranks, genus and family, explained relatively little of the variance in COG-FC

distribution, blocks of consistent colors were evident in Fig 4, indicating that enrichments or depletions

of specific COG-FCs were conserved across each taxonomic rank in some parts of the phylogenetic tree.

This is explained by 'distantly' (i.e., non-sister clades) related clades occupying similar COG-FC trait-

space. The coherence in metabolic potential at higher taxonomic ranks may help explain the distribution

of microbial clades across ecological niches. Analyses of habitat associations (9, 16) and single traits

(1–3) also support this idea. Our analysis provides quantitative evidence to this idea by demonstrating

coherence in metabolic potential with broad-scale patterns in genomic data (Fig 1-4). The question

remains: how well do the observed COG-FC enrichments reflect expressed functional traits (i.e.,

phenotype) across these lineages? It is difficult to address this question systematically, but some of the

enrichments and depletions here appear consistent with known physiologies of clades. For instance,

Rickettsiales were depleted in nucleotide metabolism and transport, consistent with previously observed

lack of a metabolic pathway for purine synthesis among five example *Rickettsiales* (17). Another

example is the depletion in the COG-FCs for energy production and conversion, amino acid transport

and metabolism, and carbohydrate transport and metabolism within the *Bacteroidetes*, *Spirochaetes*, and

Chlamydiales clade. This clade is known to contain many host-dependent pathogens and symbionts (18–

20), which are often depleted in these COG-FCs (21).

The GTDB classification is the first fully algorithmic and quantitatively self-consistent microbial

taxonomy that can be applied across the tree of life (5). By standardizing the meaning of taxonomic

ranks, it creates an objective basis on which to compare microbial functionality to phylogeny. The data

presented here support the idea that clades in this taxonomy are functionally coherent, as well as the idea

that all taxonomic ranks provide information about the metabolic and ecological niches that individual

microbes occupy.

Materials and Methods

GENOME DATABASE CURATION

All bacterial and archaeal genomes from the RefSeq database v92 (22), uncultured bacterial and

archaeal (UBA) metagenome-assembled genomes (MAGs) reported in Parks et al. (5, 23), bacterial and

archaeal MAGs from Integrated Microbial Genomes and Microbiomes (IMG/M), and bacterial and

archaeal single amplified genomes (SAGs) from IMG/M were curated into a single database. All

genomic content within the curated database is referred to as "genome(s)" for simplicity. Genomes were

assigned taxonomy consistent with the Genome Taxonomy Database (GTDB) using the GTDB toolkit

(GTDB-Tk) v0.2.1 (5). The GTDB-Tk taxonomic assignments were consistent with reference package

GTDB r86. Lineages which did not receive a genus classification, due to the absence of a reference

lineage were excluded from analyses. Due to bias in the abundance of strains in specific clades (e.g., *E.*

coli), the lowest taxonomic rank considered during our analysis was species. The COG-FC enrichments

(see below) were averaged together for all strains within a given species. An exception was made for

lineages which shared a genus classification but lacked a species classification. In this scenario, each

genome was treated as an independent lineage. Lastly, only genomes belonging to genera with at least

ten unique species in the database were retained. This criterion ensured enough data to generate

meaningful statistics during our PERMANOVA. The final database is summarized in Table 1. The

genus-level phylogenetic tree was generated from concatenated protein sequence alignments published

in Parks et al. (5).

COG FUNCTIONAL CATEGORY IDENTIFICATION, ENUMERATION, AND NORMALIZATION

Genes were predicted from individual genomes and translated into protein sequences using

Prodigal v.2.6.3 (24). The resulting protein sequences were analyzed for COGs (8). COG position-

specific scoring matrices (PSSMs) were downloaded from NCBI's Conserved Domain Database (27

March 2017 definitions). COG PSSMs were BLASTed against protein sequences with the Reverse

Position Specific-BLAST (RPS-BLAST) algorithm (25). Following previously a reported protocol (25),

we used an E-value cutoff of 0.01 to assign COGs with RPS-BLAST. The retrieved COGs were

assigned to their respective COG functional categories (COG-FCs; 25 in total) and the abundance of

each functional category was tabulated using cdd2cog (26) for each genome. The abundance for

individual COG-FCs was normalized by the respective COG-FC standard deviation across all lineages.

For the COG-FCs, extracellular structures and nuclear structures, the standard deviation was 0.

Consequently, data could not be normalized, and thus, these two categories were discarded from all

analyses.

COG-FC abundances were normalized by their respective regression slopes of COG-FC

abundance for a given genome as a function of genome size. COG-FC abundances were modelled as a

function of genome size for individual categories using a generalized additive model (GAM) with a

smoothing term due to the pairwise response to genome size (Sup. Figure 1). We used the gam function

from the R package, mgcv (27). In some instances, regression fits were visibly skewed by high-leverage

data points. High-leverage data were filtered using the influence.gam function in the mgcv package.

Data in the 99.5% percentile for influence were excluded when performing regression analysis but were

included in all downstream analyses. All regressions were significant with $p<0.001$.

PRINCIPAL COMPONENT ANALYSIS (PCA)

We performed PCA on the normalized COG-FC abundances and enrichments. Prior to PCA,

assumptions of normality were achieved by performing a boxcox transformation on individual COG-FC

abundance and enrichment distributions with the boxcox function from the R Package, MASS (28). The

resulting distributions were then scaled by the respective COG-FC standard deviation calculated from all

genomes. PCA was performed using the princomp function from the R package, stats (29).

QUANTIFYING COG-FC VARIANCE EXPLAINED BY TAXONOMIC RANK

We performed permutational multivariate analysis of variance (PERMANOVA) using the adonis

function from the R package, vegan (30). The taxonomic ranks domain, phylum, class, order, family,

and genus as well as cultured-status were used as test categorical variables for quantifying variance in

COG-FC enrichments explained by the mean taxonomic rank centroids. The default, 999 permutations

test, was performed using each categorical variable. Distances were calculated between mean phyla

COG-FC enrichment centroids and the respective genomes within that phyla by performing an analysis

of multivariate homogeneity of groups dispersions with the betadisper function from the R package,

vegan (30). The distance matrix used for both the adonis and betadisper analyses was generated

calculating Euclidean distance on the normalized COG-FC enrichments.

The mean log_{10}-distance from phylum centroid for each phylum and modeled with the following

equation, which represents a hyperbola shifted on the x-axis to ensure that mean distance is zero when n

= 1:

$$log_{10}(Mean\ Distance) = \frac{A(log_{10}(n)-1)}{B+log_{10}(n)-1} + C \hspace{3cm} (1$$

where A, B, and C are fit coefficients and n is the total number of lineages in the given phylum. The

Akaike Information Criterion was calculated with the fit from eq 1 using the AIC function from the R

package, stats (29).

DATA AVAILABILITY

The genomes analyzed for the current study are available in NCBI's RefSeq database

(ftp://ftp.ncbi.nlm.nih.gov/refseq/release/). UBA MAGs used for the current study are available under

NCBI's BioProject PRJNA417962 and PRJNA348753. JGI IMG/M acquired from Chad Burdyshaw.

Associated genome accessions are available in Dataset S1.

Acknowledgments

Funding for this project was provided by a C-DEBI graduate fellowship to TR and an-kind grant

of resources from the University of Tennessee / Oak Ridge National Lab Joint Institute for

Computational Sciences (JICS) to ADS. We thank Chad Burdyshaw of JICS for help obtaining the

genomes used in this project. This is C-DEBI contribution number [to be assigned upon manuscript

acceptance].

References

1.   Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A

292   phylogenetic perspective. Science (80-) 350:aac9323.

293   2.   Martiny AC, Treseder K, Pusch G. 2013. Phylogenetic conservatism of functional traits in

294   microorganisms. ISME J 7:830–838.

295   3.   Zimmerman AE, Martiny AC, Allison SD. 2013. Microdiversity of extracellular enzyme genes

296   among sequenced prokaryotic genomes. ISME J 7:1187–1199.

297   4.   Federhen S. 2012. The NCBI Taxonomy database. Nucleic Acids Res 40:136–143.

298   5.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P.

299   2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the

300   tree of life. Nat Biotechnol 36:996–1004.

301   6.   Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel Uncultured

302   Microbial Cells Dominate Earth Microbiomes. mSystems 3:1–12.

303   7.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,

304  Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R,

305  Thomas BC, Banfield JF. 2016. A new view of the tree of life. Nat Microbiol 1:1–6.

306 8. Tatusov RL, Galperin MY, Natale DA, Koonin E V. 2000. The COG database: a tool for genome-

307  scale analysis of protein functions and evolution. Nucleic Acids Res 28:33–36.

308 9. Philippot L, Andersson SGE, Battin TJ, Prosser JI, Schimel JP, Whitman WB, Hallin S. 2010.

309  The ecological coherence of high bacterial taxonomic ranks. Nat Rev Microbiol 8:523–529.

310 10. Danczak RE, Johnston MD, Kenah C, Slattery M, Wrighton KC, Wilkins MJ. 2017. Members of

311  the Candidate Phyla Radiation are functionally differentiated by carbon- and nitrogen-cycling

312  capabilities. Microbiome 5:112.

313 11. Emerson D, Fleming EJ, McBeth JM. 2010. Iron-Oxidizing Bacteria: An Environmental and

314  Genomic Perspective. Annu Rev Microbiol 64:561–583.

315 12. Singh AH, Doerks T, Letunic I, Raes J, Bork P. 2009. Discovering functional novelty in

316  metagenomes: Examples from light-mediated processes. J Bacteriol 91:32–41.

317 13. Mendler K, Chen H, Parks DH, Lobb B, Hug LA, Doxey AC. 2019. AnnoTree: visualization and

318  exploration of a functionally annotated microbial tree of life. Nucleic Acids Res 4567:1–7.

319 14. Soucy SM, Huang J, Gogarten JP. 2015. Horizontal gene transfer: Building the web of life. Nat

320  Rev Genet 16:472–482.

321 15. Schloter M, Lebuhn M, Heulin T, Hartmann A. 2000. Ecology and evolution of bacterial

322  microdiversity. FEMS Microbiol Rev 24:647–660.

323 16. Koeppel AF, Wu M. 2012. Lineage-dependent ecological coherence in bacteria. FEMS Microbiol

324  Ecol 81:574–582.

17.   Min CK, Yang JS, Kim S, Choi MS, Kim IS, Cho NH. 2008. Genome-based construction of the

metabolic pathways of Orientia tsutsugamushi and comparative analysis within the Rickettsiales

order. Comp Funct Genomics 2008.

18.   Wolgemuth CW. 2015. Flagellar motility of the pathogenic spirochetes. Semin Cell Dev Biol

46:104–112.

19.   Dreyer M, Aeby S, Oevermann A, Greub G. 2015. Prevalence and diversity of Chlamydiales in

Swiss ruminant farms. Pathog Dis 73:1–4.

20.   Wexler HM. 2007. Bacteroides: The good, the bad, and the nitty-gritty. Clin Microbiol Rev

20:593–621.

21.   Moran NA. 2002. Microbial Minimalism: Genome Reduction in Bacterial Pathogens. Cell

108:583–586.

22.   O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse B,

Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova O, Brover V,

Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM, Goldfarb T, Gupta T, Haft D, Hatcher E,

Hlavina W, Joardar VS, Kodali VK, Li W, Maglott D, Masterson P, McGarvey KM, Murphy

MR, O'Neill K, Pujar S, Rangwala SH, Rausch D, Riddick LD, Schoch C, Shkeda A, Storz SS,

Sun H, Thibaud-Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C, Webb D, Wu W, Landrum

MJ, Kimchi A, Tatusova T, DiCuccio M, Kitts P, Murphy TD, Pruitt KD. 2016. Reference

sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and functional

annotation. Nucleic Acids Res 44:D733–D745.

23.   Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, Hugenholtz P,

Tyson GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes substantially

expands the tree of life. Nat Microbiol 2:1533–1542.

24.   Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic

gene recognition and translation initiation site identification. BMC Bioinformatics 11:119.

25.   Marchler-Bauer A, Bryant SH. 2004. CD-Search: Protein domain annotations on the fly. Nucleic

Acids Res 32:W327–W331.

26.   Leimbach A, Poehlein A, Witten A, Wellnitz O, Shpigel N, Petzl W, Zerbe H, Daniel R, Dobrindt

U. 2016. Whole-Genome Draft Sequences of Six Commensal Fecal and Six Mastitis-Associated

Escherichia coli Strains of Bovine Origin: TABLE 1. Genome Announc 4:e00753-16.

27.   Wood S. 2017. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML Smoothness

Estimation.

28.   Venables WN, Ripley BD. 2002. Modern Applied Statistics with S. 7.3.49. Springer, New York.

29.   R Core Team. 2018. R: A Language and Environmen for Statistical Computing. Vienna, Austria.

30.   Philip D. 2003. Computer program review VEGAN , a package of R functions for community

ecology. J Veg Sci 14:927–930.

Tables

Table 1. A summary of the custom-curated genome database used in this work.

Unique Domains	Unique Phyla	Unique Classes	Unique Orders	Unique Families	Unique Genera	Unique Lineages	Cultured Lineages	Uncultured Lineages
	Actinobacteriota	3	9	22	50	2286	2115	171
	Bacteroidota	3	7	19	50	1606	741	865
	Campylobacterota	1	1	6	8	270	203	67
	Cyanobacteriota	2	3	4	7	119	84	35
	Deinococcota	1	1	2	2	44	44	0
	Desulfobacterota	2	2	2	4	43	23	20
	Elusimicrobiota	1	1	1	1	22	0	22
	Fibrobacterota	1	1	1	1	34	22	12
	Firmicutes	3	10	23	48	1543	1356	187
	Firmicutes A	2	7	10	31	600	304	296
Bacteria	Firmicutes B	1	1	1	1	22	11	11
	Firmicutes C	1	2	3	4	53	32	21
	Fusobacteriota	1	1	2	2	40	40	0
	Marinisomatota	1	1	1	1	10	0	10
	Nitrospirota	2	2	2	2	30	6	24
	Nitrospirota A	1	1	1	1	14	2	12
	Patescibacteria	6	16	26	36	707	0	707
	Proteobacteria	3	25	59	163	5589	3952	1637
	Spirochaetota	3	4	4	6	153	89	64
	Synergistota	1	1	1	1	19	2	17
	Thermotogota	1	1	2	2	23	15	8
	Verrucomicrobiota	2	4	5	7	84	16	68
	Crenarchaeota	1	1	1	2	68	7	61
Archaea	Euryarchaeota	2	2	2	2	45	26	19
	Halobacterota	4	5	7	9	164	97	67
	Thermoplasmatota	1	1	2	9	147	0	147
Total	26	50	110	209	450	13,735	9187	4548

Table 2. Phylum highly enriched (>85[th] percentile) or depleted (<15[th] percentile) in COG-FCs and depletion. All reported categories are statistically significant.

Phylum	Enriched COG-FC	Depleted COG-FC	COG-FC	Table Key
Actinobacteriota	--	4,6	Cytoskeleton	1
Bacteroidota	--	11,18,22	RNA Processing and Modification	2
Campylobacterota	4,6,10,15,19	8,12	Chromatin Structure and Dynamics	3
Cyanobacteriota	19	12	Cell Motility	4
Deinococcota	--	15	Secondary Metabolites Biosynthesis, Transport, and Catabolism	5
Desulfobacterota	13	7	Intracellular Trafficking, Secretion, and Vesicular Transport	6
Elusimicrobiota	3,10,15	14,16	Lipid Transport and Metabolism	7
Fibrobacterota	--	7,11,12,13,16, 18,20,22	Carbohydrate Transport and Metabolism	8
Firmicutes	9,12,21	--	Defense Mechanism	9
Firmicutes A	9	5,7,16,20	Signal Transduction Mechanisms	10
Firmicutes B	13,17,19	--	Amino Acid Transport and Metabolism	11
Firmicutes C	19	--	Transcription	12
Fusobacteriota	--	10,21	Energy Production and Conversion	13
Marinisomatota	--	12,14	Replication, Recombination, and Repair	14
Nitrospirota	6,10,17	8	Cell Wall/Membrane/Envelope Biogenesis	15
Nitrospirota A	4,6,10,15	12,17,22,23	Inorganic Ion Transport and Metabolism	16
Patescibacteria	--	7,11,16,19	Cell Cycle Control, Cell Division, and Chromosome Partitioning	17
Proteobacteria	--	--	Function Unknown	18
Spirochaetota	4	5,16,19	Coenzyme Transport and Metabolism	19
Synergistota	3,4,11,16	--	Post-translational Modification, Protein Turnover, and Chaperone	20
Thermotogota	3,4,8,10	--	Nucleotide Transport and Metabolism	21
Verrucomicrobiota	1	10,12,14,21, 23	General Function Prediction Only	22
Crenarchaeota	3,13,11,19,7, 12,16,5,22	9,10,14,15,17	Translation Ribosomal Structure and Biogenesis	23
Euryarchaeota	2,3,13,18,22	5,7,10,14,15		
Halobacterota	3,19,22	15,16		
Thermoplasmatota	1,2,7,13	4,6,8,9,10,14, 15,16		

366
367
368

Figures

Fig 1. PCA plots of COG-FC abundance (A,B) and enrichment (C,D) of the lineages

summarized in Table 1. Individual data points are colored by genome size (A,C) and phylum (B, D).

Panels A and B were not normalized by genome size while panels C and D were normalized by genome

size. For panels A and B, PC1 explained 71% and PC2 explained 7.0% of variance. For panels C and D,

PC1 explained 21% and PC2 explained 16% of variance.

378 **Fig 2.** The average variance in COG-FC enrichment explained by different taxonomic ranks

379 (bars) and the cumulative variance explained by taxonomic ranks (lines). All variance explained by

380 taxonomic ranks was significant ($p<0.001$). The F-value for domain, phylum, class, order, family, and

381 genus, was 726.0, 128.8, 38.76, 34.4, 11.2, and 5.1, respectively.

Fig 3. Violin plots showing the distribution in $\log_{10}$-distance for lineages from their respective phyla

mean COG-FC centroid (A) and average $\log_{10}$-distance to phyla mean COG-FC centroid as a function of

number of lineages in a given phylum (B). Coefficients in panel B correspond to fit parameters from eq

1. The * symbols denote significance as defined in the text. We note three outliers: the Crenarhaeota are

characterized by unusually high diversity of COG-FCs distributions, and the Synergistota and

Fibrobacterota are characterized by an unusually low diversity of COG-FC distributions.

 Fig 4. A heat map showing the average COG-FC enrichment for all archaeal (A) and bacterial

391 (B) genera. Categories were arranged from left to right along the x axis in order of decreasing total

392 variance in enrichment across all lineages. Clades were organized along the y axis using phylogenetic

393 relatedness based on the reported concatenated protein sequence alignments in Parks et al. (1).