1 **Partitioning of microbial function among taxonomic ranks across the tree of life**
2
3
4

5 Taylor M. Royalty
6 Andrew D. Steen
7
8 Department of Earth and Planetary Sciences, University of Tennessee, Knoxville

Concatenated protein sequence alignments are a reliable tool to determine phylogenetic relationships among microbes[1] but do not necessarily imply microbial function[2]. Microbiomes influence their environments via the collective action of microbial metabolisms, so it is important to develop a quantitative understanding of the degree to which taxonomy provides information about the function of microbes. However, the overwhelming majority of microbial lineages are resistant to culturing[3], leading to difficulties characterizing their metabolisms. Here we quantify the enrichment of clusters of orthologous gene functional categories (COG-FCs)[4], a proxy for metabolic potential, within the genomes of 3207 cultured and uncultured microbe lineages from a custom-curated genome database. Results show that, on average, 36.7% of the variation in COG-FC enrichment is explained by taxonomic rank, with domain, phylum, class, order, family, and genus explaining, on average, 2.3%, 12.9%, 2.6%, 9.4%, 6.1%, and 3.5% of the variance, respectively ($p<0.001$ for all). Furthermore, taxonomic rank explained 81.5% of the variance when a only 18 of the 23 COG-FC categories were considered for analysis. To our knowledge, this is the first work which quantifies the variance in metabolic potential from individual taxonomic ranks. A qualitative comparison between the COG-FC enrichments and order-level phylogenies, generated from concatenated protein sequence alignments, further supports the idea that metabolic potential is taxonomically coherent at higher taxonomic ranks. The coherence in metabolic potential at higher taxonomic ranks provides support to hypotheses which suggest that microbial lineages may be biased to occupy specific ecological or metabolic niches[5].

High-performance sequencing and related bioinformatic techniques have led to many newly identified microbial lineages and a reorganization of the tree of life[1,3]. However, understanding of the distribution of functional traits among lineages still lags[7]. The challenge of identifying novel genes and experimentally determining the function of their products is partially to blame. Analyses measuring the conservation of functional traits across lineages often trace single genes through phylogenies[2,8]. This approach works well for carefully-chosen genes within specific lineages but cannot be applied across

the tree of life. As such, the role that phylogeny has in influencing the distribution of functional traits

across microbial lineages is at best, poorly unconstrained, and at worst, incompletely constrained.

Here we explored the distribution of metabolic potential across 3207 microbial lineages from a

custom-curated database. The databased was built with genomes from RefSeq[9] and metagenome-

assembled genomes published by Parks et al.[1,10] (Table 1; Supplementary Dataset 1). Variations in the

abundance of COG-FCs were used as a proxy for trait variation. COG-FCs have previously been used

to characterize broad-scale trends across microbial lineages, for instance, identifying last universal

common ancestor's metabolism[11], the rapid emergence of metabolic diversity during the Archaean

Explosion[12], and the construction of "functional phylogenies"[13]. In the dataset described here, the

abundance of COG-FC within a genome correlated positively with genome size (Supplementary Figure

1). The inter-relationship between COG-FC abundance, genome size, and phylogeny was also clear in

non-metric multidimensional scaling (NMDS) plots of the scaled COG-FC abundances (Figure 1a,b).

The observation that genome size is conserved at higher taxonomic ranks is not novel[5]. Our objective

was to measure how metabolic potential partitions among taxonomic ranks, so for downstream

analyses, COG-FC abundances required genome size normalization. Variations in the explanatory

power of genome size for COG-FC abundances implied that other extrinsic variables influence the

preponderance of COG-FCs, so normalizing by genome size may increase noise rather than reducing it.

To avoid this problem, we normalized each scaled COG-FC abundance, for each genome, by a

generalized additive model (GAM) prediction of annotation abundance as a function of genome size

(see methods). This generated COG-FC enrichments relative to the expected COG-FC abundance for

given a genome size. A second NMDS of COG-FC enrichments, opposed to abundances, leads to the

loss of the strong, inter-phyla gradient in genome size while conserving the spatial relationship of

phylogeny (Figure 1 c,d). The complete removal of a gradient in genome size is not expected, as

genome size depends on phylogeny[5]. Still, the removal of the inter-phyla genome size gradient

indicates that our normalization removed the first-order effect of genome size.

To quantify the degree to which taxonomic rank explains the distribution of COG-FC

enrichments among individual genomes, we performed a permutation multivariate ANOVA

(PERMANOVA) using the following taxonomic ranks: domain, phylum, class, order, family, and

genus, as well as culture-status (cultured versus uncultured). The rank of species was excluded from the

analysis as every lineage was unique, and thus, species would explain 100% of the data. Every rank

significantly ($p<0.001$) influenced the distribution of COG-FC enrichments, but surprisingly, phylum

explained the most variance (12.9%), followed by order (9.4%). In total, an organism's full taxonomy

to the rank of genus explained 36.7% of the variance in COG-FC enrichment (Figure 2). In contrast,

culture-status failed to reject the null hypothesis ($p\leq0.05$), consistent with "unculturedness" not being

associated with enrichment of COG-FCs. The underrepresentation of uncultured lineages may have

contributed to this observation (3089 cultured lineages versus 128 uncultured lineages after data

filtering and averaging COG-FC content of strains of the same species; see methods). In order to

determine whether the explanatory power of taxonomic rank was biased by COG-FCs with low

genome size dependency, we performed a second PERMANOVA excluding five COG-FCs for which

the GAM explained <50% of COG-FC abundance. According to this analysis, a microbes's full

taxonomy explains ~81.5% variance based on 18 COG-FC categories. The relative importance of

taxonomic ranks did not change in this analysis, and all the additional explanatory power was

partitioned to the ranks of phylum, order, family, and genus. This analysis indicates that domain and

class offer essentially no information about an organism's metabolic potential while a full taxonomy

can predict the relative abundance of COG-FCs within a genome relatively accurately.

Of all the taxonomic ranks, phylum was the most powerful predictor of COG-FC abundance,

consistent with the observation that phylum is typically informative of microbial function. We therefore

quantified the average enrichment for all COG-FCs across all taxonomic orders to visualize this effect.

These enrichments were then qualitatively compared to an order-level phylogenetic tree (Figure 3) built

from concatenated ribosomal protein sequences published by Parks et al.[1]. Numerous coherent features

appear in COG-FC enrichments. For example, a prominent feature for the clade consisting of

Bacteroidetes, Spirochaetes, and Verrucomicrobia is a paucity of the less variable COG-FCs, including

energy production and conversion, amino acid transport and metabolism, and carbohydrate transport

and metabolism, among others. Another prominent feature is the near-ubiquitous enrichment of cell

motility, secondary metabolites biosynthesis, transport, and catabolism, lipid transport and metabolism,

and intracellular trafficking, secretion, and vesicular transport COGs in Proteobacteria. Interestingly, a

notable dichotomy in the enrichment of RNA processing and modification mirrors the division of the

two largest clades within the proteobacteria. Not surprisingly, the three Archaea orders have similar

COG-FC enrichments. Overall, enrichment data qualitatively appears consistent with phylogenetic

relationships. The coherence in metabolic potential at higher taxonomic ranks may help explain the

distribution of microbial clades across ecological niches. Analyses of habitat associations[5,15] and single

traits[2,7,8] also support this idea. Our analysis provides concrete evidence to this idea by demonstrating

coherence in metabolic potential with broad-scale patterns in genomic data (Figure 1-3).

The question remains: how well do the observed COG-FC enrichments (Figure 3) reflect

expressed functional traits (i.e., phenotype) across these lineages? Broad assertions are difficult to

validate based on the literature due to the recent development of the Genome Taxonomy Database

(GTDB) taxonomic classification[1] (adopted here) and the sheer magnitude of the data in Figure 3.

Nonetheless, we investigated some anomalies in our data. For instance, Rickettsiales were depleted in

nucleotide metabolism and transport, consistent with previously observed lack of a metabolic pathway

for purine synthesis among five example Rickettsiales[16]. Another example is the depletion in the COG-

FCs for energy production and conversion, amino acid transport and metabolism, and carbohydrate

transport and metabolism within the Bacteroidetes, Spirochaetes, and Chlamydiales clade. This clade is

known to contain many host-dependent pathogens and symbionts[17–19], which are often deplete in these

COG-FCs[20].

Lastly, the adoption of a new taxonomic classification system presents an opportunity for

recharacterizing the metabolic tendencies of taxonomic groups. The qualitative and quantitative data

presented here supports coherence in metabolic potential across higher taxonomic orders using the

GTDB taxonomic classification while also implicating a possible bias in niche occupation by certain

clades.

Methods Summary:

COG Functional Category Identification, Enumeration, and Normalization

Genes were predicted from individual genomes and translated into protein sequences using Prodigal

v.2.6.3[21]. The resulting protein sequences were analyzed for COGs[4]. COG position-specific scoring

matrices (PSSMs) were downloaded from NCBI's Conserved Domain Database (27 March 2017

definitions). COG PSSMs were BLASTed against protein sequences with the Reverse Position

Specific-BLAST (RPS-BLAST) algorithm[22]. Following previously a reported protocol[23], we used an E-

value cutoff of 0.01 to assign COGs with RPS-BLAST. The retrieved COGs were assigned to their

respective COG functional categories (COG-FCs; 25 in total) and the abundance of each functional

category was tabulated using cdd2cog[24] for each genome. The abundance for individual COG-FCs was

normalized by the respective COG-FC standard deviation across all lineages. For the COG-FCs,

extracellular structures and nuclear structures, the standard deviation was 0. Consequently, data could

not be normalized, and thus, these two categories were discarded from all analyses.

COG-FC abundances were normalized by their respective regression slopes of COG-FC

abundance for a given genome as a function of genome size. COG-FC abundances were modelled as a

function of genome size for individual categories using a generalized additive model (GAM) with a

smoothing term due to the pairwise response to genome size (Sup. Figure 1). We used the gam function

from the R package, mgcv[25]. All regressions were significant with $p<0.001$.

COG-FC Enrichments Significance

We tested the COG-FC enrichment significance by comparing the 95% confidence interval for a given

COG-FC enrichment of a given taxonomic order against the 95% confidence interval for the same

COG-FC enrichment across all lineages. The 95% confidence intervals were generated by a 10^5

iteration bootstrap analysis. If 95% confidence intervals overlapped for a given COG-FC, then the

COG-FC enrichment was not considered significant.

Non-metric Multidimensional Scaling (NMDS)

We performed NMDS on the normalized COG-FC abundances and enrichments. NMDS was

performed using the metaMDS function from the R package, vegan[26]. Euclidean distance was selected

as the distance metric for NMDS. Three dimensions were considered for NMDS analysis (only NMDS

1 and NMDS 2 shown) as to reduce stress to ~0.1.

Permutational Multivariate Analysis of Variance (PERMANOVA)

We performed PERMANOVA using the adonis function from the R package, vegan[26]. The taxonomic

ranks domain, phylum, class, order, family, and genus as well as cultured-status were used as test

categorial variables for quantifying variance based on a default, 999, permutation test for each variable.

A distance matrix was generated calculating Euclidean distance on the normalized COG-FC

enrichments.

Genome Database Curation

All bacterial and archaeal genomes from the RefSeq database v86[9] and uncultured bacterial and

archaeal (UBA) metagenome-assembled genomes (MAGs) reported in Parks et al.[1,10] were curated into

a single database. UBA MAGs were filtered to only include genomes with >90% completion. All

genomic content within the curated database is referred to as "genomes." Genomes were assigned

taxonomy consistent with the Genome Taxonomy Database[1]. Due to bias in the abundance of strains in

specific clades, the lowest taxonomic rank considered during our analysis was species. The COG-FC

enrichments were averaged together for all strains within a given species. We filtered all genera with

<10 unique species. This criterion ensured enough data to generate meaningful statistics during our

PERMANOVA. The final database is summarized in Table 1. The order-level phylogenetic tree was

generated from concatenated protein sequence alignments published in Parks et al.[1] and graphically

generated with Interactive Tree of Life v2[27].

References

1. Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. *Nat. Biotechnol.* **36,** 996–1004 (2018).
2. Zimmerman, A. E., Martiny, A. C. & Allison, S. D. Microdiversity of extracellular enzyme genes
among sequenced prokaryotic genomes. *ISME J.* **7,** 1187–1199 (2013).
3. Hug, L. A. *et al.* A new view of the tree of life. *Nat. Microbiol.* **1,** 16048 (2016).
4. Tatusov, R. L., Galperin, M. Y., Natale, D. A. & Koonin, E. V. The COG database: a tool for
genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* **28,** 33–36 (2000).
5. Philippot, L. *et al.* The ecological coherence of high bacterial taxonomic ranks. *Nat. Rev.*
Microbiol. **8,** 523–529 (2010).
6. Quince, C., Walker, A. W., Simpson, J. T., Loman, N. J. & Segata, N. Shotgun metagenomics,
from sampling to analysis. *Nat. Biotechnol.* **35,** 833–844 (2017).
7. Martiny, J. B. H., Jones, S. E., Lennon, J. T. & Martiny, A. C. Microbiomes in light of traits: A
phylogenetic perspective. *Science (80-.).* **350,** aac9323 (2015).
8. Martiny, A. C., Treseder, K. & Pusch, G. Phylogenetic conservatism of functional traits in
microorganisms. *ISME J.* **7,** 830–838 (2013).
9. O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: Current status, taxonomic
expansion, and functional annotation. *Nucleic Acids Res.* **44,** D733–D745 (2016).
10. Parks, D. H. *et al.* Recovery of nearly 8,000 metagenome-assembled genomes substantially
expands the tree of life. *Nat. Microbiol.* **2,** 1533–1542 (2017).
11. Weiss, M. C. *et al.* The physiology and habitat of the last universal common ancestor. *Nat.*
Microbiol. **1,** 1–8 (2016).
12. David, L. A. & Alm, E. J. Rapid evolutionary innovation during an Archaean genetic expansion.
Nature **469,** 93–96 (2011).
13. Segata, N. & Huttenhower, C. Toward an efficient method of identifying core genes for
evolutionary and functional microbial phylogenies. *PLoS One* **6,** e24704 (2011).
14. Morris, J. J., Lenski, R. E. & Zinserc, E. R. The Black Queen Hypothesis: Evolution of
Dependencies through Adaptive Gene Loss. *MBio* **3,** 1–7 (2012).
15. Koeppel, A. F. & Wu, M. Lineage-dependent ecological coherence in bacteria. *FEMS Microbiol.*
Ecol. **81,** 574–582 (2012).
16. Min, C. K. *et al.* Genome-based construction of the metabolic pathways of Orientia
tsutsugamushi and comparative analysis within the Rickettsiales order. *Comp. Funct. Genomics*
2008, (2008).
17. Wolgemuth, C. W. Flagellar motility of the pathogenic spirochetes. *Semin Cell Dev Biol.* **46,**
104–112 (2015).
18. Dreyer, M., Aeby, S., Oevermann, A. & Greub, G. Prevalence and diversity of Chlamydiales in
Swiss ruminant farms. *Pathog. Dis.* **73,** 1–4 (2015).
19. Wexler, H. M. Bacteroides: The good, the bad, and the nitty-gritty. *Clin. Microbiol. Rev.* **20,**
593–621 (2007).
20. Moran, N. A. Microbial Minimalism: Genome Reduction in Bacterial Pathogens. *Cell* **108,** 583–
586 (2002).
21. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11,** 119 (2010).
22. Marchler-Bauer, A. & Bryant, S. H. CD-Search: Protein domain annotations on the fly. *Nucleic*
Acids Res. **32,** W327–W331 (2004).
23. Mavromatis, K. *et al.* The DOE-JGI Standard Operating Procedure for the Annotations of
Microbial Genomes. *SIGS* **1,** 63–67 (2009).
24. Leimbach, A. *et al.* Whole-Genome Draft Sequences of Six Commensal Fecal and Six Mastitis-

209   Associated *Escherichia coli* Strains of Bovine Origin: TABLE 1. *Genome Announc.* **4,** e00753-
210   16 (2016).
211 25. Wood, S. mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML Smoothness
212   Estimation. (2017).
213 26. Philip, D. Computer program review VEGAN , a package of R functions for community
214   ecology. *J. Veg. Sci.* **14,** 927–930 (2003).
215 27. Letunic, I. & Bork, P. Interactive Tree of Life v2: Online annotation and display of phylogenetic
216   trees made easy. *Nucleic Acids Res.* **39,** W475–W478 (2011).
217

Tables

220 **Table 1.** A summary of the custom-curated genome database used in this work.

Unique Domains	Unique Phyla	Unique Classes	Unique Orders	Unique Families	Unique Genera	Unique Species	Cultured Lineages	Uncultured Lineages
	Actinobacteria	1	5	11	19	889	880	9
	Bacteroidetes	1	5	9	13	338	306	32
	Deinococcota	1	1	2	2	35	35	0
	Epsilonbacterota	1	1	2	2	26	25	1
	Firmicutes	2	5	13	25	492	486	6
Bacteria	Firmicutes A	2	4	4	4	58	56	2
	Firmicutes B	1	1	1	1	11	9	2
	Fusobacteria	1	1	1	1	11	11	0
	Proteobacteria	2	12	25	55	1214	1147	67
	Spirochaetes	2	3	3	3	39	31	8
	Verrucomicrobia	1	1	1	1	10	10	0
Archaea	Euryarchaeota	1	1	1	1	20	20	0
	Halobacterota	2	2	3	5	74	73	1
Total	13	18	42	76	132	3207	3089	128

222   **<u>Figures</u>**

Figure 1. NMDS plots of COG-FC abundance (a,b) and enrichment (c,d) of the lineages summarized

in Table 1. Individual data points are colored by genome size (a,c) and phylum (b, d). Panels a,b were

not normalized by genome size while panels c,d were normalized by genome size. Note the difference

in scaling for the axes.

Figure 2. The average variance in COG-FC enrichment explained by different taxonomic ranks (bars)

and the cumulative variance explained by taxonomic ranks (lines). Blue corresponds to PERMANOVA

performed on all COG-FCs while grey corresponds to PERMANOVA performed on all COG-FCs

excluding: cytoskeleton, RNA processing and modification, chromatin structure and dynamics, cell

motility, and intracellular trafficking, secretion, and vesicular transport (see text for details). All

variance explained by taxonomic ranks was significant ($p<0.001$). The F-value (blue) for domain,

phylum, class, order, family, and genus, was 75.1, 47.7, 40.7, 28.8, 20.9, and 13.7, respectively. The F-

value (grey) for domain, phylum, class, order, family, and genus, was 61.7, 130.6, 102.44, 109.3, 94.5,

and 103.4, respectively.

238
239 **Figure 3.** A heat map showing the average COG-FC enrichment for all orders in Table 1. Categories

240 were arranged along the x axis based on the total variance in enrichment across all lineages. Clades

241 were organized along the y axis using phylogenetic relatedness based on the reported concatenated

242 protein sequence alignments in Parks et al.[1]. Minus signs correspond to enrichments not consider

243 significantly different than 1 (see methods).

Genome Size (Mbp)

0.5   1.2   2.7   6.4   15

a — Genome Size Unnormalized
b — Taxonomy Unnormalized
c — Genome Size Normalized
d — Taxonomy Normalized

NMDS 2 / NMDS 1

Phylum

- Actinobacteria
- Bacteroidetes
- Proteobacteria
- Deinococcota
- Epsilonbacterota
- Spirochaetes
- Euryarchaeota
- Firmicutes
- Verrucomicrobia
- Firmicutes A
- Firmicutes B
- Fusobacteria
- Halobacterota