1

Multiple phase-variable mechanisms, including capsular polysaccharides, modify

bacteriophage susceptibility in *Bacteroides thetaiotaomicron*

*Nathan T. Porter[1], *Andrew J. Hryckowian[2], Bryan D. Merrill[2], Jackson O. Gardner[2], Shaleni

Singh[1], Justin L. Sonnenburg[2], and Eric C. Martens[1]

[1]*Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI 48109,*

USA

[2]*Department of Microbiology and Immunology, Stanford University School of Medicine,*

Stanford, CA 94305

*These authors contributed equally to this work

Correspondence to: emartens@umich.edu

Abstract

A variety of cell surface structures, including capsular polysaccharides (CPS), dictate the

interactions between bacteria and elements of their environment including their viruses

(bacteriophages). Members of the prominent human gut Bacteroidetes characteristically produce

several phase-variable CPS, which have been demonstrated as key determinants in interacting

with the host immune system. However, the contributions of Bacteroidetes CPS to bacteriophage

interactions are unknown. We used engineered strains of the human symbiont *Bacteroides*

thetaiotaomicron, which differ only in the CPS they express, to isolate bacteriophages from two

locations in the United States. Testing each of 71 bacteriophages against a panel of strains that

express wild-type phase-variable CPS, one of eight different single CPS, or no CPS at all,

revealed that each infects only a subset of strains. Deletion of infection-permissive CPS from *B.*

thetaiotaomicron was sufficient to abolish infection for several individual bacteriophages.

Likewise, infection of wild-type *B. thetaiotaomicron* with one bacteriophage from this collection

selected for a cell population expressing non-permissive CPS. Surprisingly, acapsular *B.*

thetaiotaomicron also escapes complete killing by this bacteriophage, but surviving bacteria

increased expression of a family of 9 phase-variable lipoproteins. When constitutively expressed,

one of these lipoproteins enhances resistance to this bacteriophage. Our results reveal distinct

roles for *Bacteroides* CPS in mediating bacteriophage susceptibility. Beneath this vanguard

protective layer, additional mechanisms exist to protect these bacteria from being eradicated by

bacteriophage. Given the diversity of CPS and other phase-variable surface structures encoded

by gut-dwelling Bacteroidetes, our results provide important insight into the adaptations that

allow these bacteria to persist despite bacteriophage predation and hold important implications

for using bacteriophages therapeutically to target gut symbionts.

Introduction

The community of cellular microorganisms in the human intestinal tract is dominated by

a diverse population of bacteria, with hundreds of different species and even more strains

typically present within one person[1]. In the face of frequent perturbations from a variety of

environmental pressures, including diet changes, antibiotics and host immunity, the relative

abundances of different bacteria may vary greatly within an individual over short time periods.

However, the microbiota members within a host generally remain stable over long time

periods[2,3], suggesting that individual bacteria have evolved strategies that enable them to be

resilient in the face of challenges. One potential adaptive mechanism that has clearly been

diversified in gut resident Bacteroidetes is the ability of individual strains to alternately produce

several different capsular polysaccharides (CPS), which are extracellular matrix components

encoded by *cps* biosynthetic loci. Many *cps* loci in the Bacteroidetes are under the control of

phase variable promoters, allowing for substantial phenotypic heterogeneity within an isogenic

population. Furthermore, there is much broader diversification of *cps* loci among different strains

within a species[4-7]. While a few studies have shown that these CPS take part in evasion or

modulation of host immunity[4,8-11], the sheer diversity and number of CPS synthesis loci in the

Bacteroidetes suggests that they could also fill other fundamental roles[4,6,12,13].

Bacterial viruses or bacteriophages (herein, phages), like the bacteria on which they prey,

vary greatly across individual gut microbiomes and are even responsive to host dietary changes

or disease states[14-17]. Compared to its bacterial counterpart, far less is understood about the

human gut virome, especially the mechanisms governing phage-bacteria interactions.

Specifically, while phages that target several species of *Bacteroides* (the prominent genus of the

Bacteroidetes phylum in humans) have been shown to exhibit species- or strain-specificity[18-21],

little is known about the molecular interactions that drive bacterial susceptibility[22] or the

mechanisms by which these bacteria persist despite an abundance of phages in the

gastrointestinal tract.

68   Given the observations that *Bacteroides* CPS are extremely variable, even within

members of a single species[4,12], and employ complex regulatory mechanisms that diversify

expression in members of a population[4,23], CPS are ideal candidates for modulating *Bacteroides*-

phage interactions. Thus, we sought to test the hypothesis that CPS are direct positive or negative

mediators of *Bacteroides*-phage interactions. To accomplish this, we employed a panel of

engineered strains of the model symbiont *Bacteroides thetaiotaomicron* that each constitutively

expresses a different single CPS or none at all. While our results clearly support the conclusion

that specific CPS may either block or be required for phage infection, they also reveal that *B.*

thetaiotaomicron possesses additional phage-evasion strategies that function in the absence of

CPS. For one phage, CPS-independent survival involves altered expression of nutrient receptors

and increased expression of phase-variable surface lipoproteins by the surviving bacteria. One

hypothesis is that these latter functions also encode active resistance mechanisms. This idea is

supported by locking on expression of one of these lipoproteins, which we show to be an

additional determinant of phage tropism, conferring increased resistance to this phage. Our

results provide a mechanistic glimpse into the intricacy of bacterial-phage interactions that exist

in the human gut and provide a foundation for future work aimed at leveraging these interactions

as a facet of targeted manipulation of the gut microbiome.

Results

Individual bacteriophages target *B. thetaiotaomicron* in a CPS-dependent fashion

To test the hypothesis that variable *Bacteroides* surface CPS mediate interactions with

phages, we isolated phages that infect *B. thetaiotaomicron* VPI-5482 (ATCC 29148). To

maximize our chances of collecting phages that could differ in their interactions with CPS, we

used the wild-type strain of *B. thetaiotaomicron* that expresses 8 different, phase-variable CPS,

each encoded by a different multi-gene *cps* locus; an isogenic panel of 8 single CPS-expressing

strains (designated "*cps*1" through "*cps*8")[25]; and an acapsular strain in which all *cps* genes from

all 8 loci are deleted[24] as independent hosts for phage isolation. Primary sewage effluent from

two cities within the United States (Ann Arbor, Michigan and San Jose, California; separated by

approximately 3,300 kilometers) was used as the source for these phages (for further details on

phage isolation, see *Methods* and **Table S1**). All phages were plaque purified at least 3 times and

high titer lysates were generated for each of the 71 phages. Plaque morphologies varied greatly

among the strains, ranging in size from <1 mm to 3 mm or greater and in opacity from very

turbid to clear (**Figures 1, S1**).

To determine if phages isolated on each individual host strain are restricted in their ability

to infect other strains, we systematically tested each phage against each of the 10 *B.*

thetaiotaomicron host strains that varied only in the CPS they are capable of expressing (n=3).

Hierarchical clustering of the host infection profiles revealed a cladogram with 3 main branches

that each encompasses phages from both collection sites, although substantial variation in host

tropism exists for phages within each branch (**Figure 1**). Furthermore, individual phages within

each branch displayed a range of plaque morphologies (**Figures 1, S1**), suggesting additional

diversity in the collection that is not captured by this assay. Finally, host range assays were

robust when performed by different experimenters at different research sites (**Figure S2**).

Phages in Branch 1 generally exhibited robust infection of the acapsular strain, although

3 of these phages did not form plaques on this host. Furthermore, phages in Branch 1 generally

exhibited robust infection on strains expressing CPS7 or CPS8 alone, although a separate subset

of 3 phages did not form plaques on the CPS8 expressing strain. Some Branch 1 phages also

displayed less efficient infection of other strains with the exception of *cps*4, which was not

infected by any phages in this group. Interestingly, ARB154 exclusively infected *cps*8, an

uncommon CPS among *B. thetaiotaomicron* strains that appears to be contained in a mobile

element[4]. Phages in Branch 2 generally exhibited robust infection of all strains except *cps*2, *cps*3

and *cps*4. However, subsets of this group were unable to infect *cps*1 or *cps*6. Finally, Branch 3

tended to exhibit strong infection of wild-type, *cps*1, *cps*2, and *cps*3, with some variations. Some

Branch 3 phages also exhibited the ability to infect the *cps*7 and acapsular strains but were the

only branch that poorly infected *cps*8. A subset of phages on Branches 1 and 3 failed to infect the

acapsular strain, suggesting that they may require the presence of CPS for infection. Taken

together, the observed variations in phage infection of *B. thetaiotaomicron* strains expressing

different CPS, or none at all, provide initial support for our hypothesis that these surface

structures are a key mediator of *B. thetaiotaomicron*-phage interactions.

Elimination of specific CPS subsets alters bacterial susceptibility to phages

One explanation for the differences in host infectivity described above is that there are

distinct, CPS-dependent mechanisms of phage adsorption. For example, several phages robustly

infect the acapsular strain, indicating that a capsule-independent cell surface receptor mediates

infection for these examples. However, each of these phages also infect subsets of the single

CPS-expressing strains, suggesting that some "non-permissive" CPS could block access to cell

surface receptors, while other "permissive" CPS fail to do so. For phages that do not efficiently

infect the acapsular strain, one or more CPS may serve as a direct phage receptor(s) or as a

required co-receptor.

To further define the roles of specific CPS during phage infection, we investigated a

subset of 6 phages (ARB72, ARB78, ARB82, ARB101, ARB105, and ARB25). All 6 of these

phages can infect the wild-type *B. thetaiotaomicron* strain that variably expresses its 8 different

CPS and 5 of them infect the acapsular strain poorly or not at all (**Figure 1**). We tested our

hypothesis that some CPS are required as receptors or co-receptors by deleting only the subsets

of CPS biosynthetic genes encoding permissive capsules based on our prior experiments with

single CPS-expressing strains. For ARB72, which most robustly infects the CPS1- and CPS3-

expressing strains, simultaneous elimination of both of these capsules from wild-type *B.*

thetaiotaomicron reduced infection below the limit of detection (**Figure 2A**). Likewise,

elimination of the most permissive CPS for four other phages (ARB78, ARB82, ARB101 and

ARB105) significantly reduced *B. thetaiotaomicron* infection by these phages, in some cases in

the presence of permissive CPS (**Figure 2B-E**).

For ARB25, which infected 7 of the 10 strains tested in our initial plaque assays (**Figure**

1), some single and compounded *cps* gene deletions significantly reduced infection rates or

reduced them below the limit of detection. The most noteworthy of these was deletion of the

*cps*4 (initially determined in **Figure 1** to be non-permissive for ARB25) in combination with

deleting the permissive *cps*1, which completely eliminated detectable infection (**Figure 2F**).

While individual deletion of four other permissive CPS (*cps*1,6,7,8) led to partially reduced

infection, so did single deletions of two CPS initially determined to be non-permissive (*cps*3 and

*cps*4), suggesting the possibility of more complicated regulatory interactions, which are known

to occur with *Bacteroides* CPS[23,26]. Interestingly, strains lacking either *cps4* or *cps1/cps4*

together compensated by significantly increased relative expression of the non-permissive *cps2*

locus, which could contribute to ARB25 resistance (**Figure 2G**).

A strain expressing only two of the non-permissive CPS (CPS2 and CPS3) could not be

detectably infected by ARB25 (**Figure 2F**, "2,3 only" condition). However, a strain expressing

CPS2,3,4 regained some susceptibility (**Figure 2F**, "2,3,4 only" condition), indicating that CPS4

is capable of mediating some infection by this phage (addressed further below). In contrast to

sole expression of *cps2* and *cps3* promoting resistance to ARB25, deletion of the *cps2* and *cps3*

loci led to dominant expression of *cps1* and *cps4* genes, which increased infection efficiency and

led to the production of clearer plaques (**Figure 2F,H**). Additional support for the idea that loss

of *cps4* alone modifies ARB25 susceptibility comes from plaque morphologies arising from

infection of the Δ*cps4* strain, which produced smaller and more turbid plaques, demonstrating

that when infection does occur it is less productive (**Figure 2H**). Thus, CPS4-expressing cells are

in some cases susceptible to ARB25, and loss of the genes encoding this capsule result in

increased expression of CPS2, perhaps through alleviation of the UpxZ transcription termination

mechanism, as described for capsule regulation in *B. fragilis*[23]. Because this programmed shift in

CPS expression may increase resistance to phage, it is possible that phase-variation can equip

Bacteroides populations to survive phage predation through selection of individual cell sub-

populations that are expressing non-permissive CPS.

Finally, we used ARB25 to test if purified, exogenous CPS could modify phage infection

as has been shown for other phage-CPS interactions[27]. Given that strains expressing CPS1 and

CPS2 show different susceptibility to infection by ARB25, we tested purified preparations of

both CPS types, expecting that exogenous CPS2 might inhibit ARB25 infection of the acapsular

strain if it is capable of directly interfering with recognition of a surface receptor. Arguing

against this hypothesis, pre-incubation with purified preparations of CPS2 or CPS1 (a permissive

capsule control) both resulted in no significant difference in plaquing efficiency on the acapsular

strain (**Figure S3A**), suggesting that exogenous CPS cannot block infection by phage ARB25.

***B. thetaiotaomicron* acquires transient resistance to phage infection**

Interestingly, we observed that liquid cultures of the various *B. thetaiotaomicron* strains

infected with ARB25 did not show evidence of complete lysis after 36 hours of growth, as

determined by optical density at 600 nm (OD_{600}) (**Figure 3**). Previous reports demonstrated that

B. fragilis[20] and *B. intestinalis*[21] exhibited transient resistance to phage infection that could be

"reset" through removal of the phage from the culture, although the underlying mechanism of

this transient resistance was not determined. Based on these observations, we questioned whether

similar transient resistance occurs with *B. thetaiotaomicron* and whether this resistance could be

dependent on CPS expression. Growth curves of each of the CPS-expressing strains inoculated

with active or heat-killed ARB25 confirmed our initial host range assays, except that cultures

containing the CPS4-expressing strain were sensitive to killing by this phage in liquid culture,

while cultures of the CPS6-expressing strain had no significant decrease in OD_{600} (**Figure 3**). In

these experiments, most strains deemed to be susceptible via plaque assay (**Figure 1**) exhibited

an initial lag in growth or a drop in OD_{600}. As expected, the *cps*2 and *cps*3 strains remained

resistant to ARB25 infection for the duration of the experiments. However, after initial growth

inhibition, the susceptible strains displayed either growth stagnation without complete loss of

culture density (*cps*4, *cps*5, *cps*7, *cps*8) or resumption of growth (wild-type, acapsular, *cps*1) to

near uninfected levels, the latter suggesting outgrowth of a resistant subpopulation of bacteria.

Culture supernatants taken from ARB25 post-infected, wild-type *B. thetaiotaomicron* still

contained high phage titers when exposed to naïve bacteria that had not been exposed to phage,

excluding the possibility that the phages were inactivated (**Figure S3B**).

We next determined whether strains that had survived or proliferated after exposure to

phage retained resistance after removal of phage. In order to isolate phage-free bacterial clones,

we isolated individual colonies by sequentially streaking each twice from a subset of the cultures

that gained resistance to ARB25 (WT, acapsular, *cps*1, and *cps*4) as well as the inherently

ARB25-resistant *cps*2 strain. The majority of clones isolated using this process were free from

detectable phage (see *Methods*). We then re-infected each clone with live ARB25 and monitored

susceptibility by delayed growth or drop in the culture density as compared to infection with

heat-killed phage. As expected, the *cps*2 strain remained resistant. On the other hand, the

majority of clones (42/61 total, ~69%) of the other four strains regained susceptibility (**Table 1**),

suggesting that resistance to this phage is not caused by a permanent genetic alteration in most

cases.

Phage resistant wild-type *B. thetaiotaomicron* populations exhibit altered *cps* locus

expression

Given that CPS type is correlated with resistance to phage infection (e.g., ARB25 fails to

infect strains expressing CPS2 or CPS3 under all conditions tested), we hypothesized that wild-

type *B. thetaiotaomicron* cells inherently expressing resistant capsules would be positively

selected in the presence of phage. To test this, we infected wild-type *B. thetaiotaomicron* with

ARB25 and monitored bacterial growth. For cells treated with a high MOI (MOI ≈ 1), culture

turbidity increased very slightly, declined before 3 hours after infection, and finally increased

again to ultimately achieve a high growth level as previously observed (**Figure 4A**).

Interestingly, bacterial cultures originating from different single colonies displayed variable

growth kinetics and possibly resistance frequency to ARB25, with the growth of one clone

barely delayed by treatment with live ARB25. Next, we measured if infection with ARB25

resulted in a change in CPS expression by the phage-resistant *B. thetaiotaomicron* population. In

support of our hypothesis, *B. thetaiotaomicron* exposed to heat-killed phage predominantly

expressed genes encoding CPS3 and CPS4, which we typically observe in *in vitro* culture.

Treatment with live ARB25 resulted in a dramatic loss of *cps*1 and *cps*4 expression (capsules for

which combined loss eliminates ARB25 infection, **Figure 2F**), with a concomitant increase of

expression of the non-permissive *cps*3 (**Figure 4B**). Similar growth and expression phenotypes

occurred in cultures treated with a low MOI (MOI $\approx 10^{-4}$), albeit with higher culture turbidity

before a decline and subsequent resumption of growth (**Figure S4**). Dirichlet regression (see

Methods) supported significant *cps* expression changes for *cps1*, *cps3*, and *cps4* in response to

ARB25 (p < 0.01 for experiments with both low and high MOI). Notably, the most resistant of

the three bacterial clones (as evidenced by faster outgrowth post-infection) in each of the two

experiments (low and high MOI) exhibited similar *cps* expression to the other clones after

treatment with live phage, but expressed lower levels of permissive *cps*1 and *cps4* and higher

levels of non-permissive *cps*3 in heat-killed phage treatment groups (**Figure S5**). This alteration

in expression of permissive and non-permissive CPS may contribute to the ability of these clones

to resume growth more rapidly after phage challenge because it was already skewed towards

expression of non-permissive CPS.

Multiple layers of phase-variable resistance functions equip *B. thetaiotaomicron* to survive

phage predation

The results described above support a model in which some individual cells within a *B.*

thetaiotaomicron population are pre-adapted to resist eradication by a single phage like ARB25

through expression of different CPS. Complex phase-variation mechanisms have already been

described in controlling CPS expression in *B. thetaiotaomicron* and *B. fragilis*[4,23,28] and we

demonstrate the ability of CPS to dictate interactions between phages and *B. thetaiotaomicron*

(**Figures 1, 2, 3**). However, in acapsular *B. thetaiotaomicron* infected with ARB25, we observed

significant growth after initial lysis that frequently regained susceptibility when isolated from

and subsequently re-challenged with ARB25 (**Figure 3, Table 1**), suggesting the emergence of a

transiently phage-resistant subpopulation in the absence of CPS. To determine if additional

phage resistance mechanisms are involved, we performed whole genome transcriptional profiling

by RNA-sequencing (RNA-seq) to measure transcriptional differences between ARB25 post-

infected and mock-infected *B. thetaiotaomicron*. In a wild-type *B. thetaiotaomicron* population,

in which cells retain the ability to phase-vary expression of their eight different CPS, the

transcriptional profiles of bacterial populations surviving after ARB25 infection (n=3) were

largely characterized by decreased gene expression: among a total of 56 genes that exhibited

significant expression differences ≥3-fold between *B. thetaiotaomicron* exposed to live and heat-

killed ARB25, 51 genes were decreased in post-infected bacteria (**Figure 5A, Table S2**). Most

of these genes with decreased transcription (44/51) belong to the loci encoding CPS1 and CPS4,

consistent with our findings by qPCR that lower expression of these CPS occurs after ARB25

challenge (**Figure 4B**). Correspondingly, increased expression of genes encoding CPS2 and

CPS3 was also apparent by RNA-seq, but did not reach statistical significance. Interestingly, two

additional gene clusters encoding different outer-membrane "Sus-like systems", which are well-

described Bacteroidetes mechanisms for import and degradation of carbohydrates and other

nutrients[2,29], were also decreased in post-infected bacteria (6/7 remaining non-*cps* genes). The

central features of these systems are outer membrane TonB-dependent transporters (similar to *E.*

coli TonA, or **T** **on**e phage receptor **A**; the first described phage receptor[30]), suggesting the

possibility that the proteins encoded by these genes are part of the receptor for ARB25.

To further investigate CPS-independent mechanisms that allow *B. thetaiotaomicron* to

avoid eradication by phage, we performed identical RNA-seq experiments on acapsular *B.*

thetaiotaomicron populations exposed to heat-killed or live ARB25. In this case, we

hypothesized that eliminating the ability to phase vary capsules would better reveal the

mechanism(s) that allows the acapsular strain to survive ARB25 predation, possibly by further

reducing expression of putative phage receptors like the Sus-like systems identified above.

Expression of the same two Sus-like systems (*BT2170-73, BT2365*) that were decreased in

ARB25-exposed wild-type were also decreased to similar levels in acapsular *B. thetaiotaomicron*

(**Figure 5B**). Otherwise, the transcriptional profiles of acapsular *B. thetaiotaomicron* surviving

after ARB25 infection (n=3) were largely characterized by increased gene expression: 71 of the

81 genes differentially regulated $\geq$3-fold between ARB25-infected and the heat-killed reference

showed increased expression in ARB25 post-infected cells. All but 3 of these genes were unique

to the post-infection transcriptome of acapsular *B. thetaiotaomicron* compared to wild-type *B.*

thetaiotaomicron (**Figure 5B**, **Table S2**).

Among the 71 genes with increased expression in post-infected acapsular *B.*

thetaiotaomicron, 24 genes were part of 8 different gene clusters that encode putative tyrosine

recombinases along with pairs of outer membrane lipoproteins and OmpA β-barrel proteins

(**Figure 5C**). One of these genes (*BT1927*) was previously characterized as encoding a phase-

variable, S-layer protein, which organizes into a tessellated structure on the cell surface and

when locked into the "on" orientation promoted increased *B. thetaiotaomicron* resistance to

complement-mediated killing[31]. The remaining 7 gene clusters share both syntenic organization

and homology to this original S-layer gene cluster. Closer scrutiny of the promoter regions

upstream of the 7 newly identified gene clusters encoding putative lipoproteins revealed that

each is also flanked by a pair of imperfect, 17 nucleotide palindromic repeats (**Figure 5C**). Three

of these newly identified repeats are identical to the repeats known to mediate recombination at

the *BT1927* promoter[31]. The remaining 4 sequences only varied by the sequence of a

trinucleotide located in the middle of each imperfect palindrome (colored blocks in **Figure 5C**

middle). Finally, amplicon sequencing of each promoter supported the existence of the proposed

recombination events in 5 of the 7 newly identified loci (**Figure S6)**.

Among the remaining genes that were significantly up- or down-regulated in post

ARB25-infected acapsular *B. thetaiotaomicron*, there was an additional signature of genes for

which DNA recombination may be involved in re-organizing expression of cell surface proteins.

Specifically, the expression of 3 of 4 genes in an operon (*BT1042-45*) previously implicated in

utilization of host glycans[32] were expressed an average of 6.9-fold less in ARB25-infected

acapsular cells compared to heat-killed controls. Correspondingly, 5 genes in an adjacent operon

(*BT1046-51*) with similar arrangement and predicted functions exhibited an average of 8.3-fold

increased expression in acapsular *B. thetaiotaomicron* exposed to live ARB25. Both of these

operons have been previously linked to transcriptional regulation by a nearby extra-cytoplasmic

function sigma (ECF-σ), anti-σ factor pair, such that when the single ECF-σ coding gene

(*BT1053*) is deleted, the ability to activate the adjacent operons is eliminated[33]. Based on 1) the

ARB25-dependent shift in gene expression described above; 2) previously established common

ECF-σ regulation of the *BT1042-45* and *BT1046-51* operons; 3) the observation that two genes

encoding TonB-dependent transporters (*BT1040, BT1046*) appear to be truncated at their 5' ends

compared to *BT1042* (**Figure 4**, **S7A**) and only the full-length BT1042 sequence harbors a

required anti-σ contact domain[33]; and 4) annotation of a gene encoding a putative tyrosine

recombinase (*BT1041*) located in the middle of this locus, we hypothesized that this gene cluster

possesses the ability to undergo recombination and that specific combinatorial variants are

selected under phage pressure.

To test this hypothesis, we designed PCR primer pairs (**Figure 5D**, green dumbbells) to

detect both the originally annotated sequence orientation and 3 potential alternative

recombination states derived from either moving the full-length 5' end of *BT1042* to one of two

alternative *susC*-like genes or an internal rearrangement derived from recombination of two

incomplete *susC*-like genes (**Figure 5D**, variants 1-3). In support of our hypothesis, we were

able to detect by both PCR (**Figure 5E**) and amplicon sequencing (**Figure S7B**) the presence of

all 5 predicted alternative recombination states (**Figure 5D,E** amplicons 2, 3, 4, 6, 7), plus the 3

expected from the originally published genome assembly[34]. In further support of our hypothesis,

an insertion mutation in the associated tyrosine recombinase-coding gene (*BT1041*) locked the

corresponding mutant into the native genomic architecture as determined by the presence of

amplicon 1, but the absence of amplicons 2 and 3 (**Figure 5E**). Further sequence analysis and

tracking of single nucleotide polymorphisms in the 5' ends of the three recombinationally active

susC-like genes narrowed the recombination site down to a 7 bp sequence that is flanked by an

imperfect direct repeat encompassing over 132 additional downstream bp that may also influence

recombination specificity (**Figure S7B**). Thus, three separate operons that are under the

transcriptional control of a single ECF-σ regulator and are involved in utilization of host glycans,

are also able to undergo recombinational shuffling via a tyrosine recombinase/direct repeat

mediated mechanism to vary which of the three operons is expressed to produce its

corresponding surface proteins. This strategy is similar to recombinational shufflons involving

nutrient utilization functions that have been characterized in *B. fragilis*[35,36], with the exception

that in the example described here, recombination occurs between direct repeats instead of

palindromes. One explanation for this phenomenon is that shufflons evolved to subvert phage

infection by expressing alternate cell surface receptors for important nutrients, which are also

targeted by phages. Contrary to this hypothesis that the proteins encoded by BT1042-45 might

serve as a receptor for ARB25, elimination of the genes spanning BT1033-52 did not eliminate

ARB25 infection in the acapsular strain, suggesting that an additional or different receptor(s)

exists. Interestingly, the BT1033-52 mutant exhibited variable plaquing efficiency compared to

the acapsular parent (**Figure S8**), suggesting that loss of these genes cold exert global effects that

mediate susceptibility to ARB25.

Since the gene encoding the original outer membrane S-layer protein (*BT1927*), and its

downstream gene (*BT1926*), were among the most highly activated (147- and 114-fold,

respectively) in post-infected, acapsular *B. thetaiotaomicron,* and can be alternatively locked into

the "on" or "off" orientations by mutating the recombination site upstream of the phase-variable

promoter[31], we chose to focus on the role of this single function in resisting phage infection. We

re-engineered acapsular *B. thetaiotaomicron* to contain locked "on" and locked "off" versions of

this promoter and evaluated sensitivity to ARB25 and another phage, SJC01, which has a similar

infection profile on *B. thetaiotaomicron* (**Figure 1**). Consistent with the observation that S-layer

is highly activated in acapsular *B. thetaiotaomicron* infected with ARB25, acapsular S-layer

"off" cells were more effectively killed in the presence of live phage relative to acapsular S-layer

"on" cells (**Figure 6A**). Interestingly, SJC01 showed the opposite effect, as it more effectively

killed cells with S-layer locked "on" versus cells with S-layer locked "off" (**Figure 6B**). These

data indicate that in addition to the 8 surface-exposed capsular polysaccharide types, S-layer

lipoproteins can function as positive or negative determinants of phage tropism in *B.*

thetaiotaomicron. Finally, the observation that CPS[5,6] and homologous S-layer like functions are

broadly represented in gut Bacteroidetes[31], suggest that these two mechanisms help to diversify

members of this phylum under phage-mediated selection.

Discussion

Production of multiple phase-variable CPS is a hallmark of human gut Bacteroidetes.

Previous work has revealed the importance of *Bacteroides* CPS in interactions with the host

immune system[4,8,37,38]. However, other biological roles for *Bacteroides* CPS remain relatively

unexplored. Using a panel of *B. thetaiotaomicron* strains that express individual CPS, we tested a

previously inaccessible hypothesis: that *Bacteroides*-targeting phage can be both inhibited and

assisted by the repertoire of capsules expressed by their host bacteria. Our data clearly indicate

that production of specific CPS is associated with alterations in phage susceptibility, which is

underscored by the observation that none of the 71 phages characterized here infect every CPS-

variable strain that we tested (**Figure 1**). Phage-mediated selection and interactions with the host

immune system help to explain both the extensive diversification of CPS structures in gut-

resident Bacteroidetes[4,12] and their complex phase-variable regulation mechanisms within a

given strain or species[23,25]. Surprisingly, our results also reveal that additional phase variable

markers are expressed by *B. thetaiotaomicron* under phage-mediated selection, highlighting that

other strategies exist in *Bacteroides* for surviving in the face of phage predation.

There are several mechanisms through which CPS could promote or prevent phage

infection. First, CPS may sterically mask surface receptors to block phage binding, although

additional specificity determinants must be involved because no individual phage that infects the

acapsular strain is blocked by all single *B. thetaiotaomicron* CPS. These specificity determinants

could be driven by CPS structure (physical depth on the cell surface, polysaccharide charge,

permeability) or be actively circumvented by the presence of polysaccharide depolymerases on

the phage particles, as has been described in other phage-bacterium systems (e.g., *E. coli* K1 and

phiK1-5[39]). Alternatively, certain permissive CPS could serve as obligate receptors[40] (i.e., phage

that do not infect acapsular) or more generally increase the affinity of a phage for the bacterial

cell surface – similar to what has been proposed in the "bacteriophage adhering to mucus"

model, whereby hypervariable domains on phage capsids facilitate adherence to mucus and

increase the frequency of bacteria-phage interaction[41]. This latter type of adherence to CPS might

increase the likelihood that a phage would contact its receptor by sustained interaction with the

extracellular matrix. Some combination of these possibilities is likely to explain the host range

infection profile for the majority of the phages in our collection. Collectively, our observations

provide the foundation for future mechanistic work, beginning with phage genome sequencing,

aimed at understanding the physical and chemical interactions that mediate infection of *B.*

thetaiotaomicron and other *Bacteroides* by their phages.

Using ARB25 as a representative from our larger collection, we demonstrate that

infection with this single phage does not fully eradicate presumably susceptible *B.*

thetaiotaomicron populations. Rather, resistant cells grow, often quickly, after what we interpret

to be an initial lytic event (**Figure 3**). Similar observations were previously made with

ΦCrass001, a phage that infects *B. intestinalis* [21]. Specifically, though ΦCrass001 robustly

formed plaques on lawns of *B. intestinalis*, it failed to eradicate this bacterium in liquid culture.

We hypothesize that ΦCrass001 undergoes lytic growth on a subset of *B. intestinalis* cells

expressing permissive CPS. Subsequently, clones expressing resistant CPS would grow to

dominate a culture after phage challenge, as we observed with ARB25.

Given the roles of CPS in mediating *B. thetaiotaomicron*-phage interactions, the

outgrowth of a phage resistant sub-population was especially surprising in the context of

acapsular *B. thetaiotaomicron*. While wild-type *B. thetaiotaomicron* primarily appears to survive

continued ARB25 predation by pre-adaptive CPS variation or by shifting the population to

express non-permissive CPS, the acapsular strain instead shifts to increased expression of phase-

variable sets of surface proteins, at least one of which (*BT1927-26*) confers increased resistance

to ARB25 when locked "on" in acapsular *B. thetaiotaomicron*. A previous study measured that

only 1:1000 *B. thetaiotaomicron* cells in an unchallenged population express the S-layer encoded

by *BT1927* [31]. Given that ARB25 non-permissive CPS can comprise up to 40% of the expressed

capsule population (e.g., CPS3 in **Figure 4B**), the rapid emergence of cells expressing alternative

CPS could be explained by this frequency difference, in which case S-layer expressing wild-type

bacteria might also emerge given longer phage exposure times. The original *B. thetaiotaomicron*

S-layer study also demonstrated that locking the invertible promoter for the *BT1927* S-layer into

the "on" orientation facilitated survival against complement-mediated killing[31], suggesting that

orthogonal roles for this and related proteins exist in *B. thetaiotaomicron* and likely facilitate

survival in the face of diverse environmental disturbances. Combined with our data on CPS-

mediated phage tropism, our observations that the *BT1927*-encoded S-layer confers resistance to

some phages, that 7 other homologous systems are also upregulated after exposure to ARB25,

and that a shufflon harboring three recombinationally-variable nutrient acquisition operons exists

in *B. thetaiotaomicron*, together reveal that there are at least 17 independent cell surface

structures in *B. thetaiotaomicron* that could be altered in cells exposed to phages. The fact that

almost all of these surface structures (14/17) are under the control of independent phase-variable

promoters with associated tyrosine recombinases[4,31,36] and that their products show altered

expression after phage exposure speaks to the effectiveness of this strategy in pre-adapting some

B. thetaiotaomicron cells within a population to thrive in the face of phage predation.

Phages are the most abundant biological entities in the gut microbiome[41] and interest in

the roles and identities of these gut-resident viruses is increasing as metagenomic sequencing

approaches are unveiling a more comprehensive view of their dynamics during health and

disease[16,17,42]. Although sequence-based approaches are powerful for describing the phages that

are present, they do not generate information on the definitive hosts or the mechanisms of

individual bacteria-phage interactions, which are likely to be elaborate. These limitations will

prohibit full dissection of the ecological interactions that phage exert on bacterial populations in

the gut. The approach taken here of isolating phages for a particular host of interest, with added

layers of detail like systematic variation of surface CPS when possible, will be an essential

complement to high throughput sequencing studies and will help build a foundation of

mechanistic gut bacterium-phage interactions.

Our results, with a single strain of bacteria commonly found in human gut microbiomes,

point to the existence of a very complex relationship between bacteria and phage in the gut

microbiome. Considering the possibilities that these interactions could vary over time, differ by

host species, and evolve differently within individuals or regionally distinct global populations,

the landscape becomes even more complex. Given the diverse adaptive and counter-adaptive

strategies that have apparently evolved in the successful gut symbiont *B. thetaiotaomicron* and

its relatives, our findings hold important implications for the use of phages to intentionally alter

the composition or function of the gut microbiota. While a cocktail of multiple phages could

theoretically be harnessed together to elicit more robust alteration of target populations within a

microbiome, the complexity of host tropisms and bacterial countermeasures that exist for *B.*

thetaiotaomicron imply that a deliberate selection of complementary phage would be needed. If

selections of effective phage cocktails need to be further tailored to individual microbiomes, or

elicit resistance within individuals or populations the way antibiotics do, the prospects for

effective gut microbiome-targeting phage therapy could indeed become very complicated. Given

these considerations, our findings here contribute an important early step towards building a deep

functional understanding of the bacterium-virus interactions that occur in the human gut

microbiome. As such, this work contributes to the overall goal of understanding the ecology of

this important microbial community and developing rational approaches to shape its physiology.

Methods

Bacterial strains and culture conditions.

The bacterial strains used in this study are listed in **Table S3.** Frozen stocks of these

strains were maintained in 25% glycerol at -80˚C and were routinely cultured in an anaerobic

chamber or in anaerobic jars (using GasPak EZ anaerobe container system sachets w/indicator,

BD) at 37˚C in *Bacteroides* Phage Recovery Medium (BPRM), as described previously[43]: per 1

liter of broth, 10 g meat peptone, 10 g casein peptone, 2 g yeast extract, 5 g NaCl, 0.5 g L-

cysteine monohydrate, 1.8 g glucose, and 0.12 g $MgSO_4$ heptahydrate were added; after

autoclaving and cooling to approximately 55 °C, 10 ml of 0.22 µm-filtered hemin solution (0.1%

w/v in 0.02% NaOH), 1 ml of 0.22 µm-filtered 0.05 g/ml $CaCl_2$ solution, and 25 ml of 0.22µm-

filtered 1 M Na_2CO_3 solution were added. For BPRM agar plates, 15 g/L agar was added prior to

autoclaving and hemin and Na_2CO_3 were added as above prior to pouring the plates. For BPRM

top agar used in soft agar overlays, 3.5 g/L agar was added prior to autoclaving. Hemin, $CaCl_2$,

and Na_2CO_3 were added to the top agar as above immediately before conducting experiments.

Bacterial strains were routinely struck from the freezer stocks onto agar plates of Brain Heart

Infusion supplemented with 10% horse blood (Quad Five, Rygate, Montana) (BHI-blood agar; or

for the SJC phages used in **Figure 1**, on BPRM agar) and grown anaerobically for up to 3 days.

A single colony was picked for each bacterial strain, inoculated into 5 mL BPRM, and grown

anaerobically overnight to provide the starting culture for experiments.

For the experiment described in **Figure 2G**, liquid cultures of *B. thetaiotaomicron* were

grown in BPRM using the pyrogallol method as described previously[4]. Briefly, a sterile cotton

ball was burned and then pushed midway into the tube, after which 200 μl of saturated $NaHCO_3$

and 200 μl of 35% pyrogallol solution were added to the cotton ball. A rubber stopper was used

to seal the tubes, and tubes were incubated at 37 °C.

Bacteriophage isolation from primary wastewater effluent.

The bacteriophages described in this study were isolated from primary wastewater

effluent from two locations at the Ann Arbor, Michigan Wastewater Treatment Plant and from

the San Jose-Santa Clara Regional Wastewater Treatment Facility. After collection, the primary

effluent was centrifuged at 5,500 rcf for 10 minutes at room temperature to remove any

remaining solids. The supernatant was then sequentially filtered through 0.45 μm and 0.22 μm

polyvinylidine fluoride (PVDF) filters to yield "processed primary effluent." Initial screening for

plaques was done using a soft agar overlay method[44] where processed primary effluent was

combined with 1 part overnight culture to 9 parts BPRM top agar and poured onto a BPRM agar

plate (e.g. 0.5 mL overnight culture and 4.5 mL BPRM top agar was used for standard circular

petri dishes [100 mm x 15 mm]). Soft agar overlays were incubated anaerobically at 37 ˚C

overnight. Phages were successfully isolated using three permutations of this assay: (1) Direct

plating, where processed primary effluent was directly added to overnight culture prior to

plating. (2) Enrichment, where 10 mL processed primary effluent was mixed with 10 mL

2XBPRM and 3 mL exponential phase *B. thetaiotaomicron* culture and grown overnight. The

culture was centrifuged at 5500 rcf for 10 minutes and filtered through a 0.22 μm PVDF filter.

(3) Size exclusion, where processed primary effluent was concentrated up to 500-fold via 30 or

100 kDa PVDF or polyethersulfone size exclusion columns. Up to 1 mL of processed primary

effluent, enrichment, or concentrated processed primary effluent was added to the culture prior to

adding BPRM top agar, as described above. To promote a diverse collection of phages, no more

than 5 plaques from the same plate were plaque purified and a diversity of plaque morphologies

were selected as applicable. When using individual enrichment cultures, only a single plaque was

purified.

Single, isolated plaques were picked into 100 μL phage buffer (prepared as an autoclaved

solution of 5 ml of 1 M Tris pH 7.5, 5 ml of 1 M MgSO4, 2 g NaCl in 500 ml with ddH$_2$O).

Phages were successfully plaque purified using one of two methods: (1) a standard full plate

method, where the diluted phage samples were combined with *B. thetaiotaomicron* overnight

culture and top agar and plated via soft agar overlay as described above or (2) a higher

throughput 96-well plate-based method, where serial dilutions were prepared in 96-well plates

and 1 μL of each dilution was spotted onto a solidified top agar overlay. This procedure was

repeated at least 3 times to plaque purify each phage. For more details on the phages isolated in

this work, see **Figure S1**.

High titer phage stocks were generated by flooding a soft agar overlay that plate yielded a

"lacey" pattern of bacterial growth (near confluent lysis). Following overnight incubation of each

plate, 5 ml of sterile phage buffer was added to the plate to resuspend the phage. After at least 2

hours of incubation at room temperature, the lysate was spun at 5,500 rcf for 10 minutes to clear

debris and then filter sterilized through a 0.22 μm PVDF filter.

Quantitative host range assays.

To accommodate the large number of phage isolates in our collection, we employed a

spot titer assay for semi-quantitative comparisons of infectivity on each bacterial strain. High

titer phage stocks were prepared on their "preferred host strain," which was the strain that

yielded the highest titer of phages in a pre-screen of phage host range. Lysates were then diluted

to approximately 10^6 PFU/mL, were added to the wells of a 96-well plate, and further diluted to

10^5, 10^4, and 10^3 PFU/mL using a multichannel pipettor. One microliter of each of these

dilutions was plated onto solidified top agar overlays containing single bacterial strains indicated

in each figure. After spots dried, plates were incubated anaerobically for 15-24 hours prior to

counting plaques. Phage titers were normalized to the bacterial strain that typically exhibited the

highest phage titer, which was designated as the "preferred host strain".

Images of phage plaques.

To document the morphologies of plaques formed by the purified phages, two sets of

plaque pictures were taken: the first set were taken with a Color QCount Model 530 (Advanced

Instruments) with a 0.01 second exposure. Images were cropped to 7.5 mm^2 but were otherwise

unaltered. The second set of images were taken on a ChemiDoc Touch instrument (BioRad) with

a 0.5 second exposure. Images were cropped to 7.5 mm^2 and background unnaturally high pixels

were removed (Image Lab, BioRad) to facilitate viewing of the plaques.

Incubation of phage with extracted CPS.

Approximately 50-100 PFU of ARB25 in 50 µl phage buffer were mixed with an equal

volume of H_2O or capsule (2 mg/ml) extracted by the hot water-phenol method (as described in

Reference 4) and incubated at 37 °C for 30 minutes. Samples were then plated on the acapsular

strain, and plaques were counted after 15-24 hours anaerobic incubation at 37 °C. Counts from

two replicates on the same day were then averaged, and the experiment was performed three

times.

Growth curves.

For growth curve experiments, 3 individual clones of each indicated strain were picked

from agar plates and grown overnight in BPRM. Then, for experiments in **Figures 3** and **6**, each

clone was diluted 1:100 in fresh BPRM and 100 µl was added to a microtiter plate. 10 µl of

approximately $5*10^6$ PFU/ml live or heat-killed phage were added to each well, plates were

covered with an optically clear gas-permeable membrane (Diversified Biotech, Boston, MA) and

optical density at 600 nm (OD_{600}) values were measured using an automated plate reading device

(BioTek Instruments). Phages were heat killed by heating to 95 °C for 30 minutes, and heat-

killed phage had no detectable PFU/ml with a limit of detection of 100 PFU/ml.

In **Figure S3B**, wild-type *B. thetaiotaomicron* was infected with live or heat-killed

ARB25, and bacterial growth was monitored via optical density at 600 nm (OD_{600}) on an

automated plate reader for 12 hours. At 0, 6.02, 8.36, and 11.7 hours post inoculation, replicate

cultures were vortexed in 1:5 volume chloroform, centrifuged at 5,500 rcf at 4 °C for 10 minutes,

and the aqueous phase was titered on the acapsular strain. No phages were detected in heat-killed

controls.

Generation of phage-free bacterial isolates and determination of their phage susceptibility.

To isolate phage-free bacterial clones from ARB25-infected cultures (Table 1), each

culture was streaked on a BHI-blood agar plate. Eighteen individual colonies were picked from

each plate, and each of these clones was re-streaked onto a new BHI-blood agar plate. One

colony was picked from each of these secondary plates and was inoculated into 150 μl BPRM

broth and incubated anaerobically at 37 °C for 2 days. Only one of the clones (a *cps*4 isolate)

failed to grow in liquid media. To determine whether cultures still contained viable phage, 50 μl

of each culture was vortexed with 20 μl chloroform, then centrifuged at 5,500 rcf for 10 minutes.

10 μl of the lysate was spotted on BPRM top agar containing naïve acapsular bacteria and was

incubated anaerobically overnight at 37 °C. Loss of detectable phage in the twice passaged

clones was confirmed for most of the clones (79/89, 89%) by the absence of plaques on the naïve

acapsular strain.

To determine whether the resultant phage-free plaques were resistant to ARB25 infection,

each culture was diluted 1:100 in fresh BPRM, 100 μl was added to a microtiter plate, and 10 μl

of either live or heat-killed ARB25 (approximately $5*10^6$ PFU/ml) was added. Plates were

incubated anaerobically at 37 °C for 48 hours, and OD_{600} was measured as described above.

Cultures were determined to be susceptible to ARB25 by demonstration of delayed growth or

drop in OD_{600}, as compared to heat-killed controls.

***Measurement of* cps *gene expression*.**

For **Figures 2G**, **4**, and **S4**, overnight cultures were subcultured into fresh BPRM to an

OD_{600} of 0.01. For Figure 4B, 200 µL of approximately 2 x 10^8 PFU/mL live phage or heat killed

phage were added to 5 mL of the diluted cultures. For **Figure S4**, 200 µL of approximately 2 x

10^5 PFU/mL live phage or heat killed phage were added to 5 mL of the diluted cultures. Bacterial

growth was monitored by measuring OD_{600} every 15-30 minutes using a GENESYS 20

spectrophotometer (Thermo Scientific). Cultures were briefly mixed by hand before each

measurement. For determination of relative *cps* gene expression, cultures were grown to OD_{600}

0.6-0.8, were centrifuged at 7700 rcf for 2.5 minutes, the supernatant was decanted, and the

pellet was immediately resuspended in 1 ml RNA-Protect (Qiagen). RNA-stabilized cell pellets

were stored at -80 °C.

Total RNA was isolated using the RNeasy Mini Kit (Qiagen) then treated with the

TURBO DNA-free Kit (Ambion) followed by an additional isolation using the RNeasy Mini Kit.

cDNA was then synthesized using SuperScript III Reverse Transcriptase (Invitrogen) according

to the manufacturer's instructions using random oligonucleotide primers (Invitrogen). qPCR

analyses for *cps* locus expression were performed on a Mastercycler ep realplex instrument

(Eppendorf). Expression of each of the 8 *cps* synthesis loci was quantified using primers to a

single gene in each locus (primers are listed in **Table S4**) and normalized to a standard curve of

DNA from wild-type *B. thetaiotaomicron*. Relative abundance of expression for each locus was

then calculated. A custom-made SYBR-based master mix was used for qPCR: 20 µl reactions

were made with ThermoPol buffer (New England Biolabs), and containing 2.5 mM MgSO$_4$,

0.125 mM dNTPs, 0.25 µM each primer, 0.1 µl of a 100 X stock of SYBR Green I (Lonza), and

500 U Hot Start *Taq* DNA Polymerase (New England Biolabs). 10 ng of cDNA was used for

each sample, and samples were run in duplicate. A touchdown protocol with the following

cycling conditions was used for all assays: 95 °C for 3 minutes, followed by 40 cycles of 3

seconds at 95 °C, 20 seconds of annealing at a variable temperature, and 20 seconds at 68 °C.

The annealing temperature for the first cycle was 58 °C, then dropped one degree each cycle for

the subsequent 5 cycles. The annealing temperature for the last 34 cycles was 52 °C. These

cycling conditions were followed by a melting curve analysis to determine amplicon purity.

Transcriptomic analysis of* B. thetaiotaomicron *after phage infection.

Whole genome transcriptional profiling of wild-type and acapsular *B.thetaiotaomicron*

infected with live or heat-killed ARB25 was conducted using total bacterial RNA that was

extracted the same as described above (Qiagen RNAEasy, Turbo DNA-free kit) and then treated

with Ribo-Zero rRNA Removal Kit (Illumina Inc.) and concentrated using RNA Clean and

Concentrator -5 kit (Zymo Research Corp, Irvine, CA). Sequencing libraries were prepared using

TruSeq barcoding adaptors (Illumina Inc.), and 24 samples were multiplexed and sequenced with

50 base pair single end reads in one lane of an Illumina HiSeq instrument at the University of

Michigan Sequencing Core. Demultiplexed samples were analyzed via Arraystar software

(DNASTAR, Inc.) using RPKM normalization and the default parameters. Changes in gene

expression in response to live ARB25 infection were determined by comparison to the heat-

killed reference: retained were genes with $\geq$ 3-fold expression changes up or down, Benjamini-

Hochberg corrected *P* value $\leq$ 0.05, and genes for which the lower value in the fold-change

calculation was at least 1% of the mean RPKM-normalized value for all genes in the

transcriptome. The latter cutoff was implemented to reduce the noise effects of changes in genes

with very low expression values.

PCR and sequencing of phase variable* B. thetaiotaomicron *chromosomal loci.

We found that each of the 8 chromosomal loci shown in **Figure 5C** had nearly identical

301 bp promoter sequences, including both of the imperfect palindromes that we predict to

mediate recombination and the intervening sequence at each locus. While the 8 S-layer genes

and the 7/8 of the upstream regions encoding putative tyrosine recombinases (all but the BT1927

region) shared significant nucleotide identity, we were able to design primers that were specific

to regions upstream and downstream of each invertible promoter and used these to generate an

amplicon for each locus that spanned the predicted recombination sites. After gel extracting a

PCR product of the expected size for each locus, which should contain promoter orientations in

both the "on" and "off" orientations, we performed a second PCR using a universal primer that

lies within the 301 bp sequence of each phase-variable promoter and extended to unique primers

that anneal within each S layer protein encoding gene. Bands of the expected size were excised

from agarose gels, purified and sequenced using the primer that anneals within each S layer

encoding gene to determine if the predicted recombined "on" promoter orientation can be

detected. (Note that the assembled *B. thetaiotaomicron* genome architecture places all of these

promoters in the proposed "off" orientation. We were able to detect 6/8 of these loci in the "on"

orientation in ARB25-treated cells by this method, **Figure S6**.) A similar approach was used to

determine the re-orientation of DNA fragments in the *B. thetaiotaomicron* PUL shufflon shown

in **Figure 5D**, using PCR primer amplicons positioned according to the schematic followed by

sequencing with the primer on the "downstream" end of each amplicon according to its position

relative to the shuffled promoter sequence. For a list of primers used see **Table S4**.

Construction of acapsular* B. thetaiotaomicron *S-layer 'ON' and S-layer 'OFF' mutants.

Acapsular *B. thetaiotaomicron* S-layer 'ON' and 'OFF' mutants (Δ*cps* BT1927-ON and

Δ*cps* BT1927-OFF, respectively) were created using the Δ*tdk* allelic exchange system[45]. To

generate homologous regions for allelic exchange, the primers BT_1927_XbaI-DR and

BT_1927_SalI-UF were used to amplify the BT1927-ON and BT1927-OFF promoters from the

previously-constructed BT1927-ON and BT1927-OFF strains[31] via colony PCR using Q5 High

Fidelity DNA polymerase (New England Biolabs). Candidate Δ*cps* BT1927-ON and Δ*cps*

BT1927-OFF mutants were screened by PCR using the primer pair BT1927_Diagnostic_R and

BT1927_Diagnostic_F and confirmed by Sanger sequencing using these diagnostic primers. All

plasmids and primers are listed in **Supplementary Tables S3** and **S4**, respectively.

Data representation and statistical analysis.

The heatmaps for **Figures 1** and **S2** and the dendrogram for **Figure 1** were generated in R

using the "heatmap" function. Other graphs were created in Prism software (GraphPad Software,

Inc., La Jolla, CA). Statistical significance in this work is denoted as follows unless otherwise

indicated: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. Statistical analyses other than

Dirichlet regression were performed in Prism. Dirichlet regression was performed in R using the

package "DirichletReg" (version 0.6-3), employing the alternative parameterization as used

previously[4,46]. Briefly, the parameters in this distribution are the proportions of relative *cps* gene

expression and the total *cps* expression, with *cps7* expression used as a reference. The variable of

interest used in **Figure 2G** is bacterial strain, whereas the variable of interest used in **Figure 4B**

is phage viability (live versus heat-killed phage). Precision was allowed to vary by group given

this model was superior to a model with constant precision, as determined by a likelihood ratio

test at significance level p < 0.05.

686
687   **Acknowledgements**
688

We thank Rey Honrada at the San Jose-Santa Clara Wastewater Treatment Plant and the staff at
the Ann Arbor Wastewater treatment plant for assistance in collecting primary sewage effluent.
This work was funded by NIH grants (GM099513 and DK096023 to E.C.M), an NIH
postdoctoral NRSA (5T32AI007328 to A.J.H.), a Stanford University School of Medicine
Dean's Postdoctoral Fellowship (A.J.H.), and the NIH Cellular Biotechnology Training Program
(N.T.P., T32GM008353).

695
696   **Author Contributions**
697
698   NTP, AJH, BDM, JOG, and SS performed the experiments. NTP, AJH, and ECM designed the
699   experiments, and analyzed and interpreted the data. JLS and ECM provided tools and reagents.
700   NTP, AJH and ECM prepared the display items. NTP, AJH and ECM wrote the paper. All
701   authors edited the manuscript prior to submission.
702
703   **References**

704   1   Eckburg, P. B. *et al.* Diversity of the human intestinal microbial flora. *Science (New*
705   *York, N.Y.)* **308**, 1635-1638, doi:10.1126/science.1110591 (2005).
706   2   Cockburn, D. W. & Koropatkin, N. M. Polysaccharide Degradation by the Intestinal
707   Microbiota and Its Influence on Human Health and Disease. *Journal of molecular*
708   *biology* **428**, 3230-3252, doi:10.1016/j.jmb.2016.06.021 (2016).
709   3   Hooper, L. V., Littman, D. R. & Macpherson, A. J. Interactions between the microbiota
710   and the immune system. *Science (New York, N.Y.)* **336**, 1268-1273,
711   doi:10.1126/science.1223490 (2012).
712   4   Porter, N. T., Canales, P., Peterson, D. A. & Martens, E. C. A Subset of Polysaccharide
713   Capsules in the Human Symbiont Bacteroides thetaiotaomicron Promote Increased
714   Competitive Fitness in the Mouse Gut. *Cell host & microbe* **22**, 494-506.e498,
715   doi:10.1016/j.chom.2017.08.020 (2017).
716   5   Donia, M. S. *et al.* A systematic analysis of biosynthetic gene clusters in the human
717   microbiome reveals a common family of antibiotics. *Cell* **158**, 1402-1414,
718   doi:10.1016/j.cell.2014.08.032 (2014).
719   6   Coyne, M. J. & Comstock, L. E. Niche-specific features of the intestinal bacteroidales.
720   *Journal of bacteriology* **190**, 736-742, doi:10.1128/jb.01559-07 (2008).
721   7   Xu, J. *et al.* Evolution of symbiotic bacteria in the distal human intestine. *PLoS biology* **5**,
722   e156, doi:10.1371/journal.pbio.0050156 (2007).

8   Peterson, D. A., McNulty, N. P., Guruge, J. L. & Gordon, J. I. IgA response to symbiotic
bacteria as a mediator of gut homeostasis. *Cell host & microbe* **2**, 328-339,
doi:10.1016/j.chom.2007.09.013 (2007).
9   Round, J. L. *et al.* The Toll-like receptor 2 pathway establishes colonization by a
commensal of the human microbiota. *Science (New York, N.Y.)* **332**, 974-977,
doi:10.1126/science.1206095 (2011).
10   Shen, Y. *et al.* Outer membrane vesicles of a human commensal mediate immune
regulation and disease protection. *Cell host & microbe* **12**, 509-520,
doi:10.1016/j.chom.2012.08.004 (2012).
11   Neff, C. P. *et al.* Diverse Intestinal Bacteria Contain Putative Zwitterionic Capsular
Polysaccharides with Anti-inflammatory Properties. *Cell host & microbe* **20**, 535-547,
doi:10.1016/j.chom.2016.09.002 (2016).
12   Patrick, S. *et al.* Twenty-eight divergent polysaccharide loci specifying within- and
amongst-strain capsule diversity in three strains of Bacteroides fragilis. *Microbiology*
(Reading, England) **156**, 3255-3269, doi:10.1099/mic.0.042978-0 (2010).
13   Porter, N. T. & Martens, E. C. The Critical Roles of Polysaccharides in Gut Microbial
Ecology and Physiology. *Annual review of microbiology* **71**, 349-369,
doi:10.1146/annurev-micro-102215-095316 (2017).
14   Reyes, A. *et al.* Viruses in the faecal microbiota of monozygotic twins and their mothers.
Nature **466**, 334-338, doi:10.1038/nature09199 (2010).
15   Minot, S. *et al.* The human gut virome: inter-individual variation and dynamic response
to diet. *Genome research* **21**, 1616-1625, doi:10.1101/gr.122705.111 (2011).
16   Duerkop, B. A. *et al.* Murine colitis reveals a disease-associated bacteriophage
community. *Nature microbiology*, doi:10.1038/s41564-018-0210-y (2018).
17   Manrique, P. *et al.* Healthy human gut phageome. *Proceedings of the National Academy*
of Sciences of the United States of America **113**, 10400-10405,
doi:10.1073/pnas.1601060113 (2016).
18   Booth, S. J., Van Tassell, R. L., Johnson, J. L. & Wilkins, T. D. Bacteriophages of
Bacteroides. *Reviews of infectious diseases* **1**, 325-336 (1979).
19   Cooper, S. W., Szymczak, E. G., Jacobus, N. V. & Tally, F. P. Differentiation of
Bacteroides ovatus and Bacteroides thetaiotaomicron by means of bacteriophage. *Journal*
of clinical microbiology **20**, 1122-1125 (1984).
20   Keller, R. & Traub, N. The characterization of Bacteroides fragilis bacteriophage
recovered from animal sera: observations on the nature of bacteroides phage carrier
cultures. *The Journal of general virology* **24**, 179-189, doi:10.1099/0022-1317-24-1-179
(1974).
21   Shkoporov, A. N. *et al.* PhiCrAss001 represents the most abundant bacteriophage family
in the human gut and infects Bacteroides intestinalis. *Nature communications* **9**, 4781,
doi:10.1038/s41467-018-07225-7 (2018).
22   Puig, A., Araujo, R., Jofre, J. & Frias-Lopez, J. Identification of cell wall proteins of
Bacteroides fragilis to which bacteriophage B40-8 binds specifically. *Microbiology*
(Reading, England) **147**, 281-288, doi:10.1099/00221287-147-2-281 (2001).
23   Chatzidaki-Livanis, M., Weinacht, K. G. & Comstock, L. E. Trans locus inhibitors limit
concomitant polysaccharide synthesis in the human gut symbiont Bacteroides fragilis.
Proceedings of the National Academy of Sciences of the United States of America **107**,
11976-11980, doi:10.1073/pnas.1005039107 (2010).

24   Rogers, T. E. *et al.* Dynamic responses of Bacteroides thetaiotaomicron during growth on
770   glycan mixtures. *Molecular microbiology* **88**, 876-890, doi:10.1111/mmi.12228 (2013).

771   25   Hickey, C. A. *et al.* Colitogenic Bacteroides thetaiotaomicron Antigens Access Host
772   Immune Cells in a Sulfatase-Dependent Manner via Outer Membrane Vesicles. *Cell host
773   & microbe* **17**, 672-680, doi:10.1016/j.chom.2015.04.002 (2015).

774   26   Chatzidaki-Livanis, M., Coyne, M. J. & Comstock, L. E. A family of transcriptional
775   antitermination factors necessary for synthesis of the capsular polysaccharides of
776   Bacteroides fragilis. *Journal of bacteriology* **191**, 7288-7295, doi:10.1128/jb.00500-09
777   (2009).

778   27   Pelkonen, S., Aalto, J. & Finne, J. Differential activities of bacteriophage depolymerase
779   on bacterial polysaccharide: binding is essential but degradation is inhibitory in phage
780   infection of K1-defective Escherichia coli. *Journal of bacteriology* **174**, 7757-7761
781   (1992).

782   28   Krinos, C. M. *et al.* Extensive surface diversity of a commensal microorganism by
783   multiple DNA inversions. *Nature* **414**, 555-558, doi:10.1038/35107092 (2001).

784   29   Martens, E. C., Koropatkin, N. M., Smith, T. J. & Gordon, J. I. Complex glycan
785   catabolism by the human gut microbiota: the Bacteroidetes Sus-like paradigm. *The
786   Journal of biological chemistry* **284**, 24673-24677, doi:10.1074/jbc.R109.022848 (2009).

787   30   Braun, V. FhuA (TonA), the career of a protein. *Journal of bacteriology* **191**, 3431-3436,
788   doi:10.1128/jb.00106-09 (2009).

789   31   Taketani, M., Donia, M. S., Jacobson, A. N., Lambris, J. D. & Fischbach, M. A. A Phase-
790   Variable Surface Layer from the Gut Symbiont Bacteroides thetaiotaomicron. *mBio* **6**,
791   e01339-01315, doi:10.1128/mBio.01339-15 (2015).

792   32   Martens, E. C., Chiang, H. C. & Gordon, J. I. Mucosal glycan foraging enhances fitness
793   and transmission of a saccharolytic human gut bacterial symbiont. *Cell host & microbe* **4**,
794   447-457, doi:10.1016/j.chom.2008.09.007 (2008).

795   33   Martens, E. C., Roth, R., Heuser, J. E. & Gordon, J. I. Coordinate regulation of glycan
796   degradation and polysaccharide capsule biosynthesis by a prominent human gut
797   symbiont. *The Journal of biological chemistry* **284**, 18445-18457,
798   doi:10.1074/jbc.M109.008094 (2009).

799   34   Xu, J. *et al.* A genomic view of the human-Bacteroides thetaiotaomicron symbiosis.
800   *Science (New York, N.Y.)* **299**, 2074-2076, doi:10.1126/science.1080029 (2003).

801   35   Nakayama-Imaohji, H. *et al.* Identification of the site-specific DNA invertase responsible
802   for the phase variation of SusC/SusD family outer membrane proteins in Bacteroides
803   fragilis. *Journal of bacteriology* **191**, 6003-6011, doi:10.1128/jb.00687-09 (2009).

804   36   Kuwahara, T. *et al.* Genomic analysis of Bacteroides fragilis reveals extensive DNA
805   inversions regulating cell surface adaptation. *Proceedings of the National Academy of
806   Sciences of the United States of America* **101**, 14919-14924,
807   doi:10.1073/pnas.0404172101 (2004).

808   37   Mazmanian, S. K., Liu, C. H., Tzianabos, A. O. & Kasper, D. L. An immunomodulatory
809   molecule of symbiotic bacteria directs maturation of the host immune system. *Cell* **122**,
810   107-118, doi:10.1016/j.cell.2005.05.007 (2005).

811   38   Donaldson, G. P. *et al.* Gut microbiota utilize immunoglobulin A for mucosal
812   colonization. *Science (New York, N.Y.)* **360**, 795-800, doi:10.1126/science.aaq0926
813   (2018).

39   Tomlinson, S. & Taylor, P. W. Neuraminidase associated with coliphage E that
specifically depolymerizes the Escherichia coli K1 capsular polysaccharide. *Journal of*
virology **55**, 374-378 (1985).
40   Gupta, D. S. *et al.*
41   Barr, J. J. *et al.* Bacteriophage adhering to mucus provide a non-host-derived immunity.
Proceedings of the National Academy of Sciences of the United States of America **110**,
10771-10776, doi:10.1073/pnas.1305923110 (2013).
42   Norman, J. M. *et al.* Disease-specific alterations in the enteric virome in inflammatory
bowel disease. *Cell* **160**, 447-460, doi:10.1016/j.cell.2015.01.002 (2015).
43   Tartera, C., Araujo, R., Michel, T. & Jofre, J. Culture and decontamination methods
affecting enumeration of phages infecting Bacteroides fragilis in sewage. *Applied and*
environmental microbiology **58**, 2670-2673 (1992).
44   Araujo, R. *et al.* Optimisation and standardisation of a method for detecting and
enumerating bacteriophages infecting Bacteroides fragilis. *Journal of virological methods*
93, 127-136 (2001).
45   Koropatkin, N., Martens, E. C., Gordon, J. I. & Smith, T. J. Structure of a SusD
homologue, BT1043, involved in mucin O-glycan utilization in a prominent human gut
symbiont. *Biochemistry* **48**, 1532-1542, doi:10.1021/bi801942a (2009).
46   MJ, M. DirichletReg: Dirichlet Regression in R. (2015).
833

Figure Legends

Figure 1. Host range of *B. thetaiotaomicron* phages on strains expressing different CPS.

Seventy-one bacteriophages were isolated and purified on the wild-type, Δ*cps* (acapsular), or the

8 single CPS-expressing *B. thetaiotaomicron* strains. High titer phage stocks were prepared on

their "preferred host strain," which was the strain that yielded the highest titer of phages in a pre-

screen of phage host range; phages were then tested in a quantitative host range assay. Tenfold

serial dilutions of each lysate ranging from approximately 10^6 to 10^3 plaque-forming units (PFU)

/ ml were spotted onto top agar plates containing each of the 10 bacterial strains. Plates were then

incubated overnight, and plaques on each host were counted. Titers were calculated for each host

and normalized to the titer on the "preferred host strain" for each replicate and replicates (n = 3)

were averaged. The phages were then clustered based on their plaquing efficiencies on the

different strains (see *Methods*). Each row in the heatmap corresponds to an individual phage,

whereas each column corresponds to one of the 10 host strains. Images at the far right of the

figure illustrate the range of plaque morphologies of select phages from the collection (see

Figure S1 for images of plaques for all phages). ARB25, a phage that is the focus of deeper

investigation is highlighted in red text. Scale bar = 2mm.

Figure 2. Infection by the Branch 3 phages ARB72 (A), ARB78 (B), ARB82 (C), ARB101 (D)

and ARB105 (E) is inhibited by eliminating most or all of the permissive CPS from wild-type *B.*

thetaiotaomicron. Each phage was titered on the wild-type strain, the acapsular strain, their

respective preferred host strain (blue bars), and a set of bacterial strains harboring limited CPS

locus deletions that correspond to their predetermined host range (n = 6 replicates/phage). (F)

Elimination of permissive CPS from Branch 2 phage ARB25 reduces infection, but complete

reduction of infection only occurs in the context of deleting more than one permissive CPS. The

number of replicates conducted on each strain is annotated in parentheses next to the strain name.

(G) Relative *cps* locus expression of the 8 *cps* loci in the indicated strains. (H) Representative

pictures of phage plaques on the indicated host strains. The top row of images for each phage is

unaltered; background and unnaturally saturated pixels were removed from images in in the

bottom row to facilitate plaque visualization. Scale bar = 2mm. For panels A-F, significant

differences in phage titers on the preferred host strain were calculated via Kruskal Wallis test

followed by Dunn's multiple comparisons test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, **** $p <$

0.0001. For panel G, deletion of *cps* loci resulted in significant changes in *cps* gene expression

for the missing loci; additionally, significant changes in *cps2* expression were observed in $\Delta 4$

and $\Delta 1,4$ strains ($p < 0.05$ for each change, determined by Dirichlet regression).

Figure 3. Effect of CPS and phage infection on bacterial growth. Ten strains: the wild-type

(WT), the acapsular strain (Δcps), or the eight single CPS-expressing strains were infected with

either live or heat-killed ARB25. Growth was monitored via optical density at 600 nm (OD_{600})

on an automated plate reading instrument as described in *Methods*.

Figure 4. ARB25 infection of wild-type *B. thetaiotaomicron* causes altered *cps* gene expression.

Wild-type *B. thetaiotaomicron* was infected with live or heat-killed ARB25 at an MOI of ~1. (A)

Growth was monitored by measuring OD_{600} every 15-30 minutes. (B) *cps* gene transcript

analysis was carried out by qPCR. The end of the growth curve in panel A represents the point at

which cultures were harvested for qPCR analysis (e.g. the first observed time point where culture

surpassed OD_{600} of 0.6). Significant changes in *cps1*, *cps3*, and *cps4* expression were observed

between groups treated with live or heat-killed ARB25 ($p < 0.01$ for each, determined by

Dirichlet regression).

Figure 5. Infection of acapsular *B. thetaiotaomicron* selects for the expression of multiple phase-

variable loci. (A) Wild-type *B. thetaiotaomicron* was infected with ARB25 or was alternatively

exposed to heat-killed ARB25 and cultures were grown to OD_{600}=0.6-0.7. Cells were harvested

and RNA-seq analysis was carried out as described in *Methods* (n=3 independent experiments

for each treatment group). Transcript abundance was compared between treatment groups and

fold change between ARB25-treated cells versus cells treated with heat killed ARB25 was

calculated and is plotted against the P value (T test with Benjamini-Hochberg correction) for

each fold change calculation. (B) Acapsular *B. thetaiotaomicron* was treated with ARB25 or

heat-killed ARB25 and fold change in transcript abundance was calculated, as described in panel

A. (C) Among the genes with increased expression in post-infected acapsular *B.*

thetaiotaomicron, 24 genes were part of 8 different gene clusters predicted to encode tyrosine

recombinases along with outer membrane lipoproteins and OmpA-like proteins. These gene

clusters are shown. The numbers inside the gene schematics represent the fold change in

expression in ARB25-treated cells relative to those treated with heat-killed ARB25. Flanking the

promoters of each of these loci are pairs of imperfect, 17 nucleotide palindromic repeats. qPCR

analysis and amplicon sequencing of each orientation of these 8 promoters revealed expected

increases in orientation to the "ON" position in ARB25-exposed acapsular *B. thetaiotaomicron*.

(D) Another chromosomal locus with signatures of phage-selected recombination was identified

by RNA-seq. Specifically, 3 of 4 genes in an operon (BT1042-BT1045) were significantly down-

regulated after exposure to phage and 5 genes in an adjacent operon (BT1046-BT1051) were up-

regulated an average of 8.3-fold. (E) PCR using oligonucleotides flanking direct repeats within

the BT1032-BT1053 locus (green dumbbells, panel D) were used to demonstrate locus

architecture in wild type *B. thetaiotaomicron* and in a mutant lacking the tyrosine recombinase

within this locus (*B. thetaiotaomicron* ΔBT1041).

Figure 6. S-layer differentially affects *B. thetaiotaomicron* susceptibility to ARB25 and the

related phage SJC01. Acapsular *B. thetaiotaomicron* S-layer 'ON' and 'OFF' mutants (Δ*cps*

BT1927-ON and Δ*cps* BT1927-OFF, respectively) were infected with (A) ARB25 or (B) SJC01

in liquid culture. Growth was monitored via optical density at 600 nm (OD_{600}) on an automated

plate reading instrument as described in *Methods*.

Table 1. Susceptibility of strains to infection by ARB25 after infection and passaging.

Figure S1. Representative pictures of phage plaques for all phages from this study: A) phages from Ann Arbor (ARB); B) phages from San Jose (SJC). The top row of images for each phage are unaltered; background and saturated pixels were removed from images in the bottom row to facilitate viewing of the plaques. Scale bar = 2 mm

Figure S2. Replication of subset of host range assays of *B. thetaiotaomicron*-targeting phages on strains expressing different CPS types. Ten bacteriophages isolated and purified on the wild-type, acapsular, or the 8 single CPS-expressing strains were re-tested in a spot titer assay to determine phage host range. 10-fold serial dilutions of each phage ranging from approximately 10^6 to 10^3 plaque-forming units (PFU) / ml were spotted onto top agar plates containing the 10 bacterial strains. Plates were then grown overnight, and phage titers were calculated. Titers are normalized to the titer on the preferred host strain for each replicate. Each row in the heatmap corresponds to a replicate for an individual phage, whereas each column corresponds to one of the 10 host strains. One to three replicates of the assay were conducted for each phage by the two lead authors (AJH and NTP). Assays were carried out at the same time, and each author used the same set of cultures and phage stocks. For comparison, individual replicates from **Figure 1** are included (marked with *)

Figure S3. A) Phage titers in infected cultures incubated with purified capsules. ARB25 was incubated with purified CPS1 or CPS2 (1 mg/ml) before plating on the acapsular strain, and plaques were counted after overnight incubation. Titers are normalized to mock (H_2O) treatment.

No significant differences in titers were found compared to mock treatment, as determined by

Welch's t test. B) Wild-type *B. thetaiotaomicron* was infected with live or heat-killed ARB25,

and bacterial growth was monitored via optical density at 600 nm (OD600) on an automated

plate reader for 12 hours. At 0, 6.02, 8.36, and 11.7 hours post inoculation, replicate cultures

were removed and phage levels were titered. No phage were detected in heat-killed controls.

Figure S4. Infection of wild-type *B. thetaiotaomicron* at a low multiplicity of infection and

subsequent effects on *cps* ene expression. (A) The wild-type (WT) strain was infected at a low

multiplicity of infection (MOI $= 1$ x 10^{-4}) of live or heat-killed ARB25, and bacterial growth was

monitored via OD_{600}. (B) RNA was harvested from cultures after reaching an OD of 0.6-0.7,

cDNA was generated, and relative expression of the 8 *cps* loci was determined by qPCR.

Figure S5. Single replicates of *cps* expression in heat-killed versus live phage-treated *B.*

thetaiotaomicron. Relative *cps* CPS transcript abundance in ARB25 infection experiments at

high MOI (A) and low MOI (B). In the high MOI experiment, replicate 2 showed higher starting

expression of the non-permissive CPS3 compared to others. In the low MOI experiment,

replicate 3 showed higher starting expression of the non-permissive CPS3. In both experiments,

post phage-exposed replicates displayed nearly identical CPS expression profiles characterized

by high expression of CPS3.

Figure S6. Determination of phase-variable promoter switching for six loc encoding putative S-

layer proteins. The hypothesis that the promoters associated with seven newly identified *B.*

thetaiotaomicron S-layer like lipoproteins was validated using a PCR amplicon sequencing

strategy. Because of high nucleotide identity in both the regions flanking the 7 new loci, a nested

PCR approach was required to specifically amplify and sequence each site. In the first step, a

primer lying in each S-layer gene (**Table S3** "S-layer gene" primers) was oriented towards the

promoter and used in a PCR extension to a primer in the upstream recombinase gene (**Table S3**

"recombinase gene 3" primer). The products of this PCR were purified without gel extraction

and used in a second reaction with a nested primer that lies internal to the previous recombinase

gene primer (**Table S3** "recombinase 2" primer). The expected PCR products from this reaction,

which are ~1 kb and span promoter sequences in both the ON and OFF orientations, were

excised and used for an orientation-specific PCR using the original S-layer gene primer for each

site and a universal primer (green schematic) that was designed for each promoter and is oriented

to extend upstream of the S-layer gene (e.g., OFF orientation). Resulting products from this third

reaction, which should correspond to the ON orientation if a promoter inversion has occurred in

some cells, were obtained for 5/7 of the newly identified loci and the BT1927 S-layer locus as a

control. In all cases in which an amplicon and sequence were obtained, the expected

recombination occurred between the inverted repeat site proximal to the S-layer gene start (new

DNA junction), which would orient the promoter to enable expression of the downstream S-layer

gene. The sequences shown are the consensus between forward and reverse reads for each

amplicon. The putative core promoter -7 sequence is shown in bold/red text, the coding region of

each S-layer gene is shown in bold/blue text and the S-layer gene proximal recombination site is

noted and highlighted in bold/gold text. Note that the 5'-end of the sequenced amplicon was not

resolved for the BT2486 locus.

Figure S7. Recombination between the genes BT1040, BT1042, and BT1046. (A) Pfam domain

schematics of the amino acid sequences of these three genes highlighting that BT1040 and

BT1046, as originally assembled in the *B. thetaiotaomicron* genome sequence, lack additional N-

terminal sequences that are present on BT1042. (B) Sequencing of the 8 PCR amplicons

schematized in **Figure 5C**. Amplicons 1, 5 and 8 represent the original genome architecture,

while the others represent inferred recombination events that are validated here by sequencing.

The 5' and 3' ends of the BT1042, BT1040 and BT1046 genes are color-coded to assist in

following their connectivity changes after recombination. A series of single-nucleotide

polymorphisms (SNPs) present in BT1042, downstream of the proposed recombination site, are

highlighted in yellow. The transfer of these SNPs to a fragment containing the 5' end of BT1040

(Amplicon 4) was used to narrow the recombination region to the 7 nucleotide sequence

highlighted in red. Additional SNPs that are specific to the regions upstream of this

recombination site are shown in white text for each sequence.

Figure S8. The BT1033-52 locus does not affect susceptibility of acapsular *B. thetaiotaomicron*

to ARB25. Ten-fold serial dilutions of ARB25 were spotted onto lawns of *B. thetaiotaomicron*

Δ*cps* (n=5) and *B. thetaiotaomicron* Δ*cps* ΔBT1033-52 (n=5, n=3 independent clones each).

Plaquing efficiency was determined by normalizing plaque counts on *B. thetaiotaomicron* Δ*cps*

ΔBT1033-52 relative to plaque counts on *B. thetaiotaomicron* Δcps for each replicate. Statistical

significance was determined using the Mann-Whitney test.

1002

1003 **Table S1.** Phages used in this study and details on their isolation.

1004

1005 **Table S2.** Genes that are differentially regulated in post ARB25-infected wild-type and

1006 acapsular *B. thetaiotaomicron*.

1007

1008 **Table S3.** Bacterial strains and plasmids used in this study.

1009

1010 **Table S4.** Primers used in this study.

1011

Figure 1

Figure 2

Figure 3

Time (in hours)

Figure 4

Figure 5

Figure 6

Table 1. Bacterial susceptibility to secondary ARB25 infection

Strain	# susceptible isolates after first infection	# susceptible isolates after second infection
Wild-type	9/9 (100%)	9/17 (53%)
cps1	9/9 (100%)	15/17 (88%)
cps2	0/9 (0%)	0/18 (0%)
cps4	2/9 (22%)	8/11 (73%)
Acapsular	9/9 (100%)	10/16 (63%)
Total of phage-susceptible strains[a]	**29/36 (81%)**	**42/61 (69%)**

[a]Wild-type, cps1, cps4, and acapsular