Population persistence under high mutation rate: from evolutionary rescue to lethal mutagenesis

Keywords : Evolutionary rescue, Fisher's Geometric Model, Fitness landscape, Lethal mutagenesis, Extinction dynamics, Distribution of extinction times

Abstract

Populations may genetically adapt to severe stress that would otherwise cause

their extirpation. Recent theoretical work, combining stochastic demography with

Fisher's geometric model of adaptation, has shown how evolutionary rescue becomes

unlikely beyond some critical intensity of stress. Increasing mutation rates may however

allow adaptation to more intense stress, raising concerns about the effectiveness of

treatments against pathogens. This previous work assumes that populations are rescued

by the rise of a single resistance mutation. However, even in asexual organisms, rescue

can also stem from the accumulation of multiple mutations in a single genome. Here, we

extend previous work to study the rescue process in an asexual population where the

mutation rate is sufficiently high so that such events may be common. We predict both

the ultimate extinction probability of the population and the distribution of extinction

times. We compare the accuracy of different approximations covering a large range of

mutation rates. Moderate increase in mutation rates favors evolutionary rescue.

However, larger increase leads to extinction by the accumulation of a large mutation

load, a process called lethal mutagenesis. We discuss how these results could help

design "evolution-proof" anti-pathogen treatments that even highly mutable strains

could not overcome.

Introduction

Evolutionary rescue (ER) happens when a population confronted with severe

stress avoids extinction by genetic adaptation. Understanding and predicting when and

how evolutionary rescue occurs is critical in fields as diverse as conservation biology,

invasion biology, emergence of new diseases and the management of resistance to

treatment in pests and pathogens (see reviews in Gonzalez *et al.* 2013; Carlson *et al.*

2014; Alexander *et al.* 2014; Bell 2017). In all these situations, genetic variation, be it
present before the onset of stress, or generated *de novo* after, is a key ingredient for
evolutionary rescue, as expected theoretically (e.g. Gomulkiewicz and Holt 1995) and
observed experimentally (e.g Ramsayer *et al.* 2013). Because mutation affects both
standing and *de novo* genetic variation, it comes as no surprise that a number of
evolutionary rescue models, combining stochastic evolution and demography, have
predicted that higher mutation rates are associated with higher probability of
evolutionary rescue (Orr and Unckless 2008, 2014; Martin et al. 2013; Anciaux et al.
2018). Few evolutionary rescue experiments have manipulated the mutation rate to test
these predictions (reviewed in Bell 2017). For instance, Couce *et al.* (2015) found that two
different mutator strains of bacteria with elevated rates of mutations evolved more than
100-fold resistance to antibiotic concentrations that caused the demise of control strains.
Mutator alleles are indeed often found in antibiotic resistant strains causing serious
health issues (Eliopoulos and Blázquez 2003), raising concern about pathogens escaping
our control by evolving higher mutation rates (for theoretical predictions see Taddei *et*
al. 1997; Greenspoon and Mideo 2017).

Most mathematical models of evolutionary rescue assume that the population is
rescued from extinction by the spread of a single mutant of large effect (Orr and
Unckless 2008, 2014; Martin et al. 2013; Anciaux et al. 2018) and do not describe highly
polymorphic populations where several mutations of smaller effects can combine to
allow population growth (see however the work of Uecker and Hermisson (2016) and
Uecker (2017) where sexual reproduction allows production of such rescue genotypes).
The latter situation seems in particular to be common in the evolution of herbicide
resistance, especially when the mutational target for resistance is large (Kreiner et al.
2018). Even in asexual organisms, when the mutation rate is high, evolutionary rescue
may commonly result from the cumulative effect of multiple mutations accumulating
stochastically over time in a given lineage. Such a mutation regime is particularly

relevant in highly mutable viruses, mutator strains of bacterial (e.g. Springman *et al.*

2010) or cancer cells (e.g. Loeb 2001). Our aim here is to provide theoretical predictions

for evolutionary rescue in such a regime with high mutation rates in asexual organisms,

complementing existing theory on the subject.

Several complications arise when modelling evolutionary rescue in highly

polymorphic populations with high mutation rates. First, the dynamics of allelic

frequencies at different loci interact in asexuals. For example, the selective sweep of a

given beneficial mutation is hindered by the co-segregation of other beneficial

mutations (clonal interference, Gerrish and Lenski 1998). A theoretical study by Wilson *et*

al. (2017) recently showed that, when evolutionary rescue is likely, it should most often

be driven by soft selective sweeps, where multiple resistance mutations spread through

the population simultaneously. Wilson *et al.* (2017) still assumed that each of these

lineages carried a single mutation, each with the same effect on the population growth

rate. When the mutational target is large, different lineages contributing to rescue are

however likely to carry mutations with different fitness effects. Modelling the distribution

of mutation effects (as in Martin *et al.* 2013; Anciaux *et al.* 2018) then becomes critical.

Finally, when the mutation rate is high, multiple mutations may also accumulate in each

lineage, either facilitating evolutionary rescue or impeding it, through their *cumulative*

effect. Modelling both beneficial and deleterious mutations and, critically, the epistatic

interactions between them, also becomes necessary.

Previous evolutionary rescue theory predicts that higher mutation rate allows

populations to withstand higher levels of stress (e.g. Anciaux *et al.* 2018). Yet, there are

reasons to expect this prediction not to hold above some critical mutation rate:

increased mutation rates also build-up detrimental mutation loads, thus depressing

mean fitness despite ongoing adaptation. Indeed, some previous ER models, including

both beneficial and deleterious additive effects on growth rates, have found that ER was

most likely at intermediate mutation rates (Loverdo and Lloyd-Smith 2013; Greenspoon

and Mideo 2017). Artificially increasing the mutation rate has even been proposed as a

means to weaken or even eliminate pathogen populations, by a process called lethal

mutagenesis (Loeb et al. 1999). Models of lethal mutagenesis predict that extinction of

the target population could be observed under biologically realistic sets of parameters

(Bull et al. 2007; Martin and Gandon 2010; Wylie and Shakhnovich 2012). In these

models, mean fitness dynamics and extinction stem from the deterministic effects of

selection and mutation. Alternatively, Matuszewski *et al.* (2017) discuss the continuity

between these models and models of mutational meltdown, where extinction is driven

by the interaction of genetic drift and deleterious mutation. Lethal mutagenesis has been

investigated empirically for treatment against viruses (Springman et al. 2010; Arias et al.

2014), bacteria (Bull and Wilke 2008) or cancer cells (Liu et al. 2015). In particular, the

combination of antiviral treatments with mutagenic agents is investigated as a strategy

to fight fast evolving viruses, such as influenza (Bank et al. 2016). It seems important to

improve our ability to predict whether and when such mutagenic agents will increase

treatment efficacy or, conversely, facilitate the evolution of resistance.

The population genetics of adaptation behind the rescue process, in isolated

asexual populations, roughly fall into two alternative regimes: rescue may stem (i) from

single mutations of large effect (strong selection weak mutation 'SSWM' regime) or (ii)

from multiple mutations of small effects (weak selection strong mutation 'WSSM'

regime) (reviewed in Alexander *et al.* 2014). The dichotomy between SSWM and WSSM

entails a somewhat simplistic view of adaptation regimes, at the two extremes of all

possible mutation rates. The SSWM regime of adaptation has been extensively

investigated in population genetics via "origin-fixation" models describing the average

behavior of stochastic evolutionary dynamics (McCandlish and Stoltzfus 2014) whereas

the WSSM regime has been widely analyzed via deterministic models of quantitative

genetics (Lande 1976, 1980). Corresponding evolutionary rescue models further include

a coupling of adaptation and demographic dynamics, and naturally fall into the same
two regimes (discussed in Anciaux *et al.* 2018). The SSWM regime of evolutionary rescue
is characterized by the fact that the first resistant lineage to establish (and thus cause
rescue) is only one mutational step from the predominant, sensitive 'wild-type' lineage
(e.g. Feder *et al.* 2016). Models that describe highly polymorphic dynamics (WSSM
regimes, e.g. the quantitative genetic model in Gomulkiewicz and Holt 1995) often use
the infinitesimal model assumptions (many unlinked polymorphic loci), which does not
apply to asexual populations. In the WSSM regime, the exact stochastic evolutionary
dynamics become quickly intractable, and have often been studied by simulation (e.g.
Boulding and Hay 2001). Further the latter models often consider initial standing genetic
variance as given and pay little attention to the effect of mutation rates in maintaining
this variance. They often ignore *de novo* mutations after the onset of stress, on the
argument of short timescales being most critical for evolutionary rescue (e.g.
Gomulkiewicz et al. 2010).

To make analytical progress in our understanding of the effect of mutation rates
on the process of evolutionary rescue, we build on two recent theoretical developments
(Martin and Roques 2016; Anciaux et al. 2018). Anciaux *et al.* (2018) developed a model
of evolutionary rescue in the SSWM regime using Fisher's (1930) geometric model
(hereafter "FGM") to model the distribution of mutation effects on fitness. As in Anciaux
et al. (2018), we study evolutionary rescue using the FGM. This model assumes a single
peak phenotype-fitness landscape, where fitness depends on the position, in phenotype
space, of a given genotype relative to an optimum. In the context of ER, stress may affect
this landscape in various ways (height, width or position of the peak). In this model, the
distribution of mutation effects (both beneficial and deleterious) depend on the context,
both genotypic (epistasis) and environmental (e.g. effect of stress). This context-
dependence is a key feature of the FGM; it is absent from previous ER models studying
the effect of high mutation rates, because they assume additive mutation effects on

fitness (e.g. Loverdo and Lloyd-Smith 2013; Greenspoon and Mideo 2017). The variation

in the distribution of mutation effects implied by the FGM is qualitatively, and sometimes

even quantitatively, consistent with a wealth of empirical observations (reviewed in

Tenaillon 2014). Under the assumptions of the FGM, rescue mutants become very rare as

the intensity of stress increases, because they require very large mutational steps. As a

consequence, Anciaux *et al.* (2018) predict that there is a narrow window of stress levels

where the probability of rescue shifts from being very likely to very unlikely. They also

predict that this critical level of stress, beyond which adaptation is unlikely, is increased

by higher mutation rates (Anciaux et al. 2018). Yet, predictions of this model apply to the

SSWM regime and may not hold for higher mutation rates.

We extend our previous analysis of evolutionary rescue over Fisher's geometric

model (Anciaux et al. 2018) to the more complex and more polymorphic WSSM regime.

To do so, we use the approach in Martin and Roques (2016) to model the non-

equilibrium dynamics of fitness distributions, in large asexual populations. The form of

fitness epistasis assumed may have particular impact on the results because lineages

accumulate multiple mutations at different sites over the rescue process. We use the

FGM here, which implies a particular form of epistasis that has proven consistent with

several observed patterns in fitness epistasis among mutations (Martin et al. 2007;

Perfeito et al. 2014; Blanquart and Bataillon 2016). Moreover, Martin and Roques (2016)

showed that, under the FGM, while the fitness dynamics are more complex at higher

mutation rates, they are also more predictable and less prone to stochastic fluctuations,

even in relatively small populations. To model evolutionary rescue, we still need to

describe the demographic stochasticity associated with the extinction process. In the

WSSM regime, we thus use a combination of two analytically tractable theories: a

deterministic approximation to the dynamics of mean fitness (Martin and Roques 2016)

and a diffusion approximation to the stochastic dynamics of population sizes (from

Bansaye and Simatos 2015).

Beyond a derivation of the probability of ultimate rescue or extinction, this
approach further allows tracking the rescue process over time. As stated in Gomulkiewicz
et al. (2017), tracking of transient dynamics (population size dynamics, distributions of
extinction times) are of high interest for applications of evolutionary rescue theory, yet
are not available from existing predictions, which focused mainly on ultimate outcomes.
Gomulkiewicz *et al.* (2017) studied the distribution of extinction times for populations
doomed to extinction, mostly in the absence of mutation (i.e. with a fixed arbitrary set of
competing asexual genotypes at the onset of stress). We extend this analysis to include
frequent *de novo* mutation, rescue events involving several mutational steps, a particular
form of epistasis and variable mutation effects depending on stress intensity, and an
explicit description of the dynamics of mutation load. Our approach captures the
continuum from evolutionary rescue to lethal mutagenesis, as mutation rate increases.
Interestingly, some parameter ranges prove to greatly limit evolutionary rescue at all
mutation rates, i.e. in spite of the possible mutator genotypes.

Methods

I. General framework

The present work focuses on an asexual population with independent lineages
(e.g. a population of asexual microbes without horizontal gene transfer), facing an
abrupt and stressful environmental change (e.g. an antimicrobial treatment). The
population initially consists of N_0 individuals with either a single or multiple genotypes
and is initially adapted to a non-stressful environment, where its mean growth rate is
positive. At the onset of stress, the population is shifted from the non-stressful
environment to a stressful environment, where its mean growth rate becomes negative
(definition of stress here). In such an environment, in the absence of evolution, the

population is doomed to extinction. Evolutionary rescue occurs if at least one resistant

lineage (with a positive growth rate in the new environment) establishes, in spite of

demographic stochasticity. These resistant mutant lineages can either already be present

in the population or arise *de novo* after the onset of stress. It is thus crucial to determine

how the number and growth rates of such mutants depend on the new environmental

conditions and on the parental genotypes already present in the population. We do so

using the FGM detailed below. Note that the main notations used here are summarized

in Table 1.

Fitness landscape

In the FGM, a given phenotype is a vector in a phenotypic space of n dimensions

that determine fitness (here the growth rate r). The phenotype of an individual with

genotype i, is characterized by a vector $\mathbf{z}_i \in \mathbb{R}^n$ of the breeding values (heritable

components) for the n traits, and its growth rate is r_i. In a given environment, fitness

decays as a quadratic function of the phenotypic distance to a single phenotypic

optimum, where the growth rate r_{max} is maximal at a given absolute level (height of the

fitness peak). We assume that each environment is associated with a single optimum and

fitness peak. In the scenario investigated here, in the non-stressful environment, the

population is close to the 'ancestral' optimum $\mathbf{z}_A$. When the environment changes, it is

assumed to determine a new optimum $\mathbf{z}_*$. Without loss of generality, the height of the

peak may also differ between the ancestral and new environments. However, we do

require that the n dimensions that determine fitness remain the same (in nature and

number) across environments. In the new environment, the growth rate of an individual

with genotype i is given by:

$$r(\mathbf{z}_i, \mathbf{z}_\star) = r_{max} - \frac{\|\mathbf{z}_i - \mathbf{z}_*\|^2}{2}. \qquad [1]$$

This is an isotropic version of the FGM (all directions are equivalent for selection and
mutation) where phenotypes are scaled by selective strength.

Stochastic demographic dynamics

We confine our analysis to finite haploid asexual populations. Individuals have
independent evolutionary and demographic fate (frequency or density dependence are
ignored). Each genotype i has a growth rate r_i and a reproductive variance σ_i ($r_i =$
$r(\mathbf{z_i}, \mathbf{z_*})$ in a given environment with optimum $\mathbf{z}_*$), which define its stochastic
demographic parameters in the context of a Feller diffusion approximation (Feller 1951),
as in e.g. Martin et al. (2013), Gomulkiewicz *et al.* (2017) or Anciaux *et al.* (2018). For
simplicity, we further assume here that the average stochastic variance in reproduction is
constant over time: $\bar{\sigma}(t) = 1/K_t \sum_{i=1}^{K_t} \sigma_i = \bar{\sigma}$, where the average is taken over the K_t
genotypes present at time t. This can for example be accurate whenever the σ_i are
roughly constant across genotypes i (discussed in Martin *et al.* 2013 and Anciaux *et al.*
2018).

Notation	Description	Formula
N_t, N_0	N_t: population size at time t after the onset of the stress. N_0: initial population size at the onset of the stress.	
U	Mutation rate *per* individual *per* unit time.	
n	Number of traits under stabilizing selection, or phenotypic dimensionality.	
λ	Variance of mutational effects: variance of the phenotypic effects of mutations, per trait, in a trait space scaled by the strength of stabilizing selection	$\mathbf{dz} \sim N(\mathbf{0}, \lambda \, \mathbf{I}_n)$
μ	Composite parameter : square root of mutational variance per trait	$\mu = \sqrt{U\lambda}$

$\mathbf{z}$	n-dimensional vector $\mathbf{z} \in \mathbb{R}^n$ of (breeding values for) phenotype	
$\mathbf{z}_A, \mathbf{z}_*$	$\mathbf{z}_*$: optimal phenotype in the new environment $\mathbf{z}_A$: average phenotype of the ancestral population (before the onset of stress)	
r, σ	Growth rate (r) and reproductive variance (σ) of a given genotype, in the new environment.	Eq.[1]
$\overline{r}_t, \overline{\sigma}$	Mean growth rate ($\overline{r}_t$) and mean reproductive variance ($\overline{\sigma}$) of the population, in the new environment.	Eqs. [2] and [3]
$\langle \overline{r}_t \rangle$	'ensemble mean' growth rate : expected mean growth rate $\overline{r}_t$ across stochastic replicates	Eqs. [2] and [3]
r_{max}	Maximum possible growth rate in the new environment.	$r(\mathbf{z}_*) = r_{max}$
r_D	Rate of decay of the ancestral phenotype $\mathbf{z}_A$ in the new environment.	$r(\mathbf{z}_A) = r_{max} - \|\mathbf{z}_A\|^2/2 = -r_D$
y_D	Rate of decay scaled by r_{max}.	$y_D = r_D/r_{max}$
ϵ	Ratio between the mutational load and the maximal growth rate	$\epsilon = n\,\mu/(2\,r_{max})$
$\omega_{DN}, \omega_{DN+SV}$	ω_{DN}: rate of rescue from *de novo* mutations scaled by N_0 ω_{DN+SV}: rate of rescue from both *de novo* and standing variance scaled by N_0	Eqs.[5],[6]
P_{ext}, P_R	P_{ext} : Extinction probability P_R : Rescue probability	P_{ext} see Eq.[4] $P_R = 1 - P_{ext} = 1 - \exp\left(-N_0\,\omega_R\right)$
U_c	Critical mutation rate beyond which the WSSM regime is valid.	$U_c = n^2\,\lambda/4$
U_{max}	Mutation rate beyond which certain extinction is enforced by lethal mutagenesis	$U_{max} = 4\,r_{max}^2/(n^2\,\lambda)$
U_*	Mutation rate at which the rescue probability rises to 1/2	Eq.[7]
$r_D^*(p)$	Rate of decay for which the maximal rescue probability for any U or λ is equal to p	Eq.[8]

230   ***Table 1***: *Notations*

II. Evolutionary dynamics.

In this section, we describe the model of evolutionary dynamics over the fitness

landscape (FGM of the previous section), which is embedded into the ER model. In the

following, *de novo* mutations (appearing after the onset of stress) are denoted "DN" and

mutations from standing genetic variance (mutants already present before the onset of

stress) are denoted "SV". Correspondingly, evolutionary rescue dynamics from an
isogenic population, adapting only from *de novo* mutations, are labelled "DN" and
dynamics from a polymorphic population, adapting from both *de novo* mutations and
standing genetic variance, are labelled "DN + SV".

1. Evolutionary dynamics from an isogenic population.

The population is maladapted in the new stressful environment and its growth
rate is $-r_D$, corresponding to a decay rate $r_D > 0$. Mutations arise following a Poisson
process with rate U per unit time per capita. For a given parent phenotype, each
mutation creates a random perturbation $\boldsymbol{dz}$ on phenotype, which is unbiased and
follows an isotropic multivariate Gaussian distribution, $\boldsymbol{dz} \sim \mathcal{N}(0, \lambda \, \mathbf{I}_n)$ where $\mathbf{I}_n$ is the
identity matrix in n dimensions and λ is the variance of mutational effects on traits,
standardized by the strength of selection. Mutation effects are additive on phenotype,
but not on fitness because $r(.)$ is nonlinear (epistasis on fitness and not on phenotype).

In the WSSM regime, the mean growth rate of the population shows limited
stochastic variation among replicates, even in reasonably small populations. We thus
approximate the evolutionary process by a deterministic fitness trajectory, derived in the
WSSM regime under the FGM (Martin and Roques 2016). This seemingly rough
approximation can be justified *a priori*: most of the ER process is determined by the
speed at which the population adapts at the very onset of stress. This early trajectory
takes place when the population is still large and the adaptive process proves to be
relatively deterministic, especially over this short timescale, provided that the mutation
rate is high enough compared to the mean fitness effect of random mutations (WSSM:
$U \gg U_c = n^2 \lambda / 4$, with $n \, \lambda / 2$ the mean fitness effect of random mutations). Both
analytical arguments and simulations detailed in Martin and Roques 2016, showed that

mean fitness trajectories are indeed close to the deterministic prediction (with limited

variation among replicates), provided that $U \gg U_c$ (WSSM) and $NU \gg 1$ (large mutational

input). Here, we use the deterministic fitness trajectory corresponding to these

conditions to approximate the growth rate trajectory of all replicate populations under

stress ($\bar{r}_t \approx \langle \bar{r}_t \rangle$, with $\langle . \rangle$ the expectation over replicates).

Provided $U \gg U_c$ we thus approximate the trajectory of the mean growth rate of all

replicate populations by its deterministic trajectory for the WSSM (Martin and Roques

2016):

$$\bar{r}_t^{DN} \approx \langle \bar{r}_t^{DN} \rangle \approx r_{max} - \underbrace{(r_D + r_{max})(\operatorname{sech}(\mu t))^2}_{\textbf{convergence to the optimum}} - \overbrace{\frac{n}{2} \mu \tanh(\mu t)}^{\textbf{Mutation load build} - \textbf{up}} , \quad [2]$$

where $\operatorname{sech}(z) = 2/(\exp(z) + \exp(-z))$ is the hyperbolic secant, $\tanh(z) = (\exp(z) -$

$\exp(-z))/(\exp(z) + \exp(-z))$ is the hyperbolic tangent and $\mu = \sqrt{U\lambda}$ is a composite

parameter of the mutational parameters. In Eq.[2] the first term stems from the

population nearing the phenotypic optimum, while the second term stems from the

build-up of the mutation load as the genetic variance accumulates in the initially clonal

population. Recall that r_D is the decay rate of the isogenic population and r_{max} is the

maximum fitness that can be reached in the new environment (with $r_D + r_{max}$ the fitness

distance between the parent genotype's fitness and the top of the fitness peak). The

mean growth rate in Eq.[2] reaches a plateau at infinite time of $r_{max} - n\, \mu/2$

corresponding to the maximal growth rate minus the mutational load.

2. Evolutionary dynamics from an initially polymorphic population (at mutation-selection

balance)

The evolutionary dynamics of rescue from a polymorphic population is obtained

by a similar approximation. We assume that the population is initially at mutation-

selection balance in the non-stressful environment, with an arbitrary positive mean

growth rate. The phenotypic distribution at the onset of stress is centered on a mean

phenotype $\bar{z}$, whose growth rate in the new environment is used to characterize the

harshness of the stress imposed. For consistency with the isogenic population model

above, we thus denote this growth rate $- r_D = r(\bar{z}, z_*)$, where $r_D > 0$ is the decay rate of

the central genotype as in the previous section. Resistant genotypes may already be

present in the population at the onset of stress ("SV") or appear by *de novo* mutation

("DN"), or arise as combinations of these (multiple step rescue mutants). Using the same

reasoning as in the previous subsection, we approximate the mean growth rate of all

replicate populations by the deterministic trajectory for the WSSM, i.e. whenever $U \gg U_c$

(Martin and Roques 2016):

$$\bar{r}_t^{DN+SV} \approx \langle \bar{r}_t^{DN+SV} \rangle \approx r_{max} - \underbrace{\exp{(-2\,\mu\,t)}(r_D + r_{max})}_{\textbf{convergence to the optimum}} - \overbrace{\frac{\widetilde{n}\,\mu}{2}}^{\textbf{Mutation load}}. \qquad [3]$$

In a polymorphic population at mutation-selection balance, the presence of a mutational

load implies that the mean growth rate of the population in Eq.[3] is lower than the

mean growth rate of an isogenic population in the same environmental conditions, with

the same central genotype ($\bar{r}_0^{DN+SV} = -r_D - n\,\mu/2$). Contrary to Eq.[2], the mutation load

is stable (the phenotypic variance is already at equilibrium) while the convergence to the

optimum is faster due to the presence of standing variance. The mean growth rate in

Eq.[3] ultimately reaches the same plateau as Eq. [2] (the maximal growth rate minus the

mutational load).

Note that in both Eqs.[2] and [3] the two mutational parameters U and λ play entirely

symmetric roles through the composite parameter $\mu = \sqrt{U\,\lambda}$. This is because the WSSM

regime amounts to a diffusive approximation of the effect of mutation in fitness space

(Appendix E.III of Martin and Roques 2016). Like any diffusive approximation, only the

mean and variance of the mutation kernel contribute to the dynamics: in this case $U\lambda$ is

the variance induced by mutation during one time unit.

III. Evolutionary rescue probability

As stated in introduction, we neglect evolutionary stochasticity (i.e. the variance

over replicates of the mean fitness dynamic, induced by drift and mutation) but not the

demographic stochasticity. The latter is approximated by an inhomogeneous Feller

diffusion (see Appendix I section I) with parameters $\bar{r}_t$ (the expected mean growth rate)

and $\bar{\sigma}$ (the reproductive variance averaged across segregating genotypes, which we

assume is stable over time): $\bar{r}_t$ may vary stochastically as genotype frequencies change

under the effects of drift, selection and mutation. However, as explained in the previous

subsection, we approximate each replicate's fitness trajectory by its deterministic

expectation under the WSSM regime: $\bar{r}_t \approx \langle \bar{r}_t^{DN} \rangle$ or $\langle \bar{r}_t^{DN+SV} \rangle$, using the relevant cases

from Eqs.[2] or [3] (the validity of this approximation is tested against simulations in the

Results section). Therefore, the model approximately reduces to a Feller diffusion with

constant $\bar{\sigma}$ and time-inhomogeneous deterministic growth rate $\bar{r}_t$. Under these

hypotheses, the demographic dynamics follow the stochastic differential equation:

$dN_t = \bar{r}_t\,N_t\,dt + \sqrt{\bar{\sigma}\,N_t}\,dB_t$ where B_t is a Weiner process (see Appendix I section I for

more details). We can then use the results from Theorem 1 of Appendix II (see also

Bansaye and Simatos 2015) on inhomogeneous Feller diffusions to derive the probability

that the population is extinct before time t:

$$P_{\text{ext}}(t) = \exp\left(-\frac{2\,N_0}{\bar{\sigma}\,\int_0^t \exp\left(-\rho_u\right) du}\right),$$
$$\rho_u = \int_0^u \bar{r}_v\,dv\,,$$

[4]

where $\bar{r}_t$ is given by Eqs.[2] or [3]. This expression can be evaluated numerically at any time $t > 0$. ER happens whenever evolution (change in $\bar{r}_t$) allows extinction to be avoided. Hence, the general form of the rescue probability is readily obtained as the complementary probability of the infinite time limit of Eq.[4], namely the probability of never getting extinct: $P_R = 1 - P_E$ with $P_E = P_{\text{ext}}(\infty)$, the extinction probability after infinite time. Depending on the scenarios, this rescue probability can be computed explicitly, or approximated via Laplace approximations to the integral $\int_0^\infty \exp\left(-\rho_u\right) du$, in some parameter ranges (see Results and Appendix I).

IV. Individual-based stochastic simulations: The analytical predictions are tested against exact stochastic simulations of the population size and genetic composition of populations across discrete, non-overlapping generations (**Supplementary Figures 8** shows examples of simulated dynamics of the mean fitness and the population size). Rescue probability was estimated by running 100 replicate simulations until either extinction or rescue occurred. A population was considered rescued when it reached a population size N_t and mean growth rate $\bar{r}_t$ such that its ultimate extinction probability, if it were monomorphic, would lie below 10^{-12} ($\exp(-2\,N_t\,\bar{r}_t) < 10^{-12}$). The simulation algorithm is described in Anciaux *et al.* (2018) and also detailed in Appendix I section VIII. Briefly, the number of offspring is Poisson distributed every generation with parameter e^{r_i} for genotype i, mutations occur according to a Poisson process with constant rate U per capita per generation. The phenotypic effects of the mutations are drawn from a multivariate normal distribution, with multiple mutations having additive

effects on phenotype. Fitnesses are computed according to the FGM using Eq.[1]. With

such a Poisson offspring distribution, the reproductive variance of genotype i is

$\sigma_i = 1 + r_i \approx 1$, assuming small growth rates $r_i \ll 1$, in per-generation time units. So this

particular demographic model satisfies our assumption of constant $\bar{\sigma}$ in spite of changes

in the genotypic composition of the population: here $\bar{\sigma} = 1/K_t \sum_{i=1}^{K_t} \sigma_i \approx 1$. Note also

that the analytical derivations (relying on a Feller diffusion (1951) approximation)

approximately cover other demographic models (e.g. birth death models, see Martin et

al. 2013; Gomulkiewicz et al. 2017), as long as they also satisfy $\bar{\sigma}(t) \approx \bar{\sigma}$ constant.

For *de novo* rescue, the initial population consisted of a single genotype optimal in the

environment before stress. When considering contribution from standing variance, the

initial population was at mutation selection balance in the former environment (before

the onset of stress). More precisely, 10 replicate initial equilibrium populations were

generated and 100 replicate simulations were run from each of these populations. The

overall rescue probability is then the average, over the 10 equilibrium populations, of the

rescue probabilities estimated by the proportion of rescued populations over the 100

simulations.

V. Appendices and supplementary information: Analytical derivations and

supplemental figures are described in Appendix I. Appendix II contains the derivations

for the inhomogeneous Feller diffusion. Supplementary file 1 provides the details of the

analytical derivations, the code producing the figures and the simulation code as a

Mathematica© .cdf file (MATHEMATICA v. 11.3 Wolfram Research 2018) which can be

open using the free "CDF player" available on the Wolfram website. Supplementary file 2

provides the Matlab© (MATLAB 2015a, The MathWorks, Natick, 2015) source code for

the curve fitting procedure used for Eq.[8].

Results

Rescue probability: general form. In spite of their difference in the population genetics underlying ER, the two scenarios with or without standing variance yield similar expressions for the probability of ER (as shown in Eqs.(A5) and (A11-A12) in Appendix I):

$$P_R = 1 - \exp(-N_0\,\omega) \quad \text{with } \omega = \frac{2\,\mu}{\int_0^\infty h(\tau)^{n/2} \exp(-n/(2\,\epsilon)\,f(\tau))\,d\tau}$$

$$\text{and } \begin{cases} DN: \quad f(\tau) = \tau - (1 + y_D)\tanh(\tau)\,, \quad h(\tau) = \cosh(\tau) \\ DN + SV: \quad f(\tau) = \tau - \frac{1}{2}(1 + y_D)(1 - \exp(-2\,\tau))\,, \quad h(\tau) = \exp(\tau) \end{cases}, \quad [5]$$

where the particular form of the functions $f(.)$ and $h(.)$ depend on the chosen scenario. Here, $\cosh(z) = (\exp(z) + \exp(-z))/2$ is the hyperbolic cosine and we introduce two scaled parameters: $\epsilon = n\,\mu/(2\,r_{max})$ and $y_D = r_D/r_{max}$ (recall that $\mu = \sqrt{U\lambda}$ depends on both mutation rate and effect). The parameter y_D describes how fast the initial clone decays, compared to how fast the optimal genotype grows, it gives a scaled measure of the harshness of the stress imposed (see also Anciaux *et al.* 2018). The parameter ϵ is the ratio of the mutation load (at mutation-selection balance) and the maximal absolute growth rate that can be reached in the stress. Certain extinction by lethal mutagenesis occurs whenever the load is equal to or larger than the maximal growth rate (i.e. $\epsilon \geq 1$). A small value of ϵ means that we are far from this certain extinction regime.

The quantity ω in Eq.[5] is akin to a 'per lineage rate of rescue'. Analytical expressions for this rate are either unavailable (DN) or complicated ($DN + SV$, Eq.(A12) in Appendix I).

Weak selection, intermediate mutation approximation: To get a more direct insight

into the impact of each parameter, we sought an approximate expression for ω, detailed

in Appendix I section III and IV. This approximation applies with an intermediate

mutation rate, where ER mainly depends on the early adaptation of the population to

stress, and not on the ultimate mutation load (lethal mutagenesis). More precisely, it

requires that the WSSM approximation be accurate while ϵ remains small, which implies

a small λ and intermediate U: $n^2 \lambda/4 \ll U \ll r_{max}$. This is why we call this range a 'weak

selection, intermediate mutation' approximation (WSIM in **Figure 1** and **2**).

Under this approximation, the *per capita* rate of rescue ω takes a roughly similar

form for both scenarios with or without standing variance (detailed in Eqs.(A8) through

(A14) in Appendix I):

$$\omega \underset{\epsilon \ll 1}{\approx} 2 \sqrt{\frac{r_{max}\, \mu}{\pi}} \exp\left(-\frac{r_{max}}{\mu}\, g(y_D)\right)$$
$$\text{with}\begin{cases} DN: \ g(y) = g_{DN}(y) = \sqrt{y(1+y)} - \cosh^{-1}(\sqrt{1+y}) \\ DN+SV: \ g(y) = g_{DN+SV}(y) = (y - \log(1+y))/2 \end{cases} \qquad [6]$$

In both cases, the function $g(.)$ is positive and increases (roughly log-log linearly)

with $y_D = r_D/r_{max}$. The accuracy of this approximation is illustrated in **Supplementary**

Figures 2 and **3** and **Figures 1** and **2**. Note that, whenever the approximation applies,

the ER probability is independent of dimensionality (n). This directly stems from the fact

that the growth rate trajectories in Eqs.[2] and [3] only depend on dimension via the

'mutation load' terms, whose contribution is negligible when far from the lethal

mutagenesis threshold ($U \ll r_{max}$).

Sharp decay in ER probability with increasing stress levels: A possible measure of

stress intensity in ER is the rate of decay r_D of a population after the environmental

change (see also Anciaux *et al.* 2018). However, stress might also affect other parameters

of the FGM: the height of the fitness peak r_{max}, the mutation rate U or the variance of

mutational effects: λ. We detail the two latter effects (which affect ER via the composite

parameter $\mu = \sqrt{U\,\lambda}$), in the next section, and focus here on r_{max} and r_D. In the following

we use the per *capita* rate of rescue ω_{DN} from Eq.(5) of Anciaux *et al.* (2018) in the SSWM

regime (recalled in Appendix I section VIII, Eqs.(A20)-(A21)), as a comparison to the

present results in the WSSM regime. By basic properties of the FGM described in Anciaux

et al. (2018) , increased r_D means both a faster decay (purely demographic effect of

stress) and a larger shift in optimum (which affects the whole distribution of fitness effect

of mutations), which both decrease the ER probability. On the contrary, increasing r_{max}

increases the ER probability through two effects. First, the size of the phenotypic space

of resistance increases with r_{max} (as in the SSWM regime, see Anciaux *et al.* 2018).

Second a large r_{max} counterbalances a high mutational load $n\,\mu/2$ as can be seen in

Eq.[2] (this latter effect is only captured in the WSSM approximation).

Figure 1*: ER probability against decay rate r_D for a population without standing variance*
(DN). Dots show the results of 10^2 simulations; thin plain lines: Eq.(5) from Anciaux et al.
2018 (recalled in Appendix Eq.(A21)) derived under the SSWM regime; thick plain lines:
Eq.[5] derived under the WSSM regime; dashed lines: the corresponding closed form
expression in Eq.[6] derived under the weak selection intermediate mutation regime
(WSIM). The shaded area corresponds to the extra contribution to ER from multiple
mutants, compared to single mutants. All models and simulations are shown for a high
mutation rate (blue) or a low mutation rate (brown), indicated in legend. As expected, the
model from Anciaux et al. (2018) derived in the SSWM regime captures the simulations at
low mutation rates ($U = 10^{-2}U_C$, brown), whereas Eqs.[5] and [6] captures the simulations
at high mutation rates ($U = 10^2 U_C$, blue). Other parameters are $N_0 = 10^5$, $n = 4$, $r_{max} = 1$
and $\lambda = 5 \times 10^{-3}$.

Figure 1 illustrates how the ER probability drops sharply with the decay rate $r_{D,}$

both in the SSWM regime (brown curves, $U \ll U_c$, see legend) and in the WSSM regime

(blue curves, $U \gg U_c$). This qualitatively similar behavior is *a priori* due to common

geometric constraints imposed by the FGM. The model of (Anciaux et al. 2018) and the

present model, derived under complementary approximations (SSWM *versus* WSSM),

capture the results of simulations in a different and complementary portion of the range

of possible mutation rates: compare the blue ($U = 10^2 U_c$) *vs.* brown ($U = 10^{-2} U_c$) curves

to the dots in **Figure 1**. Higher mutation rates allows higher stress levels (larger r_D) to

be endured, but it is not their only effect, as we now detail.

Non-monotonic relationship between ER probability and mutational parameters: In

the following section, we investigate the effect of mutational parameters. Both the

mutation rate U and the variance of mutational effects λ affect the system in a similar

fashion through the composite parameter $\mu = \sqrt{U\lambda}$. At small t (in Eq.[4]), an increase in μ

speeds-up the early adaptive process, thus favoring rescue but also increases the

ultimate mutation load, favoring extinction by lethal mutagenesis. These antagonistic

effects of μ create a non-monotonic relationship between the rescue probability and

mutational parameters. This is illustrated in **Figure 2**, which also shows how Eq.[5] (thick

lines) captures this effect: not neglecting the parameter $\epsilon = n\,\mu/(2\,r_{max})$ makes the

relationship between μ and P_R non-monotonic. For low stress r_D, P_R is maximal and

approximately equal to 1 over a range of mutation rates (plateau **Figure 2**, see also

Supplementary Figure 6 for higher stress values). Beyond this range, the rescue

probability in Eq.[5] drops to 0 at $U = U_{max} \equiv (4\,r_{max}^2)/(n^2\,\lambda)$. U_{max} is the mutation rate

beyond which certain extinction is enforced by lethal mutagenesis because the mutation

load ($\mu\,n/2$) is larger than the maximal growth rate that can be reached in the stress

(r_{max}). Hence, even if ER allowed the population to invade the new environment, it could

not generate a stable population once at mutation-selection balance.

Note that Eq.[6], which is only valid for intermediate μ, does not capture the decrease

in P_R close to $U = U_{max}$. It does however capture the increase in P_R as mutation rate

increases, far below U_{max}.

Figure 2: ER probability against mutation rate   for a population without standing variance (   ). Dots show the results of   simulations; thin plain lines: Eq.(5) from Anciaux et al. 2018 (recalled in Appendix Eq.(A21)) derived under the SSWM regime; thick plain lines: Eq.[5] derived under the WSSM regime; dashed lines: the corresponding closed form expression in Eq.[6] derived under the weak selection intermediate mutation regime (WSIM). The shaded area corresponds to the extra contribution to ER from multiple mutants, compared to single mutants. All models and simulations are shown for a high decay rate (purple) or a low decay rate (yellow), indicated in legend. For low mutation rates, corresponding to the SSWM regime, simulations show that populations handle only low stresses, which is captured by the model from Anciaux et al. (2018) derived in the SSWM regime. Whereas for high mutation rates, corresponding to the WSSM regime, simulations show that populations handle high stresses, which is captured by the present model derived in the WSSM regime. Other parameters are   ,   ,   and
.

Evolutionary dynamics from an initially polymorphic population (at mutation-selection balance): The results presented in the previous figures illustrate rescue from

de novo mutations. In the presence of additional standing genetic variance, rescue

mutants can arise from *de novo* mutants, from pre-existing genotypes or from a

combination of both. **Figure 3** shows the qualitative similarity between the case with and

without standing genetic variance, in their dependence on   and , as observed in

simulations and captured by Eqs.[5] and [6]. Indeed, **Figure 3** confirms that the addition

of standing genetic variance does not qualitatively modify the relationship between the

rescue probability and stress intensity (, **Figure 3a**) or mutational parameters (here ,

Figure 3b). Note that the accuracy of Eq.[5] is lower for higher   , where the

continuous time approximations become less accurate to capture discrete time

simulations (see **Supplementary Figure 4**). In the next sections, for the sake of clarity

and simplicity, we will mainly discuss the scenario of ER from *de novo* mutations only, as

the qualitative behaviors are similar with an extra contribution from standing variance.

Figure 3: ER probabilities from de novo mutants only (blue: Eq.[5], black dashed line:
Eq.[6]) or from both de novo and pre-existent mutants (green: Eq.[5], red dashed line:
Eq.[6]) against the stress level   (a) or the mutation rate   (b). Dots show the results of
   simulations (started from 10 simulated populations at mutation-selection balance for
the   scenario). The shaded area corresponds to the contribution from the standing
genetic variance to the rescue compared to de novo mutation. Other parameters are
   ,   ,   and   .   in panels (a) and   in
panel (b).

Mutation window for ER: In the previous subsections, we have shown that the ER

probability drops sharply with increasing stress and is maximal over a finite range of

mutation rates, which we denote "mutation window" for ER. The "width" (range of

mutation rates) and "height" (maximum of ER probability over the range) of this window

strongly depends on stress.

Width of the window: To characterize the mutation window, its upper and lower bounds

must be defined. The lower bound of the window (denoted U_*) corresponds to the

mutation rate at which the rescue probability rises to 1/2. Thus, this lower bound is only

defined if the height of the window lies above or at 1/2 (i.e. $\max(P_R) \geq 1/2$). The upper

bound is set to the mutation rate U_{max} beyond which certain extinction is enforced by

lethal mutagenesis. The ER probability drops off very sharply close to U_{max}, so that,

approximatively, ER is only likely within the mutation window $U_* \leq U \leq U_{max}$. These two

bounds are derived in Appendix I Eq. (A17):

$$U_{max} = \frac{4\, r_{max}^2}{n^2 \lambda}$$

$$U_* \approx \frac{1}{\lambda}\left(\frac{2\, g(y_D)\, r_{max}}{\mathcal{W}\left(\frac{8\, g(y_D)}{\pi}\left(\frac{N_0\, r_{max}}{\log(2)}\right)^2\right)}\right)^2 \underset{N_0 \to \infty}{\approx} \frac{4\, g(y_D)^2\, r_{max}^2}{\lambda \log\left(\frac{N_0^2}{\log(N_0^2)}\right)^2} \tag{7}$$

where $\mathcal{W}(.)$ is the Lambert W function, which converges to $\mathcal{W}(z) \approx \log(z/\log(z))$ as z

gets large (yielding the right hand side approximation above). Here, $g(y_D)$ is the function

of stress intensity given in Eq.[6], which describes how stress intensity ($y_D = r_D/r_{max}$)

affects ER rates. Depending on the scenario, one uses $g(y_D) = g_{DN+SV}(y_D)$ or $g(y_D) =$

$g_{DN}(y_D)$ in the presence or absence of standing genetic variance, respectively (note that

the window is always wider in the former case).

Figure 4 illustrates, for an initially clonal population, the sharpness of the transition from

ER being almost certain to being highly unlikely ($P_R = 1$ to $P_R = 0$) as a function of U

and r_D. It also shows the accuracy of the approximation for U_* in Eq.[7] (dashed black line

corresponding to the right hand side approximation of Eq.[7]), compared to its numerical

estimation from Eq.[5] (color gradient). **Supplementary Figure 5** shows a similar result

for a population starting with standing genetic variance.

Figure 4: *ER probabilities from de novo mutants only (Eq.[5]) for different values of r_D and*
U. The color gradient gives the value of P_R (see legend). The red straight line corresponds
to $U = U_{max}$ (Eq.[7]) and the black dashed line corresponds to $U = U_*$ ($P_R = 1/2$) (right
hand side approximation of Eq.[7]). For a given U, the ER probability drops sharply from
$P_R = 1$ (light yellow) to $P_R = 0$ (blue), over a short increase in r_D. For a given r_D, P_R rises
sharply as U increases about U_* and then drops sharply as U increases around U_{max}. Other
parameters are $r_{max} = 1$, $N_0 = 10^4$ and $\lambda = 5 \times 10^{-3}$. $n = 4$ in panel (a) and $n = 8$ in
panel (b).

The upper bound U_{max} is independent of initial conditions or stochasticity, as

lethal mutagenesis depends on the deterministic equilibrium state of the population,

once adapted to the stress. Therefore, U_{max} does not depend on the presence or

absence of initial standing variance, the decay rate imposed by the environmental

change (y_D) or the initial population size N_0. On the contrary, the lower bound U_*

depends on these factors as it is determined by the capacity of the population to

transiently adapt to the new conditions. It shows, however, little dependence on

dimensionality (n), as is also apparent in **Figure 4**, by the accuracy of Eq.[7], which is

independent of n. Overall, the width of the mutation window where ER is likely decreases

with increasing stress r_D (**Figure 4**) and increases with initial population size N_0 (eq. [7]:

U_* decreases with N_0, and U_{max} is unchanged). The width of the window decreases with

dimensionality n and increases with the maximum fitness in the new environment r_{max},

because both parameters affect the upper bound of the window, U_{max} (eq. [7]). Lower

dimensionality and higher maximum population growth rate reduce the mutation load,

and thus allow persisting under higher mutation rates.

Finally, note that we focused on the effect of U here, but similar results could be

obtained if λ was varied (as both parameters affect P_R as a product $\mu = \sqrt{U\,\lambda}$). This is

apparent in Eq.[7] where U and λ could be exchanged.

Height of the window and "Mutation proof extinction": within the mutation window

$(U_* \leq U \leq U_{max})$, the ER probability P_R rises above 50% and then drops back to zero. Yet,

for more extreme stresses, it cannot even reach above 50% for any value of the mutation

rate: the height of the mutation window lies below 1/2). When this height is low,

extinction is 'mutation proof', in that it is highly likely whatever the mutation rate(s) U or

the variance of mutational effects λ in the population. To illustrate this, we compute the

maximum of the ER probability $\max(P_R)$ when U varies from U_c to U_{max}, by numerically

evaluating Eq.[5] over this range. **Supplementary Figure 6** shows detailed profiles of ER

probabilities against mutation rates (illustrating how $\max(P_R)$ is found), in the presence

or absence of initial standing variance. **Figure 5** shows the maximum P_R attainable as a

function of r_D and N_0: it drops (transition from yellow to blue areas) with increasing

stress r_D and decreasing population size N_0. In this example, a large part of the

combinations of the two parameters N_0 and r_D correspond to $\max(P_R)$ lower than 10%

(blue area below the lower white dashed line in **Figure 5**). Therefore, for a given

inoculum size N_0, there is always a threshold of stress level beyond which ER is nearly

impossible, whatever the mutation rates in the population.

A closed form approximation can be obtained to describe this transition (detailed

in Appendix I section VII): denote $r_D^*(p)$ the value of r_D at which $\max(P_R) = p$ for

some $p \in [0,1]$. We obtain the following simple expression for the threshold of level of

stress beyond which $\max(P_R)$ cannot exceed some level p, independently of μ:

$$r_D^*(p) = \frac{r_{max}}{n\,\delta} \log\left(\frac{N_0\,r_{max}}{n \log(1/(1-p))}\right) \quad \text{with} \begin{cases} DN : \delta = 0.6 \\ DN + SV : \delta = 0.31 \end{cases}, \qquad [8]$$

where the values of δ come from a curve-fitting procedure (detailed in Appendix I

section VII). Setting $p \ll 1$ in Eq.[8] thus provides the stress level beyond which ER is very

unlikely, whatever the mutation rate U or the variance of mutational effects λ. This

means, in particular, that the evolution of higher mutation rates (via hypermutator

strains) or higher variance of mutational effects (larger λ) would not allow the population

to avoid extinction, when confronted to this stress level. The validity of the heuristic in

Eq. [8] is illustrated in **Figure 5**, where we see that the dashed lines ($r_D^*(p)$ with $p =$

{0.1,0.5,0.9} see legend) accurately predict the transition from high to low values

of $\max(P_R)$, computed numerically from Eq.[5].

This whole argument applies to both DN and $DN + SV$ scenarios (by choosing δ

accordingly in Eq. [8]). Interestingly, we can also see that populations initially at

mutation-selection balance (standing genetic variance) can withstand stresses twice

larger than populations only adapting from *de novo* mutations (initially clonal).

Figure 5: *Maximum ER probability reached as   is varied, for different values of   and for a population with no initial polymorphism. The color gradient gives the value of   this time (see legend). The black dashed line gives the value of   (Eq.[8] with   ) and the white dashed lines the value of   and   . The maximum of the ER probability attainable (for all possible   ) drops sharply over a short range of increasing   for a given   , or over a short range of decreasing   for a given   . Other parameters are   ,   , and   . The inset panel shows how   from Eq.[8] varies with   and   .*

Distribution of extinction times: From Eq.[4] we can derive the probability density

of this distribution, in either of the two scenarios considered (purely clonal population

     or population at mutation-selection balance   ). We get:

$$ \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \tag{9}$$

where the functions   and   depend on the scenario considered and are given

explicitly in Eq.[5]. **Figure 6** illustrates the accuracy of this result and how the distribution

of extinction times varies with stress intensity   and mutation rate   . In spite of

neglecting evolutionary stochasticity, Eq.[9] still captures the shape and scale of

extinction time distributions, in the WSSM regime.

Figure 6*: Density of the extinction probability dynamics for different values of   (panel (a)*
and (c)) and   (panel (b)). The distributions of extinction times from simulations started
with an isogenic population are shown by shaded histograms, with the corresponding
theory (Eq.[9]) given by the plain red lines. The color gradient corresponds to increasing

levels of r_D (panel (b)) or of U (lower range in panel (a) and higher range in panel (c), as
indicated on the legend). Each simulated distributions is drawn from 1000 extinct
populations among a sufficient number of replicates to observe these 1000 extinctions. The
mutation rates covered in panel (a) are $U = \{0;\ 4\ U_c;\ 8\ U_c;\ 14\ U_c\}$ and in panel (c) are
$U = \{2\ U_{max};\ 1.5\ U_{max};\ 1.1\ U_{max}\}$ and in both panel the decay rate is $r_D = 0.28$. The
decay rate covered in panel (b) are $\{0.285;\ 0.33;\ 0.4;\ 0.5\}$ and the mutation rate is
$U = 15\ U_c$. Other parameters are $N_0 = 10^5$, $n = 4$, $r_{max} = 0.1$ and $\lambda = 5 \times 10^{-3}$.

Figure 6b shows that decreasing the stress level (r_D) increases the mean

persistence time of the population and also increases the variance of this duration. This

behavior could be seen as a mere scaling: even in the absence of evolution the

population takes longer time to become extinct with a smaller r_D. On the contrary,

increasing the mutation rate, keeping it below the lethal mutagenesis threshold

($0 \leq U \leq U_{max}$, **Figure 6a**), increases the mean and the variance of the persistence time.

This likely stems from subcritical mutations ($0 > r > r_D$, beneficial but not resistant) that

can transiently invade the population, thus delaying its extinction. However, beyond the

lethal mutagenesis threshold ($U > U_{max}$, **Figure 6c**), the trend is reversed: extinctions

(which are always certain then) occur faster at high mutation rates (panel c). Therefore,

even in those cases where ER probabilities are uninformative ($P_R = 0$), the distribution of

extinction times conveys important information on the underlying adaptive or

maladaptive dynamics.

Discussion

650 We investigated the effect of an abrupt environmental change on the persistence

651 of asexual populations with a large mutational input of genetic variance (WSSM regime),

652 adapting either from *de novo* mutations arising after the environmental change (DN

653 scenario) or from both *de novo* and pre-existing mutations ($DN + SV$ scenario). In a

654 previous study (Anciaux et al. 2018), we studied evolutionary rescue when considering

adaptation over a phenotype-fitness landscape (FGM), which implies pervasive epistasis between multiple mutations and imposes a relationship between the initial decay rate of the population, the proportion and growth rate of resistance alleles, and their selective cost in the ancestral (before the stress) environment. However, we assumed that rescue resulted from rare mutations with strong effects (SSWM). The key contributions of the present model, building on our previous work, are to (i) allow for the cumulative effect of multiple mutations and (ii) provide insights into the distribution of extinction times in the presence of an evolutionary response.

Single step (SSWM regime) *vs.* multiple step (WSSM) regimes in ER: In spite of its complexity, the ER process in high mutation rate regimes can readily be captured by simple analytic approximations (Eqs.[5] and [6]), which neglect evolutionary stochasticity and only account for demographic stochasticity. Overall, the SSWM and WSSM approximations roughly capture the ER process in complementary domains of the mutation rate spectrum (**Figures 1**-**3**).

This approach shows how multiple mutations allow withstanding higher stress than what the single step approximation (SSWM in Anciaux *et al.* 2018) predicts (**Figures 1 and 3**). However, this is only true for intermediate mutation rates: a further increase in mutation rate ultimately shifts the system to a lethal mutagenesis regime (**Figures 2 to 4**). Indeed, the dependence between the ER probability and the mutation rate is not monotonic. The model shows an optimal mutation rate for the ER probability, at which the maximal ER probability may be less than 1 (depending on the stress, **Figure 5**). Beyond this rate, the ER probability drops down, to some point (U_{max}, Eq. [7]) where the mutation load is so large that absolute fitness is negative at mutation selection balance. This non-monotonic dependence reflects the continuum between ER and lethal mutagenesis along a gradient of mutation rates.

Similarities to existing ER models: Some of our key previous findings (Anciaux et al.
2018), regarding how ER depends on the parameters of the fitness landscape (FGM here)
are still valid in the more polymorphic WSSM regime. The main common features are (i)
the sharp decrease of ER probabilities with stress (decay rate r_D), (ii) their log-linear
increase with initial population size N_0, (iii) the fact that standing variance allows
withstanding higher stress, (vi) the limited effect of dimensionality n (for mutation rates
far from lethal mutagenesis, eq. [6]). The effect of initial population size ($P_R = 1 -
\exp(-N_0\,\omega)$ eq.[5]) is exactly the same as in previous models where ER stemmed from
single mutants (SSWM regime Orr and Unckless 2008, 2014; Martin *et al.* 2013; Anciaux
et al. 2018). This is expected of any model ignoring interactions between individuals, be
it evolutionary (e.g. sexual reproduction or frequency-dependent selection) or
demographic (e.g. density-dependence): each of the N_0 lineages initially present
contributes independently to ER (with some rate ω per individual). Decay rate has a
broadly similar (but quantitatively different) effect in previous ER models not based on a
fitness landscape. The other parameters (r_{max}, n, λ) are not defined outside the FGM.
More generally, the key implications of considering the FGM to model ER are detailed
more thoroughly in Anciaux *et al.* (2018).

Experimental test and parametrization: To experimentally test the predictions, the
assumptions of the WSSM regime *a priori* imply the need to use organisms with
relatively high mutation rates, such as viruses, highly mutating strains of bacteria or
possibly cancer cells. How large should the mutational parameters be for the WSSM to
apply? Empirically mutation effects and rates, at least in microbes, are typically scaled by
growth rates. In our model, if this growth rate is the maximal growth rate (that of the
wild type adapted to the lab environment), and assuming the lab environment and the

stressful environment have the same r_{max}, then, these scaled parameters would

correspond to $\bar{s} \equiv n\,\lambda/(2\,r_{max})$ and $u \equiv U/r_{max}$. If r_{max} is approximately equal to a birth

rate (i.e. the optimal 'wild-type' has a small death rate in the lab environment) $\bar{s}$ is akin to

a mean mutation effect per division, while u is akin to a mutation rate per division.

Expressed in these parameters, the WSSM approximation (Eq.[5]) applies whenever $u \gg$

$u_c \equiv \bar{s}\,n/2$, lethal mutagenesis occurs when $u = u_{max} \equiv 2/(n\,\bar{s})$ and the intermediate

mutation weak selection approximation (Eq.[6]) is valid when $u_c \ll u \ll 1$ (see the section

weak selection intermediate mutation approximation from the results). Estimates of

dimensionality n are typically not very large based on mutation fitness effects analyzed

under the FGM (Martin and Lenormand 2006; Perfeito et al. 2014). Therefore, based on

this very rough analysis, the WSSM ER model proposed here would apply in strains with

mutation rates u higher than mutation effects $\bar{s}$, measured in the same per division time

units.

For a proper experimental test of the predictions, the parameters N_0, U, r_D, r_{max}, λ

and n must be measured, in the stressful environment(s) where ER is studied. The

methods and challenges in estimating these parameters are discussed in Anciaux *et al.*

(2018). Note however that (i) the WSSM results only depend on the product $\mu = \sqrt{U\,\lambda}$

and that (ii) an error on the dimensionality may not be critical given its relatively small

effect (except in setting U_c above which the model applies). The composite parameter μ

can be estimated directly from mutation accumulation (MA) experiments as $\mu = \sqrt{U\,\lambda} =$

$\sqrt{2\Delta M/n}$ where $\Delta M = U\,E(s)$ is the change in mean fitness per unit time in the MA

experiment ($E(s) = n\,\lambda/2$ is the mean selection coefficient of random mutations).

Finally, if the model is valid, fitting an observed distribution of extinction times

(**Figure 6**) might provide estimates of r_D, r_{max}, n and possibly μ (note however that

large or low values of $\mu > \mu_c$ could produce similar distributions, see **Figure 6**). This

would allow using not only the information from rescued populations but that from
extinct ones. Such empirical test would require fine-scale time series of the population
size or at least the extinction status over time.

Treatment against pathogens, hyper-mutators and lethal mutagenesis: Our model
as well as previous ones all suggest that the effectiveness of a given treatment depends
on the mutation rate of the organism. Polymorphism for mutation rate and invasion of
hyper-mutator genotypes are thus potentially important issues for treatments against
pathogens. However, our results suggest that a sufficiently strong stress could be
effective in spite of hyper-mutator evolution. Indeed, because of the lethal mutagenesis
effect (**Figure 2**), ER is only possible within a mutation rate window: a hyper-mutator
would have to hit this window to be advantageous, and the width of the window
narrows with increased stress (**Figure 4**). At sufficiently higher stress levels, ER is unlikely
whatever the mutation rate (**Figure 5**), making these strong treatments robust to hyper-
mutator evolution. Whether this pattern is confirmed empirically and whether the
required treatment levels are then not too harmful for the treated subject remain open
questions. Note also that the same line of argument could be used, not for mutation rate
evolution, but for the evolution of the variance of mutational effects, as the end result
depends on the product $\mu^2 = U \lambda$.

Our model covers the continuum from stress induced extinction to extinction
induced by lethal mutagenesis. The latter might be an option, especially for organisms
for which no "stress treatment" exists, or whose high mutation rates (above U_* in our
model) allows them to withstand even strong stresses. Our results indeed confirm that
increasing the mutation rate in this context (above U_{max}) will allow to completely
eliminate the population. The addition of a stressor might also help in the process, as
has been suggested before (Pariente et al. 2001, 2003, 2005): indeed, the ER probability

does drop faster with increasing U (as we approach U_{max}) in the presence of a strong

stress (with high y_D), see **Supplementary Figure 6**.

Treatment duration issues: One may also wonder how long a treatment must last (be it

by stress effect or by lethal mutagenesis) for it to be efficient. From the distribution of

extinction times (**Figure 6**), it is possible to predict the duration of the treatment needed

to get rid of, say, 99% of the pathogen populations. Strong stresses (dark blue histogram

in **Figure 6b**) tend to both decrease the overall ER probability and shrink the distribution

of times to extinction. This means they are more efficient overall and require shorter

treatment durations, with less risk associated with imperfect treatment compliance.

However, when the mutation rate increases towards U_{max} (treatment by lethal

mutagenesis), although extinction becomes highly likely ($P_R \to 0$, **Figure 2**), extinction

times get more variable and the mode of the distribution is higher (**Figure 6a**). If the

mutation rate increases beyond U_{max}, the opposite pattern is observed: the distribution

of extinction times shrinks as U increases further away from U_{max} (**Figure 6c**). Hence, a

treatment by lethal mutagenesis, even if guaranteeing total extinction after infinite time,

may need to be applied for a long time to significantly decrease ER probability (if the

resulting U is close to U_{max}). Thus, observed distributions of extinction times hold

valuable information.

Limits of the model: Obviously, the different results and applications described above

are limited by the hypotheses of the model.

First, the model ignores density (discussed in Anciaux et al. 2018) or frequency-

dependent effects. Hence, our model cannot describe "competitive release" effects which

have been discussed in the context of antibiotic resistance (Read et al. 2011; Day and

Read 2016), whereby higher stresses, by eliminating sensitive genotypes more rapidly,

can "release" limiting resources for resistant ones, and ultimately increase ER probability.

However, we do not expect *a priori* the present model to show such a decrease in ER

probability at higher stress levels. In the FGM in particular, given the very sharp decay

in P_R with stress level r_D, it is highly likely that density-dependence (hence room for

competitive release) will only mitigate the result and that ER probability will still decrease

at higher stress. However, this must be studied quantitatively in a dedicated model,

which would account for density-dependence.

The approximation of the demographic dynamics by a Feller diffusion also imposes that

demographic variations remain smooth, which may be inaccurate, e.g. for some viruses

showing occasional large burst events. The Feller diffusion can also fail to predict

discrete time demography (used here in the simulations) at high growth rates per

generation ($r \geq 1$).

Second, in the demographic dynamics, we also make the approximation that σ is

roughly constant across genotypes and thus constant over time. This approximation can

be accurate if the mutant growth rates are typically small compared to their reproductive

variance $r_i \ll \sigma_i$, which is valid as long as $b_i - d_i \ll b_i + d_i$ (with b_i (resp. d_i) the birth rate

(resp. the death rate) of each genotype; discussed in Martin *et al.* 2013). However if

mutations affect substantially but dissimilarly the birth and death rates this

approximation might fail. This could be handled by considering the deterministic time

dynamics of the mean reproductive variance $\bar{\sigma}(t)$ in addition to that of the mean growth

rate $\bar{r}(t)$, in the inhomogeneous Feller diffusion.s

Third our results assume a sharp change in the environment, which does not

reflect all forms of stresses, including in the context of sudden treatment (e.g. antibiotics

can have complex pharmacokinetic patterns over time Regoes et al. 2004).

Conclusion: Recently, lethal mutagenesis treatments have received renewed interest for

their ability to provide alternative treatments against viruses resistant to other type of

drugs (Jiang et al. 2016; Escribano-Romero et al. 2017). The present model attempts to

provide a framework to predict the efficacy of such treatments. The strategy used here of

coupling deterministic evolutionary trajectories with stochastic demographic dynamics

could in principle be applied to other models, to cope with a large input of mutations

affecting the ER process. We hope that future experimental tests will evaluate its

accuracy and potential to tackle various pressing applied issues.

<u>Bibliography</u>

Alexander, H. K., G. Martin, O. Y. Martin, and S. Bonhoeffer. 2014. Evolutionary rescue: linking

theory for conservation and medicine. Evol. Appl. 7:1161–1179.

Anciaux, Y., L.-M. Chevin, O. Ronce, and G. Martin. 2018. Evolutionary Rescue over a Fitness

Landscape. Genetics 209:265–279.

Arias, A., L. Thorne, and I. Goodfellow. 2014. Favipiravir elicits antiviral mutagenesis during virus

replication in vivo. eLife 3.

Bank, C., N. Renzette, P. Liu, S. Matuszewski, H. Shim, M. Foll, D. N. A. Bolon, K. B. Zeldovich, T. F.

Kowalik, R. W. Finberg, J. P. Wang, and J. D. Jensen. 2016. An experimental evaluation of

drug-induced mutational meltdown as an antiviral treatment strategy. Evolution 70:2470–

2484.

Bansaye, V., and F. Simatos. 2015. On the scaling limits of Galton-Watson processes in varying

environments. Electron. J. Probab. 20.

Bell, G. 2017. Evolutionary Rescue. Annu. Rev. Ecol. Evol. Syst. 48:605–627.

Blanquart, F., and T. Bataillon. 2016. Epistasis and the Structure of Fitness Landscapes: Are

Experimental Fitness Landscapes Compatible with Fisher's Geometric Model? Genetics

203:847–862.

Bull, J. J., R. Sanjuán, and C. O. Wilke. 2007. Theory of Lethal Mutagenesis for Viruses. J. Virol.

81:2930–2939.

Bull, J. J., and C. O. Wilke. 2008. Lethal Mutagenesis of Bacteria. Genetics 180:1061–1070.

Carlson, S. M., C. J. Cunningham, and P. A. H. Westley. 2014. Evolutionary rescue in a changing

world. Trends Ecol. Evol. 29:521–530.

Couce, A., A. Rodríguez-Rojas, and J. Blázquez. 2015. Bypass of genetic constraints during

mutator evolution to antibiotic resistance. Proc R Soc B 282:20142698.

Day, T., and A. F. Read. 2016. Does High-Dose Antimicrobial Chemotherapy Prevent the Evolution

of Resistance? PLOS Comput. Biol. 12:e1004689.

Eliopoulos, G. M., and J. Blázquez. 2003. Hypermutation as a Factor Contributing to the

Acquisition of Antimicrobial Resistance. Clin. Infect. Dis. 37:1201–1209.

Escribano-Romero, E., N. J. de Oya, E. Domingo, and J. C. Saiz. 2017. Extinction of West Nile Virus

by Favipiravir through Lethal Mutagenesis. Antimicrob. Agents Chemother. 61:e01400-17.

Feder, A. F., S.-Y. Rhee, S. P. Holmes, R. W. Shafer, D. A. Petrov, and P. S. Pennings. 2016. More

effective drugs lead to harder selective sweeps in the evolution of drug resistance in HIV-

1. Elife 5:e10670.

Feller, W. 1951. Diffusion processes in genetics. University of California Press Berkeley, Calif.

Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Oxford University Press.

Gerrish, P. J., and R. E. Lenski. 1998. The fate of competing beneficial mutations in an asexual

population. Genetica 102:127.

Gomulkiewicz, R., and R. D. Holt. 1995. When does Evolution by Natural Selection Prevent

Extinction? Evolution 49:201.

Gomulkiewicz, R., R. D. Holt, M. Barfield, and S. L. Nuismer. 2010. Genetics, adaptation, and

invasion in harsh environments. Evol. Appl. 3:97–108.

Gomulkiewicz, R., S. M. Krone, and C. H. Remien. 2017. Evolution and the duration of a doomed

population. Evol. Appl. n/a-n/a.

Gonzalez, A., O. Ronce, R. Ferriere, and M. E. Hochberg. 2013. Evolutionary rescue: an emerging

863   focus at the intersection between ecology and evolution. Philos. Trans. R. Soc. B Biol. Sci.

864   368:20120404–20120404.

865   Greenspoon, P. B., and N. Mideo. 2017. Evolutionary rescue of a parasite population by mutation

866   rate evolution. Theor. Popul. Biol. 117:64–75.

867   Holt, R., M. Barfield, and R. Gomulkiewicz. 2004. Temporal Variation Can Facilitate Niche

868   Evolution in Harsh Sink Environments. Am. Nat. 164:187–200.

869   Holt, R. D., R. Gomulkiewicz, and M. Barfield. 2003. The phenomenology of niche evolution via

870   quantitative traits in a "black-hole" sink. Proc. R. Soc. B Biol. Sci. 270:215–224.

871   Jiang, Y.-C., H. Feng, Y.-C. Lin, and X.-R. Guo. 2016. New strategies against drug resistance to

872   herpes simplex virus. Int. J. Oral Sci. 8:1–6.

873   Kreiner, J. M., J. R. Stinchcombe, and S. I. Wright. 2018. Population Genomics of Herbicide

874   Resistance: Adaptation via Evolutionary Rescue. Annu. Rev. Plant Biol. 69:611–635.

875   Lande, R. 1976. Natural Selection and Random Genetic Drift in Phenotypic Evolution. Evolution

876   30:314.

877   Lande, R. 1980. The Genetic Covariance Between Characters Maintained by Pleiotropic Mutations.

878   Genetics 94:203–215.

879   Liu, L. L., F. Li, W. Pao, and F. Michor. 2015. Dose-Dependent Mutation Rates Determine Optimum

880   Erlotinib Dosing Strategies for EGFR Mutant Non-Small Cell Lung Cancer Patients. PLoS

881   ONE 10.

Loeb, L. A. 2001. A Mutator Phenotype in Cancer. Cancer Res. 61:3230–3239.

Loeb, L. A., J. M. Essigmann, F. Kazazi, J. Zhang, K. D. Rose, and J. I. Mullins. 1999. Lethal

mutagenesis of HIV with mutagenic nucleoside analogs. Proc. Natl. Acad. Sci. 96:1492–

1497.

Loverdo, C., and J. O. Lloyd-Smith. 2013. Evolutionary Invasion and Escape in the Presence of

Deleterious Mutations. PLoS ONE 8:e68179.

Martin, G., R. Aguilée, J. Ramsayer, O. Kaltz, and O. Ronce. 2013. The probability of evolutionary

rescue: towards a quantitative comparison between theory and evolution experiments.

Phil Trans R Soc B 368:20120088.

Martin, G., S. F. Elena, and T. Lenormand. 2007. Distributions of epistasis in microbes fit

predictions from a fitness landscape model. Nat. Genet. 39:555–560.

Martin, G., and S. Gandon. 2010. Lethal mutagenesis and evolutionary epidemiology. Philos.

Trans. R. Soc. Lond. B Biol. Sci. 365:1953–1963.

Martin, G., and T. Lenormand. 2006. The Fitness Effect of Mutations Across Environments: A

Survey in Light of Fitness Landscape Models. Evolution 60:2413–2427.

Martin, G., and L. Roques. 2016. The Non-stationary Dynamics of Fitness Distributions: Asexual

Model with Epistasis and Standing Variation. Genetics 204:1541–1558.

Matuszewski, S., L. Ormond, C. Bank, and J. D. Jensen. 2017. Two sides of the same coin: A

population genetics perspective on lethal mutagenesis and mutational meltdown. Virus

Evol. 3.

McCandlish, D. M., and A. Stoltzfus. 2014. Modeling Evolution Using the Probability of Fixation:

History and Implications. Q. Rev. Biol. 89:225–252.

Orr, H. A., and R. L. Unckless. 2008. Population Extinction and the Genetics of Adaptation. Am.

Nat. 172:160–169.

Orr, H. A., and R. L. Unckless. 2014. The Population Genetics of Evolutionary Rescue. PLOS Genet.

10:e1004551.

Pariente, N., A. Airaksinen, and E. Domingo. 2003. Mutagenesis versus Inhibition in the Efficiency

of Extinction of Foot-and-Mouth Disease Virus. J. Virol. 77:7131–7138.

Pariente, N., S. Sierra, and A. Airaksinen. 2005. Action of mutagenic agents and antiviral inhibitors

on foot-and-mouth disease virus. Virus Res. 107:183–193.

Pariente, N., S. Sierra, P. R. Lowenstein, and E. Domingo. 2001. Efficient Virus Extinction by

Combinations of a Mutagen and Antiviral Inhibitors. J. Virol. 75:9723–9730.

Perfeito, L., A. Sousa, T. Bataillon, and I. Gordo. 2014. Rates of Fitness Decline and Rebound

Suggest Pervasive Epistasis. Evolution 68:150–162.

Ramsayer, J., O. Kaltz, and M. E. Hochberg. 2013. Evolutionary rescue in populations of

Pseudomonas fluorescens across an antibiotic gradient. Evol. Appl. 6:608–616.

Read, A. F., T. Day, and S. Huijben. 2011. The evolution of drug resistance and the curious

orthodoxy of aggressive chemotherapy. Proc. Natl. Acad. Sci. 108:10871–10877.

Regoes, R. R., C. Wiuff, R. M. Zappala, K. N. Garner, F. Baquero, and B. R. Levin. 2004.

Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic

Treatment Regimens. Antimicrob. Agents Chemother. 48:3670–3676.

Springman, R., T. Keller, I. J. Molineux, and J. J. Bull. 2010. Evolution at a High Imposed Mutation

Rate: Adaptation Obscures the Load in Phage T7. Genetics 184:221–232.

Taddei, F., M. Radman, J. Maynard-Smith, B. Toupance, P. H. Gouyon, and B. Godelle. 1997. Role

of mutator alleles in adaptive evolution. Nature 387:700–702.

Tenaillon, O. 2014. The Utility of Fisher's Geometric Model in Evolutionary Genetics. Annu. Rev.

Ecol. Evol. Syst. 45:179–201.

Uecker, H. 2017. Evolutionary rescue in randomly mating, selfing, and clonal populations.

Evolution 71:845–858.

Uecker, H., and J. Hermisson. 2016. The Role of Recombination in Evolutionary Rescue. Genetics

202:721–732.

Wilson, B. A., P. S. Pennings, and D. A. Petrov. 2017. Soft Selective Sweeps in Evolutionary Rescue.

Genetics 205:1573–1586.

Wylie, C. S., and E. I. Shakhnovich. 2012. Mutation Induced Extinction in Finite Populations: Lethal

Mutagenesis and Lethal Isolation. PLOS Comput. Biol. 8:e1002609.

Notation	Description	Formula
N_t, N_0	N_t: population size at time t after the onset of the stress. N_0: initial population size at the onset of the stress.	
U	Mutation rate *per* individual *per* unit time.	
n	Number of traits under stabilizing selection, or phenotypic dimensionality.	
λ	Variance of mutational effects: variance of the phenotypic effects of mutations, per trait, in a trait space scaled by the strength of stabilizing selection	$\mathbf{dz} \sim N(\mathbf{0}, \lambda\, \mathbf{I}_n)$
μ	Composite parameter : square root of mutational variance per trait	$\mu = \sqrt{U\lambda}$
$\mathbf{z}$	n-dimensional vector $\mathbf{z} \in \mathbb{R}^n$ of (breeding values for) phenotype	
$\mathbf{z}_A, \mathbf{z}_*$	$\mathbf{z}_*$: optimal phenotype in the new environment $\mathbf{z}_A$: average phenotype of the ancestral population (before the onset of stress)	
r, σ	Growth rate (r) and reproductive variance (σ) of a given genotype, in the new environment.	Eq.[1]
$\overline{r}_t, \overline{\sigma}$	Mean growth rate ($\overline{r}_t$) and mean reproductive variance ($\overline{\sigma}$) of the population, in the new environment.	Eqs. [2] and [3]
$\langle \overline{r}_t \rangle$	'ensemble mean' growth rate : expected mean growth rate $\overline{r}_t$ across stochastic replicates	Eqs. [2] and [3]
r_{max}	Maximum possible growth rate in the new environment.	$r(\mathbf{z}_*) = r_{max}$
r_D	Rate of decay of the ancestral phenotype $\mathbf{z}_A$ in the new environment.	$r(\mathbf{z}_A) = r_{max} - \|\mathbf{z}_A\|^2/2 = -r_D$
y_D	Rate of decay scaled by r_{max}.	$y_D = r_D/r_{max}$
ϵ	Ratio between the mutational load and the maximal growth rate	$\epsilon = n\,\mu/(2\,r_{max})$
$\omega_{DN}, \omega_{DN+SV}$	ω_{DN}: rate of rescue from *de novo* mutations scaled by N_0 ω_{DN+SV}: rate of rescue from both *de novo* and standing variance scaled by N_0	Eqs.[5],[6]
P_{ext}, P_R	P_{ext} : Extinction probability P_R : Rescue probability	P_{ext} see Eq.[4] $P_R = 1 - P_{ext} = 1 - \exp(-N_0\,\omega_R)$
U_c	Critical mutation rate beyond which the WSSM regime is valid.	$U_C = n^2\,\lambda/4$
U_{max}	Mutation rate beyond which certain extinction is enforced by lethal mutagenesis	$U_{max} = 4\,r_{max}^2/(n^2\,\lambda)$
U_*	Mutation rate at which the rescue probability rises to 1/2	Eq.[7]
$r_D^*(p)$	Rate of decay for which the maximal rescue probability for any U or λ is equal to p	Eq.[8]

_Table 1__: Notations_