Progressive domain segregation in early embryonic development and underlying correlation to genetic and epigenetic changes

Hui Quan[1,†], Hao Tian[1,†],Sirui Liu[1,†], Yue Xue[1], Yu Zhang[2,3], Wei Xie[2,3], Yi Qin Gao[1,4,5*]

[1] Beijing National Laboratory for Molecular Sciences, College of Chemistry and Molecular Engineering, Peking University, Beijing, 100871, China.

[2] Center for Stem Cell Biology and Regenerative Medicine, MOE Key Laboratory of Bioinformatics, School of Life Sciences, Tsinghua University, Beijing, 100871, China.

[3] THU-PKU Center for Life Sciences, Tsinghua University, Beijing, 100871, China.

[4] Biomedical Pioneering Innovation Center (BIOPIC), Peking University, Beijing, 100871, China.

[5] Beijing Advanced Innovation Center for Genomics, Peking University, Beijing, 100871, China.

* Correspondence:

Yi Qin Gao

E-mail: gaoyq@pku.edu.cn

†These authors have contributed equally to this work.

ABSTRACT

Chromatin undergoes drastic structural organization and epigenetic reprogramming during embryonic development. We present here a consistent view of the chromatin structural change, epigenetic reprogramming and the corresponding sequence dependence in both mouse and human embryo development. The two types of domains, identified earlier as forests and prairies, become spatially segregated during embryonic development, with the exception of zygotic genome activation (ZGA) and implantation, at which notable domain mixing occurs. Structural segregation largely coincides with DNA methylation and gene expression changes. Genes located in mixed prairie domains show proliferation and ectoderm differentiation-related function in ZGA and implantation, respectively. Chromatin of ectoderm shows the weakest and endoderm the strongest domain segregation in germ layers. This chromatin structure difference between different germ layers generally enlarges in further differentiation. The systematic chromatin structure establishment and its sequence-based segregation strongly suggest DNA sequence as a possible driving force for the establishment of chromatin 3D structures which affect profoundly the expression profile. Other possible factors correlated with/influencing chromatin structures, including temperature, germ layers, and cell cycle, were discussed for an understanding of concerted chromatin structure and epigenetic changes in development.

Keywords

Sequence, Domain segregation, Epigenetic modifications, ZGA and Implantation, Temperature.

INTRODUCTION

In mammals, chromatin undergoes drastic organization and epigenetic reprogramming after fertilization(Xu and Xie, 2018; Zheng and Xie, 2019). These processes are essential for gene regulation either globally or specifically by generating chromatin environment that is permissive or repressive to gene expression(Burton and Torres-Padilla, 2014). Chromatin of mouse zygotes and two-cell embryos has obscure high-order structures, existing in markedly relaxed states, which undergoes consolidation of TADs and segregation of chromatin compartments through development(Du et al., 2017; Ke et al., 2017). TAD structure and compartmentalization are also gradually established following human fertilization(Chen et al., 2019). Along with remodeling of 3D chromatin architecture, genome-wide epigenetic reprogramming also takes place during embryonic development(Morgan et al., 2005; Burton and Torres-Padilla, 2010; Dahl et al., 2016; Liu et al., 2016; Zhang et al., 2016; Wang et al., 2018; Xu and Xie, 2018). Global hypomethylation of the genome occurs, and histone modifications globally reset, changing from a non-canonical distribution to canonical one(Smith and Meissner, 2013; Lee et al., 2014; Zheng et al., 2016) during early embryonic development, leading to the specification of the germ layers and cell differentiation.

The progression of the mammalian embryo from fertilization to germ layer formation, concurrent with transcriptional changes and cell fate transitions(Burton and Torres-Padilla, 2014), involves an ordered series of hierarchical lineage determination that ensures the establishment of a blueprint for the whole animal body. One of the most notable transcriptional changes is the zygotic genome activation(ZGA), during which the embryo changes from a state where there is little transcription to another state in which thousands of genes are transcribed(Lu et al., 2016). ZGA is mechanistically coordinated with changes in chromatin state and cell cycle(Jukam et al., 2017). Mammalian ZGA may primarily prepare for the differentiation towards the inner cell mass (ICM) and the trophectoderm (TE), which begins at the morula stage, then the TE can further develop to the extraembryonic tissue necessary for embryo implantation and receiving nutrients. After implantation, the ICM then gives rise to three germ layers (ectoderm, mesoderm and endoderm) through gastrulation, generating founder tissues for subsequent somatic development(Lawson et al., 1991; Arnold and Robertson, 2009).

Thanks to recent developments in low-input chromatin analysis technologies,
chromatin structural, epigenetic and transcriptional properties have been roundly explored
in the early embryonic development process(Zhang et al., 2018). Taken that chromatin
structural properties at the mouse pre-implantation stages have been investigated, analysis
on the latest Hi-C data describing chromatin structural properties at post-implantation
stages in mouse and embryonic development in human is expected to be able to provide a
fairly complete view of the structural change during development. On the other hand, the
relationship between global structural changes, epigenetic reprogramming and DNA
sequence remains largely unknown and therefore needs to be further investigated.

In our previous study, we analyzed the DNA sequence dependence in the formation of
3D chromatin structures for different cell types(Liu et al., 2018). Based on CpG island
(CGI) densities, the genome was divided into alternative forest (high CGI density, F) and
prairie (low CGI density, P) domains with average lengths of 1 to 3 million bases (forest
and prairie domains are therefore cell type-invariant in one species). CGI forests and
prairies effectively separate the linear DNA sequence into domains with distinctly different
genetic, epigenetic and structural properties. The segregation of the two domains tends to
intensify during embryonic development, cell differentiation and senescence as a result of
sequence based thermodynamic stabilization. However, the segregation degree of some
somatic cells is less than that of ICM, implying that chromatin structure changes in a non-
monotonic way from zygote to differentiated somatic cells. How chromatin conformation
gradually changes to establish cell identity during development is thus an interesting open
question.

In the present study, we conducted an analysis on global chromosome structural
changes during early embryonic development. Two specific stages, ZGA and implantation,
during which gene activation and lineage specification occur, respectively, were both found
to involve mixing of the two types of genomic domains, with the latter showing a more
significant change than the former. The segregation level positively correlates with the
proportion of prairie domains reside in compartment B, the larger value of which suggests
a more segregated chromatin structure. The DNA methylation distribution in early
embryonic development also correlates with the trend of domain segregation in this process,
which indicates that chromatins of endoderm and mesoderm are more segregated than that

of ectoderm. Moreover, the domain segregation levels of the earliest fate-committed germ

layers correlate with those of the differentiated cells. Detailed functions of genes residing

in more mixed domains during implantation were then analyzed, based on which we

proposed that pluripotent ICM cells may first differentiate to ectoderm-like epiblast via the

enhanced interactions of prairies with forests. Finally, we presented a consistent view for

the chromatin structural change from birth to senescence, and discussed possible factors

influencing global chromatin structure, such as temperature, germ layer, and cell division.

MATERIALS AND METHOD

Overall Segregation Ratio

Based on the Hi-C contact matrix, the inter-domain contact ratio between the same domain types was calculated as

$$R_f^s = \frac{\sum_{i,j \in F, i \neq j} D_{ij}}{\sum_{i \in F, j \in A} D_{ij}} \quad \text{and} \quad R_p^s = \frac{\sum_{i,j \in P, i \neq j} D_{ij}}{\sum_{i \in P, j \in A} D_{ij}}$$

for forests and prairies, respectively. In the above equations, D_{ij} represents the sum of contacts between the two domains i and j, which is further divided by the product of domain length of i and j. F is the collection for all forest domains, P is the collection for all prairie domains, and A is the union of sets F and P.

The inter-domain contact between different types was calculated similarly as

$$R_f^d = \frac{\sum_{i \in F, j \in P} D_{ij}}{\sum_{i \in F, j \in A} D_{ij}} \quad \text{and} \quad R_p^d = \frac{\sum_{i \in P, j \in F} D_{ij}}{\sum_{i \in P, j \in A} D_{ij}}$$

The overall segregation ratio OR_s was then defined as the ratio of inter-domain contacts between the same types and different types:

$$OR_s = \frac{R_f^s}{R_f^d} \quad \text{or} \quad OR_s = \frac{R_p^s}{R_p^d}$$

for forests and prairies, respectively. All Hi-C data in this calculation were normalized by ICE normalization(Servant et al., 2015).

3D Chromatin Structure Modelling

Detailed information can be seen in our previous work(Xie et al., 2017). Briefly, we first coarse-grained a chromosome as a string of beads. The equilibrium distance between two beads was obtained by converting the contact frequency to the spatial distance. A randomly generated initial structure was then used for further structure optimization using MD (Molecular Dynamics) simulations, until the RMSD (root-mean-square deviation) of the modeled structure became convergent.

Domain Segregation Ratio Calculation

For each forest/prairie domain i, the domain segregation ratio DR_s was defined as the ratio between its inter-domain contacts with the same types and its inter-domain contacts with different types:

$$DR_s^{i \in F} = \left.\sum_{j \in F, j \neq i} D_{ij} \middle/ \sum_{j \in P} D_{ij}\right. \quad \text{and} \quad DR_s^{i \in P} = \left.\sum_{j \in P, j \neq i} D_{ij} \middle/ \sum_{j \in F} D_{ij}\right.$$

In the above equations, D_{ij} represents the sum of contacts between the two domains i

and j, F is the collection of all forest domains, and P is the collection of all prairie

domains.

Distance-dependent Segregation Ratio Calculation

The segregation ratio at distance d was calculated as

$$R_s^f(d) = \frac{C_{f-f}}{C_{f-p}} \quad \text{and} \quad R_s^p(d) = \frac{C_{p-p}}{C_{f-p}}$$

for forests and prairies, respectively. In the above definitions, C_{f-f}, C_{f-p} and C_{p-p} are

the average Hi-C contact probabilities between forest and forest, forest and prairie, and

prairie and prairie, under genomic distance d, respectively.

Open Sea Methylation Difference Index (MDI) between Forest and Prairie

We quantified the methylation difference between adjacent forests and prairies by

$$MDI_i = \left(q_i - \frac{q_{i-1} + q_{i+1}}{2} \right) \bigg/ \left(\frac{q_{i-1} + q_i + q_{i+1}}{3} \right)$$

where q_i, q_{i-1} and q_{i+1} are the average open sea methylation levels for the ith domain

and its two flanking domains.

Overall Relative Segregation Ratios

Overall relative segregation ratios (R_{or}) between one tissue and another was identified as

$$R_{or} = ln\left(\frac{N_{ij}^o}{N_{ji}^o} \right)$$

respectively. Here N_{ij}^o is the number of domains possessing higher DR_s value in tissue

i, compared to tissue j. A positive value of R_{or} indicates that the chromatin of the former

tissue is more segregated than that of the latter. Thus, the parameter R_{or} is used to

generally reflect domain segregation behavior differences between two tissues.

Significantly More Segregated or More Mixed Domains

To identify forest or prairie domains that are significantly more (or less) segregated in one

tissue when compared to another, we first identified threshold values to distinguish more

(or less) segregated domains at certain stage. We calculated the logarithm ratio of DR_s

values between the two ICM replicates from the same laboratory. Significantly more

strongly segregated domains are taken as those whose variation of DR_s values are in the

top 2.5 percent tier of all domains, and more mixed domains in the bottom 2.5 percent. The

corresponding threshold values of DR_s variation $T_{DR_s}^{log}$ for segregation and mix are then

0.1469 and -0.1469, respectively.

We calculated the logarithm ratio of DR_s between early and late 2-cell embryos for ZGA,

ICM and E6.5 epiblast for implantation, respectively. Significantly more strongly

segregated domains were identified as those with the logarithm ratio of DR_s higher than

0.1469, and significantly more mixed domains were defined as those with the logarithm

ratio of DR_s lower than -0.1469. For convenience, we denoted more strongly segregated

and more mixed domains in forest and prairie as F_{seg}, P_{seg}, F_{mix}, and P_{mix}, respectively.

Compartment Index Calculation

To quantify the degree of compartmentalization, we defined a compartment index (C-index)

I_i for 200-kb bin i as the logarithm ratio of the average contact between this bin and all

A compartments over that between this bin and all B compartments:

$$I_i = ln \frac{(\sum_j C_{ij} \delta_j)/N_A}{(\sum_j C_{ij}(1 - \delta_j))/N_B}$$

$$\delta_j = \begin{cases} 1, \text{if bin } j \text{ is in compartment A} \\ 0, \text{if bin } j \text{ is in compartment B} \end{cases}$$

where C_{ij} is the distance-normalized Hi-C contact probability (normalized by dividing

each contact by the average contact probability at that genomic distance) between bins i

and j. N_A and N_B are the total number of bins belonging to compartments A and B,

respectively. For each region, a higher value of I_i indicates a more compartment A-like,

and a lower one, compartment B-like environment.

Gene Function Analysis

We analyzed the functional enrichment of genes located in various selected regions by

ClusterProfiler and DAVID (https://david.ncifcrf.gov), which yielded similar results. The

results were demonstrated via GO terms with p-values. The functional annotation
clustering was analyzed using DAVID, and the results were shown via GO terms with
enrichment scores.

Identification of Lineage Specific Genes

To identify lineage specific genes, we used a Shannon-entropy-based method (Xie et al.,
2013). Genes with entropy score less than 1.7 were selected as candidates for stage-specific
genes. Among them, we selected E6.5 epiblast specific genes satisfying the following
conditions: the gene is highly expressed in E6.5 epiblast (FPKM>=1), its relative
expression is higher than 1/7, and its expression level in epiblast is higher than that in ICM.
These genes were then used for further analysis.

RESULTS

Domain Segregation in Early Embryonic Development

Previous studies on chromatin structures during early embryonic development were
mainly about changes of structural elements, such as compartment and TADs(Du et al.,
2017; Ke et al., 2017). Here we present a structural analysis based on two types of genomic
domains to show domain segregation behaviors in different scales. We analyzed chromatin
structural changes by calculating overall segregation ratio OR_s (one parameter used to
globally reflect chromatin segregation behavior), domain segregation ratio DR_s (which
was used to reflect segregation behavior of individual domain), segregation ratio along
genomic distances $R_s(d)$ and F-F (forest-forest) /F-P (forest-prairie) /P-P (prairie-prairie)
spatial interaction ratio at each stage.

In our previous study(Liu et al., 2018), based on the division of the mammalian
genomes into forest and prairie domains, OR_s was defined as the inter-domain contact
ratio between domains of the same types and different types, regarding each domain as one
unit (see "Methods"). A higher OR_s for a sample indicates a stronger segregation. Here
we calculated OR_s using Hi-C data(Du et al., 2017; Zhang et al., 2018; Chen et al., 2019)
of 21 mouse cells and 5 human cells (see Additional file 2: Data sources), which allow us
to investigate the chromatin structure changes following the early embryonic development.
The calculated OR_s for each stage is given in Figure 1A and 1B. We also constructed the

modelled 3D chromatin structures following our previous work(Xie et al., 2017) to show

visually the degree of segregation at each stage (see "Methods").

As seen from Figure 1A, OR_s increases during the development of early mouse

embryonic cells except for two dips. Such a trend suggests the increased segregation of

forest and prairie domains from each other, in line with previous observation that more

inter-compartment interactions occur at early stages than late stages(Du et al., 2017). The

two dips observed along the OR_s curve correspond to the early to late 2-cell, and ICM to

E6.5 epiblast transitions, respectively. The decrease during the former transition is seen in

both alleles (Figure S1A). For the latter transition, the modelled chromatin structure clearly

changes from a domain-separated state to a forest-prairie (F-P) mixed state (Figure 1A).

Interestingly, these two processes (that is, early to late 2-cell, and ICM to E6.5 epiblast)

exactly coincide with ZGA(Svoboda et al., 2015) and implantation(Bedzhov and Zernicka-

Goetz, 2014), respectively. The domain-level F-P chromatin mixing suggested by decrease

of OR_s is possibly related to LAD (lamina-associated domains) mixing at ZGA stage as

well as germ layer differentiation following implantation, which is analyzed below.

The overall chromatin segregation follows a similar trend in human embryonic

development (Figure 1B). OR_s generally increases except for the blastocyst to 6-week

transition, accordant with the observed trend of global segregation and domain-level

mixing after ICM stage in mouse sample. These changes on domain segregation are also

reflected by the 3D structures reconstructed via Hi-C data (Figure 1B). Again, these results

suggest that forest and prairie domains tend to segregate from each other before

implantation, and domain mixing is observed at the post-implantation stage, showing an

increased interaction of the prairie with forest domains.

Besides the overall structure parameter OR_s, we also calculated the domain

segregation ratio DR_s on each forest/prairie domain during embryonic development (see

"Methods", a higher value indicates the corresponding domain is more segregated), and

the variation trend is similar to OR_s (Figures 1C and 1D). Interestingly, the change in

DR_s is more significant for the ICM to E6.5 epiblast transition ($p = 3.7 \times 10^{-96}$ by

Welch's unequal variance test) than the early to late 2-cell transition ($p = 1.9 \times 10^{-16}$ by

Welch's unequal variance test) for prairie domains, while for forest domains the former one

is less pronounced (p: 2.4×10^{-5} VS 2.1×10^{-54}, Welch's unequal variance test).

Further, to investigate chromatin segregation at different genomic distances, we

calculated the segregation ratio as a function of genomic distance $R_s(d)$ (Figures 1E, S1C

and S2, see "Methods"). It can be seen from Figure 1E that in early mouse embryo

development, prairies are characterized by elevated segregation at genomic distances of

several Million bases (Mb). The segregation becomes more pronounced at long distances

after 2-cell stage for both forests and prairies, again consistent with an enhancement in

domain segregation. Besides, Figure S1B shows a slight decrease of $R_s(d)$ for forests at

the early to late 2-cell stage and a conspicuous decrease of $R_s(d)$ for prairies at the ICM

to E6.5 epiblast stage for mouse, the underlying biological significances behind these

phenomena will be discussed below. Enhanced segregation at several Mb scale also occurs

during human embryo development (Figure S1C), and an obvious drop of $R_s(d)$ for

prairies at the blastocyst to 6 weeks transition was also observed (Figure S1D).

Finally, the top 10% contact probabilities under genomic distance d (that is, the top

10% elements of one diagonal of Hi-C matrix), $TC_{10}(d)$, were extracted, within which the

proportions of F-F, F-P and P-P interactions were calculated (notably, the premise of such

a calculation is that all elements in the $TC_{10}(d)$ should be non-zero, otherwise the

proportions of F-F, F-P and P-P are assigned zero). The genomic distance, after which the

proportions of F-F/F-P/P-P remain to be zero, is named the critical genomic distance.

During the development of mouse embryo, the critical genomic distance gradually

increases (Figure 1F), showing the gradual establishment of long-range chromatin contact.

At the same time, F-P ratio descends while P-P ratio increases from PN3 to ICM stages,

again hinting that forest and prairie become more spatially segregated along with the

development (Figure 1G). These phenomena were also observed in human embryo

development (Figure S1E).

Compartment Changes Relate with Domain Segregation

By means of the Hi-C measurement, chromatin can be partitioned into two spatial

compartments (A and B)(Lieberman-Aiden et al., 2009). It was also found that forest

domains reside mainly in compartment A and prairie domains mainly in compartment B in

different cells, manifesting that DNA with similar sequence properties tends to spatially

interact (Liu et al., 2018). Based on these, we calculated the proportion for the two

sequential domains (forests and prairies) to be distributed in the two spatial compartment

A and B during development.

The corresponding four types of DNA sequences are named as Af, Bf, Ap, and Bp,

respectively, with the first letter denoting the compartments and second the forest (f) or

prairie (p) domain. One can see from Figure 2A that during mouse development the

proportion of prairies in compartment B (Bp) changes in the same trend as the overall

segregation ratio, which increases from PN3 to ICM stage, except for the early to late 2-

cell stage, then decreases from ICM to E6.5 epiblast, and slightly increases when E6.5

epiblast develops to E7.5 ectoderm. The ratio of Ap changes in a direction opposite to that

of Bp. Similar phenomenon was also observed in human embryos, where the changes of

OR_s are in accordance with the size of Bp (Figure S3A). Recent study also found that

attractions between heterochromatic regions are crucial for compartmentalization and

domain segregation of the active and inactive genome(Falk et al., 2019). Similar to it, we

found here that during embryo development, the aggregation behavior of prairie domains

(low CpG density region) strengthens and more and more prairie regions belong to

compartment B.

Since the genome compartmentalization is weak at early stages of development(Du et

al., 2017), for the robustness of compartment partition, we calculated the distribution of

compartment index (C-index, a parameter to quantify the degree of compartmentalization,

a larger value of which corresponds to a more compartment A-like environment, see

"Methods") at each stage for compartment A and B, respectively. The gradually increased

discrepancy of C-index between compartment A and B supports the global structure

segregation from zygote to ICM (Figure S3B). Then, we identified compartment A regions

with positive C-index as strict compartment A (sA), B with negative C-index as strict

compartment B (sB), and calculated the length of genome located in sAf (forest regions in

sA), sAp, sBf and sBp, respectively. The result showed a trend similar to Figure 2A, that

is, the proportion of prairies in strict compartment B (sBp) changes in the same trend as

the chromatin segregation level, which rules out the possibility that the growth of Bp is an

artifact due to compartment definition (Figure S3C).

In comparison with stable Bp regions, the prairie domains that switch the compartment

type are particularly interesting as they may be regarded as mediators of nuclear
architecture establishment during development. The analysis of compartment
transformation during preimplantation development shows that 7.7% of prairies belong to
compartment A (stable Ap), 26.3% of prairies always reside in compartment B (stable Bp),
and the left 66.0% switches compartment type at least once (switchable p) (Figure 2B).
Genes in the stable Ap and switchable p regions are enriched in immune- and ectoderm-
related functions, while those in the stable Bp regions are not (Figure 2C). Previous
analyses showed that forest and prairie domains tend to spatially segregate (and mainly
correspond to compartment A and B, respectively), but to different extent in different
cells(Liu et al., 2018). Moreover, forest-prairie spatial intermingling is cell-type specific,
which is thought to be associated with prairie tissue-specific gene activation and the
establishment of cell identity. Therefore, compartment B (heterochromatin) provides a
silent environment for prairie genes, which wait to be activated through spatial interactions
with forest/compartment A in the following differentiation stages (as seen in their spatial
contact and expression properties in differentiated cells(Tian et al., 2020)). The function of
genes in the stable Ap and switchable p regions thus supports that the gene expression in
prairie domains plays an important role in cell fate determination. Indeed, our earlier
analysis shows that prairie genes in compartment A are highly tissue-specific and by
examining the functions of the related genes, one can deduce the tissue type of the
associated sample(Liu et al., 2018). Besides, compared to forest genes, the expression of
genes in prairie domains are more likely to be highly correlated with the compartment
environment they reside in(Tian et al., 2020).

Next, to further investigate the differences between switchable prairie and stable
prairie regions, we compared the chromatin features of these two kinds of regions between
two adjacent stages. The cells we analyzed here include PN5, early 2-cell, late 2-cell, 8-
cell and ICM due to the availability of Hi-C data. For each two adjacent stages (i.e., PN5
vs early 2-cell, early 2-cell vs late 2-cell, late 2-cell vs 8-cell, 8-cell vs ICM), we identified
stable Bp regions and switchable p regions (p regions located in compartment B in the
earlier whereas switched to compartment A in the later stage). Although these two kinds of
prairie regions are both located in compartment B in the cell of the earlier stage, the PC1
value of switchable p regions was significantly more positive than stable Bp regions

(Figures 2D and S3D) and accordingly, switchable p regions are significantly closer to A-

B compartment boundaries (Figure S3E), indicating that Bp regions near the A-B boundary

are more likely to switch. The analyses on ATAC-seq signal, H3K4me3 and H3K27me3

also support this observation (Figure 2E). For example, from 8-cell to ICM, the ATAC-seq

and H3K4me3 signals of switchable p regions are significantly higher than stable Bp

regions in both 8 cell and ICM, while the H3K27me3 signal of switchable p regions is

weaker than stable Bp regions.

To further understand how DNA sequence affects chromatin compartmentalization in

embryo development, we calculated the GC contents and CpG densities during

development for the four genomic components Af, Bf, Ap and Bp (Figures 2F and S3F).

Notably, the GC content of Ap changes in the same trend as the segregation degree in both

mouse and human embryo development (Figures 2F and S3F). Since the proportion of Bp

and the segregation level are highly correlated, when the degree of segregation increases,

a higher portion of prairie domains partition into compartment B, leaving the remaining

prairie sequences in compartment A more "bivalent" (low GC content regions of forests or

high GC content regions of prairies(Liu et al., 2018)). Moreover, for mouse, both forest

components (Af and Bf) have higher GC contents than prairie components (Ap and Bp)

before the 2-cell stage, whereas compartment A DNA (Af and Ap) have higher GC contents

than compartment B (Bf and Bp) after one passes the 2-cell stage (Figure 2F). The CpG

densities vary in a similar trend (Figure S3F), which indicates that compartmentalization

becomes more prominent after the 2-cell stage in mouse embryos, with higher-order

structures being gradually established.

To investigate how domain segregation and CpG density affect gene expression, we

calculated the median expression level of genes in forests and prairies as a function of CpG

density. It can be seen from Figures 2G and S3G that in general the median gene expression

level increases with the increase of CpG density in a certain range. Interestingly, for genes

with similar exon CpG density, those in forests tend to show a higher expression than those

in prairies (Figures 2G and S3G), showing the importance of large-scale DNA sequence

property in gene expression (given that forest regions possess significantly higher CGI and

thus CpG densities than prairies, the surrounding sequence environment of forest genes is

thus CpG richer than prairie genes, suggesting a connection between their neighboring

sequences and expression level of genes). Furthermore, the expression difference between

forest and prairie genes (that is, the averaged difference between the two curves shown in

Figure 2G for ICM and Figure S3G for other cells) increases from early 2-cell to 8-cell,

then decreases from 8-cell to E7.5 ectoderm (Figure 2H), which is accordant with the

chromatin segregation behaviors along the early embryo development.

The Association between Domain Segregation and DNA Methylation

In the earlier study we showed that differences in the methylation levels between

forests and prairies correlate well with chromatin spatial packing(Liu et al., 2018, 1). In the

following, we analyzed methylation data for early embryonic cells obtained by four

different research groups (see Additional file 2: Data sources). These data all show that the

open sea (defined as the genomic regions excluding CGIs, CGI shores and CGI

shelves(Sandoval et al., 2011)) methylation differences between forest and prairie domains

in different cell types correlate well to their corresponding chromatin structural segregation

behaviors (Figures 3A and S4A-S4C). During embryonic development, the absolute value

of MDI (F-P open sea methylation difference index, see "Methods", domain possessing a

positive (negative) value indicates the methylation level of this domains is generally higher

(lower) than its two flanking domains, therefore higher absolute value indicates the larger

methylation level difference) increases from 2-cell to ICM stages, decreases at the ICM to

E6.5 epiblast stage, and increases again in the further development to the E7.5 stage

(especially for the E7.5 endoderm stage, Figure 3A). The variation of DNA methylation

difference resembles that of the chromatin structural segregation. The correspondence

between methylation difference and segregation degree during embryonic development

further supports a connection between this epigenetic mark and the chromatin structure. In

fact, such a correlation might have a simple explanation: Since forests contribute

dominantly to the more accessible chromatin regions, they are presumably more prone to

both DNA demethylation and methylation than prairies.

Following such an observation, the extent of domain segregation of three germ layers

can be inferred from the DNA methylation difference between the forest and prairie

domains. As can be seen from Figure 3A, the calculated absolute value of MDI for

endoderm is greater than that for mesoderm (The corresponding value of ectoderm is

slightly less than mesoderm). This result suggests that the forest-prairie domains of

endoderm is more segregated than those of mesoderm and ectoderm, which is indeed

accordant with the analysis on Hi-C data (unpublished results). Interestingly, domain

segregation levels of the different tissues can be seen to reflect the germ layers they

originate from, as shown below from the analyses of both DNA methylation and structural

data.

It was found based on the Hi-C data that ectoderm-derived cortex chromatin is less

segregated than that of endoderm-derived liver(Liu et al., 2018). Since the above analysis

suggests chromatin in ectoderm to be less segregated than endoderm (Figure 3A), we

decided to examine whether chromatin structures of differentiated tissues also show such

a trend, therefore exhibiting a germ layer dependence. We analyzed average methylation

difference levels of 138 mouse differentiated cells and 45 human differentiated cells (38,

34, 66 cells originate from endoderm, mesoderm and ectoderm respectively for mouse and

13, 19, 13 for human, see Additional file 2: Data sources)(The ENCODE Project

Consortium, 2012; Lister et al., 2013; Schultz et al., 2015). As shown in Figure 3B, the

absolute values of MDI for forest domains for human ectoderm-derived tissues are

significantly smaller than those for mesoderm- ($p = 4.938 \times 10^{-7}$ by two-sample t-test)

and endoderm-derived ($p = 5.696 \times 10^{-4}$ by two-sample t-test) tissues, in the same as

the segregation degree of the three embryonic germ layers. The data on prairie domains of

human tissues show the same trend, so do those on both forest and prairie domains of

mouse tissues (Figure S4D-S4F). To further validate this finding, we analyzed the Hi-C

data of 14 human tissues(Schmitt et al., 2016) (see Additional file 2: Data sources) and

compared the segregation ratio of these tissues derived from different germ layers. To

quantify the difference between two tissues, we defined and calculated the overall relative

segregated ratios (R_{or}) of one tissue over another, a positive value of which represents that

chromatin in the former is more segregated than that in the latter (see "Methods"). For

tissues derived from endoderm and mesoderm, their overall relative segregated ratios over

those originated from ectoderm (cortex and hippocampus) are generally positive (Figure

3C), indicating that chromatin of the former indeed tends to be more segregated than that

of the latter. Similarly, the overall relative segregation ratios of tissues derived from

endoderm over those derived from mesoderm also tend to be positive (Figures 3C and S4G).

Together, these results are all consistent with the notion that the segregation level of a
differentiated cell shows the corresponding germ layer signature, that is, the order of
segregation degrees for the three germ layers is in accordance with that of differentiated
tissues.

More interestingly, we found that the absolute values of MDI for brain decrease with
aging, and the trend is opposite for mature neutrophil cells, as shown in Figures 3D and 3E.
Earlier studies have found that DNA hypomethylation, which is more likely to occur in
prairie domains, correlates with the cell cycles. Partially methylated domains (PMDs) in
tumor cells are mainly composed of prairies(Xue et al., 2020) and PMD hypomethylation
increases with age, which appears to track the accumulation of cell divisions(Zhou et al.,
2018). Xuan Ming et al. also found that solo-WCGW sites display aging- and cancer-
associated hypomethylation, which exhibit low maintenance efficiency during cell
cycle(Ming et al., 2020). For chromatin structural changes, we compared the segregation
ratio in G1 stage to that in the late S ~ G2 stage, and found that forest and prairie tend to
become more separated in the late S ~ G2 stage (Xue et al., 2020). The enhanced
segregation supports that mitosis is conducive to a more segregated chromatin structure.
Therefore, we speculate that the different methylation patterns associated with aging
among different tissues may reflect their different cell division patterns: the ectoderm-
originated brain cells hardly divide, while liver cells constantly undergo cell cycles. The
observed MDI differences between these cells are consistent with their different dividing
patterns in life. Such a consistency makes an understanding of the mechanistic connection
between methylation and cell-division patterns highly desired.

ZGA and Associated 3D Genome Architecture Change

To further investigate how transcription is associated with chromatin structure, we
analyzed the chromatin structure changes from normal early 2-cell to late 2-cell at different
genomic distance scales, since ZGA typically occurs at this period. At small genomic
distances (<500 kb), the F-P ratio for late 2-cell is larger than that for early 2-cell, while P-
P ratio is smaller for the former (Figure 4A, upper two figures). Our previous work has
revealed that prairie-forest intermingling was associated with prairie gene activation(Liu
et al., 2018), therefore, we speculate that the increase of F-P ratio may be associated with

ZGA. At large distances (500 kb-~20 Mb), the P-P ratio (F-P ratio) of late 2-cell was larger (smaller) than early 2-cell (Figure 4A, upper two figures), indicating that at such scale, the forest-prairie separation is enhanced, accordant with the increase of compartmentalization degree(Du et al., 2017) (given that compartment A and B are mainly composed of forest and prairie regions, respectively). Further, compared to early 2-cell, chromatin of late 2-cell establishes more spatial interactions over longer distances (>50 Mb, Figure 4A, down two figures). However, the compartmentalization is weak at this scale (compared to the F-P/P-P ratio of ICM in the same range (i.e., >50 Mb) and late 2-cell itself in small range (i.e., < 50 Mb), Figure 4A, down two figures).

LADs, which resemble compartment B, are established quickly after fertilization. More than 75% of LAD domains are prairies across preimplantation development in mouse and the domain segregation ratio (DR_s) of prairie domains overlapping with LAD is significantly larger than those that are not (Figure 4B, $p = 2.9 \times 10^{-34}$ and 0.01 for zygote and 2-cell, respectively, Welch's unequal variance test). We further analyzed the correlation between LAD mixing and prairie/forest mixing at the ZGA stage in mouse. LAD/iLAD (inter-LAD) regions in zygote and 2-cell stage were obtained from previous studies(Borsos et al., 2019). We then identified forest/prairie domains that become significantly more intermingled or segregated (i.e., $F_{mix}/F_{seg}/P_{mix}/P_{seg}$, see "Methods") during ZGA. The number of forest domains switching from iLAD to LAD state is 139, half of which (70 domains) belong to F_{mix}. In contrast, the proportion of F_{mix} domains within forest domains remaining in the iLAD state is only 24.8% ($p = 1 \times 10^{-7}$ by Fisher's exact test). Accordingly, forests that change from iLAD to LAD regions show a significant decrease in domain segregation ratios (Figure 4C, $p = 6.7 \times 10^{-5}$ by two-sample t-test). Such a result indicates that the forests changing from iLAD to LAD gain contacts with prairies, and a correlation does exist between LAD mixing and forest mixing during ZGA.

We next examine how compartment switch is associated with ZGA. Firstly, we examined the compartment change during embryonic development. The overall length of genomic domains changing from compartment A to B are shorter than those switching from compartment B to A in ZGA (Figure 4D). Such a phenomenon was also observed in implantation (from ICM to E6.5Epi). In addition, the prairie genes that move from compartment B to A in ZGA are found to be rich in functional annotation of "defense

response", "signal transduction", "cell membrane", "negative regulation of cell

differentiation" and "positive regulation of cell proliferation," identified using DAVID

(Huang et al., 2008, 2009) (see "Methods", Figure 4E). These results indicate that the

domain mixing during ZGA is heavily related to functions of cell proliferation.

Implantation-related Domain Mixing in Differentiation

Since it was shown that the forest-prairie inter-domain interaction coincides with

tissue-specific gene activation in differentiation(Liu et al., 2018), one wonders how the

achievements of tissue-specific functions are reflected by domain mixing in implantation.

We again calculated domain segregation ratio DR_s for each forest/prairie domain at

related embryonic developmental stages (i.e., ICM and E6.5Epi), and observed that DR_s

decreases for 89.0% of prairie domains at implantation (Figure 5A). To be more specific,

we identified DNA domains that undergo significant changes of segregation, and denoted

more strongly segregated and more mixed domains in forest and prairie as F_{seg}, P_{seg},

F_{mix}, and P_{mix}, respectively (see "Methods"). In the implantation stage, 71.0% of prairie

domains belong to P_{mix}, while 21.3% of forest domains become more mixed (Figure 5B).

As for P_{mix} regions following implantation, we first examined their structural

changes by calculating the compartment index (C-index) changes, a larger value of which

corresponds to a more compartment A-like environment (see "Methods"). Along with the

ICM to E6.5 epiblast transition, 86.9% of P_{mix} domains have an increase of C-index

(entering a more compartment A-like environment), implying that genes in these regions

move to a more active environment (Figure 5C). We analyzed next the functional

enrichment for genes located in the P_{mix} domains using ClusterProfiler(Yu et al., 2012)

and DAVID(Huang et al., 2008, 2009) (see "Methods"). The associated P_{mix} genes are

characterized by "sensory perception of taste", "regulation of lactation", "keratinocyte

differentiation", "epidermal cell differentiation", "epidermis development", "skin

development", "mammary gland development" and "long-term synaptic potentiation"

(Figure 5D). Interestingly, these terms are related to mammary glands, epidermis and

nervous system, all of which are differentiated from ectoderm(Li et al., 2013), indicating

that the mixing of prairie domains with forests during implantation is highly pertinent to

ectoderm differentiation. A supporting observation is that chromatin structures of E6.5

epiblast and E7.5 ectoderm are similar as seen by their similar DR_s values (Figure 5E). Such an observation suggests that chromatin enters an ectoderm-like state shortly after implantation, consistent with earlier studies which showed the primed ESCs, representing a post-implantation epiblast cell state, to subsequently develop towards the ectoderm(Ji et al., 2016). In addition, during the implantation process, the expression level of 309 prairie genes increases, while 162 prairie genes show decreased expression level. Here we further found that prairie domains harboring genes with elevated expression levels tend to become more forest-prairie mixed from ICM to E6.5Epi, compared to genes with lowered expression (Figure S5A, $p = 4.163 \times 10^{-4}$ by two-sample t-test), consistent with our previous finding on the vital role of forest-prairie spatial interactions in prairie gene activation(Liu et al., 2018). Moreover, we found that P_{mix} genes with increased expression tend to be enriched in neuro-related functions (Figure S5B), indicating that neuroectoderm may differentiate ahead of epidermal ectoderm. Finally, to confirm the roles of domain segregation change on biological functions, we analyzed the structural changes of domains containing lineage-specific genes. We identified 552 lineage-specific genes for E6.5 epiblast stage (see "Methods", i.e., genes specifically and highly expressed in E6.5 epiblast stage), among which 474 genes locate in forest domains and 78 genes in prairie domains. Ninety-seven percent of the prairie lineage-specific genes located in P_{mix}, suggesting that the mixing of prairies into forests does associate with lineage specification during implantation. Meanwhile, the prairie genes that move from compartment B to A during implantation are also functionally enriched in ectoderm differentiation and utero embryonic development (Figure S5C), in accordance with above analysis on P_{mix} regions.

DISCUSSION

Sequence-based Chromatin Domain Segregation

In the present study, we described the chromatin structural changes during embryonic development from zygote to the post-implantation stages and, in particular, how DNA sequence affects the chromatin structural changes. We also investigated the relationship between DNA domain segregation and the genetic/epigenetic properties. Earlier studies have shown gradual compartmentalization in embryonic development(Du et al., 2017). Here we connect the 3D structure change to DNA sequence properties and perform

functional analysis of these structural changes. The sequence-structure-function analysis

provides a common-ground for the understanding of different biological processes. In

particular, two types of genomic domains defined earlier, forests and prairies, are shown to

generally undergo an enhanced spatial separation in early embryo development of both

mouse and human, which is also reflected in the DNA methylation and gene expression

difference between the two domain types. Interestingly, noticeable domain mixing does

occur at two important stages, ZGA, when zygotic genes begin to express, and implantation,

during which differentiation starts to form germ layers. During ZGA process, short-range

(<500 kb) forest-prairie spatial interactions increase, which was thought to be associated

with gene activation since our previous work has revealed that prairie genes tend to move

to a more forest environment for activation (Tian et al., 2020). Besides, we also investigated

in detail the chromatin structural changes during ZGA process at different scales. During

implantation, we observed a conspicuous decrease of segregation ratio $R_s(d)$ for prairies

at large genomic distances (>3 Mb). Intriguingly, almost all E6.5 epiblast-specific prairie

genes resided in P_{mix} domains, again emphasizing the intimate link between domain

mixing and gene activation (lineage specification). Genes in the prairie domains that

become more mixed in implantation are prominently related with ectoderm differentiation.

Analysis of Hi-C data in differentiated and senescent cells (Liu et al., 2018), showed

a consistently enhanced segregation of chromatin structures, with a gradual establishment

of long-range DNA contacts from zygote to senescence. At the zygotic stage, few high

order structural features exist and the chromatin is organized similarly to a random coil

(Figure 6A). As the embryo develops, local structures (loops and TADs) become more

prominent and the two different types of domains segregate from each other to form

compartments. Such a trend continues through ICM (Figure 6B). After implantation and as

differentiation starts, a subset of prairie domains tends to mix into the active environment,

activating associated genes (Figure 6C). During senescence, prairie domains congregate

further, some of which detach from nuclear membranes and cluster inside the nucleus,

permitting long-range contacts establishment(Chandra et al., 2015) (Figure 6D). Previous

analysis showed that chromatin domain segregation continues in cell differentiation and

senescence(Liu et al., 2018, 1). It appears that the overall trend of chromatin structure

change follows an establishment of long-range contacts from birth to senescence, as seen

from the segregation level change (Figures S6A-S6C). In this process, intra-domain contacts are established first, followed by long-range inter-domain contact formation, and then the establishment of lineage-specific inter-domain contacts, which relates with cell differentiation (Figures S6D-S6G).

The Association between Transcription Inhibition and Genome Architecture

To investigate how transcription inhibition affects chromatin architecture, we compared the F-P and P-P ratio within the top 10% contact probabilities under one certain genomic distance between PN3, normal late 2-cell and α-amanitin-treated-20h cell (α-amanitin-treated late 2-cell). The critical genomic distance of α-amanitin-treated-20h cell is significantly smaller than that of normal late 2-cell, and the F-P and P-P ratio of α-amanitin-treated-20h cell are between PN3 and normal late 2-cell (Figure S7A), indicating that transcription inhibition can slow down the establishment of long-range contact and forest-prairie separation. We further compared the F-P and P-P ratio between α-amanitin-treated-20h cell and α-amanitin-treated-45h cell, and found that the critical genomic distance and P-P ratio of the latter are larger than the former, the F-P ratio of the latter is smaller than the former (Figure S7B), hinting that when transcription is inhibited, long-range chromatin contacts can still be established and forest-prairie separation also progresses, although at a reduced rate.

Other Factors that May Relate to Chromatin Structure Change

Sequence-based chromatin domain segregation suggests that factors effecting domain interactions could lead to functional chromatin structure changes. Besides signaling and transcriptional factors, physical parameters could also be important. Interestingly, a correlation seems to exist between chromatin domain segregation and body temperature during early embryonic development. For human, the variation of BBT (basal body temperature) corresponds closely to the change in domain segregation. Firstly, BBT rises 0.5~1°F immediately after ovulation and remains high throughout the luteal phase, even after implantation(Su et al., 2017). The segregation ratio R_s increases along with the development from zygote to ICM, except for the early to late 2-cell stage, which coincides with a small temperature dip nearly two days after ovulation, nicely corresponding to the

mixing of chromatin domains observed at this stage. Moreover, a one-day implantation dip of BBT occurs nearly one week after ovulation. Correspondingly, a significant domain mixing is observed at this stage of development, from ICM to E6.5 epiblast. As widely known, implantation dip is effective in detecting pregnancy in humans(Refinetti and Menaker, 1992) and continuous BBT measurement reliably allows for the earliest and most accurate detection of pregnancy(Smarr et al., 2016). For mouse, researchers did find that an ultimately-successful pregnancy is characterized by significantly more min/day of relative low CBT (core body temperature) in the second day after mouse pairing which rightly corresponds to ZGA, than for pregnancies which subsequently terminated prematurely(Smarr et al., 2016). These observations give rise to the possibility that temperature shifts are closely coupled with large scale chromatin structural changes during embryonic development, in accordance with high temperatures to be favorable for segregation of the prairies, and low temperatures, for a more forest-prairie mixed chromatin structure(Liu et al., 2018; Quan et al., 2020).

Besides human and mouse, temperature is also found to be important in embryonic development for many other species. One interesting example is the BBT of hibernating bears. The body temperature of the pregnant female bears decreases at the implantation stage, but remains 3-4 degrees higher and significantly more stable than non-pregnant females or males(Friebe et al., 2014), indicating the importance of a high and steady temperature in bear embryo development. In addition, researchers also reported chromosome de-condensation in rice seedlings underwent cold stress(Liu et al., 2017). All these results are consistent with the postulation that an increase of temperature correlates with an enhanced chromatin domain segregation, and imply that temperature could be a non-negligible factor influencing global chromatin structure, thus a possible parameter for affecting chromatin reprogramming or modification.

The domain segregation of the earliest fate-committed cell types also shows correlations with those of the differentiated cells. Tissues differentiated from endoderm generally have more strongly segregated chromatin structures than those from ectoderm. Such a difference between ectoderm and endoderm can be seen as early as from an E7.5 embryo. As another example, the tissue temperatures of interscapular brown adipose tissue (IBAT), liver, and the cortical (T_{IBAT}, T_L, T_{CO}, respectively) were found to follow such an

order: $T_{CO} \ll T_{IBAT} < T_L$(Dewasmes et al., 2003). This order of temperature values is

consistent with that of the corresponding chromatin domain segregation level, which

increases in the order of cortical, IBAT and liver. Furthermore, these tissues are known to

originate from ectoderm, mesoderm and endoderm, respectively, also consistent with the

degree of domain segregation of the different germ layers. Although much more data is

needed and many more factors such as protein structures and protein-DNA interactions are

likely involved in affecting the chromatin structure(Hnisz et al., 2017), the cause of such

correlations is worth of more detailed investigations.

Cell cycle is another important factor that could influence chromatin domain

segregation. During embryonic development, the establishment of TAD structure was

reported to require DNA replication but not zygotic transcription(Du et al., 2017; Ke et al.,

2017), implying the critical role of cell division in 3D structural formation. In fact, our

analysis shows that different cell division behaviors of different tissues including those

derived from different germ layers further increase their discrepancy in structural

segregation as well as in methylation pattern. The current and earlier analyses(Liu et al.,

2018) indicate that the DNA methylation are in affect associated closely to chromatin

conformation. The age related hypomethylation appears to track closely the accumulation

of cell divisions(Zhou et al., 2018). We speculate that along with hypomethylation (leading

to the formation of partially methylated domains), prairie domains tend to segregate more

stably in the less active spatial domains of heterochromatin(DEEP Consortium et al., 2018).

CONCLUSION

In this study, we delineated chromatin structural and epigenetic reprogramming in

early embryonic development based on the sequence-based domain segregation model, and

correlated the overall chromatin structural change with functional implementation in two

vital processes, ZGA and implantation. Genes within less segregated prairie domains

during implantation show lineage-relevant functions and thus may play important roles in

corresponding processes. Therefore, detailed analyses on gene expression and regulation

of transcription factors are in need for understanding the related mechanisms.

DATA AVAILABILITY STATEMENT

All data analyzed during this study are publicly available. The detailed data information can be found in Additional file 2 Data Sources.

FUNDING

This work was supported by National Natural Science Foundation of China (92053202, 22050003).

ACKNOWLEDGEMENTS

We would like to thank Prof. Xiaoliang S. Xie for helpful discussions, Zhenhai Du for the initial processing of Hi-C data, and Yupeng Huang for helping us to draw figure 6.

AUTHOR CONTRIBUTIONS

Conceptualization, Y.Q.G., H.Q., H.T. and S.L.; Methodology, Y.Q.G., H.Q., S.L. and H.T.; Software, H.Q., S.L., Y.X. and H.T.; Validation, W.X. and Y.Z.; Formal Analysis, H.Q., S.L. Y.X. and H.T.; Investigation, W.X. and Y.Z.; Writing-Original Draft, H.Q. and H.T., Writing-Review & Editing, Y.Q.G., S.L., H.T., Y.Z. and W.X.; Visualization, H.Q., X.Y. and H.T; Supervision, Y.Q.G.; Funding Acquisition, Y.Q.G.

DECLARATION OF INTERESTS

The authors declare that they have no competing interests.

SUPPLEMENTARY MATERIALS

Additional file 1:

Figures S1-S7.

Additional file 2:

Tables S1 provides a summary of all used data.

Table S2-8 contains information of analyzed mouse Hi-C data, human Hi-C data, mouse embryo methylation data, mouse tissue methylation data, human tissue methylation data, mouse expression data and mouse epigenetic data, respectively.

Additional file 3:

CGI forest definition: 1cluster_by_nearby_dist_m.m;

distance-dependent segregation ratio calculation: 2Rs(d)_calculation.m;

MDI calculation: 3mm9_embyo_diff_pf_os.m;

overall and stringent relative segregation ratio: 4overall_stringent_Rs.m;

domain segregation ratio calculation: 5DRs_calculation.m;

compartment index calculation: 6Comp_index_implan.m

REFERENCES

Arnold, S. J., and Robertson, E. J. (2009). Making a commitment: cell lineage allocation
and axis patterning in the early mouse embryo. *Nat. Rev. Mol. Cell Biol.* 10, 91–
103. doi:10.1038/nrm2618.

Bedzhov, I., and Zernicka-Goetz, M. (2014). Self-organizing properties of mouse
pluripotent cells initiate morphogenesis upon implantation. *Cell* 156, 1032–1044.
doi:10.1016/j.cell.2014.01.023.

Borsos, M., Perricone, S. M., Schauer, T., Pontabry, J., de Luca, K. L., de Vries, S. S., et al.
(2019). Genome–lamina interactions are established de novo in the early mouse
embryo. *Nature* 569, 729–733. doi:10.1038/s41586-019-1233-0.

Burton, A., and Torres-Padilla, M.-E. (2010). Epigenetic reprogramming and development:
a unique heterochromatin organization in the preimplantation mouse embryo. *Brief
Funct Genomics* 9, 444–454. doi:10.1093/bfgp/elq027.

Burton, A., and Torres-Padilla, M.-E. (2014). Chromatin dynamics in the regulation of cell
fate allocation during early embryogenesis. *Nat. Rev. Mol. Cell Biol.* 15, 723–735.
doi:10.1038/nrm3885.

Chandra, T., Ewels, P. A., Schoenfelder, S., Furlan-Magaril, M., Wingett, S. W., Kirschner,
K., et al. (2015). Global reorganization of the nuclear landscape in senescent cells.
Cell Rep. 10, 471–483. doi:10.1016/j.celrep.2014.12.055.

Chen, X., Ke, Y., Wu, K., Zhao, H., Sun, Y., Gao, L., et al. (2019). Key role for CTCF in
establishing chromatin structure in human embryos. *Nature* 576, 306-310.
doi:10.1038/s41586-019-1812-0.

Dahl, J. A., Jung, I., Aanes, H., Greggains, G. D., Manaf, A., Lerdrup, M., et al. (2016).
Broad histone H3K4me3 domains in mouse oocytes modulate maternal-to-zygotic
transition. *Nature* 537, 548–552. doi:10.1038/nature19360.

DEEP Consortium, Salhab, A., Nordström, K., Gasparoni, G., Kattler, K., Ebert, P., et al.
(2018). A comprehensive analysis of 195 DNA methylomes reveals shared and cell-

specific features of partially methylated domains. *Genome Biol.* 19, 150.
doi:10.1186/s13059-018-1510-5.

Dewasmes, G., Loos, N., Delanaud, S., Ramadan, W., and Dewasmes, D. (2003). Liver
temperature during sleep. *Sleep* 26, 948–950. doi:10.1093/sleep/26.8.948.

Du, Z., Zheng, H., Huang, B., Ma, R., Wu, J., Zhang, X., et al. (2017). Allelic
reprogramming of 3D chromatin architecture during early mammalian
development. *Nature* 547, 232–235. doi:10.1038/nature23263.

Falk, M., Feodorova, Y., Naumova, N., Imakaev, M., Lajoie, B. R., Leonhardt, H., et al.
(2019). Heterochromatin drives compartmentalization of inverted and conventional
nuclei. *Nature* 570, 395-399. doi:10.1038/s41586-019-1275-3.

Friebe, A., Evans, A. L., Arnemo, J. M., Blanc, S., Brunberg, S., Fleissner, G., et al. (2014).
Factors affecting date of implantation, parturition, and den entry estimated from
activity and body temperature in free-ranging brown bears. *PLoS One* 9, e101410.
doi:10.1371/journal.pone.0101410.

Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K., and Sharp, P. A. (2017). A phase
separation model for transcriptional control. *Cell* 169, 13–23.
doi:10.1016/j.cell.2017.02.007.

Huang, D. W., Sherman, B. T., and Lempicki, R. A. (2008). Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* 4,
44–57. doi:10.1038/nprot.2008.211.

Huang, D. W., Sherman, B. T., and Lempicki, R. A. (2009). Bioinformatics enrichment
tools: paths toward the comprehensive functional analysis of large gene lists.
Nucleic Acids Res. 37, 1–13. doi:10.1093/nar/gkn923.

Ji, X., Dadon, D. B., Powell, B. E., Fan, Z. P., Borges-Rivera, D., Shachar, S., et al. (2016).
3D chromosome regulatory landscape of human pluripotent cells. *Cell Stem Cell*
18, 262–275. doi:10.1016/j.stem.2015.11.007.

Jukam, D., Shariati, S. A. M., and Skotheim, J. M. (2017). Zygotic genome activation in
vertebrates. *Dev. Cell* 42, 316–332. doi:10.1016/j.devcel.2017.07.026.

Ke, Y., Xu, Y., Chen, X., Feng, S., Liu, Z., Sun, Y., et al. (2017). 3D chromatin structures
of mature gametes and structural reprogramming during mammalian
embryogenesis. *Cell* 170, 367-381.e20. doi:10.1016/j.cell.2017.06.029.

Lawson, K. A., Meneses, J. J., and Pedersen, R. A. (1991). Clonal analysis of epiblast fate
during germ layer formation in the mouse embryo. *Development* 113, 891–911.

Lee, H. J., Hore, T. A., and Reik, W. (2014). Reprogramming the methylome: erasing
memory and creating diversity. *Cell Stem Cell* 14, 710–719.
doi:10.1016/j.stem.2014.05.008.

Li, L., Liu, C., Biechele, S., Zhu, Q., Song, L., Lanner, F., et al. (2013). Location of
transient ectodermal progenitor potential in mouse development. *Development* 140,
4533–4543. doi:10.1242/dev.092866.

Lieberman-Aiden, E., van Berkum, N. L., Williams, L., Imakaev, M., Ragoczy, T., Telling,
A., et al. (2009). Comprehensive mapping of long range interactions reveals folding
principles   of   the   human   genome.   *Science*   326,   289–293.
doi:10.1126/science.1181369.

Lister, R., Mukamel, E. A., Nery, J. R., Urich, M., Puddifoot, C. A., Johnson, N. D., et al.
(2013). Global epigenomic reconfiguration during mammalian brain development.
Science 341, 1237905. doi:10.1126/science.1237905.

Liu, C., Cheng, Y.-J., Wang, J.-W., and Weigel, D. (2017). Prominent topologically
associated domains differentiate global chromatin packing in rice from Arabidopsis.
Nat. Plants 3, 742–748. doi:10.1038/s41477-017-0005-9.

Liu, S., Zhang, L., Quan, H., Tian, H., Meng, L., Yang, L., et al. (2018). From 1D sequence
to 3D chromatin dynamics and cellular functions: a phase separation perspective.
Nucleic Acids Res. 46, 9367-9383. doi:10.1093/nar/gky633.

Liu, X., Wang, C., Liu, W., Li, J., Li, C., Kou, X., et al. (2016). Distinct features of
H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos.
Nature 537, 558–562. doi:10.1038/nature19362.

Lu, F., Liu, Y., Inoue, A., Suzuki, T., Zhao, K., and Zhang, Y. (2016). Establishing
chromatin regulatory landscape during mouse preimplantation development. *Cell*
165, 1375–1388. doi:10.1016/j.cell.2016.05.050.

Ming, X., Zhang, Z., Zou, Z., Lv, C., Dong, Q., He, Q., et al. (2020). Kinetics and
mechanisms of mitotic inheritance of DNA methylation and their roles in aging-
associated methylome deterioration. *Cell Res*. doi:10.1038/s41422-020-0359-9.

Morgan, H. D., Santos, F., Green, K., Dean, W., and Reik, W. (2005). Epigenetic
reprogramming   in   mammals.   *Hum.   Mol.   Genet.*   14,   R47-58.
doi:10.1093/hmg/ddi114.

Quan, H., Yang, Y., Liu, S., Tian, H., Xue, Y., and Gao, Y. Q. (2020). Chromatin structure
changes during various processes from a DNA sequence view. *Curr. Opin. Struct.*
Biol. 62, 1–8. doi:10.1016/j.sbi.2019.10.010.

Refinetti, R., and Menaker, M. (1992). The circadian rhythm of body temperature. *Physiol.*
Behav. 51, 613–637. doi:10.1016/0031-9384(92)90188-8.

Sandoval, J., Heyn, H., Moran, S., Serra-Musach, J., Pujana, M. A., Bibikova, M., et al.
(2011). Validation of a DNA methylation microarray for 450,000 CpG sites in the
human genome. *Epigenetics* 6, 692–702. doi:10.4161/epi.6.6.16196.

Schmitt, A. D., Hu, M., Jung, I., Xu, Z., Qiu, Y., Tan, C. L., et al. (2016). A compendium
of chromatin contact maps reveals spatially active regions in the human genome.
Cell Rep. 17, 2042–2059. doi:10.1016/j.celrep.2016.10.061.

Schultz, M. D., He, Y., Whitaker, J. W., Hariharan, M., Mukamel, E. A., Leung, D., et al.
(2015). Human body epigenome maps reveal noncanonical DNA methylation
variation. *Nature* 523, 212–216. doi:10.1038/nature14465.

Servant, N., Varoquaux, N., Lajoie, B. R., Viara, E., Chen, C.-J., Vert, J.-P., et al. (2015).
HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol.*
16, 259. doi:10.1186/s13059-015-0831-x.

Smarr, B. L., Zucker, I., and Kriegsfeld, L. J. (2016). Detection of successful and
unsuccessful pregnancies in mice within hours of pairing through frequency
analysis of high temporal resolution core body temperature data. *PLoS One* 11,
e0160127. doi:10.1371/journal.pone.0160127.

Smith, Z. D., and Meissner, A. (2013). DNA methylation: roles in mammalian development.
Nat. Rev. Genet. 14, 204–220. doi:10.1038/nrg3354.

Su, H.-W., Yi, Y.-C., Wei, T.-Y., Chang, T.-C., and Cheng, C.-M. (2017). Detection of
ovulation, a review of currently available methods. *Bioeng Transl Med* 2, 238–246.
doi:10.1002/btm2.10058.

Svoboda, P., Franke, V., and Schultz, R. M. (2015). Sculpting the transcriptome during the
oocyte-to-embryo transition in mouse. *Curr. Top. Dev. Biol.* 113, 305–349.
doi:10.1016/bs.ctdb.2015.06.004.

The ENCODE Project Consortium (2012). An integrated encyclopedia of DNA elements
in the human genome. *Nature* 489, 57–74. doi:10.1038/nature11247.

Tian, H., Yang, Y., Liu, S., Quan, H., and Gao, Y. Q. (2020). Toward an understanding of
the relation between gene regulation and 3D genome organization. *Quant. Biol.* 8,
295-311. doi:10.1101/2020.01.13.903872.

Wang, C., Liu, X., Gao, Y., Yang, L., Li, C., Liu, W., et al. (2018). Reprogramming of
H3K9me3-dependent heterochromatin during mammalian embryo development.
Nat. Cell Biol. 20, 620–631. doi:10.1038/s41556-018-0093-4.

Xie, W. J., Meng, L., Liu, S., Zhang, L., Cai, X., and Gao, Y. Q. (2017). Structural modeling
of chromatin integrates genome features and reveals chromosome folding principle.
Sci. Rep. 7, 2818. doi:10.1038/s41598-017-02923-6.

Xie, W., Schultz, M. D., Lister, R., Hou, Z., Rajagopal, N., Ray, P., et al. (2013).
Epigenomic analysis of multilineage differentiation of human embryonic stem cells.
Cell 153, 1134–1148. doi:10.1016/j.cell.2013.04.022.

Xu, Q., and Xie, W. (2018). Epigenome in early mammalian development: inheritance,

873 reprogramming and establishment. *Trends Cell Biol.* 28, 237–253.
874  doi:10.1016/j.tcb.2017.10.008.

875 Xue, Y., Yang, Y., Tian, H., Quan, H., Liu, S., Zhang, L., et al. (2020). Domain segregated
876  3D chromatin structure and segmented DNA methylation in carcinogenesis.
877  *bioRxiv* doi:10.1101/2020.01.13.903963.

878 Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for
879  comparing biological themes among gene clusters. *OMICS* 16, 284–287.
880  doi:10.1089/omi.2011.0118.

881 Zhang, B., Zheng, H., Huang, B., Li, W., Xiang, Y., Peng, X., et al. (2016). Allelic
882  reprogramming of the histone modification H3K4me3 in early mammalian
883  development. *Nature* 537, 553–557. doi:10.1038/nature19361.

884 Zhang, Y., Xiang, Y., Yin, Q., Du, Z., Peng, X., Wang, Q., et al. (2018). Dynamic
885  epigenomic landscapes during early lineage specification in mouse embryos. *Nat.*
886  *Genet.* 50, 96–105. doi:10.1038/s41588-017-0003-x.

887 Zheng, H., Huang, B., Zhang, B., Xiang, Y., Du, Z., Xu, Q., et al. (2016). Resetting
888  epigenetic memory by reprogramming of histone modifications in mammals. *Mol.*
889  *Cell* 63, 1066–1079. doi:10.1016/j.molcel.2016.08.032.

890 Zheng, H., and Xie, W. (2019). The role of 3D genome organization in development and
891  cell differentiation. *Nat. Rev. Mol. Cell Biol*. 20, 535-550. doi:10.1038/s41580-019-
892  0132-4.

893 Zhou, W., Dinh, H. Q., Ramjan, Z., Weisenberger, D. J., Nicolet, C. M., Shen, H., et al.
894  (2018). DNA methylation loss in late-replicating domains is linked to mitotic cell
895  division. *Nat. Genet.* 50, 591–602. doi:10.1038/s41588-018-0073-4.

FIGURE LEGENDS

Figure 1. Domain segregation in early embryonic development
(A, B) The chromatin overall segregation ratio OR_s of forest and prairie domains at each stage in mouse (A) and human (B) embryonic development. Modelled chromatin structures are also shown at corresponding stages. The upper structures are colored by sequence from blue to red, and the lower structures are colored by forest/prairie domains.
(C, D) The domain segregation ratio DR_s of forest and prairie domains at each stage in mouse (C) and human (D) embryonic development.
(E) The distance-dependent segregation ratio $R_s(d)$ of forest and prairie at each stage in mouse embryonic development.
(F) The change of critical genomic distance during mouse embryonic development.
(G) The proportions of F-P and P-P spatial interactions within top 10% contact probabilities under one certain genomic distance at each stage in mouse embryonic development.

Figure 2. Compartment changes relate with domain segregation
(A) The proportions of four sequence components, Af, Bf, Ap, Bp (calculated according to the positioning of forests and prairies in compartments A and B), for each stage in the mouse embryonic development.
(B) The dynamic partition of prairie into compartment A and B in mouse embryonic development.
(C) Function annotation clustering of genes located in switch p, stable Ap and stable Bp regions, respectively.
(D) The PC1 values of stable Bp (i.e., prairie regions constantly remaining in compartment B in both 8-cell and ICM) and switchable p (i.e., prairie regions located in compartment B in 8-cell while switching to compartment A in ICM) regions in 8-cell.
(E) From 8-cell to ICM, the ATAC-seq, H3K4me3 and H3K27me3 signals of stable Bp and switchable p regions (which were explained in Figure 2D) in 8-cell and ICM.
(F) The average GC contents of the four components in mouse embryonic development.
(G) The relation between gene expression level and gene exon CpG density. Each data point represents the median expression level of one group of genes, the exon CpG density of which falls into one interval.
(H) The expression difference between forest and prairie genes at different stages during mouse embryonic development.

Figure 3. The association between domain segregation and DNA methylation
(A) The average forest/prairie open sea methylation level difference (MDI) at different stages during embryonic development.
(B) The box plots of average MDI for human differentiated cells originating from endoderm, mesoderm and ectoderm, respectively, for forest domains.
(C) The box plots of overall relative segregation ratio R_{or} (the definition can be seen below) of one tissue over another for forests. The three categories are tissues derived from endoderm and mesoderm over those derived from ectoderm, and endoderm over mesoderm. The parameter R_{or} was used to evaluate the differences of domain segregation behaviors between two tissues. For example, as for tissues 1 and 2, N_{12} and N_{21} represent the number of domains showing higher DR_s in tissues 1 and 2, respectively. Overall relative segregation ratio for tissues 1 and 2 is then defined as $\ln(\frac{N_{12}}{N_{21}})$, a positive value of which thus indicates the chromatin structure of tissue 1 is more segregated, compared to tissue 2.
(D, E) The scatter plots of age and methylation difference for brain cells (D) and mature neutrophil cells (E) for forest domains.

947
Figure 4. ZGA and associated 3D genome architecture change
(A) The upper two figures represent the proportion of F-P and P-P spatial interactions within the
top 10% contact probabilities under one certain genomic distance, for normal early 2-cell, late 2-
cell. The lower two figures were the amplifications of the partial regions of two upper figures and
ICM was also included for comparison. Inner plot represents the critical genomic distance of early
2-cell and late 2-cell.
(B) The domain segregation ratio, DR_s, of prairie domains that are/are not in LAD regions.
(C) The domain segregation ratio of forest domains switching from iLAD to LAD during ZGA, in
early 2-cell and late 2-cell.
(D) The proportion of four components (A→A, A→B, B→A, B→B) in each transition (e.g., from
early 2-cell to late 2-cell). The length ratio of e.g., AA for PN3->PN5 indicates the ratio between
the genome length of stable A regions (between PN3 and PN5) and the entire genome length.
(E) Function annotation clustering of prairie genes switching from B to A during ZGA.
961
Figure 5. Implantation-related domain mixing in differentiation
(A) The scatter diagrams of domain segregation ratio DR_s for forests and prairies during
implantation.
(B) The distribution of the logarithm ratio of DR_s between E6.5Epi and ICM for forests and
prairies. The plot, the legend of which is "standard", represents the logarithm ratio of DR_s between
two ICM replicates from the same lab. Significantly more strongly segregated domains are taken
as those whose logarithm ratio of DR_s values exceed the 97.5th percentile of the "standard"
distribution, and accordingly, domains whose logarithm ratio of DR_s is smaller than the 2.5th
percentile of the "standard" distribution are regarded as significantly more mixed domains. The
corresponding threshold values are 0.1469 and -0.1469, respectively, which was labeled using two
black dotted lines.
(C) The change of compartment index (C-index) for less-segregated prairies, P_{mix}, from ICM to
E6.5 epiblast stage.
(D) Functional assignment of genes in the P_{mix} during implantation.
(E) The scatter plot of DR_s for forests and prairies between E6.5 epiblast and E7.5 ectoderm.
977
Figure 6. Schematic pictures of chromatin structural patterns at different stages.
(A) In the beginning zygotic stage, chromatin has little long-range structural features. Here the red
lines represent CGI forest domains, and the blue ones represent CGI prairie domains.
(B) In ICM stage, the two different types of domains segregate from each other to form
compartments.
(C) In differentiated cells, prairies containing tissue-specific genes tend to form contacts with
forests.
(D) During senescence, prairies congregate further (with an increased probability to detach from
the nuclear membrane).