Computational modeling reveals frequency modulation of calcium-cAMP/PKA pathway in dendritic spines

D. Ohadi[1], D. L. Schmitt[2], B. Calabrese[3], S. Halpain[3], J. Zhang[2], and P. Rangamani[1*]

[1]Department of Mechanical and Aerospace Engineering; [2]Department of Pharmacology; [3]Division of Biological Sciences and Sanford Consortium for Regenerative Medicine, University of California San Diego, La Jolla, CA, USA

[*]Corresponding Author, Department of Mechanical Aerospace Engineering, SME 344G, Jacobs School of Engineering, University of California San Diego, La Jolla, CA 92093-0411
Tel: +1 (858) 534-4734| Email: padmini.rangamani@eng.ucsd.edu

Abstract

Dendritic spines are the primary excitatory postsynaptic sites which act as subcompartments of signaling. Ca^{2+} is often the first and rapid signal. Downstream of calcium, the cAMP/PKA pathway plays a critical role in the regulation of spine formation, morphological modifications, and ultimately, learning and memory. While the dynamics of calcium are reasonably well-studied, calcium-induced cAMP/PKA dynamics are not fully explored. In this study, we present a well-mixed model for the dynamics of calcium-induced cAMP/PKA dynamics in dendritic spines. The model is validated against experimental measurements in the literature. Further, we measure calcium dynamics in dendritic spines of cultured hippocampal CA1 neurons to justify the choice of model inputs. Our model predicts that the various steps in this pathway act as a frequency modulator for calcium and high frequency of calcium input is filtered by AC1 and PDEs in this pathway such that cAMP/PKA only responds to lower frequencies. This prediction has important implications for noise filtering and long-timescale signal transduction in dendritic spines. A companion manuscript presents a three-dimensional spatial model for the same pathway.

Abbreviations

CaM: Calmodulin; AC1: Adenylyl cyclase 1; ATP: Adenosine triphosphate; cAMP: Cyclic adenosine monophosphate; PDE1: Phosphodiesterase 1; PDE4: Phosphodiesterase 4; PKA: Protein kinase A; IC: Initial concentration; ODE: Ordinary differential equation; LTP: Long-term potentiation; LTD: Long-term depression; AKAP: A-kinase anchoring protein; IP_3 receptor: Inositol trisphosphate receptor; FRET: Fluorescence resonance energy transfer.

Introduction

Intracellular calcium oscillations ranging from seconds to minutes are known to be responsible for distinct physiological roles in the nervous system [1–4]. Oscillatory gating dynamics of plasma membrane-embedded channels or cyclic calcium release from intracellular stores can generate calcium spikes [5]. Tightly coupled with calcium oscillations, through various signaling pathways, are the dynamics of another important second messenger, cAMP [5,6]. Changes in the amplitude and frequency of these oscillating signals along with the intracellular concentrations of these second messengers can trigger diverse cellular responses in neurons [5, 7–9]. However, the mechanisms by which these oscillations are tuned to drive particular responses in cells remain unclear.

Two types of enzymes are proposed to play important roles in driving parallel Ca^{2+} and cAMP oscillations: adenylyl cyclases (ACs) and phosphodiesterases (PDEs) [10]. ACs, which synthesize cAMP from ATP, are either directly or indirectly regulated by Ca^{2+} signaling. Different AC isoforms and their spatial organization in cells are thought to play an important role in regulating the spatiotemporal dynamics of cAMP [11]. AC1 is an isoform of AC that is highly expressed in the brain and is directly activated by Ca^{2+} through the calcium-calmodulin complex. Phosphodiesterases hydrolyze cAMP to AMP and are regulated by cAMP-dependent protein kinase-A (PKA) activity downstream of cAMP [12, 13]. The isoforms PDE1 and PDE4 are highly expressed in the hippocampus and are modulated by the calcium-calmodulin complex and phosphorylation by PKA respectively [14].

The primary effector of cAMP, PKA, is a heterotetrameric protein kinase consisting of two catalytic and two regulatory subunits. Postsynaptically, calcium influx through N-methyl-D-aspartate receptors (NMDAR) triggers the synthesis of cAMP by ACs and induces PKA activity. PKA activation plays an important role in long-term potentiation and depression [15] and neurite outgrowth [16], resulting in memory formation [17] and emotional expression, among other cognitive processes [18]. Within dendritic spines, PKA is enriched following stimulation of upstream signaling pathways [19, 20] and structural plasticity [21]. Spinal PKA activity has also been associated with maintenance of synaptic strength [20]. Despite the critical role of PKA in signal transduction and physiological response within dendritic spines, little is known about how calcium-cAMP/PKA crosstalk occurs in dendritic spines.

In this work, we built a mathematical model of calcium-induced cAMP/PKA dynamics to integrate different key steps in the modulation of signaling components in the cAMP/PKA pathway. This model was constrained using experimental data available in the literature. Our model predicts that components in this pathway serve as frequency modulators of calcium dynamics such that cAMP/PKA only picks up the lower frequency of calcium influx; that is, cAMP/PKA is a leaky integrator of calcium dynamics. This finding sheds light on the kinetic mechanisms for robustness of cAMP/PKA responses in dendritic spines. Experimental measurements of calcium oscillations in dendritic spines show that calcium oscillates spontaneously in spines and these measurements are used to generate predictions of cAMP/PKA kinetics for different calcium inputs.

Materials and Methods

Experimental methods

Plasmids: GCaMP6f was purchased from Addgene.

Ethical Approval: All animal experiments in this study were conducted in accordance with the National Institutes of Health Guide for the care and use of laboratory animals and all protocols were approved by the University of California San Diego Institutional Animal Care and Use Committee. All surgical procedures were terminal and isoflurane was used as an anesthetic to prevent any pain or discomfort.

Rat primary hippocampal neuron culture: Hippocampi were dissected from E19 Sprague-Dawley male and female rat pups (Charles River) in ice-cold HEPES-buffered saline solution (Gibco), supplemented with pen-step

(Lifetech), glucose, and additional HEPES. Hippocampi were dissociated using papain (Worthington Biochemical). Cells were washed and resuspended in phenol red-free Neurobasal Medium (Lifetech) supplemented with L-glutamine (Stem Cell Technologies) and SM1 (NeuroCult, Stem Cell Technologies). Mixed neuron and astrocyte cell suspension was plated at a concentration of 200,000 cells/mL onto glass bottom 35 mm petri dishes (CellVis) coated with poly-L-lysine (Sigma). Cells were incubated in a humidified $37\,^{\circ}$C, CO_2 incubator (HeraCell). At DIV 20, cells were transfected with GCaMP6f using Lipofectamine 2000 (Invitrogen) for 48 hours, and imaged at DIV 22.

Live-cell imaging: Neurons were imaged in full medium using a resonant-scanning A1R confocal microscope (Nikon) equipped with a 60x 1.49 NA oil immersion objective, 489.1 nm laser, 525/50 nm emission filter, 488 dichroic, and DU4 GaASP detectors. Cells were kept at $37\,^{\circ}$C and 5% CO_2 using a stage-top incubation system (Okolab). Images of basal calcium dynamics were acquired using NIS Elements AR 5.10.01 at 4 frames per second for 4-5 minutes.

Image Analysis: Initial image analysis was done using NIS Elements AR Analysis 5.10.01. To measure calcium in spines, images were deconvoluted in 2D using 20 iterations of blind deconvolution with medium noise. Individual spines were identified and intensity measured over time. Fluorescence intensities were background corrected by subtracting intensity from a region with no transfected cells from spine intensities, and represented as $\Delta F/F$ where the minimum fluorescence intensity was subtracted from the instantaneous fluorescence intensity and divided by the minimum fluorescence intensity. Further processing was done using Microsoft Excel and MATLAB. Figure preparation was done using OMERO.

Mathematical Model Development

In this section, we list the model assumptions, identification of the key components in the pathway, and the governing equations.

Model assumptions

In order to simulate the temporal dynamics of calcium-stimulated cAMP/PKA pathway in dendritic spines, we developed a well-mixed model that accounts for the dynamics of the different biochemical species shown in Figure 1. We briefly discuss the assumptions made in developing this model below.

- **Compartment size:** We model the spine as a single compartment with a volume of 0.1 femtoliter (10^{-16} liter) based on the volume of a single dendritic spine [22]. All the biochemical species including the membrane-bound ones are well-mixed across this compartment. A separate spatial model is presented in [23].

- **Time scales:** We focus on cAMP/PKA dynamics at the minute timescale. Calcium input was prescribed as sinusoidal functions with exponential decays based on experimental observations in the literature [24] and our experimental measurements. Different calcium inputs are described in detail below.

- **Reaction types:** The catalytic reactions such as enzymatic reactions, phosphorylation by kinases, and dephosphorylation by phosphatases are modeled using Michaelis-Menten kinetics. All other reactions are modeled using mass action kinetics assuming that the concentrations are present in large quantities. The reactions and corresponding reaction rates are given in Table 1.

- **Enzyme regulations by calcium-calmodulin complex:** We assumed that the enzymes regulated by calcium-calmodulin, AC1 and PDE1 are fully activated only when all four binding sites of calmodulin are bound by calcium ions [25, 26]. These fully activated enzymes are shown as AC1·CaM (AC1·Ca$_4$·CaM) and PDE1·CaM (PDE1·Ca$_4$·CaM).

- **Kinetic parameters:** Kinetic parameters for these reactions were estimated using the COPASI simulator [27] and are given in Table 2. The model was constrained such that dose-response curves from the simulations matched the experimentally reported dose-response curves. To estimate the model parameters, we primarily used data from brain tissues. For AC1 activation by Ca^{2+} we used mouse hippocampus data [28]. For cAMP-ATP dynamics data from mouse neuroblastoma (N18TG2 cells) was used [29]. For PDE1 activation and PDE1/PKA phosphorylation, we used data from bovine brain (60 kDa) [30, 31]. In cases where data from brain tissues was not available, we used other cell types. For example, for PDE4/PKA phosphorylation, we used data from SF9 cells (spodoptera frugiperda Sf21 insect cells) [32].

Based on these assumptions, we constructed a well-mixed model of the calcium-induced cAMP/PKA pathway in dendritic spines.

Figure 1: Schematic of the neuronal cAMP-PKA pathway. The main biochemical components in the pathway modeled are shown. Each arrow depicts the corresponding reactions between the different biochemical components of the model. Dashed arrows represent enzymatic reactions modeled using Michaelis-Menten kinetics and solid arrows represent reactions modeled using mass action kinetics. The reaction numbers on each arrow correspond to the reactions listed in Table 1.

Key components and governing equations of the model

After identifying the key components of the cAMP pathway, using the existing experimental data in the literature, we designed the biochemical reactions. Based on the interactions between different signaling molecules, these reactions can be activation or inhibition and binding or unbinding with various types of kinetics. Some of these reactions are binding reactions that are modeled using either mass action or cooperative kinetics. Other are catalytic reactions also known as enzymatic activation or inhibition that are modeled using Michaelis-Menten kinetics. These reactions are designed in the six following modules:

1. **Ca^{2+}/CaM complex formation:** Calcium can enter the spine through NMDAR-mediated influx at the plasma membrane or through the activation of ryanodine and IP_3 receptors from internal stores [33]. Resting calcium is in the range of 0.05-0.1 μM and can rise to 10-100-fold when neurons are activated [34]. Calmodulin is a small protein involved in the regulation of many calcium-dependent events [35–37]. Calmodulin has two calcium binding sites at each N-terminus and C-terminus lobe with different Ca^{2+} affinities [38]. Calcium has a higher affinity for C-lobes than the N-lobes [39] but the N-lobe binds calcium faster [40]. Based

on these observations, we modeled the formation of the calcium-calmodulin complex using two sequential binding reactions with mass action kinetics. For simplicity, we did not consider the cooperative binding within each of these individual lobes [41]. Equation (1) accounts for binding of the first two calcium ions to the C-lobe resulting in the formation of $Ca_2 \cdot CaM$ and Equation (2) accounts for the binding of third and fourth calcium ions to the N-lobe resulting in the formation of $Ca_4 \cdot CaM$. Kinetic parameters of these reactions (kf_1, kb_1, kf_2, kb_2) were obtained by fitting experimental data [39] to the model in COPASI (Table 2, Figure 2A).

$$2\,Ca^{2+} + CaM \underset{kb_1}{\overset{kf_1}{\rightleftharpoons}} Ca_2 \cdot CaM, \qquad R_1 = k_{f1}[Ca^{2+}]^2[CaM] - k_{b1}[Ca_2 \cdot CaM]. \tag{1}$$

$$2\,Ca^{2+} + Ca_2 \cdot CaM \underset{kb_2}{\overset{kf_2}{\rightleftharpoons}} Ca_4 \cdot CaM, \qquad R_2 = k_{f2}[Ca^{2+}]^2[Ca_2 \cdot CaM] - k_{b2}[Ca_4 \cdot CaM]. \tag{2}$$

2. **AC1 activation by Ca^{2+}/CaM complex:** Calcium-calmodulin stimulated adenylyl cyclase 1 (AC1) plays an important role in learning, memory formation, and long-term potentiation by coupling calcium activity to cAMP in neurons [42]. Unlike other members of AC family, AC1 is not stimulated by G_s-coupled receptors unless it is activated by intracellular calcium [43]. AC1 responds to NMDA-mediated calcium elevations (and not calcium released from intracellular stores) and is mostly located close to the post-synaptic densities of dendritic spines [44]. AC1 is the predominant and neuro-specific calcium-stimulated adenylyl cyclase which is mostly expressed in dendrites, axons and at synapses [45]. At very high concentrations of calcium ($> 10\ \mu M$) AC1 is inhibited [46]. Our model takes this effect into account by fitting experimental data from [28] to the AC1 activity for steady state Ca^{2+} (Figure 2B). AC1 synthesizes cAMP from ATP and has been shown to be associated with hippocampal-dependent learning abilities [44]. We modeled AC1 activation by calcium as a two-step process. First, AC1 binds to $Ca_2 \cdot CaM$ by mass action kinetics to form $AC1 \cdot Ca_2 \cdot CaM$. Second, two other Ca^{2+} bind to $AC1 \cdot Ca_2 \cdot CaM$ in a cooperative binding reaction to form $AC1 \cdot Ca_4 \cdot CaM$. Only fully activated $AC1 \cdot Ca_4 \cdot CaM$ can catalyze cAMP production. The first term in R_4 accounts for cooperative kinetics, while the second term accounts for basal degradation. The parameters used in this module are listed in Table 2 and comparison against experimental data is shown in Figure 2B.

$$AC\,1 + Ca_2 \cdot CaM \underset{kb_3}{\overset{kf_3}{\rightleftharpoons}} AC\,1 \cdot Ca_2 \cdot CaM$$
$$R_3 = k_{f3}[AC1][Ca_2 \cdot CaM] - k_{b3}[AC1 \cdot Ca_2 \cdot CaM] \tag{3}$$

$$2\,Ca^{2+} + AC1 \cdot Ca_2 \cdot CaM \underset{kb_4}{\overset{kf_4}{\rightleftharpoons}} AC\,1 \cdot Ca_4 \cdot CaM$$
$$R_4 = \frac{k_{f4}[AC1 \cdot Ca_2 \cdot CaM][Ca^{2+}]}{K_{m4} + [Ca^{2+}]} - k_{b4}[AC1 \cdot Ca_4 \cdot CaM] \tag{4}$$

3. **PDE1 activation by Ca^{2+}/CaM complex:** Phosphodiesterase 1 (PDE1) is the other key enzyme that regulates calcium/cAMP interactions [26]. PDE1A (the PDE1 subtype used in our model) is widely expressed at high levels in the hippocampus [47]. PDE1 is the only type of phosphodiesterase that is calcium-calmodulin-activated [48, 49]. PDE1 becomes activated by intracellular calcium level increase and has low cAMP affinity in comparison with other members of phosphodiesterase family [50]. The phosphorylation of PDE1A (brain 60 kDa PDE1) by PKA increases the calcium concentration required for activation of PDE1A by decreasing the enzyme's affinity for calmodulin [51]. Unlike AC1, PDE1A does not discriminate between different sources of calcium [52]. We modeled PDE1 activation by calcium as a two-step process – first, PDE1 binds to $Ca_2 \cdot CaM$ by mass action kinetics to form $PDE1 \cdot Ca_2 \cdot CaM$. Second, another two Ca^{2+} bind to $PDE1 \cdot Ca_2 \cdot CaM$ in a cooperative binding reaction to form $PDE1 \cdot Ca_4 \cdot CaM$. Only fully activated

PDE1·Ca$_4$·CaM can hydrolyze cAMP into AMP. The first term in R_6 accounts for cooperative kinetics, while the second term accounts for basal degradation. The parameters used in this module are listed in Table 2 and comparison against experimental data from [30] is shown in Figure 2E.

$$\mathrm{PDE\,1} + \mathrm{Ca}_2 \cdot \mathrm{CaM} \xrightleftharpoons[\mathrm{kb}_5]{\mathrm{kf}_5} \mathrm{PDE\,1} \cdot \mathrm{Ca}_2 \cdot \mathrm{CaM}$$
$$R_5 = k_{f5}[PDE1][Ca_2 \cdot CaM] - k_{b5}[PDE1 \cdot Ca_2 \cdot CaM] \tag{5}$$

$$2\,\mathrm{Ca}^{2+} + \mathrm{PDE1} \cdot \mathrm{Ca}_2 \cdot \mathrm{CaM} \xrightleftharpoons[\mathrm{kb}_6]{\mathrm{kf}_6} \mathrm{PDE\,1} \cdot \mathrm{Ca}_4 \cdot \mathrm{CaM}$$
$$R_6 = \frac{k_{f6}[PDE1 \cdot Ca_2 \cdot CaM][Ca^{2+}]}{K_{m6} + [Ca^{2+}]} - k_{b6}[PDE1 \cdot Ca_4 \cdot CaM] \tag{6}$$

4. **cAMP production:** Adenylyl cyclases catalyze the conversion of ATP into cAMP. cAMP is implicated in both LTP and LTD by activating PKA [53–56]. In our model, fully activated AC1, AC1·Ca$_4$·CaM, catalyzes the formation of cAMP from ATP. This reaction is modeled by Michaelis-Menten kinetics with a basal degradation term. The enzyme is shown in bold type. The steady-state amount of generated cAMP with ATP is evaluated using experimental data from [29] shown in Figure 2C.

$$\mathrm{ATP} + \mathbf{AC1} \cdot \mathbf{Ca_4} \cdot \mathbf{CaM} \xrightarrow{\mathrm{Kcat}_7} \mathrm{cAMP}$$
$$R_7 = \frac{K_{cat7}[AC1 \cdot Ca_4 \cdot CaM][ATP]}{K_{m7} + [ATP]} - K_{r7}[cAMP] \tag{7}$$

5. **PKA activation:** PKA has two catalytic subunits and a regulatory dimer that binds to the catalytic subunits [57, 58]. RIIβ is the dominant PKA isoform expressed in the brain [59]. cAMP binds to its effector, PKA, and regulates the activity of regulatory subunits. Then, PKA phosphorylates different kinases responsible for neuron growth and differentiation, such as LKB1 and GSK-3β in the hippocampus [50]. Activation of PKA by cAMP is modeled in three steps inspired by [59] with some modifications in the order of A- and B-domain binding of cAMP that was observed in experiments by [60]. First, two cAMP molecules bind to the A-domain (Equation (8)), then the other two cAMP molecules bind to the B-domain (Equation (9)). Finally, when all four cAMP molecules are bound to A- and B-domains, the two catalytic subunits are released (Equation (10)). These reaction schemas are evaluated by experimental data from [60] for steady state cAMP concentration and PKA formation shown in Figure 2D and kinetic parameters are given in Table 2.

$$\mathrm{R}_2\mathrm{C}_2 + 2\,\mathrm{cAMP} \xrightleftharpoons[\mathrm{kb}_8]{\mathrm{kf}_8} \mathrm{R}_2\mathrm{C}_2 \cdot \mathrm{cAMP}_2$$
$$R_8 = k_{f8}[R_2C_2][cAMP]^2 - k_{b8}[R_2C_2 \cdot cAMP_2] \tag{8}$$

$$\mathrm{R}_2\mathrm{C}_2 \cdot \mathrm{cAMP}_2 + 2\,\mathrm{cAMP} \xrightleftharpoons[\mathrm{kb}_9]{\mathrm{kf}_9} \mathrm{R}_2\mathrm{C}_2 \cdot \mathrm{cAMP}_4$$
$$R_9 = k_{f9}[R_2C_2 \cdot cAMP_2][cAMP]^2 - k_{b9}[R_2C_2 \cdot cAMP_4] \tag{9}$$

$$\mathrm{R}_2\mathrm{C}_2 \cdot \mathrm{cAMP}_4 \xrightleftharpoons[\mathrm{kb}_{10}]{\mathrm{kf}_{10}} 2\,\mathrm{PKAc} + \mathrm{R}_2 \cdot \mathrm{cAMP}_4$$
$$R_{10} = k_{f10}[R_2C_2 \cdot cAMP_4] - k_{b10}[PKAc][R_2 \cdot cAMP_4] \tag{10}$$

6. **PDE phosphorylation by PKA:** Both PDE1A and PDE4A can be phosphorylated by PKA. PKA phosphorylation of PDE1A alters its affinity toward both calcium and calmodulin [30]. PDE4 is a cAMP-specific phosphodiesterase and its activity is increased two to six-fold upon PKA phosphorylation [50]. Different PDE4 isoforms (PDE4A, PDE4B, PDE4D) are targeted to synapses to regulate the cAMP level and modulate synaptic plasticity in learning and memory [50]. PDE4A (the PDE4 subtype used in our model) is highly expressed in the CA1 subregions of the rat hippocampus and is the major PDE4 subtype involved in mediating memory [47]. The phosphorylation reactions of the two different PDE isoforms are modeled by Michaelis-Menten kinetics with a basal degradation, where PKA acts as an enzyme in both of these reactions (Equation (11) and Equation (12)). These reactions are evaluated at steady state to show how PKA phosphorylation affects the activity of PDE1A and PDE4 in comparison with experiments from [30] and [32] in Figure 2E and Figure 2F respectively.

$$\mathrm{PDE\,1} + \mathbf{PKAc} \xrightarrow{\text{Kcat}_{11}} \mathrm{PDE\,1\,P} \qquad R_{11} = \frac{K_{cat11}[PDE1][PKAc]}{K_{m11} + [PDE1]} - K_{r11}[PDE1P] \qquad (11)$$

$$\mathrm{PDE\,4} + \mathbf{PKAc} \xrightarrow{\text{Kcat}_{12}} \mathrm{PDE\,4\,P} \qquad R_{12} = \frac{K_{cat12}[PDE4][PKAc]}{K_{m12} + [PDE4]} - K_{r12}[PDE4P] \qquad (12)$$

7. **cAMP degradation:** We assumed that cAMP can be degraded to AMP by PDE1·Ca$_4$·CaM, PDE4, and PDE4P (PDE4 phosphorylated by PKA). All cAMP inhibition reactions are modeled by Michaelis-Menten kinetics with a basal degradation term, in which phosphodiesterases act as enzymes (Equations (13) to (15)). cAMP hydrolysis by PDE4 and PDE4P evaluated against experimental data from [32] is shown in Figure 2F.

$$\mathrm{cAMP} + \mathbf{PDE1} \cdot \mathbf{Ca_4} \cdot \mathbf{CaM} \xrightarrow{\text{Kcat}_{13}} \mathrm{AMP}$$
$$R_{13} = \frac{K_{cat13}[PDE1 \cdot Ca_4 \cdot CaM][cAMP]}{K_{m13} + [cAMP]} - K_{r13}[AMP] \qquad (13)$$

$$\mathrm{cAMP} + \mathbf{PDE4} \xrightarrow{\text{Kcat}_{14}} \mathrm{AMP} \qquad R_{14} = \frac{K_{cat14}[PDE4][cAMP]}{K_{m14} + [cAMP]} - K_{r14}[AMP] \qquad (14)$$

$$\mathrm{cAMP} + \mathbf{PDE4P} \xrightarrow{\text{Kcat}_{15}} \mathrm{AMP} \qquad R_{15} = \frac{K_{cat15}[PDE4P][cAMP]}{K_{m15} + [cAMP]} - K_{r15}[AMP] \qquad (15)$$

The list of the above reactions and estimated parameters are shown in Table 1 and Table 2, respectively.

Ordinary differential equations

The temporal dynamics of each species, c, is defined as

$$\frac{dc_i}{dt} = J_i \qquad (16)$$

where c_i, $i \in 1, 2, ..., 21$, represents the concentration of the i^{th} species as a function of time and J_i is the net reaction flux for the i^{th} species. These ordinary differential equations (ODEs) and the corresponding initial concentration of the different species are shown in Table 3. The sensitivity analysis with respect to reaction parameters and initial conditions of the model for cAMP and PKA are shown in the Supplemental Material (Figure S1-S4).

Figure 2: Steady-state parameter scans to evaluate the model based on available experimental data in the literature: (A) Saturation curve of C- and N-lobes of calmodulin, predicted by our model, and comparison with experimental data from [39]. (B) Ca^{2+}-stimulated adenylyl cyclase activity, predicted by our model, and comparison with experimental data from [28]. (C) cAMP production at increasing concentrations of ATP, predicted by our model, and comparison with experimental data from [29]. (D) Activation of RIIβ holoenzymes by cAMP in A- and B-domains, predicted by our model, and comparison with experimental data from [60]. (E) Activation of phosphodiesterase isoenzyme PDE1A by different concentrations of Ca^{2+} before and after the phosphorylation of the isoenzyme by PKA, predicted by our model, and comparison with experimental data from [30]. (F) PDE4A-catalyzed cAMP hydrolysis at increasing cAMP concentration before and after PKA phosphorylation, predicted by our model, and comparison with experimental data from [32]. It must be noted that none of the experimental datasets in the literature reported error bars or standard deviations for the measured data points.

Ca^{2+} input signals as the stimulus

The four different types of input calcium are modeled as:

1. Sinusoidal oscillations (ω_1=0.5 Hz) with exponential decay that is repeated every 5 minutes (ω_2=0.003 Hz):

$$Ca^{2+}(t) = (0.45 \sin \pi(t - 0.5) + 0.45)(1 - 0.01)^t + [Ca^{2+}]_{rest} \qquad [\mu M] \qquad (17)$$

2. Calcium bursts with exponential decay that is being repeated every 5 minutes (ω_2=0.003 Hz):

$$Ca^{2+}(t) = 0.95(1 - 0.01)^t + \frac{[Ca^{2+}]_{rest}}{2} \qquad [\mu M] \qquad (18)$$

Table 1: cAMP-PKA pathway reactions, reaction types, and reaction rates used in the model

	List of Reactions	Reaction Type	Reaction Rate
Module 1:	Ca^{2+}/CaM Complex Formation		
1	$2\,Ca^{2+} + CaM \underset{kb_1}{\overset{kf_1}{\rightleftharpoons}} Ca_2{\cdot}CaM$	mass action	$R_1 = k_{f1}[Ca^{2+}]^2[CaM] - k_{b1}[Ca_2CaM]$
2	$2\,Ca^{2+} + Ca_2{\cdot}CaM \underset{kb_2}{\overset{kf_2}{\rightleftharpoons}} Ca_4{\cdot}CaM$	mass action	$R_2 = k_{f2}[Ca^{2+}]^2[Ca_2CaM] - k_{b2}[Ca_4CaM]$
Module 2:	Enzyme Activations By CaCaM Complex		
3	$AC1 + Ca_2{\cdot}CaM \underset{kb_3}{\overset{kf_3}{\rightleftharpoons}} AC1{\cdot}Ca_2{\cdot}CaM$	mass action	$R_3 = k_{f3}[AC1][Ca_2CaM] - k_{b3}[AC1Ca_2CaM]$
4	$2\,Ca^{2+} + AC1{\cdot}Ca_2{\cdot}CaM \underset{kb_4}{\overset{kf_4}{\rightleftharpoons}} AC1{\cdot}Ca_4{\cdot}CaM$	cooperative binding	$R_4 = \frac{k_{f4}[AC1Ca_2CaM][Ca^{2+}]}{K_{m4}+[Ca^{2+}]} - k_{b4}[AC1Ca_4CaM]$
5	$PDE1 + Ca_2{\cdot}CaM \underset{kb_5}{\overset{kf_5}{\rightleftharpoons}} PDE1{\cdot}Ca_2CaM$	mass action	$R_5 = k_{f5}[PDE1][Ca_2CaM] - k_{b5}[PDE1Ca_2CaM]$
6	$2\,Ca^{2+} + PDE1{\cdot}Ca_2{\cdot}CaM \underset{kb_6}{\overset{kf_6}{\rightleftharpoons}} PDE1{\cdot}Ca_4{\cdot}CaM$	cooperative binding	$R_6 = \frac{k_{f6}[PDE1Ca_2CaM][Ca^{2+}]}{K_{m6}+[Ca^{2+}]} - k_{b6}[PDE1Ca_4CaM]$
Module 3:	cAMP Production		
7	$ATP + \mathbf{AC1{\cdot}Ca_4{\cdot}CaM} \xrightarrow{Kcat_7} cAMP$	Michaelis-Menten	$R_7 = \frac{K_{cat7}[AC1Ca_4M][ATP]}{K_{m7}+[ATP]} - K_{r7}[cAMP]$
Module 4:	PKA Formation		
8	$R_2C_2 + 2\,cAMP \underset{kb_8}{\overset{kf_8}{\rightleftharpoons}} R_2C_2{\cdot}cAMP_2$	mass action	$R_8 = k_{f8}[R_2C_2][cAMP]^2 - k_{b8}[R_2C_2cAMP_2]$
9	$R_2C_2{\cdot}cAMP_2 + 2\,cAMP \underset{kb_9}{\overset{kf_9}{\rightleftharpoons}} R_2C_2{\cdot}cAMP_4$	mass action	$R_9 = k_{f9}[R_2C_2cAMP_2][cAMP]^2 - k_{b9}[R_2C_2cAMP_4]$
10	$R_2C_2{\cdot}cAMP_4 \underset{kb_{10}}{\overset{kf_{10}}{\rightleftharpoons}} 2\,PKAc + R_2{\cdot}cAMP_4$	mass action	$R_{10} = k_{f10}[R_2C_2cAMP_4] - k_{b10}[PKAc][R_2cAMP_4]$
Module 5:	PDE Phosphorylation By PKA		
11	$PDE1 + \mathbf{PKAc} \xrightarrow{Kcat_{11}} PDE1P$	Michaelis-Menten	$R_{11} = \frac{K_{cat11}[PDE1][PKAc]}{K_{m11}+[PDE1]} - K_{r11}[PDE1P]$
12	$PDE4 + \mathbf{PKAc} \xrightarrow{Kcat_{12}} PDE4P$	Michaelis-Menten	$R_{12} = \frac{K_{cat12}[PDE4][PKAc]}{K_{m12}+[PDE4]} - K_{r12}[PDE4P]$
Module 6:	cAMP Inhibition		
13	$cAMP + \mathbf{PDE1{\cdot}Ca_4{\cdot}CaM} \xrightarrow{Kcat_{13}} AMP$	Michaelis-Menten	$R_{13} = \frac{K_{cat13}[PDE1Ca_4CaM][cAMP]}{K_{m13}+[cAMP]} - K_{r13}[AMP]$
14	$cAMP + \mathbf{PDE4} \xrightarrow{Kcat_{11}} AMP$	Michaelis-Menten	$R_{14} = \frac{K_{cat14}[PDE4][cAMP]}{K_{m14}+[cAMP]} - K_{r14}[AMP]$
15	$cAMP + \mathbf{PDE4P} \xrightarrow{Kcat_{15}} AMP$	Michaelis-Menten	$R_{15} = \frac{K_{cat15}[PDE4P][cAMP]}{K_{m15}+[cAMP]} - K_{r15}[AMP]$

*Enzymes are in bold type

Table 2: Reaction parameters calculated for the model

Reaction rate	k_f	unit	k_b	unit	K_{cat}	unit	K_m	unit	K_r	unit
R_1	0.1	$\mu M^{-2}\cdot s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_2	0.005	$\mu M^{-2}\cdot s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_3	45.46	$\mu M^{-1}\cdot s^{-1}$	1	-	-	-	-	-	-	-
R_4	6.89	s^{-1}	0.39	s^{-1}	-	-	93.02	μM	-	-
R_5	87.65	$\mu M^{-1}\cdot s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_6	49.80	s^{-1}	1.09	s^{-1}	-	-	6.95	μM	-	-
R_7	-	-	-	-	7.41	s^{-1}	193.73	μM	0.44	s^{-1}
R_8	0.19	$\mu M^{-1}s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_9	27.89	$\mu M^{-1}s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_{10}	0.21	$\mu M^{-1}s^{-1}$	1	s^{-1}	-	-	-	-	-	-
R_{11}	-	-	-	-	0.10	s^{-1}	0.42	μM	0.12	s^{-1}
R_{12}	-	-	-	-	7.67	s^{-1}	1.34	μM	0.02	s^{-1}
R_{13}	-	-	-	-	1*	s^{-1}	35*	μM	0	s^{-1}
R_{14}	-	-	-	-	8.66	s^{-1}	1.21	μM	0.10	s^{-1}
R_{15}	-	-	-	-	1.59	s^{-1}	0.84	μM	0	s^{-1}

*All the kinetic parameters are calculated by fitting the experimental data to the kinetic equations. For reaction 13, the kinetic parameters are directly extracted from [31].

3. Non-oscillating calcium:

$$Ca^{2+}(t_0) = 10\,[Ca^{2+}]_{rest} \qquad [\mu M] \tag{19}$$

in which $[Ca^{2+}]_{rest} = 0.1\ \mu M$ is the resting concentration of calcium in the cytoplasm.

4. Experimental calcium: Spontaneous calcium oscillations measured in neurons are shown in Figure 5. The measured calcium is in the range of 0.1 to 10 μM with oscillations in the seconds timescale.

Table 3: Initial concentrations* and ordinary differential equations

Species	IC (μM)	ODE
Ca^{2+}	1	$\frac{d[Ca^{2+}]}{dt} = -2R_1 - 2R_2 - 2R_4 - 2R_6$
CaM	1.24	$\frac{d[CaM]}{dt} = -R_1$
$Ca_2 \cdot CaM$	0	$\frac{d[Ca_2CaM]}{dt} = R_1 - R_2 - R_3 - R_5$
$Ca_4 \cdot CaM$	0	$\frac{d[Ca_4CaM]}{dt} = R_2$
AC1	5	$\frac{d[AC1]}{dt} = -R_3$
$AC1 \cdot Ca_2 \cdot CaM$	0	$\frac{d[AC1Ca_2CaM]}{dt} = R_3 - R_4$
$AC1 \cdot Ca_4 \cdot CaM$	0	$\frac{d[AC1Ca_4CaM]}{dt} = R_4$
PDE1	3.80	$\frac{d[PDE1]}{dt} = -R_5 - R_{11}$
$PDE1 \cdot Ca_2 \cdot CaM$	0	$\frac{d[PDE1Ca_2CaM]}{dt} = R_5 - R_6$
$PDE1 \cdot Ca_4 \cdot CaM$	0	$\frac{d[PDE1Ca_4CaM]}{dt} = R_6$
ATP	2000	$\frac{d[ATP]}{dt} = -R_7$
cAMP	0	$\frac{d[cAMP]}{dt} = R_7 - 2R_8 - 2R_9 - R_{13} - R_{14} - R_{15}$
R_2C_2	0.22	$\frac{d[R_2C_2]}{dt} = -R_8$
$R_2C_2 \cdot cAMP_2$	0	$\frac{d[R_2C_2cAMP_2]}{dt} = R_8 - R_9$
$R_2C_2 \cdot cAMP_4$	0	$\frac{d[R_2C_2cAMP_4]}{dt} = R_9 - R_{10}$
$R_2 \cdot cAMP_4$	0	$\frac{d[R_2cAMP_4]}{dt} = R_{10}$
PKAc	0	$\frac{d[PKAc]}{dt} = 2R_{10}$
PDE1P	0	$\frac{d[PDE1P]}{dt} = R_{11}$
PDE4	1	$\frac{d[PDE4]}{dt} = -R_{12}$
PDE4P	0	$\frac{d[PDE4P]}{dt} = R_{12}$
AMP	0	$\frac{d[AMP]}{dt} = R_{13} + R_{14} + R_{15}$

*Initial concentrations are estimated in COPASI using the experimental data fitted to the model parameters.

Numerical Methods

The system of ordinary differential equations for this model is solved for a time course of 1800 seconds and interval size of 0.1 (sec) through a deterministic simulation in COPASI [61]. Using a deterministic algorithm (LSODA), the system of ODEs is solved with a dense or banded Jacobian when the problem is stiff and automatically selects between non-stiff (Adams) and stiff (BDF) methods by initially using the non-stiff method and dynamically monitoring data in order to decide which method to use [62]. Parameter estimations are done in COPASI using the Evolutionary Programming (EP) method [63–65]. Sensitivity analyses were conducted in COPASI by numerical differentiation using finite differences [66].

Results

Large frequency modulation of immediate downstream effectors of calcium: The first module of the model focused on the formation of the calcium-calmodulin complex. Calcium input was modeled as a sinusoidal function with a 2-second oscillation period with an exponential decay every 5-minutes [24] (Figure 3A). From here onwards, we refer to the larger frequency as ω_1 for the oscillations in the seconds timescale and the smaller frequency as ω_2 for the oscillations in the minutes timescale. The dynamics of calmodulin consumption leading to the formation of

Ca$_2$·CaM respond only to the low-frequency oscillations of calcium in minute-scale (Figure 3A). Ca$_2$·CaM demonstrates both low frequency (minute-scale) and high frequency (second-scale) oscillations (Figure 3). In the next step, two more Ca^{2+} bind to Ca$_2$·CaM to form Ca$_4$·CaM. Ca$_4$·CaM also responds to second-scale oscillations of calcium; however, the oscillation amplitudes are almost 10-fold lower in comparison with Ca$_2$·CaM (Figure 3A). Another key difference between the observed oscillation patterns of Ca$_2$·CaM and Ca$_4$·CaM is that the Ca$_2$·CaM oscillation pattern seems to be mainly driven by CaM on the minute-scale, while the Ca$_4$·CaM oscillation pattern looks almost exactly like calcium patterns with less significant oscillations on the second scale.

AC1 and PDE1 are activated by the Ca^{2+}/CaM complex. Ca$_2$·CaM binds to AC1 and forms AC1·Ca$_2$·CaM. Both AC1 and AC1· Ca$_2$·CaM show second and minute-scale calcium oscillations (Figure 3B) and their oscillation pattern seem to be mostly deriven by Ca$_2$·CaM oscillation pattern shown in Figure 3C with less significant second-scale oscillations. In the second step, AC1· Ca$_2$·CaM fully activates AC1 by forming AC1· Ca$_4$·CaM with the other two Ca^{2+}. In comparison to AC1·Ca$_2$·CaM, AC1· Ca$_4$CaM shows less significant second-scale oscillations (almost 10-fold lower) and the minute-scale oscillations seems to be driven by the calcium oscillation pattern (Figure 3B). AC1· Ca$_4$·CaM dynamics mostly follow Ca$_4$·CaM dynamics with less sensitivity to Ca^{2+}. The dynamics of PDE1 activation by Ca$_2$·CaM is very similar to AC1. However, both PDE1·Ca$_2$·CaM and PDE1·Ca$_4$·CaM show much higher sensitivity to second-scale oscillations of Ca^{2+} (Figure 3C). This higher activity and increased sensitivity of PDE to Ca^{2+}/CaM in comparison with AC, which makes the amplitude of the oscillations of the PDE activity larger than the amplitude of the oscillations of the AC activity, has been reported by [67].

Figure 3: Ca^{2+}/CaM complex formation and Ca^{2+}/CaM-activated enzymes time course predicted by our model: (A) Calcium input with sinusoidal oscillations (0.5 Hz) and exponential decay (0.003 Hz) as the stimulus in the model. Free calmodulin concentration over a 15-minute and a 10-second time course showing only minute-scale oscillations. The first step of calcium-calmodulin binding of two calcium ions to C-lobe of calmodulin over a 15-minute and a 10-second time course. Ca^{2+}·CaM shows both minute-scale (0.003 Hz) and second-scale (0.5 Hz) oscillations. The second step of calcium-calmodulin binding of another two calcium ions to N-lobe of calmodulin over a 15-minute and a 10-second time course. Ca$_4$CaM also shows both minute-scale (0.003 Hz) and second-scale (0.5 Hz) oscillations. (B) AC1 partial activation by Ca$_2$·CaM to from AC1·Ca$_2$·CaM followed by binding of two extra calcium ions to fully activate AC1 and form AC1·Ca$_4$·CaM. (C) PDE1 partial activation by Ca$_2$·CaM initially followed by further binding of two calcium ions to fully activate PDE1 and form PDE1·Ca$_4$·CaM.

Small frequency modulation of cAMP/PKA and PDE phosphorylation: After enzyme activation by calcium-calmodulin complex, cAMP is synthesized by AC1·CaM from ATP. Then PKA is activated by cAMP and activated PKA phosphorylates PDE1 and PDE4. Finally, cAMP is degraded by PDE1, PDE4, and PDE4P and shows minute-scale oscillations and leaky oscillations on the second-scale (Figure 4A). PKA is activated by cAMP and as a result,

it only shows minute-scale oscillations (Figure 4B). While PDE1-P (phosphorylated PDE1) shows minute-scale oscillations (Figure 4C), PDE4P (phosphorylated PDE4) barely shows any oscillations (Figure 4D). Degradation of cAMP by PDE4 and PDE4P, which are indirectly affected by calcium oscillations might filter the second-scale oscillations induced by AC1·CaM in the cAMP synthesis and change the second-scale oscillations to leaky oscillations (Figure 4A). Thus, our model predicts that phosphodiesterases may filter the large frequency (ω_1) oscillations of the upstream components driven by calcium dynamics and capture only the smaller frequency (ω_2).

Figure 4: cAMP and PKA formation time course and PDE1 and PDE4 phosphorylation by PKA predicted by our model: (A) cAMP dynamics predicted by our model show mainly minute-scale oscillations leaky oscillations on the second-scale shown in the figure inset (B) PKA dynamics predicted by our model shows only minute-scale oscillations and no second-scale oscillations (figure inset). (C) Phosphorylation of PDE1 by PKA and formation of phosphorylated PDE1 (PDE1P) that oscillates only on minute-scale. (D) PDE4 and phosphorylated PDE4 (PDE4P) formed by PKA that is indirectly affected by calcium and shows almost neither of the second- and minute-scale oscillations.

cAMP/PKA pathway filtering of Ca^{2+} signals: The response of cAMP/PKA to different calcium patterns raises the following questions: do spines experience vastly varying calcium patterns? Can experimentally-measured calcium patterns be used to generate predictions from our model? In order to answer these questions, we measured basal calcium fluctuations in dendritic spines of hippocampal neurons. Cultured neurons have been previously reported to form neuronal networks *in vitro* exhibiting spontaneous action potentials [68], which can be measured using reporters of neuronal activity, such as calcium sensors. We expressed the genetically encoded calcium biosensor GCaMP6f [69] in primary rat hippocampal neurons. To allow for useful parameters to be determined, we assigned calcium concentrations to correspond to changes in the measured fluorescence intensity (ΔF/F), based on previously reported intracellular calcium concentrations [34]. Using the measured calcium in three spines from three neurons as the stimulus in the model, we found our model predicts different types of cAMP/PKA dynamics based on the calcium input (Figure 5). We chose these three spines to reflect the different calcium patterns tested in Figure 6. The first spine measured had minimal changes in fluorescence intensity and displays a low range of predicted calcium concentrations (0.1 to 0.2 μM) (Figure 5A). There seem to be no apparent calcium spikes, and the calcium concentration is barely enough to fully activate AC1 and PDE1 and show steady-state cAMP and PKA. AC1·CaM and PDE1·CaM integrate some calcium peaks; however, cAMP and PKA do not exhibit significant peaks. The second spine measured shows higher fluorescence intensity which is almost 10-fold higher than the first data set (Figure 5B). In the second spine, AC1, PDE1, cAMP, and PKA reach a steady state. AC1 and PDE1 seem to reflect most of the calcium major peaks, while cAMP appears to filter out some of the less significant peaks, and PKA almost shows none of these peaks. The third spine measured showed many calcium spikes and a very high range of fluorescence intensity, which we correlated to a large range of calcium concentrations (Figure 5C). The slopes at the first 60 seconds of simulation show that the activation of AC1 and PDE1, and formation of cAMP and PKA occurs much faster in the third spine (Figure 5C) in comparison to the other two (Figure 5A and B). With

high ranges of calcium transients, AC1, PDE1, and cAMP seem to integrate most of the calcium peaks and even PKA has nominal peaks (Figure 5C). Thus, we find that for experimentally derived calcium transients cAMP/PKA serve to filter out the larger frequency (ω_1).

Figure 5: Model predictions for AC1·CaM, PDE1·CaM, cAMP, and PKA for experimental calcium input measured from GCaMP6f expressed in DIV 22 primary rat hippocampal neurons. Fluorescence intensity measurements shown for calcium have an arbitrary starting time and the model system at time zero (start of the recording) is already in equilibrium. The calcium concentration is estimated based on the relative increases and decreases in calcium to the minimum intensity measured. (A) The calcium input is at low range, almost on the range of resting calcium concentration. This calcium amount slows down the rate of AC1 and PDE1 activation by calcium-calmodulin complex and as a result, only a low amount of cAMP is produced. AC1·CaM and PDE1·CaM seem to pick up only calcium peaks, while these calcium dynamics seem to barely affect the cAMP and PKA dynamics over the time course of recorded calcium (4 minutes). (B) The second set of measured calcium in the spine shows a higher amount of calcium which accelerates the rate of AC1 and PDE1 activation and cAMP and PKA formation. Due to the higher range of calcium oscillations in this data set, AC1·CaM, PDE1·CaM, and cAMP seem to be affected more by calcium transients. (C) This set of measured calcium data shows the highest recorded calcium. Because of the high range of calcium oscillations (10 μM), the induced calcium transients are significant in this case and even PKA shows a subtle effect of the most prominent calcium spikes. The data are representative of n=10 neurons from three independent experiments. Representative images at the indicated time points are shown for each spine selected for analysis. Scale bar, 10 μm.

Ca^{2+} signal frequency modulates cAMP/PKA dynamics: If it is true that cAMP/PKA filters out ω_1 dynamics to pick ω_2 dynamics, then this must hold true for different values of these two frequencies. To further test our

predictions, we designed three different calcium inputs (Figure 6) – first, no oscillations of calcium; second, ω_1 is zero and only ω_2 is present; and third, both ω_1 and ω_2 are present. In the first set of simulations, Ca^{2+} input has an initial constant concentration of 1 μM with no oscillations or calcium bursts during the 15 minutes of oscillations (Figure 6A). From AC1·CaM and PDE1·CaM to cAMP and PKA, it is evident that there is only one peak of the concentrations for all these species, with a 40-60 seconds delay between AC1·CaM and PDE1·CaM and cAMP/PKA. This behavior has also been observed by [21], where PKA returns to basal level about 10 minutes after stimulation by a pulse of calcium. For the second set of simulations, starting from the same initial calcium concentration (1 μM), we simulated a calcium burst every five minutes (Figure 6B). These calcium bursts at every five minutes are reflected in AC1·CaM, PDE1·CaM, cAMP, and PKA dynamics. The range of concentrations for all of these molecules is the same as the concentration seen in the first simulation (Figure 6A). The third set of simulations has sinusoidal oscillations with two-second periods in addition to the calcium bursts at every five minutes (Figure 6C). AC1·CaM and PDE1·CaM show both second- and minute-scale oscillations while cAMP and PKA only show the minute-scale oscillations (Figure 6C). In comparison to the second simulation (Figure 6B), the concentration of AC1·CaM, PDE1·CaM, cAMP, and PKA are lower (Figure 6C). We also chose different ω_2 values and confirmed that cAMP only picks up minute time scales (Figure S5).

Discussion

cAMP/PKA activity triggered by Ca^{2+} is an essential biochemical pathway for synaptic plasticity, regulating spine structure, and long-term potentiation [21, 70]. In this study, we have developed a mathematical model, constrained by existing experimental data (Figure 2), to simulate the dynamics of cAMP/PKA pathway in response to calcium input. Analysis of this model allowed us to make the following predictions: *first*, for a given calcium input, AC1, and PDE1 kinetics reflect both the large and the small frequencies albeit with different amplitudes; *second*, cAMP/PKA dynamics reflect only the small frequency of the calcium input; and *finally*, cAMP/PKA acts as a leaky integrator of calcium because of frequency attenuation by the intermediary steps (Figure 7). We also found that spines generated different spontaneous calcium frequencies in cultured neurons (Figure 5), thus motivating the need to study multiple time scales. These findings have implications for cAMP/PKA signaling in dendritic spines in particular and neuronal signal transduction in general.

The ability to modulate the output frequency in response to a signal input is an important feature in biochemical signal transduction; it ensures robustness and noise-filtering. As a result, the output signal, cAMP/PKA in this case, will respond to calcium input for strong signals but not to weak signals (Figure 6). In general, the combination of activation and inactivation kinetics along with feedback loops, many of which oscillate, may be responsible for noise attenuation in signal regulation [71, 72]. Noise attenuation has a significant role in regulating biological switches. These switches must sense changes in signal concentration while buffering against signal noise [73]. Our model predicts that depending on the sensitivity of regulating enzymes to Ca^{2+}, the signaling input can be modified to the desired signaling output with a different range of frequency. For example, starting with a stimulus Ca^{2+} signal with a combination of ω_1 and ω_2 oscillation frequencies, the cAMP/PKA pathway can modify the initial signal and filter it so that components downstream in the pathway, such as PKA receive a signal with an oscillation frequency of ω_2 (Figure 7).

As a result, cAMP/PKA can be thought of as a leaky integrator of calcium signals (Figure 4A and B). This signal modification is important in the context of how cAMP/PKA directly impacts transcription factor activation (e.g. CREB and Epac) [17, 74]. While these transcription factors are not directly considered in our study, it is possible that the frequency modulation exhibited by the pathway we modeled plays a critical role in contributing to neuronal robustness to small fluctuations while retaining the ability to rapidly respond to large calcium influxes.

The calcium input to our model was inspired by measurements of spontaneous calcium dynamics in dendritic spines (Figure 5). Ideally, we would like to use imaging to simultaneously measure spontaneous calcium and cAMP/PKA activity in dendritic spines in order to test our model predictions. However, despite the recent developments in flu-

Figure 6: Effect of calcium stimulation patterns on AC1·CaM, PDE1·CaM, cAMP, and PKA dynamics: (A) Non-oscillating calcium with an initial concentration of 1 μM induces only one peak in each of AC1·CaM, PDE1·CaM, cAMP, and PKA. (B) Calcium bursts with exponential decays with 0.003 Hz frequency (ω_2) induces minute-scale oscillations in each of AC1·CaM, PDE1·CaM, cAMP, and PKA. (C) The combination of sinusoidal calcium oscillations of 0.5 Hz with exponential decays of 0.003 Hz induces both second-scale and minute-scale oscillations in both AC1·CaM and PDE1·CaM as the plot insets show. While cAMP barely shows a leaky oscillation on the second-scale, PKA merely shows minute-scale oscillations.

orescence biosensing methods, imaging and co-imaging of multiple second messengers, kinases, and phosphatase activities in micron-sized compartments remains challenging [75]. This is also true for simultaneously imaging the dynamics of both cAMP and calcium in single dendritic spines [21, 76]. Recently, Tang *et al.* measured PKA activity in dendritic spines in response to glutamate uncaging using new FRET (fluorescence resonance energy transfer) probes that they developed [21]. The timescale of PKA dynamics from our model (Figure 6A) are in good agreement with the time scale of approximately 10 min (Figure 6 of Tang et al [21]). This indicates our developed model closely mimics reported PKA activity in subcompartments, in this case, dendritic spines.

Despite the predictive capability of our model, there are certain limitations that must be acknowledged. First, we only consider a deterministic model and have currently ignored stochastic dynamics in the system. It is possible that the frequency modulation effects presented here will be somewhat altered by including stochasticity but we anticipate that the overall conclusions will not be affected on average if the same network is considered. An additional limitation of our work emerges in the form of challenges associated with co-imaging two second messengers si-

multaneously in the same spine. We are working on developing such probes to overcome these challenges. Finally, spatial regulations of cAMP signaling are other important factors that need to be considered [77–79]. Furthermore, enzyme regulation seems to play a significant role in cAMP/PKA pathway dynamics and signal filtration [80–82]. These enzymes can also be localized by A-kinase anchoring proteins (AKAPs) and change cAMP/PKA dynamics further [83–85]. The effect of spine size and enzyme localizations are considered in a companion study [23].

Figure 7: The frequency control in the modeled cAMP/PKA pathway from calcium stimulation through PKA. Calcium oscillates with a combination of frequencies (ω_1=0.5 Hz and ω_2=0.003 Hz). From calcium-calmodulin complex activation of AC1 and PDE1 enzymes through cAMP production and degradation, the effect of ω_1 becomes trivial. Ultimately, through the downstream pathway, ω_2 becomes the only significant oscillatory frequency.

Acknowledgments

The authors wish to thank Miriam Bell, Justin Laughlin, and Allen Leung for their valuable comments. DO and PR were supported by AFOSR MURI grant number FA9550-18-1-0051. DLS was supported by NIH/NCI T32 CA009523. Confocal imaging was done at the UC San Diego Nikon Imaging Center with the assistance of Eric Griffis. DLS would like to thank AJ Slepain for her assistance with neuronal culture.

Author contributions

DO and PR conceived the study; DO developed the model and conducted the simulations; DLS conducted neuronal culture experiments; BC aided in experimental design; all authors analyzed the data and wrote the manuscript.

References

[1] P. Uhlen and N. Fritz, "Biochemistry of calcium oscillations," *Biochemical and Biophysical Research Communications*, vol. 396, pp. 28–32, may 2010.

[2] G. Dupont, L. Combettes, G. S. Bird, and J. W. Putney, "Calcium Oscillations," *Cold Spring Harbor Perspectives in Biology*, vol. 3, pp. a004226–a004226, mar 2011.

[3] E. Smedler and P. Uhlen, "Frequency decoding of calcium oscillations," *Biochimica et Biophysica Acta (BBA) - General Subjects*, vol. 1840, pp. 964–969, mar 2014.

[4] J. Sneyd, J. M. Han, L. Wang, J. Chen, X. Yang, A. Tanimura, M. J. Sanderson, V. Kirk, and D. I. Yule, "On the dynamical structure of calcium oscillations," *Proceedings of the National Academy of Sciences*, vol. 114, pp. 1456–1461, feb 2017.

[5] M. Zaccolo and T. Pozzan, "cAMP and Ca2+ interplay: a matter of oscillation patterns," *Trends in Neurosciences*, vol. 26, pp. 53–55, feb 2003.

[6] T. A. Dunn, C.-T. Wang, M. A. Colicos, M. Zaccolo, L. M. DiPilato, J. Zhang, R. Y. Tsien, and M. B. Feller, "Imaging of cAMP Levels and Protein Kinase A Activity Reveals That Retinal Waves Drive Oscillations in Second-Messenger Cascades," *Journal of Neuroscience*, vol. 26, pp. 12807–12815, dec 2006.

[7] J. S. O'Neill and A. B. Reddy, "The essential role of cAMP/Ca 2+ signalling in mammalian circadian timekeeping," *Biochemical Society Transactions*, vol. 40, pp. 44–50, feb 2012.

[8] S. Ujita, T. Sasaki, A. Asada, K. Funayama, M. Gao, K. Mikoshiba, N. Matsuki, and Y. Ikegaya, "cAMP-Dependent Calcium Oscillations of Astrocytes: An Implication for Pathology," *Cerebral Cortex*, vol. 27, p. bhv310, jan 2016.

[9] F. Siso-Nadal, J. J. Fox, S. A. Laporte, T. E. Hébert, and P. S. Swain, "Cross-Talk between Signaling Pathways Can Generate Robust Oscillations in Calcium and cAMP," *PLoS ONE*, vol. 4, p. e7189, oct 2009.

[10] D. Willoughby, W. Wong, J. Schaack, J. D. Scott, and D. M. F. Cooper, "An anchored PKA and PDE4 complex regulates subplasmalemmal cAMP dynamics.," *The EMBO journal*, vol. 25, pp. 2051–61, may 2006.

[11] M. L. Halls and D. M. F. Cooper, "Regulation by Ca2+-Signaling Pathways of Adenylyl Cyclases," *Cold Spring Harbor Perspectives in Biology*, vol. 3, pp. a004143–a004143, jan 2011.

[12] N. K. Tulsian, S. Krishnamurthy, and G. S. Anand, "Channeling of cAMP in PDE-PKA Complexes Promotes Signal Adaptation," *Biophysical Journal*, vol. 112, pp. 2552–2566, jun 2017.

[13] C. Sette and M. Conti, "Phosphorylation and Activation of a cAMP-specific Phosphodiesterase by the cAMP-dependent Protein Kinase," *Journal of Biological Chemistry*, vol. 271, pp. 16526–16534, jul 1996.

[14] A. Cardinale and F. R. Fusco, "Inhibition of phosphodiesterases as a strategy to achieve neuroprotection in Huntington's disease," *CNS Neuroscience & Therapeutics*, vol. 24, pp. 319–328, apr 2018.

[15] J. W. Hell, "How Ca 2+ -permeable AMPA receptors, the kinase PKA, and the phosphatase PP2B are intertwined in synaptic LTP and LTD," *Science Signaling*, vol. 9, pp. pe2–pe2, apr 2016.

[16] N. J. Batty, K. K. Fenrich, and K. Fouad, "The role of cAMP and its downstream targets in neurite growth in the adult nervous system," *Neuroscience Letters*, vol. 652, pp. 56–63, jun 2017.

[17] E. R. Kandel, "The molecular biology of memory: cAMP, PKA, CRE, CREB-1, CREB-2, and CPEB," *Molecular Brain*, vol. 5, p. 14, may 2012.

[18] T. Nagai, J. Yoshimoto, T. Kannon, K. Kuroda, and K. Kaibuchi, "Phosphorylation Signals in Striatal Medium Spiny Neurons," *Trends in Pharmacological Sciences*, vol. 37, pp. 858–871, oct 2016.

[19] S. E. Tillo, W.-H. Xiong, M. Takahashi, S. Miao, A. L. Andrade, D. A. Fortin, G. Yang, M. Qin, B. F. Smoody, P. J. Stork, and H. Zhong, "Liberated PKA Catalytic Subunits Associate with the Membrane via Myristoylation to Preferentially Phosphorylate Membrane Substrates," *Cell Reports*, vol. 19, pp. 617–629, apr 2017.

[20] H. Zhong, G.-M. Sia, T. R. Sato, N. W. Gray, T. Mao, Z. Khuchua, R. L. Huganir, and K. Svoboda, "Subcellular Dynamics of Type II PKA in Neurons," *Neuron*, vol. 62, pp. 363–374, may 2009.

[21] S. Tang and R. Yasuda, "Imaging ERK and PKA Activation in Single Dendritic Spines during Structural Plasticity," *Neuron*, vol. 93, pp. 1315–1324.e3, mar 2017.

[22] R. Yasuda, "Studying signal transduction in single dendritic spines," *Cold Spring Harbor Perspectives in Biology*, vol. 4, oct 2012.

[23] D. Ohadi and P. Rangamani, "Geometric control of frequency modulation of cAMP oscillations due to Ca2+-bursts in dendritic spines," *bioRxiv*, p. 520643, jan 2019.

[24] Y. V. Gorbunova and N. C. Spitzer, "Dynamic interactions of cyclic AMP transients and spontaneous Ca2+ spikes," *Nature*, vol. 418, pp. 93–96, jul 2002.

[25] N. Masada, S. Schaks, S. E. Jackson, A. Sinz, and D. M. F. Cooper, "Distinct Mechanisms of Calmodulin Binding and Regulation of Adenylyl Cyclases 1 and 8," *Biochemistry*, vol. 51, pp. 7917–7929, oct 2012.

[26] T. A. Goraya and D. M. Cooper, "Ca2+-calmodulin-dependent phosphodiesterase (PDE1): Current perspectives," *Cellular Signalling*, vol. 17, pp. 789–797, jul 2005.

[27] S. Hoops, R. Gauges, C. Lee, J. Pahle, N. Simus, M. Singhal, L. Xu, P. Mendes, and U. Kummer, "COPASI - A COmplex PAthway SImulator," *Bioinformatics*, vol. 22, pp. 3067–3074, dec 2006.

[28] S. T. Wong, J. Athos, X. A. Figueroa, V. V. Pineda, M. L. Schaefer, C. C. Chavkin, L. J. Muglia, and D. R. Storm, "Calcium-Stimulated Adenylyl Cyclase Activity Is Critical for Hippocampus-Dependent Long-Term Memory and Late Phase LTP," *Neuron*, vol. 23, pp. 787–798, aug 1999.

[29] C. Klein, R. K. Sunahara, T. Y. Hudson, T. Heyduk, and A. C. Howlett, "Zinc Inhibition of cAMP Signaling," *Journal of Biological Chemistry*, vol. 277, pp. 11859–11865, apr 2002.

[30] R. K. Sharma and J. H. Wang, "Regulation of cAMP concentration by calmodulin-dependent cyclic nucleotide phosphodiesterase," *Biochemistry and Cell Biology*, vol. 64, pp. 1072–1080, nov 1986.

[31] R. K. Sharma and J. Kalra, "Characterization of calmodulin-dependent cyclic nucleotide phosphodiesterase isoenzymes," tech. rep., 1994.

[32] F. Laliberté, S. Liu, E. Gorseth, B. Bobechko, A. Bartlett, P. Lario, M. J. Gresser, and Z. Huang, "In vitro PKA phosphorylation-mediated human PDE4A4 activation," *FEBS Letters*, vol. 512, pp. 205–208, feb 2002.

[33] N. L. Rochefort and A. Konnerth, "Dendritic spines: From structure to in vivo function," aug 2012.

[34] R. Yasuda, "Biophysics of Biochemical Signaling in Dendritic Spines: Implications in Synaptic Plasticity," *Biophysical Journal*, vol. 113, pp. 2152–2159, nov 2017.

[35] M. Zhang, C. Abrams, L. Wang, A. Gizzi, L. He, R. Lin, Y. Chen, P. J. Loll, J. M. Pascal, and J.-f. Zhang, "Structural Basis for Calmodulin as a Dynamic Calcium Sensor," *Structure*, vol. 20, pp. 911–923, may 2012.

[36] A. R. Means, "Calcium, calmodulin and cell cycle regulation," *FEBS Letters*, vol. 347, pp. 1–4, jun 1994.

[37] G. C. Faas, S. Raghavachari, J. E. Lisman, and I. Mody, "Calmodulin as a direct detector of Ca2+ signals," *Nature Neuroscience*, vol. 14, pp. 301–304, mar 2011.

[38] D. B. Halling, B. J. Liebeskind, A. W. Hall, and R. W. Aldrich, "Conserved properties of individual Ca 2+ -binding sites in calmodulin," *Proceedings of the National Academy of Sciences*, vol. 113, pp. E1216–E1225, mar 2016.

[39] M. Lai, D. Brun, S. J. Edelstein, and N. Le Novère, "Modulation of Calmodulin Lobes by Different Targets: An Allosteric Model with Hemiconcerted Conformational Transitions," *PLOS Computational Biology*, vol. 11, p. e1004063, jan 2015.

[40] A. B. Sorensen, M. T. Søndergaard, and M. T. Overgaard, "Calmodulin in a Heartbeat," in *FEBS Journal*, vol. 280, pp. 5511–5532, nov 2013.

[41] N. V. Valeyev, D. G. Bates, P. Heslop-Harrison, I. Postlethwaite, and N. V. Kotov, "Elucidating the mechanisms of cooperative calcium-calmodulin interactions: a structural systems biology approach," *BMC Systems Biology*, vol. 2, p. 48, jun 2008.

[42] G. C. Chan, S. Tonegawa, and D. R. Storm, "Hippocampal neurons express a calcineurin-activated adenylyl cyclase," *J Neurosci*, vol. 25, pp. 9913–9918, may 2005.

[43] G. A. Wayman, S. Impey, Z. Wu, W. Kindsvogel, L. Prichard, and D. R. Storm, "Synergistic activation of the type I adenylyl cyclase by Ca2+ and Gs-coupled receptors in vivo," *Journal of Biological Chemistry*, vol. 269, pp. 25400–25405, oct 1994.

[44] D. M. F. Cooper, N. Mons, and J. W. Karpen, "Adenylyl cyclases and the interaction between calcium and cAMP signalling," *Nature*, vol. 374, pp. 421–424, mar 1995.

[45] G. Antunes, A. C. Roque, and F. M. Simoes De Souza, "Modelling intracellular competition for calcium: Kinetic and thermodynamic control of different molecular modes of signal decoding," *Scientific Reports*, vol. 6, p. 23730, apr 2016.

[46] D. Willoughby and D. M. F. Cooper, "Organization and Ca 2+ Regulation of Adenylyl Cyclases in cAMP Microdomains," *Physiological Reviews*, vol. 87, pp. 965–1010, jul 2007.

[47] N. J. Brandon and A. R. West, eds., *Cyclic-Nucleotide Phosphodiesterases In The Central Nervous System*. Hoboken, New Jersey: John Wiley & Sons, Inc., may 2014.

[48] R. Kakkar, R. V. Raju, and R. K. Sharma, "Calmodulin-dependent cyclic nucleotide phosphodiesterase (PDE1).," *Cellular and molecular life sciences : CMLS*, vol. 55, pp. 1164–86, jul 1999.

[49] R. K. Sharma, "Diversity of calcium action in regulation of mammalian calmodulin-dependent cyclic nucleotide phosphodiesterase.," *Indian journal of biochemistry & biophysics*, vol. 40, pp. 77–91, apr 2003.

[50] H.-T. Zhang, Y. Xu, and J. M. O'Donnell, eds., *Phosphodiesterases: CNS Functions and Diseases*, vol. 17 of *Advances in Neurobiology*. Cham: Springer International Publishing, 2017.

[51] R. K. Sharma, S. B. Das, A. Lakshmikuttyamma, P. Selvakumar, and A. Shrivastav, "Regulation of calmodulin-stimulated cyclic nucleotide phosphodiesterase (PDE1): Review," *INTERNATIONAL JOURNAL OF MOLECULAR MEDICINE*, vol. 18, pp. 95–105, jul 2006.

[52] T. A. Goraya, N. Masada, A. Ciruela, and D. M. F. Cooper, "Sustained entry of Ca2+ is required to activate Ca 2+-calmodulin-dependent phosphodiesterase 1A," *Journal of Biological Chemistry*, vol. 279, pp. 40494–40504, sep 2004.

[53] E. Fourcaudot, F. Gambino, Y. Humeau, G. Casassus, H. Shaban, B. Poulain, and A. Luthi, "cAMP/PKA signaling and RIM1 mediate presynaptic LTP in the lateral amygdala," *Proceedings of the National Academy of Sciences*, vol. 105, pp. 15130–15135, sep 2008.

[54] T. Tzounopoulos, R. Janz, T. C. Südhof, R. A. Nicoll, and R. C. Malenka, "A Role for cAMP in Long-Term Depression at Hippocampal Mossy Fiber Synapses," *Neuron*, vol. 21, pp. 837–845, oct 1998.

[55] E. D. Roberson and J. D. Sweatt, "Transient Activation of Cyclic AMP-dependent Protein Kinase during Hippocampal Long-term Potentiation," *Journal of Biological Chemistry*, vol. 271, pp. 30436–30441, nov 1996.

[56] T. Abel and P. V. Nguyen, "Regulation of hippocampus-dependent memory by cyclic AMP-dependent protein kinase.," *Progress in brain research*, vol. 169, pp. 97–115, 2008.

[57] R. Walker-Gray, F. Stengel, and M. G. Gold, "Mechanisms for restraining cAMP-dependent protein kinase revealed by subunit quantitation and cross-linking approaches," *Proceedings of the National Academy of Sciences*, vol. 114, pp. 10414–10419, sep 2017.

[58] R. M. Gibson, Y. Ji-Buechler, and S. S. Taylor, "Interaction of the Regulatory and Catalytic Subunits of cAMP-dependent Protein Kinase," *Journal of Biological Chemistry*, vol. 272, pp. 16343–16350, jun 1997.

[59] D. Kulasiri and Y. He, *Computational Systems Biology of Synaptic Plasticity*, vol. 10 of *Series on Advances in Bioinformatics and Computational Biology*. WORLD SCIENTIFIC (EUROPE), aug 2017.

[60] P. Zhang, E. V. Smith-Nguyen, M. M. Keshwani, M. S. Deal, A. P. Kornev, and S. S. Taylor, "Structure and allostery of the PKA RIIβ tetrameric holoenzyme," *Science*, vol. 335, pp. 712–716, feb 2012.

[61] L. Petzold, "Automatic Selection of Methods for Solving Stiff and Nonstiff Systems of Ordinary Differential Equations," *SIAM Journal on Scientific and Statistical Computing*, vol. 4, pp. 136–148, mar 1983.

[62] P. Mendes, S. Hoops, S. Sahle, R. Gauges, J. Dada, and U. Kummer, "Computational Modeling of Biochemical Networks Using COPASI," pp. 17–59, Humana Press, 2009.

[63] D. B. Fogel, "An Overview of Evolutionary Programming," pp. 89–109, Springer, New York, NY, 1999.

[64] T. Baeck, D. B. Fogel, and Z. Michalewicz, *Handbook of evolutionary computation*. Institute of Physics Pub., 1997.

[65] T. Baeck and H.-P. Schwefel, "An Overview of Evolutionary Algorithms for Parameter Optimization," *Evolutionary Computation*, vol. 1, pp. 1–23, mar 1993.

[66] S. Sahle, P. Mendes, S. Hoops, and U. Kummer, "A new strategy for assessing sensitivities in biochemical models," *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, vol. 366, pp. 3619–3631, oct 2008.

[67] X. Yu, J. H. Byrne, and D. A. Baxter, "Modeling Interactions Between Electrical Activity and Second-Messenger Cascades in Aplysia Neuron R15," *Journal of Neurophysiology*, vol. 91, pp. 2297–2311, may 2004.

[68] J. van Pelt, I. Vajda, P. S. Wolters, M. A. Corner, and G. J. Ramakers, "Dynamics and plasticity in developing neuronal networks in vitro," in *Progress in Brain Research*, vol. 147, pp. 171–188, Elsevier, jan 2005.

[69] T.-W. Chen, T. J. Wardill, Y. Sun, S. R. Pulver, S. L. Renninger, A. Baohan, E. R. Schreiter, R. A. Kerr, M. B. Orger, V. Jayaraman, L. L. Looger, K. Svoboda, and D. S. Kim, "Ultrasensitive fluorescent proteins for imaging neuronal activity," *Nature*, vol. 499, pp. 295–300, jul 2013.

[70] D. D. Murphy and M. Segal, "Morphological plasticity of dendritic spines in central neurons is mediated by activation of cAMP response element binding protein," *Proceedings of the National Academy of Sciences*, vol. 94, pp. 1482–1487, feb 1997.

[71] L. Wang, J. Xin, and Q. Nie, "A Critical Quantity for Noise Attenuation in Feedback Systems," *PLoS Computational Biology*, vol. 6, p. e1000764, apr 2010.

[72] O. Brandman and T. Meyer, "Feedback Loops Shape Cellular Signals in Space and Time," *Science*, vol. 322, pp. 390–395, oct 2008.

[73] M. Chen, L. Wang, C. C. Liu, and Q. Nie, "Noise Attenuation in the ON and OFF States of Biological Switches," *ACS Synthetic Biology*, vol. 2, pp. 587–593, oct 2013.

[74] M. P. Delghandi, M. Johannessen, and U. Moens, "The cAMP signalling pathway activates CREB through PKA, p38 and MSK1 in NIH 3T3 cells," *Cellular Signalling*, vol. 17, pp. 1343–1351, nov 2005.

[75] C. R. Hackley, E. O. Mazzoni, and J. Blau, "cAMPr: A single-wavelength fluorescent sensor for cyclic AMP," *Science Signaling*, vol. 11, p. eaah3738, mar 2018.

[76] Y. Ohta, T. Furuta, T. Nagai, and K. Horikawa, "Red fluorescent cAMP indicator with increased affinity and expanded dynamic range," *Scientific Reports*, vol. 8, p. 1866, dec 2018.

[77] S. R. Neves, P. Tsokas, A. Sarkar, E. A. Grace, P. Rangamani, S. M. Taubenfeld, C. M. Alberini, J. C. Schaff, R. D. Blitzer, I. I. Moraru, and R. Iyengar, "Cell Shape and Negative Links in Regulatory Motifs Together Control Spatial Information Flow in Signaling Networks," *Cell*, vol. 133, pp. 666–680, may 2008.

[78] M. Zaccolo, "Spatial control of cAMP signalling in health and disease.," *Current opinion in pharmacology*, vol. 11, pp. 649–55, dec 2011.

[79] A. E. Cowan, I. I. Moraru, J. C. Schaff, B. M. Slepchenko, and L. M. Loew, "Spatial modeling of cell signaling networks.," *Methods in cell biology*, vol. 110, pp. 195–221, 2012.

[80] M. D. HOUSLAY and D. R. ADAMS, "PDE4 cAMP phosphodiesterases: modular enzymes that orchestrate signalling cross-talk, desensitization and compartmentalization," *Biochemical Journal*, vol. 370, pp. 1–18, feb 2003.

[81] D. M. F. COOPER, "Regulation and organization of adenylyl cyclases and cAMP," *Biochemical Journal*, vol. 375, pp. 517–529, nov 2003.

[82] V. De Arcangelis, S. Liu, D. Zhang, D. Soto, and Y. K. Xiang, "Equilibrium between Adenylyl Cyclase and Phosphodiesterase Patterns Adrenergic Agonist Dose-Dependent Spatiotemporal cAMP/Protein Kinase A Activities in Cardiomyocytes," *Molecular Pharmacology*, vol. 78, pp. 340–349, sep 2010.

[83] C. W. Dessauer, "Adenylyl Cyclase-A-kinase Anchoring Protein Complexes: The Next Dimension in cAMP Signaling," *Molecular Pharmacology*, vol. 76, pp. 935–941, nov 2009.

[84] M. S. Kapiloff, M. Rigatti, and K. L. Dodge-Kafka, "Architectural and functional roles of A kinase–anchoring proteins in cAMP microdomains," *The Journal of General Physiology*, vol. 143, pp. 9–15, jan 2014.

[85] K. Gorshkov, S. Mehta, S. Ramamurthy, G. V. Ronnett, F.-Q. Zhou, and J. Zhang, "AKAP-mediated feedback control of cAMP gradients in developing hippocampal neurons," *Nature Chemical Biology*, vol. 13, pp. 425–431, feb 2017.