1
2
3
The Dynamic Shift Detector: an algorithm to identify changes in parameter values governing populations

Short title: Dynamic Shift Detector: an algorithm to identify population changes

Christie A. Bahlai[1] and Elise F. Zipkin[2]

1 Department of Biological Sciences, Kent State University. Kent, Ohio, United States of America. Email
for correspondence: cbahlai@kent.edu

2 Department of Integrative Biology; Program in Ecology, Evolutionary Biology and Behavior, Michigan
State University, East Lansing, Michigan, United States of America.

Abstract

Environmental factors interact with internal rules of population regulation, sometimes perturbing systems to alternate dynamics though changes in parameter values. Yet, pinpointing when such changes occur in naturally fluctuating populations is difficult. An algorithmic approach that can identify the timing and magnitude of parameter shifts would facilitate understanding of abrupt ecological transitions with potential to inform conservation and management of species.

The "Dynamic Shift Detector" is an algorithm to identify changes in parameter values governing temporal fluctuations in populations with nonlinear dynamics. The algorithm examines population time series data for the presence, location, and magnitude of parameter shifts. It uses an iterative approach to fitting subsets of time series data, then ranks the fit of break point combinations using model selection, assigning a relative weight to each break. We examined the performance of the Dynamic Shift Detector with simulations and two case studies. Under low environmental/sampling noise, the break point sets selected by the Dynamic Shift Detector contained the true simulated breaks with 70-100% accuracy. The weighting tool generally assigned breaks intentionally placed in simulated data (i.e., true breaks) with weights averaging >0.8 and those due to sampling error (i.e., erroneous breaks) with weights averaging <0.2. In our case study examining an invasion process, the algorithm identified shifts in population cycling associated with variations in resource availability. The shifts identified for the conservation case study highlight a decline process that generally coincided with changing management practices affecting the availability of hostplant resources.

When interpreted in the context of species biology, the Dynamic Shift Detector algorithm can aid management decisions and identify critical time periods related to species' dynamics. In an era of rapid global change, such tools can provide key insights into the conditions under which population parameters, and their corresponding dynamics, can shift.

Author Summary

Populations naturally fluctuate in abundance, and the rules governing these fluctuations are a result of both internal (density dependent) and external (environmental) processes. For these reasons, pinpointing when changes in populations occur is difficult. In this study, we develop a novel break-point analysis tool for population time series data. Using a density dependent model to describe a population's underlying dynamic process, our tool iterates through all possible break point combinations (i.e., abrupt changes in parameter values) and applies information-theoretic decision tools (i.e. Akaike's Information Criterion corrected for small sample sizes) to determine best fits. Here, we develop the approach, simulate data under a variety of conditions to demonstrate its utility, and apply the tool to two case studies: an invasion of multicolored Asian ladybeetle and declining monarch butterflies. The Dynamic Shift Detector algorithm identified parameter changes that correspond to known environmental change events in both case studies.

Introduction

Abrupt and persistent changes in ecological processes, and methods to detect them, have long interested ecologists [1–5]. Changes to the rules governing system dynamics are often associated with substantial changes to biodiversity and ecosystem functions. Understanding when, and how these changes occur is critically important to broader evaluation of system behaviors. The study of abrupt changes, discontinuities, and regime shifts is highly interdisciplinary, and has been examined for a variety of processes related to climate [6,7], ecology [8], and economic and social systems [9,10].

Population dynamics are determined by internal, biotic rules and also external abiotic factors, leading to both stochastic and deterministic forces affecting abundance through time [11]. External perturbations can lead to shifts in population dynamics, such that the parameters governing population abundance transition to other values [12,13]. In the context of this study, we define the set of parameters controlling dynamics as a population's *dynamic rule*, and an abrupt shift in these parameter values as a *dynamic shift*. We use the term *break point* to describe the location in time series data where the dynamic shift occurs.

Although theoretically straightforward, identifying dynamic shifts in noisy ecological systems is challenging using real-world data due to a lack of systematic, adaptable tools [2]. Arguably, the most common approach to identify break points is through the use of segmented regressions [12,14,15]. However, these models don't adequately account for nonlinearities, and uncertainties in the existence and location of breaks are typically not quantified [1,2]. Break points are often applied to time series data *ad hoc*, based on data visualization or specific hypotheses surrounding factors affecting population changes [12,14–17], creating the potential for biases in break point selection.

Several break point detection methods have been developed to address issues associated with *ad hoc* approaches. Such dynamic shift analysis tools use a variety of statistical optimization strategies, including linear and moving average methods [18–21]. For example, climatological and econometrics time series data are frequently examined for stepwise statistical deviations from the mean or variance [6,7,22]. To fit the periodicity of population time series data, wavelet analyses have also been used to detect break points [23] but this method does not mechanistically account for density-dependent processes such that changes in parameter values are not easily interpretable [24]. Dynamic shift detection methods that explicitly account for non-linear population processes may be less likely to yield false positives than methods based on summary statistics [25].

Break point detection methods based on statistical measures tend to rely on null hypothesis testing (i.e., that no dynamic shift occurred) and thus they have low sensitivity in situations where statistical power is limited. Additionally, such methods do not provide a means for assessing uncertainty in the existence and magnitude of break points [1,26]. In a 2009 review, Andersen and colleagues note that if common break point methods were used on typical ecological time series with 20-40 time steps, only the most extreme transitions occurring near the midpoint of the time series would be deemed 'significant' [1]. They concluded that break point analyses could be enhanced with respect to both sensitivity and parsimony by use of model selection procedures. Thus, to address these limitations in the ability to identify dynamic shifts in population processes, it is necessary to develop rigorous tools that allow users to accommodate non-linear population models and quantify uncertainties associated with the existence of potential break points.

In this paper, we develop a generalizable algorithm, the Dynamic Shift Detector (DSD), to identify
dynamic shifts in populations with density-dependent growth using time series data. The DSD algorithm
uses an iterative approach, grounded in information theoretic model selection. We illustrate the DSD
using the Ricker model because of this model's simplicity and high performance under a variety of
realistic environmental scenarios. Density-dependent population models such as the Ricker provide a
convenient, generalizable proxy for a variety of ecological processes because of their relatively simple
parameterization and potential to explain complex dynamics [27]. Although deterministic approaches to
population modelling have largely fallen out of favor for more complex structures and stochastic
elements [28–30], simple dynamic models remain useful due to their easily interpretable and
ecologically meaningful parameters [31]. Further, the techniques described in our paper can be readily
adapted to other model structures, including more complicated processes such as seasonal periodicity
or lag effects.

We describe the basic structure of our DSD algorithm and how it can be used to evaluate the presence,
location, and magnitude of dynamic shifts in population parameters (i.e., break points in time series
abundance data). We demonstrate the utility of our algorithm through a series of simulations and apply
the algorithm to empirical case studies of two populations of economic and conservation concern. First,
we examine the invasion process of the multicolored Asian ladybeetle (*Harmonia axyridis*), a
cosmopolitan invasive, in the two decades following its arrival in Midwestern US agricultural
ecosystems. Then, we examine the declining eastern monarch butterfly (*Danaus plexippus*) population
using census data collected on its overwintering grounds in Mexico over a similar two-decade period. In
our ladybeetle case study, the DSD algorithm identified dynamic shifts associated with known variation
in prey availability, with moderately high weight for a break coinciding with prey arrival and moderately
low weight for a break coinciding with management actions aimed to control the prey. The results for
the monarch population were more ambiguous, with greater uncertainty about the number and location
of breaks in the time series. Several equivalently performing break point combinations had divergent
weights for various break points, suggesting that multiple, super-imposed biological processes drive the
dynamics of this population.

Models

The Dynamic Shift Detector algorithm

Although any time series population model can be used with our tool, we illustrate the DSD algorithm
with a Ricker population model. To do this, we assume that the population size in time t+1, N_{t+1}, is
dependent on the population size in time t, N_t and regulated by two parameters: the carrying capacity
of the system, K, and the annual intrinsic growth rate, r [27]:

124
$$N_{t+1} = N_t \exp\left(r\left(1 - \frac{N_t}{K}\right)\right) + \varepsilon_t.$$

We further assume that observed annual population abundance is partially stochastic and may be
influenced by environmental variation and sampling error. We include an error term ε_t to represent this
noise, which follows a normal distribution centered around zero with a variance of σ^2. The parameters
K, r, and ε_t are estimated from the population time series data (N_1, N_2, … N_t). The Ricker model is a
useful starting point for break point analyses because 1) it does not rely on any external information
(abundance in time *t* is a function of only abundance in time *t*-1); 2) only three parameters (including the

error) need to be estimated, and those parameters have ecologically meaningful interpretations; and 3)
it is an extremely flexible distribution, taking a variety of forms, from linear to compensatory to over-
compensatory, and thus has a wide range of applications across a variety of taxa [32,33]. Subsequent
applications of the DSD algorithm can incorporate other population models if the life history of the
target organism requires a different structure.

To build the DSD algorithm, we use an iterative, model-selection process to determine if, and when,
shifts in parameter values occur within a given time series. To achieve this, we first fit the Ricker model
to the entire time series of available data. Then the population time series is subdivided into all possible
combinations of 2, 3, …, n subsets of sequential data points (hereafter, 'break point combinations') and
the Ricker model is fit to each of the subsets produced for each break point combination. To avoid
overfitting, we constrain break point combinations to include only subsets with a minimum of four
sequential data points.

After fitting all break point combinations, we evaluate the candidate set of models by calculating the
Akaike Information Criteria for small sample sizes (AICc) for each segment and summing them
accordingly [34]. Fits for break point combinations with comparatively lower AICc values are considered
to have better performance. For a given time series, the DSD algorithm produces a set of top performing
break point combinations for cases in which model fits produce equivalent AICc values (i.e. within 2
units of the best-performing fit; [35]). To evaluate the strength of evidence for an identified break in the
time series, we use the relative variable importance method [35]. We compute the Akaike weight w_i (a
measure of the relative likelihood of a break point combination, given the data and the set of break
point combinations being tested) for every identified break point across all combinations. Commonly
used in model averaging, the w_1, w_2,… w_n are interpreted as the respective conditional probabilities for
each model in a set of n models [36]. Break weight (i.e., relative variable importance, *sensu* [35]) is
computed as the sum of the Akaike weights for all break point combinations, where that break point
appears. Break point combinations with weights <0.001 were excluded to increase computational
efficiency.

We selected AICc as our information criterion for model selection within the DSD algorithm because it
provides a balance of specificity and sensitivity. However, we also completed a parallel analysis with an
identical procedure using AIC as the information criterion for decision-making, which is documented in
Appendix S1. AICc is a function of AIC with a correction for small sample bias, which is appropriate for
the sample sizes typical to contemporary population time series data (i.e., 15-30 years/data points) and
is designed to minimize the risk of overfitting during model selection [35]. However, use of AIC for
model selection may be desirable when increased algorithmic sensitivity to dynamic shifts is desired.

The DSD algorithm is implemented as a series of R functions to enable a user to quickly generate a list of
potential break points for a population time series dataset. The algorithm (and all subsequent
simulations and case studies) were scripted and run in R Version 3.3.3 [37]. For fitting the Ricker model,
we used the Levenberg-Marquardt nonlinear least-squares algorithm as implemented in the package
minpack.LM [38]. All data manipulations, analyses and figure scripts, including the complete
development history, are publicly available in a Github repository at
https://github.com/cbahlai/dynamic_shift_detector [39]. We summarize the role of each function used
in the algorithm within Appendix S2.

Simulation study

We conducted a series of simulations to test the accuracy of the DSD algorithm under a variety of
plausible parameter spaces. For all scenarios, we fix N_1 = 3000, and K = 2000 in the initial conditions, as
the Ricker model is most reliably fit for populations fluctuating around their carrying capacity. As the
dynamic observed in a Ricker population is driven primarily by the relationship of other parameters to K
than by the absolute value of K itself, we held the starting value of K constant for all simulations. For
each set of simulations, we held the variables not being varied at "base values" defined as: starting value
of r = 2, change in r = ±25%, change in K = ±75%, 2% noise (τ; described below), with time series length
of 20 years. We examined the effect of the size of initial r on algorithm performance by creating
scenarios with different starting values of r = 0.5, 1, 1.5, 2. For each value of initial r, we modified the
percent change in r at break points from the starting values (± no change, 10%, 25%, 50%, 75%) while
holding all other parameters at base values. We then ran a set of simulations examining the percent
change in K at break points from its starting value (± no change, 10%, 25%, 50%, 75%) while holding all
other parameters (including r) at base values. This led to a total of 40 scenarios (four starting values of r
with five percent changes in r and five percent changes in K).

We further evaluated how the magnitude of stochasticity in the system (as measured by the error term
ε_t) influenced algorithm performance. For generalizability of our simulation results, we simulated error
as a percentage of the mean population size, rather than as absolute value (as described in the model
above that we used for fitting the DSD). For each annual population size in the simulated dataset, a
random value was selected from a normal curve of mean 0 and standard deviation of τ (where τ = 1%,
2%, 5%, 10%, 15%) and multiplied by the expected population size generated from the deterministic
portion of the model.

We ran these simulations with all noise (τ) levels across all percent change values for r and K (with other
parameters held at base values) for a total of 50 additional scenarios (five percent noise values with five
percent changes in r and five percent changes in K). Finally, we tested the impact of time series length
by modifying the length of the simulated time series at five-year intervals over a range from 15 – 30
years (as the number of break point allows) while holding all other parameters constant, for four
additional scenarios. We generated 250 simulated datasets for each of the 94 possible scenarios
assuming breakpoint combinations with 0, 1, 2 and 3 breaks. Break point locations were randomly
selected from within the set of possible time points. In total, we generated 93,572 data sets that we
examined with our DSD algorithm. (Note that 94,000 simulations were run but simulations for higher
numbers of break points in shorter time series occasionally failed to converge; results for such
combinations are not presented).

We evaluated the DSD algorithm's performance for all test scenarios by examining its ability to identify
the true break points within the set of the best fitting break point combinations (i.e. the top ranked
break point combination and those break point combinations whose AICc values fell within two units of
the top ranked). We also examined the performance of the break-point weighting tool by calculating the
average weightings of all true and erroneous break points identified in the top performing model(s)
across all runs of a given scenario.

Results

Simulation Study

The scenario with the correct number of breaks and their locations was detected within the top
performing break point combination sets with >70% accuracy under nearly all parameterizations (Fig. 1).
Accuracy was generally lowest in time series with three break points but above 70% for most scenarios.
These results remained roughly consistent regardless of the value of the variance (σ^2) determining the
annual amount of environmental/sampling noise (Fig. 1 A). Results were similar across all r values tested
but performance of the DSD declined slightly when initial r was large (>2.0; Fig. 1 B). The DSD algorithm
had the highest accuracy with larger shifts in K (≥25%; Fig 1. C) and relatively smaller changes to r (≤25%;
Fig. 1 D). This result is somewhat counter-intuitive, as we would generally expect large shifts in all
parameters to be more easily detected. However, because nonlinear models produce chaotic dynamics
with high population growth rates (e.g., r > 2.3 in the Ricker model), a large shift in parameters could
potentially result in a situation where multiple break point fits would perform equally well. Finally, the
accuracy of the DSD algorithm decreased as scenario length increased, likely because of the factorial
increase in potential break point combinations with additional data in the time series (Fig. 1 E). Accuracy
was also lower in cases where the number of break points was high relative to the time series length
(e.g., 20 years and three breaks).

The breakpoint weighting analysis revealed that in the vast majority of cases, the average weight of a
true break exceeded a value of 0.8 (Fig. 2A-E), whereas the weight of erroneous breaks averaged less
than 0.2 in weight. The notable exception occurred when true breaks resulted from very small shifts in K
(Fig. 2 C). Thus, our results suggest the following decision rules to evaluate strength of evidence for a
break occurring at a given time point: when a weight of >0.8 is indicated for a break found by the DSD
algorithm, we can reasonably conclude this is a true break, and likewise, a break with a weight of <0.2
can reasonably assumed to be erroneous. Weight values intermediate to those two thresholds can be
interpreted as a quantification of the strength of evidence that a break occurred.

Case study applications

We tested the performance of the DSD algorithm with two case studies using population time series
data from field observations. Both case studies involve approximately two decades of observations of
economically or culturally important insect species: one examines an invasion process and the other
examines a population decline, both occurring over the same time period in recent history.

<u>Multicolored Asian ladybeetles in southwestern Michigan</u>

The 1994 invasion of multicolored Asian ladybeetles to southwestern Michigan, United States was
documented in monitoring data collected on agriculturally-important Coccinellidae (ladybeetles) in
landscapes dominated by field crops. Population density of ladybeetles is monitored in ten plant
communities weekly over the growing season using yellow sticky card glue traps starting in 1989 at the
Kellogg Biological Station at Michigan State University. We used data on the captures of adults at the
site from 1994-2017, culled at day of year 222 (August 10) to minimize the effect of year-to-year
variation in the sampling period. We then calculated the average number of adults captured per trap,
across all traps deployed within a sampling year, and used this value in our analysis. Detailed sampling
methodology is available in previous work [40–42].

Two break points, one occurring after 2000 and one occurring after 2005, were observed in the top
performing break point combination (Fig. 3 A, AICc=-18.02). However, the DSD algorithm indicated that
two additional break point combinations, a single break after 2000 (AICc=-17.46), and a no break series

(AICc=-17.64), had equivalent performance. Break weight analysis suggested a weight of 0.56 for the
2000 break, and a weight of 0.29 for the break after 2005. As these weights fall into a range
intermediate to the 0.2 and 0.8 decision rules, we conclude that there is moderately strong evidence of
a shift in dynamic rule after 2000, and moderate-weak evidence for a shift after 2005. The shift in 2000
is characterized by substantial increases in the values of K and r, with approximate increases of 75% and
40% over their initial estimates, respectively (Table 1). The shift in 2005 is characterized by a return to
parameter estimates that were nearly identical to those observed at the beginning of the time series
(Table 1, Fig. 3 B).

Our results can be explained in the context of the known ecology of this ladybeetle. Dynamics of the
ladybeetle invasion appear to be closely coupled with prey availability [41,43–45], which, in turn, is
driven by documented pest management practices (neonicotinoid insecticide use; [42]) leading to a
relatively simple pulsed change. The first shift in the dynamics of the Asian ladybeetle, after 2000,
corresponds to the well documented arrival and establishment of soybean aphid to North America, a
preferred prey item from the ladybeetle's native range [46,47]. The invasion of this aphid dramatically
increased resources available to the ladybeetle in habitats where the beetles were already well-
established [40], supporting both a higher carrying capacity and a greater intrinsic growth rate. The
second shift, after 2005, was less strongly supported, but coincides with the introduction and uptake of
a management strategy for aphids that incompletely controlled the prey item. Landscape-scale use of
neonicotinoid insecticides decreased prey numbers, particularly during the spring when aphids colonize
new hosts, which could limit early season reproduction of ladybeetles [42]. Indeed, in this case, we
would expect a weaker shift in dynamics as the prey item is incompletely controlled, and control tactics
were not uniformly adopted across the prey's range simultaneously.

Monarch butterflies in Mexican overwintering grounds

The eastern population of the North American monarch butterfly (*Danaus plexippus*) is migratory, with
the majority of individuals overwintering in large aggregations in Oyamel fir forests within the
transvolcanic mountains in the central region of Mexico [48]. Monarchs are highly dispersed over their
breeding season, occupying landscapes throughout the agricultural belt in central and eastern United
States and southern Canada [49]. As such, estimates of the overwintering population size can provide a
convenient and inclusive annual metric of the size of the eastern migratory population [50]. This
population of monarchs has been in dramatic decline in recent decades, although the degree and cause
of this decline is hotly debated [51]. We used data on the total area occupied by monarchs from 1995-
2017 (based on early winter surveys conducted in December) compiled by the World Wildlife Fund
Mexico (available at MonarchWatch; Lovett 2017).

The DSD algorithm estimated that the best break point combination fit for the monarch overwintering
data was a single break after 2003 (Fig. 4; AICc=120.18). However, the algorithm indicated that two
additional break point combinations, a single break after 2006 (AICc=121.87) and a two-break
combination of 2003 and 2008 (AICc=-121.86), had equivalent performance. The weight analysis
computed weights of 0.49, 0.14, and 0.26, for 2003, 2006, and 2008 respectively, suggesting that the
break at 2006 is erroneous and providing intermediate support for the 2003 and 2008 breaks. As with
our ladybeetle case study, the strength of evidence was strongest for the first break, and weaker for the
second break. The shift corresponds with a >50% reduction in K in 2003, and, if the secondary break is
taken at 2008, a further reduction of K nearing 50% again at that point (Table 1; Fig. 4 B).

The patterns we observe are consistent with a leading hypothesis to explain monarch population
decline. Loss of milkweed hostplants due to changing agricultural practices on Midwestern breeding
grounds [53,54] is hypothesized to be a major driver in the dynamics of this species. Changing herbicide
practices in central North America have largely eliminated milkweed hostplants from agricultural field
crops, with fairly consistent, low levels of milkweed on the landscape starting from about 2003-2005
[55]. Although glyphosate tolerant soybeans and maize were introduced to the US market in 1996 and
1998 respectively [56], actual glyphosate use lagged behind, with dramatic increases in use of the
pesticide in 1998-2003 in soybean, and 2007-2008 in maize [57].

However, additional drivers likely play a role in monarch processes given the uncertainty in our results.
Abiotic drivers of monarch population dynamics are complex and can interact at local, regional, and
continental scales [58]. Other studies have implicated climate [59], extreme weather events [60],
changing habitat availability on wintering grounds [61], and mortality during the fall migration [62,63] as
possible factors influencing monarch population dynamics. With many super-imposed drivers, monarch
dynamics are likely driven by both press and pulsed processes, making the detection of discrete break
points associated with dynamic shifts complicated.

Discussion

The DSD algorithm provides a novel tool for evaluating shifts in parameter values that govern density-
dependent populations, such as carrying capacity and population growth rates. As illustrated with our
simulations and case studies, the DSD algorithm can not only identify and quantify parameter changes
but also assess uncertainty in potential break points and help detect time frames where additional
research should be focused. We recommend that the model selection approach be used to identify a list
of potential break points and break point combinations, and the weighting tool be used to evaluate the
strength of evidence for each potential break, providing a clear direction to focus downstream research
on changing dynamic processes. Characterizing dynamic transitions in population time series data has
been hindered by a lack of a common, accessible, and empirical approaches [2]. Most assessments of
break points in ecological research are *ad hoc* [12,14–17], introducing the potential for bias in break
point selection, particularly in cases where nonlinear dynamics are likely to occur. Our DSD algorithm
directly addresses this gap in tools, using an information-theoretic, model selection approach and a
framework that can incorporate a variety nonlinear population models to assess ecological processes.

The core novelty of our tool lies in the model selection procedure used within the DSD algorithm, which
allows for greater accuracy than common break point detection models [1]. The DSD additionally allows
users to assess the confidence in a given break point, as well as providing a measure of which break
points are likely to appear together, if multiple dynamic shifts have occurred. Information-theoretic
approaches such as model selection using AIC, may be prone to over-fitting, particularly when data are
limited [35]. Thus, we used AICc, the Akaike Information Criterion corrected for small sample sizes, as
the selection criterion within the DSD algorithm. AICc is generally recommended in place of AIC in
situations where small samples sizes are being examined as it incorporates a penalty term that is
inversely related to the number of observations. As sample size increases, the penalty for model
complexity is reduced and AICc approaches AIC [35]. Break point detection methods must incorporate a
compromise between sensitivity and penalty for over-parameterization [1]. We examined the
performance of both AICc (here) and AIC (in Appendix S1) and found that using AICc performed best for
our simulated data. The DSD algorithm aides in the interpretation of break points by incorporating a

metric based on Akaike weights, which allows users to assess the relative 'strength' of multiple breaks.
Where many tools aim to identify points at which parameter changes occur, the DSD algorithm provides
a measure of the confidence in each break, as well as a measure of how differing break sets perform in
explaining variation in the data relative to each other.

The performance of the DSD algorithm was fairly stable among the break point simulations we tested.
We found that the amount of environmental/sampling noise (ranging from 1-15% of the population size)
had little effect on algorithm performance (Fig. 1 A). Other input conditions had relatively greater
impacts on the performance of the DSD algorithm, depending on which parameter was changed and by
how much. Large shifts in a population's carry capacity were more easily detected than smaller shifts
(Fig. 1 C). However, large changes in population growth rate were harder to detect, but this effect was
most pronounced when simulated data contained multiple breaks (Figs. 1, 2 D). Although larger shifts in
regression parameters would, intuitively, lead to a higher likelihood of detection, large shifts in growth
rate are also more likely to induce variations in transient dynamics in the years immediately following
the shift, potentially making the timing of shifts more difficult to pinpoint. Similarly, longer time series
yielded results that were more error prone (Fig. 1-2 E). This is likely because there were simply more
possible break-point combinations for the algorithm to select from and because the penalty for
increasing parameterization (i.e. AICc) decreases as sample sizes grow (leading to increasing likelihood
of identifying extra, erroneous breaks).

In our case studies, we found interpretation of the ladybeetle example was straightforward (Fig. 3). Our
top break point combination and the equivalently-performing set did not contain contradictory
information: each candidate set was simply a subset of breakpoints from the most complex set, and only
two break points were found. Both of these break points were associated with moderate or greater
weights, although the values of these break weights were in the intermediate range (i.e., between 0.2
and 0.8), suggesting breaks in natural systems may not be as well behaved as those in simulated data.
The monarch butterfly case study results were slightly more ambiguous, as the model selection tool
identified a break that the weighting tool suggested was erroneous (Fig. 4). Weights of the two most
strongly-supported breaks were numerically similar to those of the ladybeetle case study, and are also
interpretable with knowledge of the study system. However, the model selection results suggest
additional, superimposed processes may be affecting monarch population dynamics, creating a noisier
signal.

The DSD algorithm is readily adaptable to other population models and, indeed, potentially to other
nonlinear processes. Density-dependent population growth has the potential to mask transition points
because of its inherent nonlinear structure. For example, transient dynamics occurring immediately
after a temporary disturbance can result in a change in population size, but not necessarily in the rules
governing dynamics. We used the Ricker model as the core population model within the algorithm
because it had a number of useful characteristics, namely its simple parameterization and realistic
behavior [27]. However, simple density-dependent population models including Ricker, Beverton-Holt,
and logistic models have similar performance in predicting outbreaks of insects driven by food limitation
[64]. Indeed, in early iterations examining the DSD, we fit both logistic and Ricker models to real data
and found that the two models ranked break point combinations nearly identically, even while the
Ricker model generally provided a better fit for the data [42]. Thus, we expect the DSD algorithm would
have similar performance across models with similar structures, but performance may vary with other
model structures, particularly those that incorporate additional terms.

We recommend users carefully consider the strengths and limitations of the DSD algorithm in the
context of their own data. For example, if changes to parameter values occur frequently (e.g., less than
3-4 years or time periods), the frequency of shifts would violate the constraints placed on the DSD to
prevent overfitting. We also observed that the likelihood of identifying erroneous break points increased
as time series length increased. Thus, in cases where a long time series exists, but a particular time
period is of interest, the DSD algorithm could be used on the time period of interest alone to minimize
the likelihood of distracting or erroneous results. The DSD algorithm functions as a method for
identifying break points within time series data and quantifying the strength of evidence for each
potential break point. When interpreted in the context of species biology, the DSD algorithm has the
potential to aid management decisions, identify critical drivers of change in species' dynamics, and help
determine where best to focus additional research efforts.

<u>Acknowledgements</u>

398   **References**

399   1.   Andersen T, Carstensen J, Hernández-García E, Duarte CM. Ecological thresholds and regime shifts:
400   approaches to identification. Trends in Ecology & Evolution. 2009;24: 49–57.
401   doi:10.1016/j.tree.2008.07.014

402   2.   Bestelmeyer BT, Ellison AM, Fraser WR, Gorman KB, Holbrook SJ, Laney CM, et al. Analysis of
403   abrupt transitions in ecological systems. Ecosphere. 2011;2: art129. doi:10.1890/es11-00216.1

404   3.   Scheffer M, Carpenter S, Foley JA, Folke C, Walker B. Catastrophic shifts in ecosystems. Nature.
405   2001;413: 591–596. doi:10.1038/35098000

406   4.   Scheffer M, Carpenter SR. Catastrophic regime shifts in ecosystems: linking theory to observation.
407   Trends in Ecology & Evolution. 2003;18: 648–656. doi:10.1016/j.tree.2003.09.002

408   5.   May RM. Thresholds and breakpoints in ecosystems with a multiplicity of stable states. Nature.
409   1977;269: 471–477. doi:10.1038/269471a0

410   6.   Ducré-Robitaille J-F, Vincent LA, Boulet G. Comparison of techniques for detection of
411   discontinuities in temperature series: DETECTING DISCONTINUITIES IN TEMPERATURE SERIES.
412   International Journal of Climatology. 2003;23: 1087–1101. doi:10.1002/joc.924

413   7.   Rodionov SN. A sequential algorithm for testing climate regime shifts: ALGORITHM FOR TESTING
414   REGIME SHIFTS. Geophysical Research Letters. 2004;31: n/a-n/a. doi:10.1029/2004GL019448

415   8.   Beaugrand G., Conversi A., Chiba S., Edwards M., Fonda-Umani S., Greene C., et al. Synchronous
416   marine pelagic regime shifts in the Northern Hemisphere. Philosophical Transactions of the Royal
417   Society B: Biological Sciences. 2015;370: 20130272. doi:10.1098/rstb.2013.0272

418   9.   Zampolli F. Optimal monetary policy in a regime-switching economy: The response to abrupt shifts
419   in exchange rate dynamics. Journal of Economic Dynamics and Control. 2006;30: 1527–1567.
420   doi:10.1016/j.jedc.2005.10.013

421   10.   Holling CS. Understanding the Complexity of Economic, Ecological, and Social Systems. Ecosystems.
422   2001;4: 390–405. doi:10.1007/s10021-001-0101-5

423   11.   Bjørnstad ON, Grenfell BT. Noisy Clockwork: Time Series Analysis of Population Fluctuations in
424   Animals. Science. 2001;293: 638. doi:10.1126/science.1062226

425   12.   Hare SR, Mantua NJ. Empirical evidence for North Pacific regime shifts in 1977 and 1989. Progress
426   in Oceanography. 2000;47: 103–145. doi:10.1016/S0079-6611(00)00033-1

427   13.   Carpenter SR, Brock WA, Cole JJ, Kitchell JF, Pace ML. Leading indicators of trophic cascades.
428   Ecology Letters. 2008;11: 128–138. doi:10.1111/j.1461-0248.2007.01131.x

429   14.   Weimerskirch H, Inchausti P, Guinet C, Barbraud C. Trends in bird and seal populations as
430   indicators of a system shift in the Southern Ocean. Antarctic Science. 2003;15: 249–256.

431   15.   Berryman A, Lima M. Deciphering the effects of climate on animal populations: diagnostic analysis
432   provides new interpretation of Soay sheep dynamics. The American Naturalist. 2006;168: 784–795.

16.　Toms JD, Lesperance ML. Piecewise regression: a tool for identifying ecological thresholds.
Ecology. 2003;84: 2034–2041. doi:10.1890/02-0472

17.　Knapp AK, Smith MD, Hobbie SE, Collins SL, Fahey TJ, Hansen GJA, et al. Past, present, and future
roles of long-term experiments in the LTER Network. Bioscience. 2012;62: 377–389.
doi:10.1029/2008gb003336

18.　Braun JV, Muller H-G. Statistical Methods for DNA Sequence Segmentation. Statistical Science.
1998;13: 142–162.

19.　Zeileis A, Leisch F, Hornik K, Kleiber C. strucchange. An R package for testing for structural change
in linear regression models. 2001.

20.　Killick R, Eckley I. changepoint: An R package for changepoint analysis. Journal of Statistical
Software. 2014;58: 1–19.

21.　Priyadarshana W, Sofronov G. Multiple break-points detection in array CGH data via the cross-
entropy method. IEEE/ACM Transactions on Computational Biology and Bioinformatics (TCBB).
2015;12: 487–498.

22.　Mauget SA. Multidecadal Regime Shifts in U.S. Streamflow, Precipitation, and Temperature at the
End of the Twentieth Century. Journal of Climate. 2003;16: 3905–3916. doi:10.1175/1520-
0442(2003)016<3905:MRSIUS>2.0.CO;2

23.　Jenouvrier S, Weimerskirch H, Barbraud C, Park Y-H, Cazelles B. Evidence of a shift in the cyclicity
of Antarctic seabird dynamics linked to climate. Proceedings of the Royal Society B: Biological
Sciences. 2005;272: 887–895. doi:10.1098/rspb.2004.2978

24.　Cazelles B, Chavez M, Berteaux D, Ménard F, Vik JO, Jenouvrier S, et al. Wavelet analysis of
ecological time series. Oecologia. 2008;156: 287–304. doi:10.1007/s00442-008-0993-2

25.　Carl Boettiger, Alan Hastings. Early warning signals and the prosecutor's fallacy. Proceedings of the
Royal Society B: Biological Sciences. 2012;279: 4734–4739. doi:10.1098/rspb.2012.2085

26.　Royall R. The Likelihood Paradigm for Statistical Evidence. in ML Taper and SR Lele, eds. The nature
of scientific evidence. University of Chicago Press, Chicago; 2004.

27.　Turchin P. Complex population dynamics: a theoretical/empirical synthesis. Princeton University
Press; 2003.

28.　May RM. Simple mathematical models with very complicated dynamics. Nature. 1976;261: 459–
467. doi:10.1038/261459a0

29.　Barraquand F, Louca S, Abbott KC, Cobbold CA, Cordoleani F, DeAngelis DL, et al. Moving forward
in circles: challenges and opportunities in modelling population cycles. Ecology letters. 2017;20:
1074–1092.

30.　Boettiger C. From noise to knowledge: how randomness generates novel phenomena and reveals
information. Ecology Letters. 2018;21: 1255–1267. doi:10.1111/ele.13085

31.   Gadrich T, Katriel G. A Mechanistic Stochastic Ricker Model: Analytical and Numerical
Investigations. Int J Bifurcation Chaos. 2016;26: 1650067. doi:10.1142/S021812741650067X

32.   Ricker WE. Stock and Recruitment. J Fish Res Bd Can. 1954;11: 559–623. doi:10.1139/f54-039

33.   Brook BW, Bradshaw CJ. Strength of evidence for density dependence in abundance time series of
1198 species. Ecology. 2006;87: 1445–1451.

34.   Hall AR, Osborn DR, Sakkas N. Inference on Structural Breaks using Information Criteria. The
Manchester School. 2013;81: 54–81. doi:10.1111/manc.12017

35.   Burnham KP, Anderson DR. Model selection and multimodal inference: a practical information-
theoretic approach. 2nd ed. New York: Springer Science + Business Media, LLC; 2002.

36.   Wagenmakers E-J, Farrell S. AIC model selection using Akaike weights. Psychonomic Bulletin &
Review. 2004;11: 192–196. doi:10.3758/BF03206482

37.   R Development Core Team. R: A Language and Environment for Statistical Computing 3.3.3. R
Foundation for Statistical Computing. 2017. Available: http://www.R-project.org

38.   Elzhov TV, Mullen KM, Spiess A-N, Bolker B. minpack. lm: R Interface to the Levenberg-Marquardt
Nonlinear Least-Squares Algorithm Found in MINPACK. 2016.

39.   Bahlai C. cbahlai/dynamic_shift_detector: The Dynamic Shift Detector. Zenodo; 2019.
doi:10.5281/zenodo.3368486

40.   Bahlai CA, Colunga-Garcia M, Gage SH, Landis DA. Long term functional dynamics of an
aphidophagous coccinellid community are unchanged in response to repeated invasion. PLoS One.
2013;8: e83407. doi:10.1371/journal.pone.0083407

41.   Bahlai C, Colunga-Garcia M, Gage S, Landis D. The role of exotic ladybeetles in the decline of native
ladybeetle populations: evidence from long-term monitoring. Biol Invasions. 2015;17: 1005–1024.
doi:10.1007/s10530-014-0772-4

42.   Bahlai CA, vander Werf W, O'Neal M, Hemerik L, Landis DA. Shifts in dynamic regime of an invasive
lady beetle are linked to the invasion and insecticidal management of its prey. Ecological
Applications. 2015 [cited 14 May 2015]. doi:10.1890/14-2022.1

43.   Bahlai CA, Sears MK. Population dynamics of Harmonia axyridis and Aphis glycines in Niagara
Peninsula soybean fields and vineyards. Journal of the Entomological Society of Ontario. 2009;140:
27–39.

44.   Heimpel G, Frelich L, Landis D, Hopper K, Hoelmer K, Sezen Z, et al. European buckthorn and Asian
soybean aphid as components of an extensive invasional meltdown in North America. Biological
Invasions. 2010;12: 2913–2931. doi:10.1007/s10530-010-9736-5

45.   Rhainds M, Yoo HJS, Kindlmann P, Voegtlin D, Castillo D, Rutledge C, et al. Two-year oscillation
cycle in abundance of soybean aphid in Indiana. Agricultural and Forest Entomology. 2010;12:
251–257.

46.   Ragsdale DW, Voegtlin DJ, O'Neil RJ. Soybean aphid biology in North America. Annals of the
Entomological Society of America. 2004;97: 204–208.

47.   Wu Z, Schenk-Hamlin D, Zhan W, Ragsdale DW, Heimpel GE. The soybean aphid in China: a
historical review. Annals of the Entomological Society of America. 2004;97: 209–218.

48.   Urquhart FA, Urquhart NR. Autumnal migration routes of the eastern population of the monarch
butterfly (Danaus p. plexippus L.; Danaidae; Lepidoptera) in North America to the overwintering
site in the Neovolcanic Plateau of Mexico. Can J Zool. 1978;56: 1759–1764. doi:10.1139/z78-240

49.   Flockhart DTT, Brower LP, Ramirez MI, Hobson KA, Wassenaar LI, Altizer S, et al. Regional climate
on the breeding grounds predicts variation in the natal origin of monarch butterflies overwintering
in Mexico over 38 years. Glob Change Biol. 2017;23: 2565–2576. doi:10.1111/gcb.13589

50.   Prysby MD, Oberhauser KS. Temporal and geographic variation in monarch densities: citizen
scientists document monarch population patterns. The monarch butterfly: Biology and
conservation. 2004; 9–20.

51.   Sarkar S. What Is Threatening Monarchs? BioScience. 2017;67: 1080–1080.
doi:10.1093/biosci/bix120

52.   Lovett J. Monarch Population Status. Monarch Watch; 2017. Available:
http://monarchwatch.org/blog/2017/02/11/monarch-population-status-30/

53.   Hartzler RG. Reduction in common milkweed (Asclepias syriaca) occurrence in Iowa cropland from
1999 to 2009. Crop Protection. 2010;29: 1542–1544. doi:10.1016/j.cropro.2010.07.018

54.   Pleasants JM, Oberhauser KS. Milkweed loss in agricultural fields because of herbicide use: effect
on the monarch butterfly population. Insect Conservation and Diversity. 2013;6: 135–144.
doi:10.1111/j.1752-4598.2012.00196.x

55.   Zaya DN, Pearse IS, Spyreas G. Long-Term Trends in Midwestern Milkweed Abundances and Their
Relevance to Monarch Butterfly Declines. BioScience. 2017;67: 343–356.
doi:10.1093/biosci/biw186

56.   Powles SODSB. Glyphosate-resistant crops and weeds: now and in the future. 2010.

57.   Baker NT. Estimated annual agricultural pesticide use by crop group for states of the conterminous
United States, 1992-2014. National Water Quality Assessment Program. 2017.
doi:10.5066/F7NP22KM

58.   Saunders SP, Ries L, Oberhauser KS, Thogmartin WE, Zipkin EF. Local and cross-seasonal
associations of climate and land use with abundance of monarch butterflies Danaus plexippus.
Ecography. 2017; n/a-n/a. doi:10.1111/ecog.02719

59.   Zipkin EF, Ries L, Reeves R, Regetz J, Oberhauser KS. Tracking climate impacts on the migratory
monarch butterfly. Glob Change Biol. 2012;18: 3039–3049. doi:10.1111/j.1365-2486.2012.02751.x

60. Brower LP, Kust DR, Rendon-Salinas E, Serrano EG, Kust KR, Miller J, et al. Catastrophic winter
storm mortality of monarch butterflies in Mexico during January 2002. The Monarch butterfly:
biology and conservation. 2004; 151–166.

61. Vidal O, Rendón-Salinas E. Dynamics and trends of overwintering colonies of the monarch butterfly
in Mexico. Biological Conservation. 2014;180: 165–175. doi:10.1016/j.biocon.2014.09.041

62. Davis AK, Altizer S. New Perspectives on Monarch Migration, Evolution, and Population Biology.
Monarchs in a Changing World: Biology and Conservation of an Iconic Butterfly. 2015; 203.

63. Saunders SP, Ries L, Neupane N, Ramírez MI, García-Serrano E, Rendón-Salinas E, et al. Multiscale
seasonal factors drive the size of winter monarch colonies. Proc Natl Acad Sci USA. 2019;116:
8609. doi:10.1073/pnas.1805114116

64. Abbott KC, Dwyer G. Food Limitation and Insect Outbreaks: Complex Dynamics in Plant-Herbivore
Models. Journal of Animal Ecology. 2007;76: 1004–1014.

Supporting Information Legends

S1- Additional Simulations using AIC for break point selection

S2- Description of functions used in the Dynamic Shift Detector (DSD) algorithm

Table 1: Ricker model parameter values for each phase between break points resulting from fitting
population data of 1) multicolored Asian ladybeetles from Michigan, USA (1994-2017), and 2) the area
occupied by monarch butterflies in their winter habitat in central Mexico (1995-2017). The parameter *r*
is the per capita yearly intrinsic rate of increase and *K* is the carrying capacity (e.g., average number of
adult ladybeetles captured per trap annually and hectares occupied by monarchs annually).

Species	Years in subset	*r* (± SE)	*K* (± SE)
Ladybeetle	1994-2000	1.3 ± 0.3	0.31 ± 0.02
Harmonia axyridis	2001-2005	2.3 ± 0.3	0.43 ± 0.03
	2006-2017	1.6 ± 0.3	0.27 ± 0.03
Monarch	1995-2003	1.0 ± 0.5	10.1 ± 1.9
Danaus plexippus	2004-2008	1.6 ± 0.2	5.6 ± 0.3
	2009-2017	1.2 ± 0.4	2.8 ± 0.5

Figure captions

Figure 1: Performance of the Dynamic Shift Detector (DSD) algorithm under varying parameter values.
Proportion of simulation results in which the true break scenario was detected within the top break point combinations as identified by the DSD implemented with an underlying Ricker model with varied A) noise (in the form of normally distributed error), B) starting values of the r parameter, C) percent changes in the K parameter, D) percent changes in r, and E) simulated time series length. Sets of 0, 1, 2 and 3 break points were randomly generated from within the set of possible values, and 250 datasets were simulated for each scenario. In each panel, other variables (that were not being varied) were held constant at their base values (i.e., noise=2%; starting value of r = 2; change in r = ±25%; change in K = ±75%; time series length = 20 years). Trends within a set of scenarios (grey lines) are illustrated with a third-order GAM smoothing line.

Figure 2: Average break weight of break points detected under varying parameterization conditions.
Average weights of break points identified by the Dynamic Shift Detector algorithm reflecting true parameterization conditions (diamonds) or erroneous breaks suggested by the algorithm (triangles) under varied A) noise (in the form of normally distributed error), B) starting values of the r parameter, C) percent changes in the K parameter, D) percent changes in r, and E) simulated time series length. Sets of 0, 1, 2 and 3 break points were randomly generated from within the set of possible values, and 250 datasets were simulated for each scenario. In each panel, other variables (that were not being varied) were held constant at their base values (i.e., noise=2%; starting value of r = 2; change in r = ±25%; change in K = ±75%; time series length = 20 years). Trends within a set of scenarios (grey lines) are illustrated with a third-order GAM smoothing line.

Figure 3: Dynamic Shift Detector breaks and Ricker model fits for an invasive species. Population data documenting the invasion of multicolored Asian ladybeetle in Michigan, USA from 1994-2017. A) Time series data showing the average number of adults captured, per trap, per year. Vertical blue lines indicate years in which dynamic shifts occurred, as estimated by the Dynamic Shift Detector algorithm. B) Ricker fits of time series data segments. Ladybeetle art by M. Broussard, used under a CC-BY 3.0 license.

Figure 4: Dynamic Shift Detector breaks and Ricker model fits for a species of conservation concern.
Population data documenting the area occupied by monarch butterflies in their winter habitat in central Mexico from 1995-2017. A) Time series data showing the total area occupied by overwintering monarchs each year in December. Vertical blue lines indicate years in which dynamic shifts occurred, as estimated by the Dynamic Shift Detector algorithm. B) Ricker fits of time series data segments. Butterfly art by D. Descouens and T.M. Seesey, used under a CC-BY 3.0 license.