Exercise-stimulated muscle ROS production and glucose uptake requires NADPH oxidase 2

Carlos Henríquez-Olguin[1,2], Jonas R. Knudsen[1], Steffen H. Raun[1], Zhencheng Li[1], Emilie Dalbram[3], Jonas T. Treebak[3], Lykke Sylow[1], Rikard Holmdahl[4], Erik A. Richter[1], Enrique Jaimovich[2], and Thomas E. Jensen[1,*]

1. Department of Nutrition, Exercise and Sports, University of Copenhagen, Denmark

2. Center of Exercise, Metabolism and Cancer, Biomedical Sciences Institute, Universidad de Chile.

3. Novo Nordisk Foundation Center for Basic Metabolic Research, Research Program in Integrative Metabolism and Environmental Influence, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

4. Section for Medical Inflammation Research, Department of Medical Biochemistry and Biophysics, Karolinska Institute, Stockholm, Sweden.

* Corresponding author and person to who reprint requests should be addressed:

Thomas E. Jensen, PhD

Department of Nutrition, Exercise, and Sports (NEXS)

University of Copenhagen

Universitetsparken 13, DK-2100 Copenhagen, Denmark

Phone: +45 3532 1757

Email: TEJensen@NEXS.ku.dk

ABSTRACT

Reactive oxygen species (ROS) act as intracellular compartmentalized second messengers mediating metabolic stress-adaptation. In skeletal muscle fibers, ROS have been suggested to stimulate glucose transporter 4 (GLUT4)-dependent glucose transport during artificially evoked contraction *ex vivo* but whether myocellular ROS production is stimulated by *in vivo* exercise to control metabolism is unclear. Here, we combined exercise in humans and mice with fluorescent dyes, genetically-encoded biosensors, and NADPH oxidase 2 (NOX2) loss-of-function models to demonstrate that NOX2 is the main source of ROS during moderate-intensity exercise in skeletal muscle. Furthermore, two NOX2 loss-of-function mouse models lacking either p47phox or Rac1 presented striking phenotypic similarities, including greatly reduced exercise-stimulated glucose uptake and GLUT4 translocation. These findings indicate that NOX2 is a major myocellular ROS source regulating glucose transport capacity during moderate intensity exercise.

INTRODUCTION

A single bout of exercise prompts a rapid adaptive increase in energy substrate metabolism in contracting skeletal muscle to meet the increased demand for ATP production. Understanding the molecular signal transduction pathways, that orchestrate these metabolic responses in muscle has broad implications, from optimizing athletic performance, to understanding fundamental stress-adaptive responses at the cellular level, to prevention and treatment of aging- and lifestyle-related diseases in muscle and other tissues [1].

The production of oxygen-derived free radicals and derivatives thereof, collectively referred to as reactive oxygen species (ROS), increases in skeletal muscle during exercise and has long been considered to mediate adaptive responses to both acute bouts of exercise and chronic exercise training [2]. Although historically viewed as a by-product of oxidative metabolism in mitochondria, it has been suggested that ROS may be produced enzymatically by several potential sources in exercising muscle, including NADPH oxidase (NOX) and xanthine oxidase [3]. In contrast, mitochondria are likely not the major source of exercise-induced ROS [4, 5]. The major ROS source in the context of physiological *in vivo* exercise, however, remains uncertain due to the difficulty in measuring and quantifying exercise-stimulated ROS production [6].

Research in the past 60 years has established that glucose transport *in vivo* is controlled by a coordinated increase in glucose delivery, glucose transport into muscle fibers and intracellular metabolism [7, 8]. Among these events, the insulin-independent translocation of glucose transporter protein 4 (GLUT4) from intracellular storage depots to the cell surface to facilitate glucose entry appears to be a key molecular event [9, 10]. Studies in

isolated *ex vivo* incubated skeletal muscles from rodents have shown that glucose

transport in response to electrically stimulated contraction and mechanical stress is

lowered by antioxidants [11, 12], suggesting that ROS signal to increase glucose transport

[13]. Interestingly, the small GTPase Rac1 is activated by muscle contraction and passive

stretch and is necessary for both stimuli to increase glucose uptake in isolated mouse

muscles [14, 15]. Rac1 is best known to bind and orchestrate regulators of actin

remodeling[16], but the recruitment of GTP-loaded active Rac GTPase isoforms in

conjunction with a complex of other NOX2 regulatory proteins (p67phox, p47phox and

p40phox) is also essential to stimulate superoxide (O_2^-)-production by the membrane-

bound NOX2 complex[17]. However, whether NOX2-induced ROS production regulates

glucose uptake in vivo and if the reduced exercise-stimulated glucose uptake observed

in Rac1 KO muscles[18] is due to reduced ROS production, is currently unknown.

In the present study, we took advantage of recent methodological approaches in the

ROS signaling field, including a method for preservation of *in vivo* ROS modifications

and genetically-encoded ROS biosensors [19]. This allowed us to measure both general

and localized NOX2-specific ROS production in wild-type (WT) mice and mice lacking

NOX2 activity due to the absence of either the Rac1 or p47phox regulatory subunits.

Using these approaches, we show that NOX2 is activated by moderate intensity exercise

and is the predominant source of ROS production under these conditions. Moreover, a

large reduction in exercise-stimulated glucose uptake and GLUT4 translocation were

shared features between the two NOX2 loss-of-function models. Collectively, these

results imply that NOX2 is a major source of ROS production during moderate-intensity

exercise and that Rac1 is required for GLUT4 translocation and glucose uptake due to

its essential role in NOX2 activation.

RESULTS

Moderate intensity exercise increase ROS in human muscle

Plasma redox markers are increased during both moderate- and high-intensity exercise

in humans[20]. However, whether muscle ROS production increases during moderate-

intensity exercise in humans is unknown. To test this, we estimated total oxidant

production in skeletal muscle from healthy young men before and after a single 30 min

bout of bicycle ergometer exercise at a moderate 65% peak power output exercise

intensity using the redox-sensitive dye dichlorodihydrofluorescein diacetate (DCF). We

found that exercise stimulated a 86% increase in DCF oxidation in human skeletal

muscle (Fig. 1A), which was accompanied by an expected increase in the

phosphorylation state of known exercise-responsive proteins (Fig. 1B). This shows that

moderate-intensity exercise increases ROS in human skeletal muscle.

ROS production during exercise requires NOX2 activity in mice

To dissect the ROS source during exercise, we used a previously described p47phox-

mutated mice (*ncf1**) harboring a loss-of-function mutation in the regulatory NOX2

subunit, p47phox [21]. An acute moderate-intensity treadmill exercise bout (65% maximal

running speed) for 20 min increased DCF oxidation (+ 45%) in WT mice, a response

that was completely abolished in p47phox-deficient *ncf1** mice (Fig. 1C). This

observation was found to be independent of alterations in antioxidant enzyme

abundance in TA muscles from *ncf1** mice compared to WT mice (Fig. S1).

To substantiate that NOX2 activity was required for exercise-stimulated ROS

production, we used inducible muscle-specific Rac1 knockout mice (Rac1 imKO),

which are predicted to lack functional NOX2. Similar to the *ncf1** mice, exercise-

stimulated DCF oxidation was completely absent in Rac1 imKO compared to WT

littermates (Fig. S-2C). To directly demonstrate NOX2 activation during *in vivo*

exercise, we adapted a recently described redox histology method [19] that enabled the

preservation and visualization of the redox state of a transfected oxidation-sensitive

NOX2 activity reporter, p47-roGFP [22], in mouse muscle sections. In WT mice,

electrical stimulation *in vitro* induced p47-roGFP oxidation in FDB fibers (Fig. S2B), as

did treadmill exercise in TA muscle (Fig.1D). In contrast, exercise or electrical

stimulation failed to increase p47-roGFP oxidation in Rac1 imKO. Importantly, the lack

of response in Rac1 imKO mice occurred without changes in antioxidant enzymes

abundance (Fig. S2B). Collectively, these results strongly suggest that NOX2 is

activated and constitutes a source of ROS production during endurance-type exercise in

skeletal muscle.

NOX2 is required for exercise-stimulated glucose uptake

Given that Rac1 imKO mice display a severe reduction in treadmill exercise-stimulated

glucose uptake and GLUT4 translocation [18], we reasoned that if Rac1 was acting via

NOX2 then *ncf1** mice should show a similar reduction. We first conducted a general

characterization of *ncf1** mice. Similar total body weight but reduced fat mass was

observed in *ncf1** mice compared to age-matched WT mice (Fig. S3). Respiratory

exchange ratio (RER) was lower in *ncf1** mice during the light phase compared to WT

mice (Fig. 2A), despite a similar habitual activity observed between genotypes (Fig.

2B). Thus, *ncf1** mice, similar to Rac1 imKO mice, display a shift towards fat oxidation

at rest and during fasting.

Next, we determined whether NOX2-dependent ROS generation is required for

exercise-stimulated glucose uptake. Each mouse was exercised at a relative work load

corresponding to 65% of its maximum running capacity for 20 min. Exercise-induced 2-

[^{3}H] deoxyglucose (2DG) uptake was markedly attenuated in Quad, SOL and TA

muscles from *ncf1** mice compared to WT (Fig. 2C-E). Importantly, the reduction of

exercise-stimulated glucose uptake in *ncf1* mice* was not due to differences in maximal

running capacity (Fig. 2F), blood glucose levels (Fig. 2G) or intramuscular energetics

evaluated as NAD+ and NADH levels (Fig. 2H-I). However, plasma lactate levels were

increased by exercise in WT but not in *ncf1** mice (Fig. 2J), consistent with a reduced

reliance on glycolysis for energy production.

NOX2 is required for exercise-stimulated GLUT4 translocation

Next, we tested if the reduction in exercise-stimulated glucose uptake in *ncf1** mice

could be explained by reduced GLUT4 translocation as seen in the Rac1 imKO [18].

Consistent with this notion, the exercise-stimulated increase in surface-membrane

GLUT4-GFP-*myc* in TA muscle from WT was virtually abolished in *ncf1** mice (Fig.

3A). Importantly, total protein abundance of GLUT4 and HKII (Fig. 3B-D and Fig. S4)

did not differ between genotypes nor did protein abundance of the mitochondrial

electron transport chain complexes (Fig. 3E-F and Fig. S5). Taken together, these data

demonstrate that *ncf1** mice phenocopy the Rac1 imKO mice in terms of pronounced

impairments in exercise-stimulated muscle glucose uptake and GLUT4 translocation[18].

This strongly suggests that the common denominator between p47phox and Rac1,

NOX2 activation, is required for exercise-stimulated glucose uptake and GLUT4

translocation.

Exercise-stimulated p38 MAPK$^{Thr180/Tyr182}$ phosphorylation is reduced in *ncf1*

quadriceps muscle

Exercise-induced ROS production has been suggested to activate a number of kinases

linked to glucose uptake-regulation in skeletal muscle[23]. Interestingly, p-p38

MAPK$^{Thr180/Tyr182}$ levels were lower in quadriceps muscle of *ncf1** mice compared to

WT mice after exercise, but not in soleus or TA muscles (Fig. 4). The phosphorylation

of ERK $^{Thr202/Tyt204}$, AMPK^{Thr172}, and its substrate ACC^{Ser79} did not differ significantly

between genotypes (Fig. 4) nor did total $\alpha2$ AMPK, p38 MAPK, and ERK1/2

expression in any of the studied muscles (Fig. S6). Taken together, this makes reduced

signaling by these kinases an unlikely explanation of the consistent reduction in

exercise-stimulated glucose uptake in *ncf1** vs. WT mice across muscles.

DISCUSSION

The source of ROS during exercise and the role of ROS in the acute and chronic

adaptations to exercise, in particular in the context of non-damaging moderate intensity

in vivo exercise, has remained undetermined for more than 30 years [24]. The current

study observed that moderate-intensity endurance exercise acutely increased ROS levels

in both humans and mice. Strikingly, functional NOX2 was required for the exercise-

stimulated ROS increase and is thus a major source of ROS during moderate intensity

exercise. Furthermore, NOX2 activity was a major requirement for exercise-stimulated

GLUT4 translocation and glucose uptake in skeletal muscle, providing a likely

molecular explanation of why Rac1 imKO mice have a strong impairment of exercise-

stimulated GLUT4 translocation and glucose uptake.

We and others have shown that the small rho family GTPase Rac1 is necessary for

insulin-stimulated, muscle-contraction and passive stretch-induced skeletal muscle

glucose uptake $^{14, 15, 25}$. We have previously found this Rac1-dependency of glucose

uptake and GLUT4 translocation to be more pronounced during *in vivo* treadmill

exercise compared to in vitro contractions [18]. The GLUT4 trafficking field has mostly focused on Rac1 as an orchestrator of the dynamic reorganization of the cortical actin cytoskeleton consisting of the cytoplasmic β and γ actin-isoforms [16]. This actin-reorganization was shown to be crucial for insulin-stimulated GLUT4 translocation in muscle cells [26]. However, the cytoplasmic β and γ actin-isoforms are both downregulated during muscle differentiation and whether dynamic reorganization occurs in adult skeletal muscle fibers similarly to in cells is presently unclear [27]. Our present data suggest that NOX2 is a vital downstream mediator of the effect of Rac1 on exercise-induced glucose uptake and GLUT4 translocation since the Rac1 imKO phenotype is shared by mice lacking another essential regulatory subunit of NOX2, p47phox. In addition, our *in vivo* glucose uptake data provide a likely physiological mechanism explaining why exogenous antioxidants inhibit *ex vivo* contraction- and stretch-stimulated glucose uptake [11, 12]. The relative dependence on actin and NOX2 and whether NOX2 and actin interact in muscle to regulate glucose uptake, as suggested in other cell types[28], should be clarified in future studies.

Due to the uncertainties regarding the downstream events mediating NOX2 signal transduction, we utilized a classical, non-ROS-specific redox-sensitive fluorescent DCF which, upon oxidation, is converted into fluorescent DCF [29]. DCF fluorescence should thus be regarded as a general measure of total cellular oxidant levels rather than a specific measure of a particular ROS[30]. We observed an increase in DCF fluorescence in exercising mice and humans. Remarkably, despite the non-specific nature of DCF oxidation, we found that the increase in exercise-stimulated DCF fluorescence was virtually abolished in mice lacking either Rac1 or p47phox components of NOX2. Regardless of the exact ROS measured, this indicates that NOX2 is a predominant source of ROS in skeletal muscle of exercising mice.

In the current study, the exercise-stimulated phosphorylation of p-p38MAPK was

reduced in *ncf1** mice compared to WT, in particular in quadriceps muscle. As p38

MAPK has been proposed as a regulator of muscular glucose uptake [12], we cannot

exclude that reduced p38 MAPK contributed to the reduced glucose uptake we observed

in *ncf1** mice. However, the reduction in exercise-stimulated kinase signaling is less

consistent than the reduction in glucose uptake in the different muscles analyzed in the

*ncf1** mice. Moreover, Rac1 imKO mice share the same glucose uptake phenotype

without showing reductions in stretch-stimulated signaling, including p38 MAPK[18].

Instead, we speculate that the shared reduction in exercise-stimulated glucose uptake in

*ncf1** and Rac1 imKO mice might relate to a shared and yet undetermined redox-

sensitive signaling mechanism. Worth noting, we believe NOX2 to signal independently

of the AMPK-TBC1D1 signaling axis, based on the previous work by us and others [14,

31, 32, 33, 34]. Future studies should work to identify the shared cell signaling traits between

different NOX2-deficient mouse models using e.g. redox proteomics [35].

In conclusion, this study showed for the first time that NOX2 is activated during

moderate-intensity exercise in human and mouse skeletal muscle and is a major source

of ROS under such conditions. Furthermore, our comparison of two mouse models

lacking regulatory NOX2 subunits showed that lack of ROS generation during exercise

strongly impaired muscle glucose uptake and GLUT4 translocation during exercise.

This indicates that NOX2 is a major source of ROS generation during exercise and that

NOX2-dependent ROS production is an important signal for increasing muscle glucose

uptake during exercise.

MATERIAL AND METHODS

Animals. All experiments were approved by the Danish Animal Experimental

Inspectorate (2015−15−0201−00477). Inducible muscle-specific male Rac1 mice

(imKO) and littermate control mice were generated by crossbreeding Rac1$^{fl/fl}$ mice [36]

with mice expressing Cre recombinase from a tetracycline-controlled transactivator

coupled to the human skeletal actin promoter [37]. For muscle-specific Rac1 deletion, all

mice were treated with doxycycline in the drinking water around 10-14 weeks of age

(1g/L; Sigma Aldrich) for 3 weeks followed by a wash out period of 3 weeks.

Male B10.Q wild-type (WT) and B10.Q. p47phox mutated (*ncf1**) mice contain a point

mutation in exon 8 [21]. All mice were group-housed maintained on a 12:12-h light-dark

cycle, at 21º C, and received standard rodent chow diet (Altromin no. 1324; Chr.

Pedersen, Denmark) and water ad libitum.

In vivo gene transfer in adult skeletal muscle. For the *in vivo* transfection experiments,

mice were anesthetized with 2-3% isoflurane. Hyaluronidase (Sigma-Aldrich) dissolved

in sterile saline solution (0.36 mg/ml) was injected subcutaneously in FDB and

intramuscularly in TA muscle, followed by 20 and 40 μg plasmid injections 1 hour

later, respectively. FDB muscle electroporation was performed by delivering 10

electrical pulses at an intensity of 75 V/cm, 10-ms pulse duration, 200-ms pulse interval

using acupuncture needles (0.20 x 25 mm, Tai Chi, Lhasa OMS) connected to an ECM

830 BTX electroporator (BTX Harvard Apparatus). TA electroporation was performed

following the same protocol but raising the intensity to 100 V/cm (#45-0101 Caliper

Electrode, BTX Caliper Electrodes, USA). The p47-roGFP construct used to determine

NOX2 activity was a kind gift from Dr. George G. Rodney [22]. The GLUT4-myc-GFP

construct was a kind gift from Dr. Jonathan Bogan (Addgene plasmid # 52872).

Metabolic Chambers. O_2 uptake (VO_2) and CO_2 production (VCO_2) were measured

using a CaloSys apparatus (TSE Systems, Bad Homburg, Germany). The data presented

is the average for light and dark periods measured over 2 consecutive days in WT and

ncf1* mice. The respiratory exchange ratio (RER) was calculated as VCO_2

production/VO_2 uptake.

Maximal running tests. Mice were acclimated to the treadmill three times (15 min at

0.16 m/s) (Treadmill TSE Systems) a week prior to the maximal running tests. The

maximal running test started at 0.16 m/s for 300 s with 10 % incline, followed by a

continuing increase (0.2 m/s) in running speed every 60s until exhaustion. Exhaustion

was defined as the point at which instead of running on the treadmill, the mice fall back

on the grid 3 times within a 30 sec period. Maximal running speed was determined as

the last completed stage during the incremental test.

FDB fiber isolation. Muscle fibers were obtained by enzyme digestion of the whole

muscle with collagenase type II (1.5 mg/ml) (Worthington, Lakewood, NJ) for 90 min

at 37°C followed by mechanical dissociation with fire polished Pasteur pipettes. Isolated

fibers were seeded in ECM Gel-coated (Sigma-Aldrich) cell culture dishes in DMEM

supplemented with 10% fetal bovine serum. After 16 h of seeding, the fibers were used

for experimentation. Electrical stimulation protocol (10 V, 1 Hz, 0.1 ms duration) was

delivered to isolated FDBs for 15 min via field electric field stimulation device (RC-

37FS, Warner instruments, USA).

Redox changes in muscle cryosections. TA muscles were dissected and embedded in

optimum cutting temperature (OCT) medium from Tissue Tek, frozen in melting

isopentane and kept at -80 °C until processing. Total oxidant levels were estimated as

previously described [13]. Briefly, 10 μm thickness muscle cryosections were incubated

with 5 μM of 2',7'-dichlorodihydrofluorescein diacetate (DCF; Molecular Probes,

Eugene) and allowed to dry overnight at room temperature in dark. Muscle membranes

were visualized using Texas red labeled wheat germ agglutinin (WGA; Molecular

Probes). The redox histology was performed as previously described [19]. p47roGFP-

transfected muscle cryosections were incubated with PBS containing 50 mM N-

Ethylmaleimide (NEM) for 10 min at 4°C, followed by fixation using 4%

paraformaldehyde for 10 min at room temperature.

Running-stimulated muscle 2-deoxyglucose (DG) uptake. Muscle-specific 2DG uptake

was measured as previously described[31]. [^{3}H]2DG (PerkinElmer) was injected

intraperitoneally (as a bolus in saline, 10 μL/g body weight) containing 0.1 mmol/L

2DG and 50 μCi/mL [^{3}H]2DG corresponding to ~12 μCi/mouse) into fed mice

immediately before the exercise bout. Blood glucose was measured before and

immediately after 20 min of exercise. Mice were euthanized by cervical dislocation, and

muscle tissues and plasma samples quickly frozen in liquid nitrogen and stored at

−80°C until analysis. Samples were deproteinated using 0.1 mol/L Ba (OH)$_2$ and 0.1

mol/L ZnSO$_4$. The total muscle [^{3}H]2DG tracer activity found in 2DG-6-phosphate was

divided by the area under the curve of the specific activity at time points 0 and 20 min

multiplied with the average blood glucose at time points 0 and 20 min. This was related

to muscle weight and the time to obtain the tissue-specific 2DG uptake as micromoles

per gram per hour.

In vivo GLUT4 translocation. Immediately after exercise, TA muscles were fixed by

immersion in ice-cold 4% in paraformaldehyde in PBS for 4 h. Individual fibers were

teased from fixed muscle with fine forceps under a dissection microscope. Non-

permeabilized isolated muscle fibers were incubated in 1% BSA in PBS for 1 h min and

then incubated with an anti-myc antibody (rabbit polyclonal) overnight (CST, #2278).

The next day fibers were incubated with a secondary antibody conjugated with Alexa

Fluor 568 (Invitrogen, UK). The muscle fibers were mounted in Vectashield mounting

medium.

Western blot analyses. Tissue was homogenized for 1 min at 30 Hz using a Tissue

Lyser in ice cold lysis buffer (0.05 mol/L Tris Base pH 7.4, 0.15 mol/L NaCl, 1 mmol/

L EDTA and EGTA, 0.05mol/L sodium flouride, 5 mmol/L sodium pyrophosphate, 2

mmol/L sodium orthovanadate, 1 mmol/L benzamidine, 0.5% protease inhibitor

cocktail (P8340, Sigma Aldrich), and 1% NP- 40). After rotating end-over-end for 30

min, lysate supernatants were collected by centrifugation (18,327 g) for 20 min at $4^{o}C$.

Lysate protein concentrations were determined using BSA standards and bicinchoninic

acid assay reagents (Pierce). Total protein and phosphorylation levels of relevant

proteins were determined by standard immunoblotting techniques, loading equal

amounts of protein. The primary antibodies used p-AMPK^{Thr172} (Cell Signaling

Technology (CST)), #2535S, p-p38 MAPK$^{Thr180/Tyr182}$ (CST, #9211), Hexokinase II

(CST, #2867), GLUT4 p-ACC2 Ser^{212} (Millipore, 03-303), (ThermoFisher Scientific,

PA-23052), Rac1 (BD Biosciences, #610650), NOX2 (Abcam, #Ab129068), Catalase

(SCBT, sc-271803), MnSOD (Millipore, 06-984), TRX2 (SCBT, sc-50336), actin

(CST, #4973) total p38 MAPK (CST, #9212), alpha2 AMPK (a gift from D. Grahame

Hardie, University of Dundee), total ERK 1/2 (CST, #9102), and TBC1D1^{ser231}

(Millipore #07-2268), OXPHOS cocktail (Abcam, #ab110413) and Hexokinase II

(CST, #2867). All antibodies were optimized for signal linearity. Membranes were

blocked for 1 h at room temperature in TBS-Tween 20 containing either 2% skimmed

milk or 2% BSA and incubated overnight at 4°C with primary antibody. The

membranes were incubated in the corresponding horseradish peroxidase-conjugated

secondary antibody for 1 h at room temperature and washed in TBST before

visualization of the proteins (ChemiDoc^{TM}MP Imaging System, Bio Rad).

Imaging and imaging analysis. For all live-imaging experiments confocal images were

collected using a 63x 1.4 NA oil immersion objective lens on an LSM 780 confocal

microscope (Zeiss) driven by Zen 2011. FDB fibers were imaged in phenol red-free

growth medium (5 % horse serum) while kept at 95 % O_2 and 5 % CO_2 and 37°C using

a Pecon Lab-Tek S1 heat stage.

Human experiments. Three young (Age 29 ± 3.56y BMI 24.8 ± 1.76 kg/m^2), healthy

men gave their written, informed consent to participate in the study approved by the

Regional Ethics Committee for Copenhagen (H-16040740) and complied with the

ethical guidelines of the Declaration of Helsinki II. The volunteers visited the laboratory

on two separate days. The first day, the subjects completed an incremental test on a

Monark Ergomedic 839E cycle ergometer (Monark, Sweden) to determine peak power

output (PPO). On the second day, each subject performed a 30-min exercise trial at the

intensity corresponding to ~65% of their individual PPO in fasted state. Muscle biopsies

were obtained from the m. vastus lateralis under local anesthesia [~3ml xylocaine

(20mg ml−1lidocaine), Astra, Stockholm, Sweden] before and after exercise using a 5

mm Bergstrom needle with suction. The muscle biopsies were embedded in optimal

cutting temperature compound (tissue-tek) and frozen in liquid nitrogen-cooled

isopentane and stored at −80°C for further analysis.

Statistical analyses. Results are shown as individual values or means ± S.E.M.

Statistical testing was carried out using *t*-tests or two-way (repeated measures when

appropriate) ANOVA as applicable. Tukey's *post hoc* test was performed to pinpoint

significant differences between groups. Statistical analyses were performed using

GraphPad Prism.

Acknowledgements. ZL was supported by Chinese Scholarship Council PhD stipends.

LS was supported by a Danish Research Council grant 4004-00233 and Novo Nordisk

Foundation Excellence grant NNF18OC0032082. CHO was supported by a CONICYT

PhD travel grant. JRK was supported by Danish Diabetes Academy PhD stipends. EJ

was supported by FONDECYT grant 1151293. TEJ was supported by a Novo Nordisk

Foundation Excellence project grant (#15182). EAR was supported by the Danish

Council for Independent Research/Science 6108-00203. We thank Kim Anker

Sjøberg for pre-testing the human subjects and Betina Bolmgren for technical

assistance with measuring 2DG uptake. Imaging data were collected at the

Center for Advanced Bioimaging and the Core Facility for Integrated

Microscopy, University of Copenhagen, Denmark.

Author Contribution. Conceptualization, CHO, EJ and TEJ. Methodology, CHO and

TEJ. Investigation, CHO, JRK, ZL, ED, JTT, SHR, RH, EAR and LS. Formal

analysis, CHO. Visualization, CHO. Writing – Original draft, CHO and TEJ. Writing

– Review and Editing, CHO with input from all authors. Funding Acquisition, TEJ.

Supervision, EJ and TEJ.

REFERENCES

1. Egan B, Zierath JR. Exercise metabolism and the molecular regulation of
skeletal muscle adaptation. *Cell metabolism* **17**, 162-184 (2013).
402

403 2. Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms
404   and impact on muscle force production. *Physiol Rev* **88**, 1243-1276 (2008).
405

406 3. Powers SK, Radak Z, Ji LL. Exercise-induced oxidative stress: past, present and
407   future. *J Physiol* **594**, 5081-5092 (2016).
408

409 4. Sakellariou GK*, et al.* Studies of mitochondrial and nonmitochondrial sources
410   implicate nicotinamide adenine dinucleotide phosphate oxidase(s) in the
411   increased skeletal muscle superoxide generation that occurs during contractile
412   activity. *Antioxidants & redox signaling* **18**, 603-621 (2013).
413

414 5. Michaelson LP, Shi G, Ward CW, Rodney GG. Mitochondrial redox potential
415   during contraction in single intact muscle fibers. *Muscle Nerve* **42**, 522-529
416   (2010).
417

418 6. Jackson MJ. Recent advances and long-standing problems in detecting oxidative
419   damage and reactive oxygen species in skeletal muscle. *J Physiol* **594**, 5185-
420   5193 (2016).
421

422 7. Richter EA, Hargreaves M. Exercise, GLUT4, and skeletal muscle glucose
423   uptake. *Physiol Rev* **93**, 993-1017 (2013).
424

425 8. Sylow L, Kleinert M, Richter EA, Jensen TE. Exercise-stimulated glucose
426   uptake - regulation and implications for glycaemic control. *Nat Rev Endocrinol*
427   **13**, 133-148 (2017).
428

429 9. Kristiansen S, Hargreaves M, Richter EA. Exercise-induced increase in glucose
430   transport, GLUT-4, and VAMP-2 in plasma membrane from human muscle. *Am
431   J Physiol* **270**, E197-201 (1996).
432

433 10. Zisman A*, et al.* Targeted disruption of the glucose transporter 4 selectively in
434   muscle causes insulin resistance and glucose intolerance. *Nat Med* **6**, 924-928
435   (2000).
436

437 11. Sandstrom ME*, et al.* Role of reactive oxygen species in contraction-mediated
438   glucose transport in mouse skeletal muscle. *J Physiol* **575**, 251-262 (2006).
439

440 12. Chambers MA, Moylan JS, Smith JD, Goodyear LJ, Reid MB. Stretch-
441   stimulated glucose uptake in skeletal muscle is mediated by reactive oxygen
442   species and p38 MAP-kinase. *J Physiol* **587**, 3363-3373 (2009).
443

444 13. Merry TL, Steinberg GR, Lynch GS, McConell GK. Skeletal muscle glucose
445   uptake during contraction is regulated by nitric oxide and ROS independently of
446   AMPK. *Am J Physiol Endocrinol Metab* **298**, E577-585 (2010).
447

14.   Sylow L, *et al.* Rac1 is a novel regulator of contraction-stimulated glucose uptake in skeletal muscle. *Diabetes* **62**, 1139-1151 (2013).

15.   Sylow L, Moller LL, Kleinert M, Richter EA, Jensen TE. Stretch-stimulated glucose transport in skeletal muscle is regulated by Rac1. *J Physiol* **593**, 645-656 (2015).

16.   Chiu TT, Jensen TE, Sylow L, Richter EA, Klip A. Rac1 signalling towards GLUT4/glucose uptake in skeletal muscle. *Cell Signal* **23**, 1546-1554 (2011).

17.   Bedard K, Krause KH. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* **87**, 245-313 (2007).

18.   Sylow L, *et al.* Rac1 governs exercise-stimulated glucose uptake in skeletal muscle through regulation of GLUT4 translocation in mice. *J Physiol* **594**, 4997-5008 (2016).

19.   Fujikawa Y, *et al.* Mouse redox histology using genetically encoded probes. *Science signaling* **9**, rs1 (2016).

20.   Margaritelis NV, *et al.* Adaptations to endurance training depend on exercise-induced oxidative stress: exploiting redox interindividual variability. *Acta physiologica (Oxford, England)* **222**, (2018).

21.   Hultqvist M, Olofsson P, Holmberg J, Backstrom BT, Tordsson J, Holmdahl R. Enhanced autoimmunity, arthritis, and encephalomyelitis in mice with a reduced oxidative burst due to a mutation in the Ncf1 gene. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 12646-12651 (2004).

22.   Pal R, Basu Thakur P, Li S, Minard C, Rodney GG. Real-time imaging of NADPH oxidase activity in living cells using a novel fluorescent protein reporter. *PLoS One* **8**, e63989 (2013).

23.   Ji LL, Kang C, Zhang Y. Exercise-induced hormesis and skeletal muscle health. *Free radical biology & medicine* **98**, 113-122 (2016).

24.   Davies KJ, Quintanilha AT, Brooks GA, Packer L. Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun* **107**, 1198-1205 (1982).

25.   Ueda S, *et al.* Crucial role of the small GTPase Rac1 in insulin-stimulated translocation of glucose transporter 4 to the mouse skeletal muscle sarcolemma. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* **24**, 2254-2261 (2010).

26.   Chiu TT, Patel N, Shaw AE, Bamburg JR, Klip A. Arp2/3- and cofilin-coordinated actin dynamics is required for insulin-mediated GLUT4 translocation to the surface of muscle cells. *Mol Biol Cell* **21**, 3529-3539 (2010).

27.   Madsen AB, *et al.* beta-actin shows limited mobility and is only required for
supraphysiological insulin-stimulated glucose transport in young adult soleus
muscle. *Am J Physiol Endocrinol Metab*, (2018).

28.   Wilson C, *et al.* A Feed-Forward Mechanism Involving the NOX Complex and
RyR-Mediated Ca2+ Release During Axonal Specification. *J Neurosci* **36**,
11107-11119 (2016).

29.   Keller A, Mohamed A, Drose S, Brandt U, Fleming I, Brandes RP. Analysis of
dichlorodihydrofluorescein and dihydrocalcein as probes for the detection of
intracellular reactive oxygen species. *Free Radic Res* **38**, 1257-1267 (2004).

30.   Chen X, Zhong Z, Xu Z, Chen L, Wang Y. 2',7'-Dichlorodihydrofluorescein as a
fluorescent probe for reactive oxygen species measurement: Forty years of
application and controversy. *Free Radic Res* **44**, 587-604 (2010).

31.   Sylow L, *et al.* Rac1 and AMPK Account for the Majority of Muscle Glucose
Uptake Stimulated by Ex Vivo Contraction but Not In Vivo Exercise. *Diabetes*
66, 1548-1559 (2017).

32.   Jensen TE, *et al.* Contraction-stimulated glucose transport in muscle is
controlled by AMPK and mechanical stress but not sarcoplasmatic reticulum
Ca(2+) release. *Mol Metab* **3**, 742-753 (2014).

33.   Dokas J, *et al.* Conventional knockout of Tbc1d1 in mice impairs insulin- and
AICAR-stimulated glucose uptake in skeletal muscle. *Endocrinology* **154**, 3502-
3514 (2013).

34.   Szekeres F, *et al.* The Rab-GTPase-activating protein TBC1D1 regulates
skeletal muscle glucose metabolism. *Am J Physiol Endocrinol Metab* **303**, E524-
533 (2012).

35.   McDonagh B, Sakellariou GK, Jackson MJ. Application of redox proteomics to
skeletal muscle aging and exercise. *Biochem Soc Trans* **42**, 965-970 (2014).

36.   Chrostek A, *et al.* Rac1 is crucial for hair follicle integrity but is not essential for
maintenance of the epidermis. *Molecular and cellular biology* **26**, 6957-6970
(2006).

37.   Rao P, Monks DA. A tetracycline-inducible and skeletal muscle-specific Cre
recombinase transgenic mouse. *Dev Neurobiol* **69**, 401-406 (2009).

FIGURE LEGENDS

Figure 1. Moderate-intensity exercise increases ROS production in a NOX2-

dependent manner. A) Dichlorofluorescein (DCF) oxidation and B) exercise signaling

in muscle biopsies before and after a bout of moderate-intensity cycling (30 min, 65%

maximal power output) in young-healthy volunteers (n=3). C) Representative images

and quantification of the exercise-stimulated (20 min, 65% maximal running speed)

DCF oxidation in tibialis anterior (TA) muscle from WT and *ncf1** mice (n=5). D)

Representative image and quantification of the *in vivo* running-induced p47roGFP

oxidation in tibialis anterior (TA) muscle in WT and Rac1 imKO mice. Scale bar=50

uM. For A) Unpaired t-test was performed for statistical analysis, * denotes P< 0.05

compared to resting condition. For C-D) Two-way ANOVA was performed to test for

effects of exercise (exer) genotype (geno), and interaction (Int), followed by Tukey's

post hoc corrected for multiple comparisons. #, ### denotes P< 0.05 and p< 0.001,

respectively compared to the WT group. Values are mean ± SEM.

Figure 2. Exercise-stimulated glucose uptake is reduced in *ncf1** mice. Different

parameters were compared between p47phox mutated mice (*ncf1**) and wild-type (WT)

mice. A) Respiratory exchange ratio (RER) during light and dark cycle and B) habitual

activity. C-E) *In vivo* running-induced (20 min, 65% maximal running speed) 2-deoxy-

glucose uptake (2DG) uptake in WT and *ncf1** mice (n=12-16). F) Maximal running

speed (n=15 per group), G) blood glucose concentration after rest/exercise (n=6-8), H)

NAD+, I) NADPH levels in quadriceps muscle (n=6-8), and J) Plasma lactate

concentration. Unpaired t-test (F) and two-way ANOVA were performed to test for

effects of exercise (Exer) genotype (Geno), and interaction (Int), followed by a Tukey's

post hoc corrected for multiple comparisons. Data is shown as mean and SEM. #, ###

denotes P< 0.05 and p< 0.001, respectively compared to the WT group (n=8 per group).

Figure 3. Exercise-stimulated GLUT4 translocation requires p47phox. A) The
GLUT4-myc-GFP construct was electroporated into tibialis anterior (TA) muscles of
both WT and *ncf1** mice. Non-permeabilized muscle fibers from exercised (20 min,
65% maximal running speed) or resting mice were stained with anti-*myc* antibody and
imaged by confocal microscopy (n=3-5). Total endogenous GLUT4 and HK II were
determined by western blot in the following muscles B) quadriceps, C) soleus, and D)
TA muscles in WT and *ncf1** mice (n=14). Total oxphos proteins were determined in
E) quadriceps and F) Soleus muscle. Unpaired t-test (B-D) and two-way ANOVA were
performed to test for effects of exercise (Exer) genotype (Geno), and interaction (Int),
followed by a Tukey's *post hoc* corrected for multiple comparisons. Values are mean ±
SEM.

Figure 4. Exercise-stimulated intracellular signaling in *ncf1** muscles. Activation of
intracellular kinases after moderate-intensity treadmill exercise (20 min, 65% maximal
running speed) in muscle lysates of A) quadriceps, B) Soleus, and C) tibialis anterior
from WT and p47phox-mutated (*ncf1**) mice (n=6-8). Two-way ANOVAs was
performed to test for effects of exercise (Exer) genotype (Geno), and interaction (Int),
followed by Tukey's *post hoc* corrected for multiple comparisons. # denotes P< 0.05
compared to the WT group. Values are mean ± SEM.

Figure S1. Similar abundance of antioxidant enzymes in *ncf1** muscles. *Tibialis*
anterior muscle lysates from both *ncf1** and WT mice. Values are mean ± SEM.

Figure S2. Rac1 imKO mice fail to increase ROS production *in vitro* and *in vivo*. A)
ROS generation induced by electrical stimulation in FDB Rac1 imKO fibers (*n*=5) b)

redox-regulated proteins in FDB fibers are similar between WT and Rac1 imKO muscle

(*n*=5). C) DCF oxidation after 20 min of exercise in WT and Rac1 imKO *tibialis*

anterior muscle (*n*=4). Two-way ANOVA were performed to test for effects of exercise

(Exer) genotype (Geno), and interaction (Int), followed by a Tukey's *post hoc* corrected

for multiple comparisons. # denotes P< 0.05 respectively compared to the WT group.

Values are mean ± SEM.

Figure S3. Body weight and composition of *ncf1* mice.* a) Similar body weight

despite tendency to b) lower body fat and c) higher lean mass and d) equal oxygen

consumption in *ncf1** compared to WT mice. The midline represents the median; box

represents the IQR between the first and third quartiles, and whiskers represent the

lowest or highest values within 1.5 times IQR from the first or third quartiles. Unpaired

t-test (F) and two-way ANOVA were performed to test for effects of exercise (Exer)

genotype (Geno), and interaction (Int), followed by a Tukey's *post hoc* corrected for

Figure S4. Similar endogenous glucose handling-related protein abundance in

***nfc1* and WT muscles.** Quantification of protein levels showed in figure 3B-D in a)

quadriceps b) soleus, and c) *tibialis anterior* muscle. Values are mean ± SEM.

Figure S5. Oxphos protein content in *ncf1* muscles are similar to WT.*

Quantification of protein levels showed in figure 3E-F in a) quadriceps and b) soleus

muscle. Values are mean ± SEM.

Figure S6. Total AMPK, p38 MAPK and ERK protein in *ncf1** muscles.

Quantification of protein levels showed in figure 4A-C in a) quadriceps and b) soleus

and *tibialis anterior* muscle. Values are mean ± SEM.

Figure S1

Figure S2

A

B

C

Figure S3

Figure S4

Figure S5

Figure S6