1

Adapting genotyping-by-sequencing and variant calling for heterogeneous stock rats

Jianjun Gao[*,‖], Alexander F. Gileta[* †‖], Hannah V. Bimschleger[*], Celine L. St. Pierre[*], Shyam Gopalakrishnan[#], Abraham A. Palmer[*‡]

* Department of Psychiatry, University of California San Diego, La Jolla, California, 92093

† Department of Human Genetics, University of Chicago, Chicago, Illinois, 60637

Department of Biology, University of Copenhagen, 2200 København N, Denmark

‡ Institute for Genomic Medicine, University of California San Diego, La Jolla, California, 92093

15   ‖ These authors contributed equally to this work.

Running title: GBS and variant calling in HS rats

Key words: genotyping-by-sequencing, heterogeneous stock, rat, imputation

Corresponding author: Abraham A. Palmer

Mailing address: 9500 Gilman Drive #0667, La Jolla, CA, 92093

Phone number: 858-534-2093

Email: aapalmer@ucsd.edu

ABSTRACT

The heterogeneous stock (**HS**) is an outbred rat population derived from eight inbred rat strains.

The population is maintained with the goal of minimizing inbreeding and maximizing the genetic

diversity of the stock. A few genotyping microarrays have been created for rats; however, they

were expensive and are no longer in production. To obtain high-density genome-wide marker

data for genetic mapping, we have adapted genotype-by-sequencing (**GBS**) for use in rats. Here,

we outline the laboratory and computational steps we took to design and optimize an efficient

double digest genotype-by-sequencing (**ddGBS**) protocol for rats. To analyze the ddGBS

sequencing data, we evaluated multiple existing computational tools and designed a workflow

that allowed us to call and impute over 3.7 million SNPs genome-wide in the HS. We also

compared various rat genetic maps for use in imputation, including a recently developed map

specific to the HS. Using the pipeline, we obtained concordance rates of 99% with data from a

rat genotyping array. The computational pipeline that we have developed can be easily adapted

for use in other species. We include a detailed protocol to perform ddGBS in rats.

41   INTRODUCTION

Advances in next-generation sequencing technology over the past decade have enabled the

discovery of high-density, genome-wide single nucleotide polymorphisms (**SNPs**) in model

systems. Comprehensive assays of the standing genetic variation in these organisms has allowed

for the identification of quantitative trait loci and the application of numerous population genetic

and phylogenetic methods. However, due to the high degree of linkage disequilibrium (**LD**) in

many structured breeding populations, sequencing whole genomes is not necessary. SNPs are

frequently in strong LD with adjacent loci, effectively 'tagging' nearby variation, and thereby

reducing the number of sites that need to be genotyped.   Several reduced-representation

sequencing approaches that take advantage of LD structure have been previously described

(Miller et al. 2007; van Orsouw et al. 2007; Van Tassell et al. 2008; Baird et al. 2008; X. Huang

et al. 2009; Davey et al. 2011; Elshire et al. 2011; Poland et al. 2012; Peterson et al. 2012; Sun et

al. 2013; Scheben, Batley, and Edwards 2017). Thousands of SNPs can be identified in large

numbers of samples for a fraction of the price of other genotyping methods (Chen et al. 2013; He

et al. 2014). The advantages of these methods are especially attractive when applied to less

commonly utilized species or strains for which genotyping microarrays are not available.

Of the existing reduced-representation protocols, the genotyping-by-sequencing (**GBS**)

approach developed by Elshire et al. (Elshire et al. 2011) has been frequently modified to

accommodate non-model species, such as: soybean (Sonah et al. 2013), rice (Furuta et al. 2017),

oat (Fu and Yang 2017), chicken (Pértille et al. 2016; Wang et al. 2017), mouse (Parker et al.

2016), fox (Johnson et al. 2015), and cattle (De Donato et al. 2013), among others. The greatly

varying genomic composition among organisms necessitates a diverse and customized set of

approaches for obtaining high-quality genotypes. As such, both the GBS protocol and

computational pipeline require modifications when being applied to a new species. Recent work

from our group showed that GBS can be effectively applied to outbred mice (Parker et al. 2016;

Gonzales et al. 2017; Zhou et al. 2018) and rats (Fitzpatrick et al. 2013). However, those

publications used protocols that had not been optimized, leaving significant room for

improvement in genotype quality and marker density. Additionally, although several tools and

workflows for the analysis of GBS data have been described, including Stacks (Catchen et al.

2013), IGST-GBS (Sonah et al. 2013), TASSEL-GBS (Glaubitz et al. 2014), Fast-GBS

(Torkamaneh et al. 2017), and GB-eaSy (Wickland et al. 2017), the majority were developed and

optimized for use in plant species and given the lack of well-developed genomic resources in

these species, do not include an imputation step. Here we describe the customized computational

and laboratory protocols for applying GBS to HS rats.

The HS is an outbred rat population created in 1984 using eight inbred strains and has

been maintained since then with the goal of minimizing inbreeding and maximizing the genetic

diversity of the colony (Johannesson et al. 2008; Woods and Mott 2017). After more than 80

generations of accumulated recombination events, their genome has become a fine-scale mosaic

of the inbred founders' haplotypes. The breeding scheme and the number of accumulated

generations has made the HS colony attractive for genetic studies. Additionally, extensive deep

sequencing data exists for the eight progenitor strains, allowing for accurate imputation from

sites directly captured by GBS to millions of additional SNPs.

Detailed here are the steps we have taken to optimize a rat GBS protocol and

computational pipeline. Drawing on existing protocols (Elshire et al. 2011; Peterson et al. 2012;

Poland et al. 2012; Parker et al. 2016) as models, we redesigned our GBS approach and have

developed a novel, reference-based, high-throughput workflow to accurately and cost-effectively

call and impute variants from low-coverage double digest GBS (**ddGBS**) data in HS rats. This

publication is intended as a resource for others who might wish to perform GBS in rats and

should provide a roadmap for adapting GBS for use in new species. We demonstrate that with a

suitable reference panel, applying reduced representation approaches and imputation in model

systems can provide high-confidence genotypes on millions of genome-wide markers.

92   MATERIALS AND METHODS

93 **Tissue samples and DNA extraction**

Samples for this study originated from three sources: an inhouse advanced intercross line (**AIL**) derived from LG/J and SM/J mice (Gonzales et al. 2018), Sprague Dawley (**SD**) rats from Charles River Laboratories and Harlan Sprague Dawley, Inc. (Gileta et al. 2018), and an HS rat colony (Woods and Mott 2017; Chitre et al. 2018). Early ddGBS optimization utilized AIL genomic DNA extracted from spleen by a standard salting-out protocol. Later optimization steps were performed using genomic DNA from SD rats extracted from tail tissue using the PureLink Genomic DNA Mini Kit (Thermo Fisher Scientific, Waltham, MA). Optimization of the ddGBS sequencing data analysis pipeline was performed on various subsets of our final sample of 4,973 HS rats. HS rat DNA was extracted from spleen tissue using the Agencourt DNAdvance Kit (Beckman Coulter Life Sciences, Indianapolis, IN). All genomic DNA quality and purity was assessed by NanoDrop 8000 (Thermo Fisher Scientific, Waltham, MA). Interestingly, we have observed that rat genomic DNA appears to degrade faster than mouse genomic DNA following extraction; therefore, we recommend storing rat genomic DNA at -20° and using it within months of extraction whenever possible.

In silico digest of rat genome

We used *in silico* digests to aid in the selection of restriction enzymes, with the goal of maximizing the portion of the genome captured at sufficient depth to make confident genotype calls. We used the *restrict* function within EMBOSS (version 6.6.0) (Rice, Longden, and Bleasby 2000) in conjunction with the REBASE database published by New England BioLabs (NEB; version 808) (Roberts and Macelis 1999) to perform *in silico* digest of rn6, the current release of the Norway brown rat reference genome. For the primary restriction enzyme, we chose PstI. We performed the digest with PstI alone and then with PstI paired with each of 7 secondary enzymes: AluI, BfaI, DpnI, HaeIII, MluCI, MspI, and NlaIII. We only considered fragments with

one PstI cut site and one cut site from the secondary enzyme because the adapter and primer sets

are designed to only allow these fragments to amplify. The final choice of enzyme (NlaIII) was

determined empirically and is detailed in the Results.

Restriction enzyme selection

Initial criteria for selecting a secondary restriction enzyme were: a 4bp recognition sequence, no

ambiguity in the recognition sequence (i.e. N's), compatibility with the NEB CutSmart Buffer,

and an incubation temperature of 37°C. The list of enzymes meeting these criteria at the time

included: AluI, BfaI, DpnI, HaeIII, MluCI, MspI, and NlaIII. Using the *in silico* digest data, we

looked to maximize the portion of the genome contained within a fragment size range of 125-

275bp (250-400bp with annealed adapters and primers) (Figure 1; Table 1). We excluded

enzymes that produced blunt ends, both because it would be more difficult to anneal adapters to

blunt ended fragments and because our adapters would then also anneal to blunt ends produced

by DNA shearing. We also excluded methylation-sensitive enzymes, as we did not want to limit

the breadth of our sequencing efforts, accepting the possibility of read pileup in repetitive

regions. Based on these criteria, NlaIII, BfaI, and MluCI were selected for further testing.

ddGBS library preparation and sequencing

The full ddGBS protocol is available in File S1. In brief, approximately 1µg of DNA is used per

sample. Sample DNA, PstI barcoded adapters, and NlaIII Y-adapter are combined in a 96-well

plate and allowed to evaporate at 37°C overnight. Sample DNA and adapters are re-eluted on day

two with a PstI/NlaIII digestion mix and incubated at 37°C for two hours to allow for complete

digestion. Ligation reagents are then added and incubated at 16°C for one hour to anneal the

adapters to the DNA fragments, followed by a 30-minute incubation at 80°C to inactivate the

restriction enzymes. Sample libraries are purified using a plate from a MinElute 96 UF PCR

Purification Kit (QIAGEN Inc., Hilden, Germany), vacuum manifold, and ddH$_2$O. Once re-

eluted, libraries are quantified in duplicate with Quanit-iT PicoGreen (Thermo Fisher Scientific,

Waltham, MA) and pooled to the desired level of multiplexing (i.e. 12, 24, or 48 samples per

library). Pooled libraries are concentrated to obtain the desired volume for use in the Pippin Prep.

The concentrated pool is quantified to ensure the gel cassette will not be overloaded with DNA

(>5µg). The pool is then loaded into the Pippin Prep for size selection (300-450bps) using a 2%

agarose gel cassette on a Pippin Prep (Sage Science, Beverly, MA). Size-selected libraries were

then PCR amplified for 12 cycles to increase the quantity of DNA, concentrated, and checked for

quality on an Agilent 2100 Bioanalyzer with a DNA 1000 Series II chip (Agilent Technologies,

Santa Clara, CA)., Bioanalyzer results were used to assure sufficient DNA concentration and to

identify excessive primer dimer peaks.

An initial 96 HS samples were sequenced, 12 samples per library, at Beckman Coulter

Genomics (now GENEWIZ) on an Illumina HiSeq 2500 with v4 chemistry and 125bp single-end

reads. Subsequently, we began using a set of 48 unique barcoded adapters (File S2) to multiplex

48 HS samples per ddGBS library. Each library was run on a single flow cell lane on an Illumina

HiSeq 4000 with 100bp single-end reads at the IGM Genomics Center (University of California

San Diego, La Jolla, CA).

161 **Figure 2. ddGBS sequencing data analysis workflow.** Each step of the workflow is described
162 in the text.

163
164

Evaluation of ddGBS pipeline performance

We present the steps required to call and impute genotypes from ddGBS data in Figure 2. During

optimization of the pipeline, performance was assessed by two primary metrics: (1) the number

of variants called and (2) genotype concordance rates for calls made in 96 HS rats that had both

ddGBS genotypes and array genotypes from the Axiom MiRat microarray. After each step of the

pipeline, the variant count and concordance rates were computed again. There were two

checkpoints in the GBS pipeline where genotype quality (measured by concordance rate) was

assessed: after internal imputation within Beagle (Browning and Browning 2009, 2016) and

again after imputation to the reference panel with IMPUTE2 (B. N. Howie, Donnelly, and

Marchini 2009; B. Howie et al. 2012). A third, additional metric we checked was the transition to

transversion ratio ($T_S T_V$), which is expected to be ~2.

Demultiplexing

The PstI adapter barcodes were used to demultiplex FASTQ files into individual sample files.

Three demultiplexing software packages were tested: FASTX Barcode Splitter (Hannon Lab

2010), GBSX (Herten et al. 2015), and an in-house Python script (Parker et al. 2016). Reads that

could not be matched with any barcode (maximum of 1 mismatch allowed), or that lacked the

appropriate enzyme cut site, were discarded. Samples with less than two million reads after

demultiplexing were discarded. Data concerning demultiplexing are shown in Table S1 are from

a single HS rat sequenced in a 12-sample library on one lane after demultiplexing and

adapter/quality trimming.

Barcode, adapter, and quality trimming

186 Read quality was assessed using FastQC v0.11.6 (Andrews 2017). We compared the efficacy of

187 two rapid, lightweight software options for trimming barcodes, adapters, and low-quality bases

188 from the NGS reads: Cutadapt (Martin 2011) and the FASTX Clipper/Trimmer/Quality Trimmer

189 tools (Hannon Lab 2010) (Table S2). A base quality threshold of 20 was used and reads trimmed

190 to below 25bps were discarded.

Read alignment and indel realignment

192 Rattus norvegicus genome assembly rn6 was used as the reference genome for read alignment

193 with the Burrows-Wheeler Aligner (BWA) (H. Li and Durbin 2009) using the *mem* algorithm.

194 We then used GATK IndelRealginer (McKenna et al. 2010) to improve alignment quality by

195   locally realigning reads around a reference set of known indels in 42 whole-genome sequenced

196   inbred rat strains, including the eight HS progenitor strains (Hermsen et al. 2015).

Variant calling and imputation

Variants were called using ANGSD v0.911, for which we employed the SAMtools model

(Korneliussen, Albrechtsen, and Nielsen 2014; Durvasula et al. n.d.). Beagle v4.1 (Browning and

Browning 2009, 2016) was used to impute missing genotypes at sites that had calls for only a

subset of individuals (no external reference panel was used). Before settling on the combination

of ANGSD and Beagle for genotype calling, we also tried GATK's UnifiedGenotyper and

HaplotypeCaller (McKenna et al. 2010) with various parameter settings, but we did not obtain

satisfactory performance. Using ANGSD, we inferred the major and minor alleles (-

domajorminor 1) from the genotype likelihoods, kept only high confidence polymorphic sites (-

snp_pval 1e-6), and estimated the allele frequencies based on the inferred alleles *(-domaf* 1). We

discarded sites missing read data in more than 4% of samples (*–minInd*). Additionally, we tested

altering thresholds for minimum base quality score (*-minQ*) and mapping quality (*-minMapQ*).

QC and pre-phasing for imputation using an external reference panel

Prior to imputing SNPs from the 42 inbred strain reference panel (Hermsen et al. 2015), we

checked concordance rates for the 96 HS animals with array genotypes, examined the T_ST_V ratio,

and assessed whether the sex as recorded in the pedigree records agreed with the sex empirically

determined by the proportion of reads on the X-chromosome out of the total number of reads

(Figure S1). We also identified Mendelian errors using the --mendel option in *plink* and known

pedigree information for 1,136 trios from 214 families within the HS sample. Using the fraction

of the trios that were informative for a given SNP and the formula $1-(1-2p(1-p))^3$, where p

represents the minor allele frequency of the allele, we formed curves for the distributions of the

expected number of Mendelian errors for both SNPs and samples and chose the inflection points

as thresholds for the number of Mendelian errors allowed. Lastly, we only retained variants

previously identified in the 8 HS founders (Ramdas et al. 2018) for use in imputation to the 42

genomes (Hermsen et al. 2015).

To improve imputation accuracy and computational efficiency, we employed a pre-

phasing step prior to reference imputation. A flowchart outlining the pre-phasing protocol is

presented in Figure S2. We tested three methods of converting genotype data to phased .hap

files: VCFtools (--*impute*) (Danecek et al. 2011), SHAPEIT (Delaneau, Marchini, and Zagury

2011), and IMPUTE2 (-*prephase_g*) (B. N. Howie, Donnelly, and Marchini 2009).

Genetic maps

Genetic maps are required for phasing and imputation with IMPUTE2. When we began this

project, no strain-specific recombination map was available for HS rats. Thus, we considered a

sparse genetic map for SHRSPxBN (Steen et al. 1999). We also tested two types of linearly

interpolated genetic maps, with recombination rates set at either 1cM/Mb or the chromosome

specific averages for rats, as reported by Jensen-Seaman et al. (Jensen-Seaman 2004). Lastly,

late in the course of this project, we experimented with an HS-specific genetic map developed by

Littrell et al. the Medical College of Wisconsin (Littrell et al. 2018).

Imputation to reference panel

We used existing sequencing and array data from the HS rat founder and other inbred laboratory

rat strains (Hermsen et al. 2015) as reference panel for imputation. Genotype data underwent QC

and were phased by Beagle into single chromosome haplotype files. Haplotype files were then

created using the workflow detailed in Figure S2. Imputation by IMPUTE2 was performed in

5Mb using the aforementioned reference panels and genetic maps.

Data availability

242 Data Genotype data will be available at https://doi.org/10.6084/m9.figshare.7597697 and the

243 code necessary to run the steps outline in the publication provided here

244 https://doi.org/10.6084/m9.figshare.7597694. Additional data is available upon request.

245   RESULTS

ddGBS optimization

Previous projects utilizing GBS in mice and rats (Fitzpatrick et al. 2013; Parker et al. 2016;

Gonzales et al. 2018) often encountered an issue where certain regions of the genome

experienced high pileups of reads per sample (>100x), while other regions were covered by just

1-2 reads. This read distribution imbalance can be caused in part by PCR amplification bias,

where a subset of fragments are preferentially amplified until they dominate the final library

(Kanagawa 2003; Aird et al. 2011). These previous protocols utilized 18 cycles of amplification.

We tested reducing this to 8, 10, 12, or 14 cycles and found that only 12 cycles were necessary to

yield sufficient PCR product for sequencing; the reduction in the number of PCR cycles was

expected to reduce PCR bias.

Another concern regarding previous sequencing results was an excess of long fragments

(>700bps as determined by *in silico* digest), which do not provide sufficient reads to make

confident genotype calls (< 5 reads per sample) and are therefore wasteful. We tested multiple

methods of combatting this issue, including: reducing the PCR extension time, increasing the

digestion length or enzyme concentration, increasing the selectivity of the primers, performing

size selection on the libraries, or using a two-enzyme restriction digest.

We began by testing shorter extension times (15s/20s vs. 30s) to try and reduce the

average fragment size of our sequencing libraries by preventing fragments with larger inserts

from completing amplification. However, we did not see noticeable improvement in the final

library composition at shorter extension times and determined the original length of 30 seconds

was appropriate.

We also considered the possibility that the restriction enzyme digests might not be

running to completion. To address this possibility, we increased the length of the digestion from

2 hours to 3 or 4 hours. We also tried increasing the number of units of PstI enzyme added, to

ensure complete digest. Neither of these modifications impacted the final fragment length

distribution of the library, indicating that the digest was reaching completion after 2 hours using

the original concentration of PstI.

We also explored reducing the complexity of our library as a means of obtaining more

reads at fewer sites. We accomplished this using the methods described by Sonah et al. 2013

(Sonah et al. 2013). This method involved adding an additional 1-2 nucleotides to the sequencing

PCR primers (A or AG). The addition of these nucleotides increases the selectivity of the primers

and reduces the number of amplifiable fragments, sacrificing sequencing breadth for read depth.

However, this did not ameliorate the issue of reads being wasted on long fragments but did

appear to increase the PCR bias (data not shown). Therefore, we chose to retain our original

primer design.

Our previous GBS protocol did not have an explicit library fragment size selection step.

The final library was purified using a MinElute PCR Purification Kit (QIAGEN Inc., Hilden,

Germany), which isolates PCR products 70bp-4kb in length, leaving a wide range of fragment

sizes in the final library, under the assumption that only shorter fragments would bridge amplify

on the flow cell. This method was imprecise and had low reproducibility, negatively impacting

our ability obtain reads at consistent sites across libraries. Rather than attempt size selection by

gel extraction, we chose to utilize a Pippin Prep, which automates the elution of DNA libraries of

desired fragment size ranges. By using this automated size selection, we reduced the proportion

of the genome targeted for sequencing, and because restriction enzymes make the consistent cuts

across samples, ensure the same fragments are sequenced in the majority of libraries. Since the

clustering process involves a bridge amplification step that preferentially amplifies library

fragments with shorter insert sizes (Illumina, Inc. 2014), we kept the size selection window

narrow (250-400bps) to avoid introducing a bias in which fragments were sequenced.

To increase the proportion of the genome captured within the fragment size window, we pursued

a double digest of the genome using a secondary enzyme with a more frequently occurring

recognition sequence. When used alone, *in silico* digest of the rn6 reference genome by PstI

(Figure 1; Table 1) showed that only ~0.5% of the genome would have fallen within a 150bp

fragment size window selected on the Pippin Prep. Given the LD structure of other outbred

rodent lines such as DO and CFW mice (Parker et al. 2016), we speculated this small fraction

would not yield sufficient SNPs to tag the majority of variation in the genome. Additionally, we

were concerned about potential biases in coverage, heterozygosity, and the minor allele

frequency (**MAF**) spectrum that may be introduced by incomplete capture of the genome

(Flanagan and Jones 2018).

The number of fragments with one of each of the cut sites were summed for all observed

lengths and the results summarized in Figure 1 and Table 1. BfaI, MluCI, and NlaIII were

chosen for further testing due to their compatibility with PstI digestion reagents and

temperatures, sticky ends, and the proportion of the genome falling in the size selection window.

We ruled out BfaI because it only had a 2bp overhang after cleavage, which led to a high

concentration of adapter dimer in the sequencing libraries. NlaIII was chosen because it

contained the greatest portion of the genome in the size selection window.

Restriction Enzyme(s)	Recognition sequence	Length of Overhang (bp)	% Genome in 250-400bp Region[+]	% Genome in 300-450bp Region[+]
PstI	CTGCA^G	4	0.48%	0.56%
PstI + AluI	AG^CT	0	3.06%	2.88%
PstI + BfaI	C^TAG	2	3.10%	3.25%
PstI + DpnI*	GA^TC	0	2.69%	3.00%
PstI + HaeIII	GG^CC	0	2.71%	2.79%
PstI + MluCI	^AATT	4	3.32%	3.21%
PstI + MspI	C^CGG	2	1.16%	1.24%
PstI + NlaIII	CATG^	4	3.45%	3.31%

312   **Table 1. Restriction enzyme options for double digest.**

313   The percent genome in region columns indicate the percentage of the genome that falls within
314   the provided fragment size ranges and can therefore be captured by GBS.

316   * Restriction enzyme is methylation sensitive.

317   [+] Calculated using rn6 genome length of 2,870,182,909bps.

Figure 1. *In silico* **digest fragment distributions for PstI and potential secondary restriction**
enzymes.

Each panel represents an independent digest of rn6 with the listed enzyme(s). Regions
highlighted in blue are fragments that would be selected by the Pippin Prep (125-275bp) after
annealing adapters and primers. These regions are quantified in Table 1 by multiplying the
length of the fragments by the number of fragments to estimate the portion of the genome
captured.

In our previous GBS protocol, all fragments were cut on both ends by PstI. By using a

substantially lower concentration of the barcoded PstI adapter than the common PstI adapter, we

ensured the barcoded adapter would be the limiting reagent and the majority of fragments with

an annealed barcoded adapter would have a common adapter on the other end. This is crucial, as

having one of each of the adapters is required for proper amplification of the fragments on the

flow cell. However, when using both PstI and NlaIII, the library is predominantly composed of

fragments cut on both sides by NlaIII, which will amplify during PCR with a common adapter,

but not on the flow cell. Therefore, we employed a Y-adapter (Poland et al. 2012) to control the

direction of the first round of PCR and prevent two-sided NlaIII fragments from dominating the

final sequencing library (File S2).

We tested numerous quantities of PstI and NlaIII adapters in an attempt minimize the

amount used and avoid adapter dimers in the final libraries. For the barcoded PstI adapters, we

tested 120pmol, 40pmol, 20pmol, 4pmol, 3pmol, 1.5pmol, 0.6pmol, and 0.2pmol and for the

NlaIII Y-adapter, 16pmol, 8pmol, 4pmol, 2pmol, 1pmol, and 0.5pmol. We found that 0.2pmol of

PstI adapter and 4pmol of NlaIII Y-adapter yielded sufficient library and avoided adapter dimers.

We sequenced a trial flow cell with 8 pooled ddGBS libraries of 12 SD rat samples each

(96 total) on a HiSeq 2500 (Illumina, San Diego, CA) with 125bp reads and v3 chemistry,

obtaining an average of 15.3 million reads per sample. Given the NlaIII *in silico* digest results

suggested we were capturing ~3.4% of the genome and that we were using 125bp reads, this

corresponded to approximately 20x coverage of captured sites. We subsequently increased the

number of samples to 48 per library for the HS rats because we hypothesized 5x would be

sufficient coverage per sample when utilizing imputation to a reference panel. We also

discovered that a portion of the reads contained sequence fragments of the NlaIII adapter

sequence, indicating there were fragments with insert sizes smaller than 125bps in the final

library. To avoid this, we increased the fragment size range to 300-450bps (Table 1), which

corresponds to a 175-325bp insert size once the adapters and primers are accounted for. Due to

the high concentrations of our libraries after pooling, the library size distribution obtained from

the Pippin Prep was uniformly shifted towards higher fragment lengths (Figure S3).

The final ddGBS protocol can be found in File S1 and the necessary primer and adapter

sequences in File S2. This protocol was used for the sequencing of all HS rats included in the

following computational optimization steps.

Demultiplexing

The number of base pairs of sequencing data retained after demultiplexing was fairly consistent

across demultiplexing software (Table S1). We ultimately decided to use FASTX Barcode

Splitter because it yielded the greatest number of reads after quality/adapter trimming and had

faster run times, though this may be predominantly attributable to outputting uncompressed

FASTQ files. An average of 330 million 100bp reads were obtained per library, resulting in ~7

million reads per sample. Figure S4 shows the distribution of reads counts for all samples after

demultiplexing, but prior to sample filtering based on read count.

Adapter and quality trimming

Read quality was substantially elevated after trimming the barcode and adapter sequences and

low-quality base pairs at the ends of reads (Figure S5). Overall read counts were only marginally

reduced by quality trimming (Table S1). We observed that the number of called variant sites and

the genotyping rate were both greater when using reads initially processed by Cutadapt (Martin,

2011) than reads processed by the FASTX_Toolkit (Table S2). Importantly, a large portion of

the additional identified variants were known variant sites from the 42 inbred strains reference

set (Figure S6), indicating the elevated call rate was at least in part due to capturing more true

variant sites. We viewed this as sufficient support for proceeding with Cutadapt for quality

trimming.

Read alignment quality

The number of called variants and genotype call rates were identical at read mapping quality

(mapQ) thresholds of either 20 or 30 (Table S3) within ANGSD. As the ANGSD mapQ

threshold was raised to 45, there was a small reduction in the number of called variants, and then

much greater losses at thresholds of 60 or 90. Fortunately, genotype concordance rates at both

low and high mapQ thresholds were stable, despite the putatively higher quality of the

alignments (data not shown). This permitted us to select a lower mapQ threshold (mapQ = 20),

maximizing the number of variants called without sacrificing genotype quality.

Variant calling

Neither GATK's UnifiedGenotyper nor HaplotypeCaller (McKenna et al. 2010; DePristo et al.

2011) called an adequate number of SNPs for fine-mapping in the HS. Panel A of Figure 3

shows that the combination of HaplotypeCaller and Beagle had lower rates of genotype

concordance with array data than calls made by ANGSD and Beagle. We tried several

adjustments to the GATK default thresholds for variant quality, heterozygosity, graph pruning,

dangling branch length, and number of reads sharing an alignment start to declare an active

region; however, none of these changes significantly improved the results (data not shown). As

shown in Panel B of Figure 3, ANGSD/Beagle yielded a substantially greater number of SNPs at

all genotyping error rate thresholds. This observation held when variants were limited only to

biallelic sites and SNPs with an MAF > 0.05.

Figure 3. Genotype discordance rates between array data and variants called by GATK or
ANGSD.

The two panels compare the number variants called by combination of ANGSD and Beagle or
GATK HaplotypeCaller and Beagle at various thresholds of genotype discordance with array
data. Calls were made using the 96 HS rats with array data. (A) The x-axis represents the
genotype discordance rate thresholds and the y-axis is the number of variants that surpass that
threshold for each genotype calling method. (B) Additional filters were applied to the original
SNP sets and the plot zooms in on a smaller range of acceptable discordance rates. Blue lines
represent the unfiltered SNP set. Yellow lines have been filtered for singletons. Red lines have
further excluded SNPs with an MAF < 0.05. Each line contains the same number of points.

ANGSD supports four different models for estimating genotype likelihoods: SAMtools,

GATK, SOAPsnp and SYK. We compared the methods to determine which produced the most

SNPs with the lowest error rates. The SOAPsnp model demonstrated an advantage in genotype

accuracy and number of variants called post-imputation with Beagle (Figure S7). However,

SOAPsnp requires considerably more computational resources, outweighing its marginal benefits

when applied to a large sample set (>2,000). The GATK model showed a greater number of

variants for more lenient discordance rate threshold, but as stringency increased, the number of

variants converged across the remaining 3 models. We proceeded with the SAMtools model for

genotype likelihood estimation due to its previous support in the GBS literature (Torkamaneh et

al. 2017), accepting a nominal decrease in highly concordant variants for a large gain in

efficiency.

Imputation to reference panel

Imputation is use in two ways in our protocol. As described above, we use imputation to assign

missing genotypes at SNPs called in some but not all individuals. In addition, we use imputation

in this section to call genotypes at site where GBS did not provide a call for any individuals.

Thus, our second application (described here) is similar to the human genetics application in

which imputation using 1000 Genomes increases the number of SNPs beyond those included on

a given microarray platform.

Before starting this imputation step, we observed an inflated transition/transversion ratio

(Table S4) in our ANGSD/Beagle SNPs. This issue was ameliorated when filtered the SNP set

for only "known" variants that were previously identified in either the 42 inbred strains

(Hermsen et al. 2015) or the 8 deep-sequenced HS founders (Ramdas et al. 2018). For

imputation, we therefore only provided IMPUTE2 with previously identified variant sites from

our ANGSD/Beagle output. Prior to running IMPUTE2, we also filtered the variants for biallelic

sites with a genotype call in more than one individual. Using pedigree data for the HS rats, we

further removed samples and SNPs showing uncharacteristically high levels of Mendelian errors.

There were 4 samples and 4,179 SNPs removed from subsequent analyses. Lastly, we removed

any samples where the sex chromosome read ratio was contrary to their reported sex (Figure S1).

To determine which reference set to use for imputation, we tested six different possible

combinations of available reference data (Table 2). The most accurate imputation was observed

for the reference set containing only the 8 deep-sequenced HS founder strains from UMich

(Ramdas et al. 2018); however, imputation to this set had the lowest genotyping rate of all

panels. In contrast, using the 42 rat inbred strains displayed a balance of high accuracy and low

missingness, leading us to choose this as our reference set. To better understand the role of the 8

founder strains, who were part of the 42 strains reference panel, we created a reference panel that

included only the 34 non-HS founder strains. As expected, discordance rates were much higher

when only considering non-founders. However, genotyping rates were greater for the 34 than the

8 founders, suggesting a combination of the two was the optimal set.

Table 2. Imputation accuracy based on different variant reference panels for IMPUTE2.

The table includes six different possible reference panels for imputation. The 42 inbred strains, 34 non-founder inbred strains, and 8 HS founders from the 42 inbred strains all were derived from Hermsen et al. 2015 (Hermsen et al. 2015). The UMich 8 HS founders were obtained from Ramdas et al. 2018 (Ramdas et al. 2018). The final set of 8 HS founder was taken from Baud et al. 2013 (Rat Genome Sequencing and Mapping Consortium et al. 2013).

		Chr1	Chr2
42 inbred strains	**Discordance rate**	0.011	0.010
	# Variants	790,659	882,993
	Genotyping Rate	0.85	0.81
UMich 8 founders + 42 inbred strains	**Discordance rate**	0.007	0.011
	# Variants	864,670	898,621
	Genotyping Rate	0.63	0.64
All 34 non-founder inbred strains	**Discordance rate**	0.035	0.030
	# Variants	812,550	912,749
	Genotyping Rate	0.84	0.80
8 HS founders only from the 42 inbred strains	**Discordance rate**	0.012	0.011
	# Variants	805,424	902,061
	Genotyping Rate	0.57	0.53
UMich 8 HS founders only	**Discordance rate**	0.0059	0.008
	# Variants	865,514	898,621
	Genotyping Rate	0.42	0.41
Baud et. al 2013 8 HS founders only	**Discordance rate**	0.0095	0.0096
	# Variants	507,909	540,844
	Genotyping Rate	0.43	0.40

IMPUTE2 requires a genetic map. As described in the methods section, we considered

four different genetic maps, two that were empirically derived and two that were linear

extrapolations based on the physical map. All genetic map performed similarly (Table S5).

Surprisingly, the linear genetic maps performed just as well as the HS-specific map (Littrell et al.

2018). Thus, for simplicity, we chose to use the chromosome-specific values initially published

by Jensen-Seaman (Jensen-Seaman 2004).

Pre-phasing with VCFtools and IMPUTE2 performed similarly, therefore we used

IMPUTE2. SHAPEIT was not used because it showed a higher rate of genotype discordance

compared to VCFtools and IMPUTE2 (data not shown).

To obtain our final set of ~3.7 million variants, a final round of variant filtering was

performed after imputation to the 42 strain reference panel. We removed SNPs with MAF <

0.005, a post-imputation genotyping rate < 90%, and SNPs that violated HWE with $p<1\times10^{-10}$.

DISCUSSION

The use of microarrays and WGS for genotyping large samples in model organisms remains

cost-prohibitive. There is therefore an urgent and wide-spread need for a high-performance and

economical methods of obtaining genome-wide genotype data. While reduced-representation

approaches have been utilized in numerous species of plants and animals, including rodents

(Peterson et al. 2012; Fitzpatrick et al. 2013; Parker et al. 2016; Gonzales et al. 2017; Zhou et al.

2018), there has yet to be a published protocol optimized specifically for rats. Prior to

sequencing thousands of HS samples with GBS for our mapping efforts, we wanted to ensure we

were capturing the greatest possible number of high-quality variants at the lowest possible cost.

The protocol we present here is the culmination of careful testing and optimization of each step

of the GBS protocol for rats. We have now applied the approach to 4,973 HS rats, as well as

4,608 Sprague Dawley rats (Gileta et al. 2018).

Our previous GBS protocol (Parker *et al*, 2016), which was designed for use with CFW

mice, was unsuitable for our current genotyping efforts in HS rats, due to the much higher levels

of genetic diversity in the HS population. There are multiple reasons we chose to develop our

own computational pipeline for GBS rather than using existing workflows. Foremost, the

prominent GBS analysis pipelines were developed and optimized for use in crop species (Sonah

et al. 2013; Catchen et al. 2013; Glaubitz et al. 2014; Torkamaneh et al. 2017; Wickland et al.

2017), which are polyploid and have differing levels of variation and LD than outbred rodent

populations. Additionally, there were elements of each pipeline that did not meet our needs or

lacked customizability. For instance, TASSEL-GBS v2 (Glaubitz et al. 2014) trims all reads to

92 base pairs; however, other projects underway in our lab utilized up to 125bp reads, leading to

a ~20% reduction in data. TASSEL-GBS also ignores read base quality scores, which are

informative in probabilistic frameworks for estimating uncertainty in alignments and variant

calls (H. Li, Ruan, and Durbin 2008; DePristo et al. 2011; Nielsen et al. 2011), and uses a naïve

binomial likelihood ratio method for calling SNPs. Stacks has previously shown poor

demultiplexing (Herten et al. 2015; Torkamaneh et al. 2017) and does not make use of the

reference genome for priors when calling SNPs (Catchen et al. 2013). Fast-GBS relies on

Platypus for variant calling (WGS500 Consortium et al. 2014; Torkamaneh et al. 2017), which

employs a Bayesian method of constructing candidate haplotypes that works poorly with low-

pass sequencing data and does not scale well (Z. Li, Wang, and Wang 2018). Lastly, none of

these pipelines included an imputation step, which is crucial for filling in missing genotypes in

GBS data and can provide millions of additional SNPs given an appropriate composite reference

panel (B. Howie, Marchini, and Stephens 2011; G.-H. Huang and Tseng 2014).

Though we have not explicitly tested each alternate GBS pipeline for the purposes of this

publication, this has been recently done by Wickland et al. (Wickland et al. 2017). Their pipeline

GB-eaSy, which ours most closely resembles, was found to be superior by a number of metrics

to Stacks, TASSEL-GBS, IGST, and Fast-GBS. Similar to GB-eaSy, our pipeline utilizes a

double-digest GBS protocol, aligns reads to the reference genome with *bwa-mem*, and uses the

SAMtools genotype likelihood model for calling SNPs (H. Li 2011). The combination of bwa-

mem and SAMtools algorithm was independently shown to have the best performance for calling

SNPs from Illumina data (Hwang et al. 2015), further supporting our choice of these programs

for read alignment and variant calling. Additionally, using the ANGSD wrapper provided us with

the ability to convert the posterior genotype probabilities into genotype dosages for mapping

studies (Korneliussen, Albrechtsen, and Nielsen 2014).

A minor difference between GB-eaSy and our pipeline is the use of Cutadapt (Martin

2011) rather than GBSX (Herten et al. 2015) for demultiplexing, though both performed equally

well (Table S1). The primary improvement is our extension of the pipeline with the

implementation of effective internal and reference-based imputation steps using the 42 inbred rat

genomes (Hermsen et al. 2015) and 8 deep-sequenced HS founders from UMich (Ramdas et al.

2018). There are two stages of imputation in our pipeline: the first one is accomplished by

Beagle, and aims to fill in missing genotypes at called variants using information from other

samples; this raises the genotype call rate to 100% but may also introduce errors if there is low

confidence in the existing genotype calls, emphasizing the need for careful filtering steps. The

second stage of imputation made use of IMPTUE2 and an external reference panels of variants

called from WGS data on the 8 inbred HS founders, as well as 34 additional inbred rat strains.

We decided to include the 34 additional strains because of the elevated genotyping rate we

observed upon their inclusion in the IMPUTE2 reference panel. We attribute this to the presence

of haplotypes that exist in both the 8 the HS founder strains and a subset of the 34 additional

strains in this panel.

In summary, we have adapted a GBS protocol and genotyping and imputation pipeline to

obtain dense genotypes on genome-wide markers in highly-multiplexed HS rats. After quality

filtering on the level of SNP and sample, over 3.7 million were called with a concordance rate of

99%. The ddGBS protocol and bioinformatic methods used to produce this data are publicly

available, easy to handle, and cost-effective. The presented workflow could be feasibly followed

with marginal modifications for application in other species.

550   ACKNOWLEDGEMENTS:

551   This work was supported by P50 DA037844, R21 DA036672, T32 GM07197, and

552   F31 DA039638.

553   LITERATURE CITED

Aird, Daniel, Michael G Ross, Wei-Sheng Chen, Maxwell Danielsson, Timothy Fennell, Carsten Russ,
David B Jaffe, Chad Nusbaum, and Andreas Gnirke. 2011. "Analyzing and Minimizing PCR
Amplification Bias in Illumina Sequencing Libraries." *Genome Biology* 12 (2): R18.
https://doi.org/10.1186/gb-2011-12-2-r18.
Andrews, Simon. 2017. *FastQC* (version 0.11.6).
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
Baird, Nathan A., Paul D. Etter, Tressa S. Atwood, Mark C. Currey, Anthony L. Shiver, Zachary A. Lewis,
Eric U. Selker, William A. Cresko, and Eric A. Johnson. 2008. "Rapid SNP Discovery and Genetic
Mapping Using Sequenced RAD Markers." Edited by Justin C. Fay. *PLoS ONE* 3 (10): e3376.
https://doi.org/10.1371/journal.pone.0003376.
Browning, Brian L., and Sharon R. Browning. 2009. "A Unified Approach to Genotype Imputation and
Haplotype-Phase Inference for Large Data Sets of Trios and Unrelated Individuals." *The
American Journal of Human Genetics* 84 (2): 210–23.
https://doi.org/10.1016/j.ajhg.2009.01.005.
Browning, Brian L., and Sharon R. Browning. 2016. "Genotype Imputation with Millions of Reference
Samples." *The American Journal of Human Genetics* 98 (1): 116–26.
https://doi.org/10.1016/j.ajhg.2015.11.020.
Catchen, Julian, Paul A. Hohenlohe, Susan Bassham, Angel Amores, and William A. Cresko. 2013. "Stacks:
An Analysis Tool Set for Population Genomics." *Molecular Ecology* 22 (11): 3124–40.
https://doi.org/10.1111/mec.12354.
Chen, Qiang, Yufang Ma, Yumei Yang, Zhenliang Chen, Rongrong Liao, Xiaoxian Xie, Zhen Wang, et al.
2013. "Genotyping by Genome Reducing and Sequencing for Outbred Animals." Edited by
Shuhong Zhao. *PLoS ONE* 8 (7): e67500. https://doi.org/10.1371/journal.pone.0067500.
Chitre, Apurva S, Oksana Polesskaya, Katie Holl, Jianjun Gao, Riyan Cheng, Angel Martinez, Tony George,
et al. 2018. "Genome Wide Association Study of Body Weight, Body Mass Index, Adiposity, and
Fasting Glucose in 3,173 Outbred Rats," September. https://doi.org/10.1101/422428.
Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, et al. 2011.
"The Variant Call Format and VCFtools." *Bioinformatics* 27 (15): 2156–58.
https://doi.org/10.1093/bioinformatics/btr330.
Davey, John W., Paul A. Hohenlohe, Paul D. Etter, Jason Q. Boone, Julian M. Catchen, and Mark L.
Blaxter. 2011. "Genome-Wide Genetic Marker Discovery and Genotyping Using next-Generation
Sequencing." *Nature Reviews Genetics* 12 (7): 499–510. https://doi.org/10.1038/nrg3012.
De Donato, Marcos, Sunday O. Peters, Sharon E. Mitchell, Tanveer Hussain, and Ikhide G. Imumorin.
2013. "Genotyping-by-Sequencing (GBS): A Novel, Efficient and Cost-Effective Genotyping
Method for Cattle Using Next-Generation Sequencing." Edited by James C. Nelson. *PLoS ONE* 8
(5): e62137. https://doi.org/10.1371/journal.pone.0062137.
Delaneau, Olivier, Jonathan Marchini, and Jean-François Zagury. 2011. "A Linear Complexity Phasing
Method for Thousands of Genomes." *Nature Methods* 9 (December): 179.
DePristo, Mark A, Eric Banks, Ryan Poplin, Kiran V Garimella, Jared R Maguire, Christopher Hartl,
Anthony A Philippakis, et al. 2011. "A Framework for Variation Discovery and Genotyping Using
Next-Generation DNA Sequencing Data." *Nature Genetics* 43 (5): 491–98.
https://doi.org/10.1038/ng.806.
Durvasula, Arun, Paul J Hoffman, Tyler V Kent, Chaochih Liu, Thomas J Y Kono, Peter L Morrell, and
Jeffrey Ross-Ibarra. n.d. "ANGSD-Wrapper: Utilities for Analyzing next Generation Sequencing
Data." Accessed September 5, 2018. https://doi.org/10.7287/peerj.preprints.1472v2.

Elshire, Robert J., Jeffrey C. Glaubitz, Qi Sun, Jesse A. Poland, Ken Kawamoto, Edward S. Buckler, and Sharon E. Mitchell. 2011. "A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species." Edited by Laszlo Orban. *PLoS ONE* 6 (5): e19379. https://doi.org/10.1371/journal.pone.0019379.

Fitzpatrick, Christopher J., Shyam Gopalakrishnan, Elizabeth S. Cogan, Lindsay M. Yager, Paul J. Meyer, Vedran Lovic, Benjamin T. Saunders, et al. 2013. "Variation in the Form of Pavlovian Conditioned Approach Behavior among Outbred Male Sprague-Dawley Rats from Different Vendors and Colonies: Sign-Tracking vs. Goal-Tracking." Edited by Patrizia Campolongo. *PLoS ONE* 8 (10): e75042. https://doi.org/10.1371/journal.pone.0075042.

Flanagan, Sarah P., and Adam G. Jones. 2018. "Substantial Differences in Bias between Single-Digest and Double-Digest RAD-Seq Libraries: A Case Study." *Molecular Ecology Resources* 18 (2): 264–80. https://doi.org/10.1111/1755-0998.12734.

Fu, Yong-Bi, and Mo-Hua Yang. 2017. "Genotyping-by-Sequencing and Its Application to Oat Genomic Research." In *Oat*, edited by Sebastian Gasparis, 1536:169–87. New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-6682-0_13.

Furuta, Tomoyuki, Motoyuki Ashikari, Kshirod K. Jena, Kazuyuki Doi, and Stefan Reuscher. 2017. "Adapting Genotyping-by-Sequencing for Rice F2 Populations." *G3: Genes|Genomes|Genetics* 7 (3): 881–93. https://doi.org/10.1534/g3.116.038190.

Gileta, Alexander F., Christopher J. Fitzpatrick, Apurva S. Chitre, Celine L. St. Pierre, Elizabeth V. Joyce, Rachael J. Maguire, Africa M. McLeod, et al. 2018. "Genetic Characterization of Outbred Sprague Dawley Rats and Utility for Genome-Wide Association Studies," September. https://doi.org/10.1101/412924.

Glaubitz, Jeffrey C., Terry M. Casstevens, Fei Lu, James Harriman, Robert J. Elshire, Qi Sun, and Edward S. Buckler. 2014. "TASSEL-GBS: A High Capacity Genotyping by Sequencing Analysis Pipeline." Edited by Nicholas A. Tinker. *PLoS ONE* 9 (2): e90346. https://doi.org/10.1371/journal.pone.0090346.

Gonzales, Natalia M., Jungkyun Seo, Ana Isabel Hernandez-Cordero, Celine L. St. Pierre, Jennifer S. Gregory, Margaret G. Distler, Mark Abney, Stefan Canzar, Arimantas Lionikas, and Abraham A. Palmer. 2017. "Genome Wide Association Study of Behavioral, Physiological and Gene Expression Traits in a Multigenerational Mouse Intercross," December. https://doi.org/10.1101/230920.

———. 2018. "Genome Wide Association Analysis in a Mouse Advanced Intercross Line," September. https://doi.org/10.1101/230920.

Hannon Lab. 2010. *FASTX-Toolkit* (version 0.0.13). http://hannonlab.cshl.edu/fastx_toolkit/index.html.

He, Jiangfeng, Xiaoqing Zhao, André Laroche, Zhen-Xiang Lu, HongKui Liu, and Ziqin Li. 2014. "Genotyping-by-Sequencing (GBS), an Ultimate Marker-Assisted Selection (MAS) Tool to Accelerate Plant Breeding." *Frontiers in Plant Science* 5 (September). https://doi.org/10.3389/fpls.2014.00484.

Hermsen, Roel, Joep de Ligt, Wim Spee, Francis Blokzijl, Sebastian Schäfer, Eleonora Adami, Sander Boymans, et al. 2015. "Genomic Landscape of Rat Strain and Substrain Variation." *BMC Genomics* 16 (1). https://doi.org/10.1186/s12864-015-1594-1.

Herten, Koen, Matthew S Hestand, Joris R Vermeesch, and Jeroen KJ Van Houdt. 2015. "GBSX: A Toolkit for Experimental Design and Demultiplexing Genotyping by Sequencing Experiments." *BMC Bioinformatics* 16 (1). https://doi.org/10.1186/s12859-015-0514-3.

Howie, Bryan, Christian Fuchsberger, Matthew Stephens, Jonathan Marchini, and Gonçalo R Abecasis. 2012. "Fast and Accurate Genotype Imputation in Genome-Wide Association Studies through Pre-Phasing." *Nature Genetics* 44 (July): 955.

Howie, Bryan, Jonathan Marchini, and Matthew Stephens. 2011. "Genotype Imputation with Thousands of Genomes." *G3: Genes|Genomes|Genetics* 1 (6): 457–70. https://doi.org/10.1534/g3.111.001198.

Howie, Bryan N., Peter Donnelly, and Jonathan Marchini. 2009. "A Flexible and Accurate Genotype Imputation Method for the Next Generation of Genome-Wide Association Studies." Edited by Nicholas J. Schork. *PLoS Genetics* 5 (6): e1000529. https://doi.org/10.1371/journal.pgen.1000529.

Huang, Guan-Hua, and Yi-Chi Tseng. 2014. "Genotype Imputation Accuracy with Different Reference Panels in Admixed Populations." *BMC Proceedings* 8 (Suppl 1): S64. https://doi.org/10.1186/1753-6561-8-S1-S64.

Huang, X., Q. Feng, Q. Qian, Q. Zhao, L. Wang, A. Wang, J. Guan, et al. 2009. "High-Throughput Genotyping by Whole-Genome Resequencing." *Genome Research* 19 (6): 1068–76. https://doi.org/10.1101/gr.089516.108.

Hwang, Sohyun, Eiru Kim, Insuk Lee, and Edward M. Marcotte. 2015. "Systematic Comparison of Variant Calling Pipelines Using Gold Standard Personal Exome Variants." *Scientific Reports* 5 (December): 17875.

Illumina, Inc. 2014. "Nextera(R) Library Validation and Cluster Density Optimization: Guidelines for Generating High-Quality Data with Nextera Library Prepartion Kits." https://www.illumina.com/documents/products/technotes/technote_nextera_library_validation.pdf.

Jensen-Seaman, M. I. 2004. "Comparative Recombination Rates in the Rat, Mouse, and Human Genomes." *Genome Research* 14 (4): 528–38. https://doi.org/10.1101/gr.1970304.

Johannesson, M., R. Lopez-Aumatell, P. Stridh, M. Diez, J. Tuncel, G. Blazquez, E. Martinez-Membrives, et al. 2008. "A Resource for the Simultaneous High-Resolution Mapping of Multiple Quantitative Trait Loci in Rats: The NIH Heterogeneous Stock." *Genome Research* 19 (1): 150–58. https://doi.org/10.1101/gr.081497.108.

Johnson, Jennifer L., Helena Wittgenstein, Sharon E. Mitchell, Katie E. Hyma, Svetlana V. Temnykh, Anastasiya V. Kharlamova, Rimma G. Gulevich, et al. 2015. "Genotyping-By-Sequencing (GBS) Detects Genetic Structure and Confirms Behavioral QTL in Tame and Aggressive Foxes (Vulpes Vulpes)." Edited by William J. Murphy. *PLOS ONE* 10 (6): e0127013. https://doi.org/10.1371/journal.pone.0127013.

Kanagawa, Takahiro. 2003. "Bias and Artifacts in Multitemplate Polymerase Chain Reactions (PCR)." *Journal of Bioscience and Bioengineering* 96 (4): 317–23. https://doi.org/10.1016/S1389-1723(03)90130-7.

Korneliussen, Thorfinn Sand, Anders Albrechtsen, and Rasmus Nielsen. 2014. "ANGSD: Analysis of Next Generation Sequencing Data." *BMC Bioinformatics* 15 (1). https://doi.org/10.1186/s12859-014-0356-4.

Li, H. 2011. "A Statistical Framework for SNP Calling, Mutation Discovery, Association Mapping and Population Genetical Parameter Estimation from Sequencing Data." *Bioinformatics* 27 (21): 2987–93. https://doi.org/10.1093/bioinformatics/btr509.

Li, H., and R. Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows-Wheeler Transform." *Bioinformatics* 25 (14): 1754–60. https://doi.org/10.1093/bioinformatics/btp324.

Li, H., J. Ruan, and R. Durbin. 2008. "Mapping Short DNA Sequencing Reads and Calling Variants Using Mapping Quality Scores." *Genome Research* 18 (11): 1851–58. https://doi.org/10.1101/gr.078212.108.

Li, Zhentang, Yi Wang, and Fei Wang. 2018. "A Study on Fast Calling Variants from Next-Generation Sequencing Data Using Decision Tree." *BMC Bioinformatics* 19 (1). https://doi.org/10.1186/s12859-018-2147-9.

Littrell, John, Shirng-Wern Tsaih, Amelie Baud, Pasi Rastas, Leah Solberg-Woods, and Michael J. Flister.
2018. "A High-Resolution Genetic Map for the Laboratory Rat." *G3:*
Genes|Genomes|Genetics, May, g3.200187.2018. https://doi.org/10.1534/g3.118.200187.
Martin, Marcel. 2011. "Cutadapt Removes Adapter Sequences from High-Throughput Sequencing
Reads." *EMBnet.Journal* 17 (1): 10. https://doi.org/10.14806/ej.17.1.200.
McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernytsky, K. Garimella, et al. 2010.
"The Genome Analysis Toolkit: A MapReduce Framework for Analyzing next-Generation DNA
Sequencing Data." *Genome Research* 20 (9): 1297–1303.
https://doi.org/10.1101/gr.107524.110.
Miller, M. R., J. P. Dunham, A. Amores, W. A. Cresko, and E. A. Johnson. 2007. "Rapid and Cost-Effective
Polymorphism Identification and Genotyping Using Restriction Site Associated DNA (RAD)
Markers." *Genome Research* 17 (2): 240–48. https://doi.org/10.1101/gr.5681207.
Nielsen, Rasmus, Joshua S. Paul, Anders Albrechtsen, and Yun S. Song. 2011. "Genotype and SNP Calling
from Next-Generation Sequencing Data." *Nature Reviews Genetics* 12 (6): 443–51.
https://doi.org/10.1038/nrg2986.
Orsouw, Nathalie J. van, René C. J. Hogers, Antoine Janssen, Feyruz Yalcin, Sandor Snoeijers, Esther
Verstege, Harrie Schneiders, et al. 2007. "Complexity Reduction of Polymorphic Sequences
(CRoPS^{TM}): A Novel Approach for Large-Scale Polymorphism Discovery in Complex Genomes."
Edited by Ivan Baxter. *PLoS ONE* 2 (11): e1172. https://doi.org/10.1371/journal.pone.0001172.
Parker, Clarissa C, Shyam Gopalakrishnan, Peter Carbonetto, Natalia M Gonzales, Emily Leung, Yeonhee J
Park, Emmanuel Aryee, et al. 2016. "Genome-Wide Association Study of Behavioral,
Physiological and Gene Expression Traits in Outbred CFW Mice." *Nature Genetics* 48 (8): 919–26.
https://doi.org/10.1038/ng.3609.
Pértille, Fábio, Carlos Guerrero-Bosagna, Vinicius Henrique da Silva, Clarissa Boschiero, José de Ribamar
da Silva Nunes, Mônica Corrêa Ledur, Per Jensen, and Luiz Lehmann Coutinho. 2016. "High-
Throughput and Cost-Effective Chicken Genotyping Using Next-Generation Sequencing."
Scientific Reports 6 (May): 26929.
Peterson, Brant K., Jesse N. Weber, Emily H. Kay, Heidi S. Fisher, and Hopi E. Hoekstra. 2012. "Double
Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and Genotyping in Model
and Non-Model Species." Edited by Ludovic Orlando. *PLoS ONE* 7 (5): e37135.
https://doi.org/10.1371/journal.pone.0037135.
Poland, Jesse A., Patrick J. Brown, Mark E. Sorrells, and Jean-Luc Jannink. 2012. "Development of High-
Density Genetic Maps for Barley and Wheat Using a Novel Two-Enzyme Genotyping-by-
Sequencing Approach." Edited by Tongming Yin. *PLoS ONE* 7 (2): e32253.
https://doi.org/10.1371/journal.pone.0032253.
Ramdas, Shweta, Ayse Bilge Ozel, Katie Holl, Myrna Mandel, Leah Solberg Woods, and Jun Z Li. 2018.
"Extended Regions of Suspected Mis-Assembly in the Rat Reference Genome," September.
https://doi.org/10.1101/332932.
Rat Genome Sequencing and Mapping Consortium, Amelie Baud, Roel Hermsen, Victor Guryev, Pernilla
Stridh, Delyth Graham, Martin W McBride, et al. 2013. "Combined Sequence-Based and Genetic
Mapping Analysis of Complex Traits in Outbred Rats." *Nature Genetics* 45 (7): 767–75.
https://doi.org/10.1038/ng.2644.
Rice, Peter, Ian Longden, and Alan Bleasby. 2000. "EMBOSS: The European Molecular Biology Open
Software Suite." *Trends in Genetics* 16 (6): 276–77. https://doi.org/10.1016/S0168-
9525(00)02024-2.
Roberts, R. J., and D. Macelis. 1999. "REBASE--Restriction Enzymes and Methylases." *Nucleic Acids*
Research 27 (1): 312–13. https://doi.org/10.1093/nar/27.1.312.

Scheben, Armin, Jacqueline Batley, and David Edwards. 2017. "Genotyping-by-Sequencing Approaches
to Characterize Crop Genomes: Choosing the Right Tool for the Right Application." *Plant*
Biotechnology Journal 15 (2): 149–61. https://doi.org/10.1111/pbi.12645.
Sonah, Humira, Maxime Bastien, Elmer Iquira, Aurélie Tardivel, Gaétan Légaré, Brian Boyle, Éric
Normandeau, et al. 2013. "An Improved Genotyping by Sequencing (GBS) Approach Offering
Increased Versatility and Efficiency of SNP Discovery and Genotyping." Edited by Zhanjiang Liu.
PLoS ONE 8 (1): e54603. https://doi.org/10.1371/journal.pone.0054603.
Steen, R. G., A. E. Kwitek-Black, C. Glenn, J. Gullings-Handley, W. Van Etten, O. S. Atkinson, D. Appel, et
al. 1999. "A High-Density Integrated Genetic Linkage and Radiation Hybrid Map of the
Laboratory Rat." *Genome Research* 9 (6): AP1-8, insert.
Sun, Xiaowen, Dongyuan Liu, Xiaofeng Zhang, Wenbin Li, Hui Liu, Weiguo Hong, Chuanbei Jiang, et al.
2013. "SLAF-Seq: An Efficient Method of Large-Scale De Novo SNP Discovery and Genotyping
Using High-Throughput Sequencing." Edited by Jan Aerts. *PLoS ONE* 8 (3): e58700.
https://doi.org/10.1371/journal.pone.0058700.
Torkamaneh, Davoud, Jérôme Laroche, Maxime Bastien, Amina Abed, and François Belzile. 2017. "Fast-
GBS: A New Pipeline for the Efficient and Highly Accurate Calling of SNPs from Genotyping-by-
Sequencing Data." *BMC Bioinformatics* 18 (1). https://doi.org/10.1186/s12859-016-1431-9.
Van Tassell, Curtis P, Timothy P L Smith, Lakshmi K Matukumalli, Jeremy F Taylor, Robert D Schnabel,
Cynthia Taylor Lawley, Christian D Haudenschild, Stephen S Moore, Wesley C Warren, and Tad S
Sonstegard. 2008. "SNP Discovery and Allele Frequency Estimation by Deep Sequencing of
Reduced Representation Libraries." *Nature Methods* 5 (3): 247–52.
https://doi.org/10.1038/nmeth.1185.
Wang, Yuzhe, Xuemin Cao, Yiqiang Zhao, Jing Fei, Xiaoxiang Hu, and Ning Li. 2017. "Optimized Double-
Digest Genotyping by Sequencing (DdGBS) Method with High-Density SNP Markers and High
Genotyping Accuracy for Chickens." Edited by Peng Xu. *PLOS ONE* 12 (6): e0179073.
https://doi.org/10.1371/journal.pone.0179073.
WGS500 Consortium, Andy Rimmer, Hang Phan, Iain Mathieson, Zamin Iqbal, Stephen R F Twigg,
Andrew O M Wilkie, Gil McVean, and Gerton Lunter. 2014. "Integrating Mapping-, Assembly-
and Haplotype-Based Approaches for Calling Variants in Clinical Sequencing Applications."
Nature Genetics 46 (8): 912–18. https://doi.org/10.1038/ng.3036.
Wickland, Daniel P., Gopal Battu, Karen A. Hudson, Brian W. Diers, and Matthew E. Hudson. 2017. "A
Comparison of Genotyping-by-Sequencing Analysis Methods on Low-Coverage Crop Datasets
Shows Advantages of a New Workflow, GB-EaSy." *BMC Bioinformatics* 18 (1).
https://doi.org/10.1186/s12859-017-2000-6.
Woods, Leah C. Solberg, and Richard Mott. 2017. "Heterogeneous Stock Populations for Analysis of
Complex Traits." In *Systems Genetics*, edited by Klaus Schughart and Robert W. Williams,
1488:31–44. New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-6427-7_2.
Zhou, Xinzhu, Celine L. St. Pierre, Natalia M. Gonzales, Riyan Cheng, Apurva S. Chitre, Greta Sokoloff, and
Abraham A. Palmer. 2018. "Genome-Wide Association Study, Replication, and Mega-Analysis
Using a Dense Marker Panel in a Multi-Generational Mouse Advanced Intercross Line," August.
https://doi.org/10.1101/387613.

782
783

Figure S1. Ratio of reads on X-chromosome to total sequencing reads.

The color of the points indicates the pedigree-recorded sex of the samples. Females are expected to have approximately twice as many reads for the X-chromosome. Samples that did not cluster with their pedigree-recorded sex were removed from the study for possible sample mix-up.

Figure S2. Data preparation workflow for imputation with IMPUTE2.

Figure S3. Programmed vs. empirical Pippin Prep fragment size range.

This plot comes from the Bioanalyzer output for a pooled HS library. The x-axis shows the library fragment sizes in base pairs, and the y-axis is in fluorescent units, which represent the quantity of the fragments on the gel chip. There is approximately a 50-75bp shift in the empirical library distribution compared to expectation due to the high quantity of fragments loaded into the Pippin Prep gel cassette.

Figure S4. Raw read counts grouped by shipment batch.

Raw read counts are on a per-sample basis after demultiplexing FASTQ files with FASTX Barcode Splitter.

Figure S5. FASTQC results pre- and post-filtering with Cutadapt.

FASTQC results are from a single sample from the original set of 96 HS samples prepared in 12-plex
and sequenced on the Illumina HiSeq 2500 with 125bp reads.

868 **Figure S6. Overlap of called SNPs with known variants after read trimming with FASTX**
869 **or Cutadapt.**

Figure S7. Number of variants by genotype discordance rates for 4 ANGSD genotype likelihood models.

893
894

Figure S8. Available rat genetic maps.

Plotted physical and genetic distances are for chromosome 12.

Table S1. Demultiplexing performance.

All methods began with the same number of reads from the original FASTQ. Final read and base pair counts are from after the reads have been trimmed of adapter, barcode, and restriction site sequences, as well as low-quality base pairs (< Q20).

	In-house Python Script	GBSX	FASTX Barcode Splitter
Reads with NlaIII adapter sequence	545,177 (3.07%)	475,581 (2.67%)	547,697 (3.07%)
Total bps processed	2,061,523,464	2,116,436,361	2,227,542,500
Total bps written to file	2,059,714,312	2,114,841,934	2,225,724,833
Proportion of bps retained	99.91%	99.92%	99.92%
Reads post-processing	17,771,754	17,786,280	17,820,340

926
927

Table S2. Comparison of variants calls after filtering with FASTX vs Cutadapt.

Data shown comes from the original set of 96 HS samples prepared in 12-plex and sequenced on the Illumina HiSeq 2500. At this step of pipeline optimization, variants were called utilizing GATK UnifiedGenotyper. Calls were unfiltered.

	FASTX Clipper	Cutadapt
Number of variants	6,075,821	6,581,115
Genotyping call rate	0.17	0.19
Mean minor allele count	3.96	4.25
Mean minor allele frequency	0.15	0.15
Number of singletons	433,960	548,975
Number monomorphic sites	807,453	773,074
Transition/transversion ratio	2.32	2.40
T_IT_V ratio for singletons	3.23	3.40
Mean variant read depth	109.56	126.35
Mean quality score	601.79	715.56

940
941

Table S3. Variant metrics resulting from reads filtered at different mapping quality thresholds.

Data shown comes from the original set of 96 HS samples prepared in 12-plex and sequenced on the Illumina HiSeq 2500. Variants were called utilizing the SAMtools model and the -minMapQ filter in ANGSD. Calls were unfiltered.

	MAPQ = 20	MAPQ = 30	MAPQ = 45	MAPQ = 60	MAPQ = 90
Number of variants	372,860	372,330	363,790	316,949	233,322
Genotyping call rate	0.64	0.64	0.64	0.61	0.75
Mean minor allele count	5.96	5.96	6.06	5.86	7.36
Mean minor allele frequency	0.18	0.18	0.18	0.18	0.19
Number of singletons	16,781 (4.50%)	16,732 (4.49%)	16,550 (4.55%)	17,352 (5.47%)	11,773 (5.05%)
Number of monomorphic sites	122,478 (32.85%)	122,188 (32.82%)	116,738 (32.09%)	100,074 (31.57%)	56,179 (24.08%)
Transition/transversion ratio	1.23	1.24	1.26	1.31	1.41
$T_I T_V$ ratio for singletons	1.27	1.28	1.28	1.31	1.38
Mean variant read depth	157.78	157.73	159.25	152.48	188.80
Mean quality score	2,547	2,548	2,556	2,461	2,954

Table S4. Transition/transversion ratio before and after known sites filtering.

The presented data comes from ANGSD/Beagle variant calls for 3,601 HS samples, prior to imputation with IMPUTE2. Known SNPs came from both the 42 inbred genomes from Hermsen et. al 2015 (Hermsen et al. 2015) and the 8 inbred HS founder strains sequenced by the University of Michigan (Ramdas et al. 2018).

	Unfiltered SNPs	Filtered for known SNPs
AC	15,157	9,166
AG	888,657	42,275
AT	15,432	7,610
CG	18,043	8,061
CT	893,653	41,938
GT	15,118	9,177
T_S	1,782,310	84,213
T_V	63,750	34,014
T_ST_V	27.96	2.48
Total # SNPs	1,846,060	118,227

961
962

Table S5. Imputation accuracy for chromosome 12 across different genetic maps.

The number of variants used for the concordance check is dependent on the overlap of the imputed variants with array data for the 96 HS rats with array genotypes. The MAF filter only removes monomorphic sites within the 96 HS rat sample used for the concordance check.

	cM/Mb = 1.00	cM/Mb = 1.16	SHRSPxPN	HS-specific
Number of variants before QC	158,452	158,452	158,452	158,452
Genotyping rate before QC	0.94	0.92	0.92	0.92
Variant removed for missingness > 10%	22,217	28,959	28,356	28,858
Variants removed for MAF < 0.005	50,380	61,270	61,592	59,812
Variants removed for HWE < 1×10^{-10}	53	56	57	56
Number of variants after QC	85,802	68,167	68,447	69,726
Genotyping rate after QC	0.93	0.91	0.92	0.91
Number of variants in concordance check	5,912	5,590	5,594	5,646
Discordance rate	0.095	0.011	0.011	0.010