Epigenetic suppression of interferon lambda receptor expression leads to an enhanced

HuNoV replication *in vitro*.

Sabastine E. Arthur*, Frédéric Sorgeloos*[#], Myra Hosmillo, and Ian G. Goodfellow[#].

Division of Virology, Department of Pathology, University of Cambridge, Addenbrooke's Hospital,

Hills Road, Cambridge, CB2 0QQ, United Kingdom

[#]Corresponding authors.

* These authors contributed equally to this work.

Email: ig299@cam.ac.uk , frederic.sorgeloos@gmail.com

Keywords: Human norovirus replicon, microarray analysis, interferon lambda receptor

(IFNLR1), DNA methylation, epigenetic reprogramming.

Running Title: Epigenetic reprogramming of HuNoV replicon harbouring cells

The authors declare no conflict of interests.

Abstract

Human norovirus (HuNoV) is the main cause of acute gastroenteritis worldwide yet no therapeutics

are currently available. Here, we utilize a human norovirus replicon in epithelial human gastric tumor

(HGT) cells to identify host factors involved in promoting or inhibiting HuNoV replication. We

observed that an IFN-cured population of replicon-harboring HGT cells (HGT-cured) was enhanced

in their ability to replicate transfected HuNoV RNA compared to parental HGT cells, suggesting that

differential gene expression in HGT-cured cells created an environment favouring the replication of

viral RNA. Microarray analysis was used to identify genes differentially regulated in HGT-NV and

HGT-cured compared to parental HGT cells. We found that the IFN lambda receptor alpha (IFNLR1)

expression was highly reduced in HGT-NV and HGT-cured cells. All three cell lines responded to

exogenous IFN-β by inducing interferon stimulated genes (ISGs), however, HGT-NV and HGT-cured

failed to respond to exogenous IFN-λ. Inhibition of DNA methyltransferase activity with 5-aza-2'-

deoxycytidine partially reactivated IFNLR1 expression in HGT-NV and IFN-cured cells suggesting

that host adaptation occurred via epigenetic reprogramming. In line with this observation, ectopic

expression of the IFN-λ receptor alpha rescued HGT-NV and HGT-cured cells response to IFN-λ. We

conclude that type III IFN is important in inhibiting HuNoV replication in vitro and that the loss of

IFNLR1 enhances replication of HuNoV. To the best of our knowledge, this study unravels for the

first time epigenetic reprograming of the interferon lambda receptor as a new mechanism of cellular

adaptation during long-term RNA virus replication and shows that an endogenous level of interferon

lambda signalling is able to control human norovirus replication.

Introduction

With the introduction of the rotavirus vaccine, human norovirus (HuNoV) is now the main etiologic

agent responsible for gastroenteritis worldwide (Fankhauser *et al.*, 1998; Lopman *et al.*, 2013; Hall *et*

al., 2013; Mawatari and Kato, 2014). The disease, characterized by diarrhoea, nausea and vomiting is

generally self-limiting in healthy adults. However, in immunocompromised, elderly patients and

young children under the age of five, the disease can become chronic and sometimes lead to death

mostly due to dehydration (Ludwig *et al.*, 2008; Patel *et al.*, 2008; Phillips *et al.*, 2010; Payne *et al.*,

2013; Belliot *et al.*, 2014). Despite this and the profound economic burden of the disease (Bartsch *et*

al., 2016), there are no therapeutics or licensed vaccines available.

Until the advent of the B lymphocyte-based and stem cell-derived human intestinal enteroid culture

systems (Jones *et al.*, 2014; Ettayebi *et al.*, 2016), HuNoV research has been hampered by the lack of

cell culture and small animal models recapitulating noroviral infection and pathogenesis. However,

limitations in robustness, cost and labour intensity associated with both methods foreshadow that

advances in HuNoV research still relies on the well-established murine norovirus (MNV) infection

model and norovirus replicon systems (Wobus *et al.*, 2004; Chang *et al.*, 2006). These systems have

been used to identify key host factors that impact the lifecycle of HuNoV, reviewed by Thorne and

Goodfellow (Thorne and Goodfellow, 2014).

Contrary to the type I IFNs (IFN-α/β) which were discovered more than an half century ago (Isaacs

and Lindenmann, 1988), type III interferons (IFN-λ) were discovered only little over a decade

(Kotenko *et al.*, 2003; Sheppard *et al.*, 2003). Although both cytokine families possess similar

functions, a few but crucial differences exist in their biology. Notably, both cytokines signal through

distinct heterodimeric receptors with type I IFN signaling through the interferon alpha/beta receptor

(IFNAR) composed of the IFNAR1 and IFNAR2 subunits and type III IFN through the interferon

lambda receptor (IFNLR) which consists of the interferon lambda receptor 1 (IFNLR1) and

interleukin 10 receptor beta (IL10RB) subunits. Unlike the type I IFN receptor that is expressed on

most cell types, expression of the IFNLR1 subunit is restricted to cells of the mucosal epithelium,

neutrophils and human hepatocytes (De Weerd *et al.*, 2007; Sommereyns *et al.*, 2008; Blazek *et al.*,

2015). Although the immune cells of the blood were also shown to express the IFNLR1 subunit, the

receptor has been reported to lack the ability to respond to IFN-λ (Witte *et al.*, 2009).

Expression of type I and III interferons is regulated at the transcriptional level and relies on the

recognition of conserved pathogen-associated molecular patterns (PAMPs). Detection of these

molecular signatures by extracellular and intracellular pattern recognition receptors triggers the

coordinated activation of distinct signaling pathways responsible for the activation of IRF-3/7 and

NF-κB transcription factors that are required for IFN gene transcription. Secreted IFN-λ then binds to

the IFNLR and is thought to activate the Janus kinase 1 and tyrosine kinase 2. Subsequently, recruited

signal transducer and activator of transcription 1 (STAT1) and STAT2 are activated through

phosphorylation leading to the expression of IFN stimulated genes (ISGs), some of which have direct

antiviral activities (Dickensheets *et al.*, 2013). Interferon lambdas are important players in both innate

and adaptive immunity and have profound antiviral effects on a variety of viruses (Ank *et al.*, 2008;

Mordstein *et al.*, 2010; Hemann *et al.*, 2017; Lee and Baldridge, 2017; Syedbasha and Egli, 2017).

Expression of IFNLR1 on epithelial cells of the small intestine and colon was shown to be important

in IFN-λ-mediated antiviral activity against persistent MNV and reovirus infection *in vivo* (Baldridge

et al., 2017). Treatment of persistently infected mice lacking adaptive immune response (Rag1$^{-/-}$)

with IFN-λ abolished viral replication, suggesting that IFN-λ can cure persistent MNV infection in

absence of adaptive immunity and this ability requires the expression of IFNLR1 (Nice *et al.*, 2015).

In line with this, the effect of antibiotics that inhibit persistent MNV infection in the gut has also been

shown to be dependent on IFNLR1 expression as well as IRF3 and STAT1 transcription factors

(Baldridge *et al.*, 2014). It was observed that AG129 sentinel mice lacking the ability to respond to

both IFN-α/β and IFN-γ housed together with MNV-infected mice developed a diarrhoea-associated

MNV infection. Overexpression of IFN-λ in sentinel mice upregulated ISG expression, inhibited

MNV replication in the small intestine and prevented them from being infected when co-housed with

MNV-infected mice (Rocha-Pereira *et al.*, 2018a).

Results

Generation and characterization of human cell lines bearing stable human norovirus replicons.

To understand the influence of viral and host factors involved in HuNoV replication, we sought to generate several human cell lines stably replicating HuNoV RNA. To this end, BHK-21 cells were transfected with capped Norwalk replicon RNA harbouring a neomycin selection marker (Chang *et al.*, 2006) and subjected to G418 selection 48 hours after transfection (Fig. 1A). Although the vast majority of the cells died within one week, individual cell colonies were observed and subjected to limiting dilution. A single high-expressing clone was selected and expanded to generate stable replicon-containing BHK-21 cells (BHK-NV). VPg-linked RNA extracted from these cells was transfected into HGT cells, a cell line of epithelial origin which was subsequently selected on the basis of their G418 resistance in order to generate human norovirus replicon cells (HGT-NV). These HGT-NV cells were either collected as a population or subjected to limiting dilution to produce HGT-NV cell clones. The HGT-NV population was further passaged 16 times in the presence of IFN□α at a concentration of 1000 U/mL in the absence of G418 selection over an 8 weeks period leading to the generation of HGT-Cured cells. These cells were subsequently cultured in presence of G418 to observe their loss of resistance to G418 confirming the complete elimination of the replicon. Detection of HuNoV RNA by RT-qPCR analysis confirmed the presence of noroviral genomes in HGT-NV cells that were absent from HGT-Cured or parental HGT cells used as control (Fig. 1B). To confirm the presence of authentic steady-state replication of Norwalk RNA, cells were subjected to immunofluorescence analysis using monoclonal antibodies directed against dsRNA, a by-product assumed to be universally generated during viral replication (Weber *et al.*, 2006; Son *et al.*, 2015). As shown in Fig. 1C, punctate structures reminiscent of replication complexes were identified in HGT-NV cells while no signal above background levels was detected in HGT-Cured or in parental cell lines. Taken together these results suggest that HuNoV VPg-linked RNA successfully replicates in HGT cells and displays all the characteristics of a replication-competent RNA.

Cured population of HGT-NV cells demonstrate enhanced HuNoV replication.

Stable viral RNA replication under drug-mediated selection leads to an environment where on the one

hand host cells face the selective pressure of the drug and, on the other hand, replicating RNA faces

innate cellular responses aimed at inhibiting replication. This results in the establishment of a

metastable equilibrium prone to adaptation from both cells and the replicating viral RNA. To test for

the appearance of such adaptive mutation(s) in the replicon, VPg-linked RNA was extracted from

BHK-NV and HGT-NV cells and subjected to consensus genome sequencing. Sequence analysis

revealed that viral RNA originating from both BHK-NV and HGT-NV cells underwent many

synonymous and non-synonymous genomic changes, suggesting the potential adaptation of HuNoV

RNA to specific cellular environments (Supporting information Table 1). To test specifically for

host cell adaptation, wild-type VPg-linked RNA extracted from BHK-NV cells was re-transfected in

both HGT and HGT-Cured cell lines that were subsequently selected for 5 days with G418. As shown

in Fig. 2A, HGT-Cured cells gave rise to a greater number of stable replicon colonies than parental

HGT cells. To quantitatively confirm this observation, each cell line was transfected with wild-type

VPg-linked RNA replicon and cells were harvested at various time points for RNA extraction and

viral RNA quantification (Fig. 2B). From 5 days post transfection, the levels of Norwalk replicon

RNA were significantly higher in HGT-Cured cells when compared to HGT cells. Noroviral RNA

levels further increased at day 6 to yield a 15-fold difference between HGT-Cured and HGT cells.

Quantification of transfected VPg-linked viral RNA at 6h post transfection revealed similar levels in

both HGT and HGT-Cured cells suggesting that IFN treatment did not alter the ability of HGT-Cured

cells to be transfected (data not shown). We thus concluded that HGT-Cured cells possess a greater

degree of permissiveness to viral replication than parental HGT cells.

An alteration in cellular environment is responsible for enhanced viral replication in HGT-

Cured cells.

Increased replication of HuNoV VPg-linked RNA in HGT-Cured cells suggests that these cells

provide a better cellular environment compared to that of parental HGT cells. To get insights into the

mechanism of this cell-derived HuNoV replicon permissiveness, genome-wide expression profiles

from HGT, HGT-NV and HGT-Cured cell lines were quantified using Illumina BeadChip microarrays

(Tables S2 and S3). We first compared the gene expression profiles of HGT-NV cells relative to their

HGT controls. As seen in Figure 3A, more than 2000 genes (HGT-NV-vs-HGT: 2074, HGT-Cured-

vs-HGT: 858) were identified for which mRNA expression was significantly modified at a FDR lower

than 0.01. Of these, a minority of 151 genes had expression changes in either direction greater than 2-

fold suggesting that HuNoV replication has a marginal effect on the whole transcriptional landscape

of replicon harbouring cells. Using the same criteria, comparison of gene expression profiles between

HGT cells and HGT-Cured cells led to identification of 101 genes for which expression was

differentially regulated between the two cell lines (Tables S4-S5). To confirm these observations and

to probe for the accuracy of gene expression measured by microarray analysis, we first compared

expression changes obtained by microarrays with that measured by RT-qPCR analysis. To this end,

ten genes differentially regulated across conditions and spanning a wide range of fold changes were

chosen for RT-qPCR validation (Fig. 3B). Globally, we observed a high correlation between the two

techniques with Pearson's correlation coefficients ranging from 0.90 to 0.95. Direct comparison of

raw microarray signal intensities with differences in cycle threshold obtained by real-time PCR

displayed the same trends suggesting that similar fold changes were primarily the consequence of

gene expression differences and were not an artefact resulting from different gene normalization

techniques. Differential gene expression at the protein level of selected genes was further confirmed

by western blot analysis (Fig. S1A-B). Taken together these data extensively confirms the reliability

of gene expression measurements by microarray analysis.

The number of genes showing similar regulation between HGT-NV and HGT-Cured is surprisingly

high and likely reflects cellular long-term adaption to replicating RNA (GO term enrichment analysis

did not reveal specific pathway enrichment in this gene set). Remarkably, we observed an opposite

regulation of several interferon-stimulated genes (ISGs) when comparing the HGT-NV transcriptional

landscape with that of parental HGT cells (Fig. 3C and Table S4). While genes coding for IFITM2,

IFITM3, IFIH1 and IFI27L2 were down-regulated, IFIT1 and IFIT2 genes were significantly up-

regulated in HGT-NV cells. Expression of IFIT1 and IFIT2 proteins is known to be IRF3-dependent

(Grandvaux *et al.*, 2002; Andersen *et al.*, 2008), while expression of IFITMs and IFIH1 genes was

shown to be mediated by ISRE binding (Friedman *et al.*, 1984; Reid *et al.*, 1989), this suggests that

while HuNoV replication induces IRF3-dependent immune responses, activation of ISRE-dependent

genes located downstream in the interferon signalling pathway is inhibited. In line with this

observation, microarray analysis identified a statistically significant down-regulation of the type III

interferon receptor (IFNLR1) expression in HGT-NV and HGT-Cured cells when both were

compared to HGT cells. To confirm and extend this observation, we compared IFNLR1 and IL10RB

gene expression by quantitative RT-PCR. Remarkably, we found more than 20-fold decrease in

IFNLR1 expression when parental HGT cells were compared to HGT-NV or HGT-Cured cells. In

contrast, no significant difference was observed when IL10RB gene expression was measured (Fig.

3D-E).

Sensing of cytosolic RNA and DNA PAMPs by the innate immune system is functional.

To examine whether HGT-NV and HGT-Cured cells were able to detect and mount an innate immune

response against cytosolic PAMPs, poly (I:C) and poly (dA:dT) were transfected in the various HGT-

derived cell lines and viperin expression levels were measured by RT-qPCR (Fig. 4A-B). Although

the cell lines exhibited upregulation of viperin mRNA to various extents, a significant increase of

viperin expression above basal levels was detected in all the three cell lines in response to both

poly(l:C) and poly(dA:dT). Given that the overall levels of viperin following treatment was

comparable across the three cell lines, evident by the relative ratio with respect to β-actin, we

concluded that PAMP sensing and downstream signalling pathways are functional in these cell lines.

The ability to respond to type III but not type I IFN in HGT-NV and HGT-Cured cells is

debilitated.

We next investigated the ability of type I and type III interferons to elicit ISGs induction. To this end,

cells were incubated with human IFN-β or IFN-λ2 for 16 hours and viperin expression levels were

measured by RT-qPCR. We observed that viperin mRNA was readily induced in all three cell lines in

response to exogenous type I IFN (IFN-β) treatment (Fig. 5A). However, whereas HGT cells

responded to exogenous type III IFN (IFN-λ2) treatment, HGT-NV and HGT-Cured cells did not

(Fig. 5B). To test whether the absence of ISG expression is linked to a defect of STAT1

phosphorylation in response to type I or type III interferons, cells were incubated with either

recombinant IFN-β or IFN-λ2 and STAT1 phosphorylation status was analysed by western blot using

anti-STAT1 phospho-specific antibodies. We observed that although STAT1 was phosphorylated in

all cell lines following IFN-β treatment (Fig. 5C), no STAT1 phosphorylation was detected in HGT-

NV and HGT-Cured cells following IFN-λ2 treatment (Fig. 5D). To ensure that the absence of

STAT1 phosphorylation in HGT-NV cells is not due to a potential clonal effect, different clones as

well as a population of the HGT-NV cells were included in the experiment. Similarly, the clones

represented here by clone 2 (C2) and the polyclonal cell population responded to IFN-β but not to

IFN-λ2 (Fig. 5E-F). Taken together, these results indicate that signal transduction induced by type I

IFN is intact while STAT1 phosphorylation induced in response to lambda interferon is inhibited in

HGT cells that sustained HuNoV replication.

IFNLR1 overexpression rescues STAT1 phosphorylation in response to IFN lambda.

To directly test whether the down-regulation of IFNLR1 expression measured in HGT-NV and Cured

cells is responsible for their unresponsiveness to lambda interferon, cells were transfected with an

IFNLR1-expressing plasmid and stimulated with IFN-λ2 the day after. We observed that IFNLR1

expression rescued STAT1 phosphorylation in response to interferon lambda suggesting that IFNLR1

down-regulation in HGT-NV and HGT-Cured cells is responsible for IFN-λ insensitivity (Fig. 6). In

addition, this experiment shows that the JAK-STAT signalling cascade comprising JAK1, Tyk2 and

possibly JAK2 (Odendall *et al.*, 2014) is functional and leads to the phosphorylation of STAT1.

Inactivation of interferon type I and III receptors increases HuNoV replication in epithelial

cells.

To directly test the influence of type I and type III IFNs on HuNoV replication, HGT cells lines

deficient either for the interferon alpha/beta receptor IFNAR1 or for the interferon lambda receptor

IFNLR1 were generated using CRISPR/Cas9-mediated genome editing. To this end, parental HGT

cells were transduced with lentiviruses expressing IFNAR1 or IFNLR1 single guide RNAs and

clonally selected in the presence of puromycin. Individual clones were screened for gene inactivation

by IFN-β or IFN-λ2 challenging followed by analysis of STAT1 phosphorylation by western-blot and

viperin induction by RT-qPCR (Fig. S2A-B). VPg-linked viral RNA was then transfected and cells

were selected in the presence of 0.5mg/mL G418 for up to 6 days. Relative to day 0 taken as

reference, we observed a significant increase in genomic viral RNA in all cell lines with the exception

of parental HGT cells corroborating the antiviral activity of both type I and type III IFNs on HuNoV

replication (Fig. 7). Similarly, an increased ability of modified cell lines to promote colony formation

induced by the HuNoV replicon was observed (data not shown).

IFNLR1 promoter is methylated in HGT-NV and HGT-Cured cells.

The dimeric interferon lambda receptor consists of the ubiquitously expressed IL10RB chain subunit

and the interferon lambda specific chain IFNLR1 whose expression is limited to cells of epithelial

origin (Sommereyns et al., 2008). Cell-type specific expression of the IFNLR1 subunit was later

shown to be inversely correlated with the methylation of its promoter (Ding et al., 2014). To examine

whether viral replication induced, or lead to the selection of, alterations of IFNLR1 gene expression

through epigenetic modifications, cells were incubated in the presence 5-aza-2'-deoxycytidine

(5azadC), a deoxynucleoside analogue that strongly inhibits DNA methyltransferase (DNMT)

activity. As shown in Fig. 8A, we observed a significant 6-fold increase of IFNLR1 mRNA levels in

HGT-NV cells when compared to cells incubated with the vehicle only. A statistically significant

increase in IFNLR1 gene expression was also observed in the case of HGT-Cured cells but to a lower

extent (2.8-fold). In contrast, no change in IL10RB expression was detected when the same cells were

incubated with 5azadC (Fig. 8B). In addition, incubation in the presence of MS-275 alone (an HDAC

inhibitor) or in combination with 5azadC did not result in increased IFNLR1 expression when

compared to 5azadC alone suggesting that the presence of replicating HuNoV did not modify the

chromatin structure of the IFNLR1 gene (data not shown). Taken together, these results suggest that

the replication of HuNoV in HGT epithelial cells induces a long-term transcriptional silencing of the

IFNLR1 gene through the methylation of its promoter.

Discussion

The purpose of this study was to compare genome-wide transcriptional responses of epithelial cells supporting autonomous HuNoV replication with those elicited by IFN-cured derivatives showing increased viral replication abilities. We found that the interferon lambda receptor is downregulated in cells that sustained HuNoV replication. Mechanistically, this down-regulation of IFNLR1 was mediated by gene methylation as treatment with 5azadC, a DNA methyltransferase inhibitor, rescued the phenotype. This suggests that the lambda IFN pathway is a key constituent of the innate intrinsic defence against human norovirus and shows that an endogenous IFN lambda signalling activity is able to modulate the replication of the virus.

Influence of lambda interferons on murine noroviruses has been highlighted in previous studies using different experimental setups (Baldridge *et al.*, 2015; Baldridge *et al.*, 2017; Rocha-Pereira *et al.*, 2018b). In the case of the human norovirus, recent work using the same replicon in Huh-7 human hepatoma cell line showed that all three types of IFN, when exogenously added, were able to inhibit HuNoV replication leading to virus clearance during long-term treatment (Dang *et al.*, 2018). However, it is not known whether physiological activation of the interferon pathways, particularly, the lambda IFN signalling pathway exerts an antiviral effect against human noroviruses. The present results add further evidence for the involvement of type III IFN in the control of human noroviruses, but extend earlier findings, by showing that physiological levels of type III IFN signalling can effectively restrict HuNoV replication. One of the major concerns with replicon systems is that gene expression landscapes might reflect clonal selection of RNA-replicating cells rather than a universal impact of viral replication on cellular gene expression. To exclude that differences in gene expression were exclusive to a specific cell type or population, we compared our dataset with previous genome wide-transcription profiles from Huh-7 hepatoma cells supporting autonomous replication of HuNoV RNAs (Chang, 2009). We found a (significant) positive correlation between the two datasets when statistically differentially expressed genes were compared (r=0.19; p=0.03; n=113) (or r=0.49; p=0.1; n=12, FC>2), suggesting that analogous cellular responses were induced in response to HuNoV replication in both cell lines.

It is interesting to note that IFNLR1 is also down-regulated in HGT-NV cells which have not been

treated with exogenous type I IFN. This indicates that in the absence of selective pressure mediated by

type I IFN, modulation of the type III IFN pathway is preferentially selected over other genes with

antiviral properties as exemplified in other studies using RNA replicons with selectable marker. For

example, in the case of HCV replicon systems, several antiviral genes including *Viperin* and *MX1*

were shown to be silenced through gene methylation, rendering Huh-7 cells permissive to HCV

replication (Naka *et al.*, 2006; Chen *et al.*, 2013).

An important observation in this study is the upregulation of several IRF3-dependent genes when

HGT-NV cells were compared to parental HGT cells. As IRF3 activation is essential for type I and III

interferon induction, upregulation of IRF3-dependent genes suggests that HuNoV replication

produces PAMPs that are sensed by RIG-I like receptor(s) which in turn activate the cell-intrinsic

immune signalling pathways. Similarly, down-regulation of ISRE-dependent genes in HGT-NV when

compared to HGT-Cured cells, which have silenced the IFNLR1 gene to the same extent, put forward

the idea that HuNoV replication induces IFN-dependent responses. In line with this, a recent study

using the same replicon identified RIG-I and MDA5 proteins as potent negative regulator of HuNoV

replication suggesting that viral RNA secondary structures can readily be detected during replication

(Dang et al., 2018). These observations are however in striking contrast to the study of Qu and

colleagues in which transfection of stool-derived HuNoV RNA readily replicated in 293T cells but

failed to induce detectable interferon responses (Qu *et al.*, 2016). Differences in experimental setups

such as including cell lines, virus strains, stable versus transient viral replication and IFN response

readouts may account for this discordant observation. Investigating whether diverse strains of HuNoV

induce interferon responses in cell-derived human intestinal enteroids will be of interest to probe the

influence of HuNoV replication on IFN induction and responses.

Increased replication of HuNoV in HGT-cured compared to either IFNAR and IFNLR1 knock-out

cells suggests that differentially regulated genes other than IFN receptors may favour virus

replication. However, manual curation of genes differentially regulated between parental and IFN-

cured cells did not reveal individual genes convincingly known to modulate viral replication other

than IFNLR1 and genes involved in the biosynthesis and trafficking of cholesterol. This suggests

either that a low but collective influence of these genes contributes to the increased replication of

HuNoV observed in HGT-Cured cells, or that some of the differentially expressed genes identified

have a potent yet unknown proviral activity towards the human norovirus.

Overall, our results provide insights into the interactions between the human norovirus and innate

cellular responses and show that endogenous levels of λ-IFNs control HuNoV replication suggesting

that they may have a therapeutic potential in the treatment of noroviral infections. In addition, the

high confidence gene expression datasets provided with this study is expected be useful for the

selection and examination of new targets aimed to antiviral therapy.

321
322
Methods

Cells and Media

Human gastric tumour (HGT) cells, human norovirus replicon-harbouring HGT (HGT-NV) cells and

IFN-α interferon-treated HGT-NV cells (HGT-Cured) were maintained in Dulbecco's minimal

essential medium supplemented with 10% fetal calf serum, 2 mM glutamine, 100 U/ml penicillin, 100

µg/ml streptomycin, 1X non-essential amino acid and 0.5 mg/ml G418 in the case of HGT-NV cells.

The DNA methyltransferase inhibitor 5-Aza-2'-deoxycytidine (Sigma-Aldrich, A3656) was diluted in

sterile water and used at a final concentration of 10 µM.

Microarray analysis

Four lineages of HGT, HGT-NV and HGT-Cured cells were grown in tissue culture flasks in the

appropriate culture media. After four successive passages, total RNA was extracted using TRIzol®

(Invitrogen). Microarray analysis was done using the HumanHT-12 v4 Expression BeadChip

(Illumina, Chesterford, UK). All microarray experiments, data normalizations and preliminary

analysis were fulfilled by the Cambridge Genomic Services, UK.

Quantitative RT-PCR analysis

Total cell RNA was extracted using a GenElute Mammalian Total RNA Miniprep kit (Sigma) and

contaminating genomic DNA was removed through RNase-free DNase I treatment (Roche). Total

RNA was then reverse transcribed using random hexamers and the M-MLV RT enzyme (Promega).

SYBR green-based quantitative PCR was performed using gene-specific primers listed in

(Supporting information Table 6). Each experimental condition was measured in biological triplicate

and results are shown as a ratio to levels detected in control cells according to the $\Delta\Delta$Ct method

(Schmittgen and Livak, 2008). Additional non-template and non-reverse transcriptase samples were

analysed as negative controls. Data were collected using a ViiA 7 Real-Time PCR System (Applied

Biosystems). Genomic viral RNA was quantified by one step RT-qPCR using GI NV-specific

primers. Viral genome copy numbers were calculated by interpolation from a standard curve

generated using serial dilutions of viral RNA transcribed from the pNV101 plasmid coding for the

full-length Norwalk genome (Fernandez-vega *et al.*, 2004).

Nucleic acid transfection and IFN treatment

Transfections of plasmid DNA or purified viral RNA (NV replicon) were carried out using

Lipofectamine 2000 (Invitrogen) according to manufacturer's instructions. Briefly, cells were seeded

in antibiotic-free growth medium at a density of 2×10^5 or 1×10^6 cells per well in 24- or 6-well plates,

respectively, and incubated overnight at 37°C. Lipofectamine 2000 reagent was diluted in Opti-

MEM® (Gibco) and incubated for 5 minutes at 25°C. After the incubation, plasmid DNA or viral

RNA, diluted in Opti-MEM® (Gibco), was mixed with the Lipofectamine: Opti-MEM mixture,

vortexed briefly and incubated at 25°C for 20 minutes. The DNA or RNA complex was subsequently

inoculated onto 80-90 % confluent cell monolayers, followed by incubation at 37°C. For viral RNA

transfection, media was replaced after 24 h with fresh complete growth medium containing G418 at a

concentration of 0.5 mg/ml. For IFN treatments, cells were seeded as described above and incubated

at 37°C for indicated time-points with or without recombinant type I IFN (IFN-β; Peprotech, Cat

N°:300-02BC), or type III IFN (IFN-λ2; Peprotech, Cat N°:300-02K) at a final concentration of 0.1

ug/ml.

Immunofluorescence microscopy

Cells were plated on 12 mm glass coverslips and allowed to adhere overnight before fixation with 4%

paraformaldehyde in PBS. Cells were then permeabilized for 5 min with PBS-Triton X-100 0.2% and

unspecific antigens were blocked for 1h using 2% normal goat serum (Sigma-S2007) in PBS-Tween-

20 0.1% (PBST). Cells were then incubated for 1h with primary mouse monoclonal J2 anti-dsRNA

antibodies in PBST at a dilution of 1:1000 (J2, SCICONS English & Scientific Consulting, Hungary).

After extensive washes with PBST, species-matched AlexaFluor-conjugated secondary antibodies

(ThermoFisher Scientific, A-11029) were added at a dilution of 1:500 in PBST for one additional

hour. Coverslips were extensively washed and mounted on slides with Mowiol supplemented with

DAPI and DABCO. Confocal micrographs were acquired on a Leica TCS SP5 confocal fitted with a

63x 1.3NA oil immersion objective using 405nm and 488nm laser excitation lines under sequential

channel scanning to prevent fluorophore bleed-through artefacts due to spectral overlap.

Western blot analysis

Cell lysates were prepared in radio-immuno precipitation assay buffer (RIPA: 150 mM NaCl, 0.5%

sodium deoxycholate, 0.1% SDS, 1mM EDTA, 1% Triton X-100, and 50 mM Tris pH8)

supplemented with protease and phosphatase inhibitors (Calbiochem; Cat N°: 539134 and 524625).

Protein concentrations were determined by BCA assay (Thermo Fisher Scientific). Equal amounts of

total proteins were resolved by SDS-PAGE and transferred to nitrocellulose membranes. Blocking of

unspecific antigens was carried out in 5% non-fat dried milk or 5% BSA in PBST for 1 h at 4°C.

Primary antibodies were diluted in blocking buffer and incubated overnight at 4°C with gentle rocking

(Supporting information Table 7). Membranes were washed three times in PBST for 5 min at RT.

Species-matched IRDye-800CW secondary antibodies were diluted in blocking buffer as before and

incubated at RT for 1 h. Membranes were washed again three times in PBST for 5 min at RT.

Fluorescent signal was detected through an Odyssey CLx infrared imaging system (Li-COR).

Generation of IFNAR1 or IFNLR1 knockout HGT cells

HGT cells knockout for IFNAR1 or IFNLR1 genes were generated using the CRISPR/Cas9 system.

Lentivirus vectors encoding single-guide RNAs against IFNAR1 or IFNLR1 were generously

provided by Dr Steeve Boulant and are described in Pervolaraki et al. (Pervolaraki *et al.*, 2017).

Vesicular stomatitis virus G-protein-pseudotyped lentiviral particles were generated by transient

transfection of 293T cells grown in 6-well plates using 1.25 µg lentiviral vector, 0.63 µg

pMDLg/pRRE (Addgene #12251), 0.31 µg pRSV-Rev (Addgene #12253) and 0.38 µg pMD2.G

(Addgene #12259) per well. Parental HGT cells were transduced with lentiviral supernatants and

incubated for 48 h. Transduced cells were then selected on the basis of their resistance to puromycin

at a concentration of 2.5 µg/mL. Clonal isolation was performed by limiting dilution into 96-well

plate at a density of 0.3 cell per well and single cell clones were selected on the basis of visual

examination. Single cell clones were expanded and tested for IFNAR1 or IFNLR1 gene disruption by

RT-qPCR measurement of viperin induction following incubation with IFN-β or IFN-λ2,

respectively. Absence of STAT1 phosphorylation following incubation with receptor-matched

interferons confirmed the gene ablation.

Statistical analysis and data sharing

Statistical significance was determined from experiments where n $\geq$ 3 using two-tailed Student *t* tests

in Prism 6.0 (GraphPad). All microarray expression data reported in this study have been deposited

into Gene Expression Omnibus (GEO, http://www.ncbi.nlm.nih.gov/geo) with the accession number

GSE111041.

<u>**Figure legends**</u>

412 **Figure 1.** Generation and characterization of stable HuNoV replicons in HGT and U2OS cell lines.

413 A. Diagram showing the steps used to generate the various cell lines carrying human norovirus

414 replicons and their IFN-cured counterparts.

415 B. Total cellular RNA was extracted from each cell line and viral RNA quantified by RT-qPCR. Viral

416 RNA copy numbers were normalized to the total input RNA and expressed as genome equivalents/μg

417 of input RNA. The error bars represent the standard error of the mean determined from four biological

418 replicates. Viral RNA copy numbers were below the limit of detection in both HGT and HGT-Cured

419 cells. The dotted line represents the low limit of detection.

420 C. Detection of viral replication complexes by confocal imaging. Representative merged confocal

421 micrographs showing the detection of dsRNA (green) in indicated cell lines. Nuclei were stained with

422 DAPI (blue). Scale bars correspond to 20μm.

423 **Figure 2.** HGT-Cured cells demonstrate enhanced HuNoV replication.

424 A. HGT and HGT-Cured cells were transfected with VPg-linked replicon RNA. After five days of

425 selection with G418 at a concentration of 0.5 mg/mL, cellular morphology was analysed by light

426 microscopy.

427 B. Cells transfected with VPg-linked RNA were harvested at various time points post-transfection for

428 total RNA extraction and viral RNA was quantified by RT-qPCR. Viral RNA levels were determined

429 by comparison to a standard curve and normalized to the RNA input. The data are presented as mean

430 and standard error of the mean from three replicates. Indicated values are expressed as fold change in

431 genome equivalent normalized to viral RNA levels at 6 hours post-transfection to control for

432 transfection efficiency. Unpaired two-tailed Student's t-test was used to evaluate the statistical

433 significance.

435 **Figure 3.** Microarray analysis and validation of differentially regulated genes between HGT-NV,

436 HGT-Cured and parental HGT cells.

A. Volcano plots of differentially expressed genes from microarray analysis comparing gene

expression in HGT-NV or HGT-Cured compared to parental HGT cells. Significantly up- or down-

regulated genes (FDR<0.01 and |log$_2$ fold change| ≥ 1) are represented in red or green, respectively.

B. Fold change correlation between microarray analysis and quantitative real-time PCR (RT-qPCR).

Scatter plots comparing log$_2$ fold changes of selected genes measured by microarray analysis and RT-

qPCR in HGT-NV or HGT-Cured compared to parental HGT cells. Error bars represent the standard

deviation of biological quadruplicate experiments analyzed in triplicate reactions. The Pearson

correlation coefficient (r) and the number of pairs analysed (n) are indicated on each graph. Dotted

lines illustrate the 95% confidence interval of the linear regression.

C. Venn diagrams representing the overlap of significantly up- and down-regulated genes between

HGT-NV or HGT-Cured cells compared to parental HGT cells. The top ten genes within each

category are shown.

D-E. IFNLR1 (D) but not IL10RB (E) gene expression is down-regulated in HGT-NV and HGT-

Cured cells. Changes in gene expression of IFNLR1 and IL10RB mRNA levels were normalized to β-

actin levels and calculated using the $\Delta\Delta C_T$ method. Relative expression was determined from one

experiment performed in biological triplicate and compared to the expression levels measured in HGT

control. Statistical significance was determined using the unpaired t test. Error bars represent the

standard deviation between biological replicates.

Figure 4. Innate immune responses to RNA and DNA PAMPs are functional.

Cells were transfected with poly I:C (A) or poly dA:dT (B) and total RNA was harvested 8h post-

transfection for quantification of viperin mRNA levels by RT-qPCR. Viperin mRNA levels were

expressed as differences of cycle threshold (C$_t$) of viperin, relative to the C$_t$ of beta-actin. The error

bars represent standard error of the mean, determined from the result of three replicates.

Figure 5. The ability to respond to type III but not type I IFN is debilitated in HGT-NV and HGT-

Cured cells.

Cells were treated with type I (IFN-β) (A) or type III interferon (IFN-λ2) (B), incubated overnight and

harvested for total RNA extraction and quantification of viperin mRNA levels by RT-qPCR. Viperin

mRNA expression levels were expressed as differences of cycle threshold (C_t) of viperin, relative to

the C_t of beta-actin. The error bars represent the standard error of the mean, determined from the

result of three replicates.

(C-F) Western-blot analysis showing STAT1 phosphorylation after stimulation with type I and type

III interferons in HGT, HGT-NV and HGT-Cured cell lines (C, E). HGT-NV Clone 2 (C2) and

population of cells (pop) were included in the experiment (D, F). GAPDH was used as loading

control.

Figure 6. Overexpression of IFNLR1 in HGT-NV and HGT-Cured cells rescues the ability to respond

to type III IFN and the loss of the receptor enhances viral RNA replication in HGT cells.

HGT, HGT-NV and HGT-Cured were transfected with IFNLR1 expressing plasmid and STAT1

phosphorylation was measured in whole cell lysate by western blot after incubating cells with type III

IFN for 15 min (A). STAT1 was used as a positive control and GAPDH was used as the loading

control. HGT IFNLR1$^{-/-}$ cells transfected with GI viral VPg-linked RNA were harvested at 0 and 5

days post-transfection for total RNA extraction and viral RNA was quantified by RT-qPCR (B). The

viral RNA levels were determined by comparing to a standard curve and normalized to the total input

RNA. The data are presented as the mean and standard error of the mean from at least 11 replicates

and are expressed as fold change in genome equivalent at day 0.

Figure 7. Genetic ablation of IFNAR1 and IFNLR1 interferon receptors promotes HuNoV

replication.

Parental HGT, IFNAR1 and IFNLR1 knock-out cells were transfected with VPg-linked replicon RNA

and submitted to G418 selection. Total cellular RNA was harvested at various time points post-

transfection and viral RNA was quantified by RT-qPCR. Relative viral replication was determined

from one experiment performed in biological triplicate and compared to the replication levels

measured in HGT control at day 0. Statistical significance was determined using the unpaired *t* test.

Error bars represent the standard deviation between biological replicates.

Figure 8. IFNLR1 promoter is methylated during long-term HuNoV replication.

HGT, HGT-NV and HGT-Cured cells were incubated in the presence of 10 uM 5azadC for 72 hours

or left untreated. Total cell RNA was extracted and IFNLR1 and IL10RB expression was measured by

RT-qPCR. Total cellular RNA was harvested 72 hours post-treatment and IFNLR1 and IL10RB

mRNA were quantified by RT-qPCR. Relative gene expression was determined from one experiment

performed in biological triplicate and compared to the expression levels measured in corresponding

untreated cells. Statistical significance was determined using the unpaired t test. Error bars represent

the standard deviation between biological replicates.

Supplementary figure legends

Figure S1. Expression change validation of selected genes at the transcription and translation

levels.

A. Direct comparison between microarray signal intensities and difference in cycle thresholds for

ANXA1 and *IFITM*3 genes.

B. Validation by western-blot analysis of ANXA1 and IFITM3 protein levels in HGT, HGT-NV and

HGT-Cured cell lines. The indicated ratios represent the ANXA1 or IFITM3 protein levels

normalized to the corresponding GAPDH loading controls relative to parental HGT cells.

Figure S2. Western blot validation of CRISPR/Cas9-induced knockout cells.

A-B. HGT wild-type cells as well as IFNAR1 (A) and IFNLR1 (B) CRISPR/Cas9 clones were treated

with type I (IFN-β) (A) or type III interferon (IFN-λ2) (B) for 15 min and STAT1 phosphorylation

(pT701) was measured by immunoblot analysis. GAPDH was used as a loading control.

Funding information

I.G. is a Wellcome Senior Fellow and this work was supported by funding from the Wellcome

Trust (Ref: 207498/Z/17/Z). F.S. was funded by a Biotechnology and Biological Sciences Research

Council (BBSRC) sLoLa grant (BB/K002465/1). S.A.E. is supported through a Cambridge Trust

Cambridge-Africa PhD studentship.

522
523   Myra?

524   Sabastine?

525   **Author contributions**
526
527   S.A.E., F.S., M.H., and I.G. designed, performed the research and analysed the data. S.A.E., F.S., and
528   I.G. wrote the manuscript.
529
530

Acknowledgements

We thank Laure Dumoutier (de Duve Institute, Universite catholique de Louvain) for the gift of

the IFNLR1 (LICR2) expressing plasmid. We thank K Green (NIH, Bethesda) and KO Chang for

the provision of pNV-Neo.

Conflicts of interest

The authors declare that they have no conflict of interest.

References

Andersen, J., VanScoy, S., Cheng, T.F., Gomez, D., and Reich, N.C. (2008) IRF-3-dependent and
augmented target genes during viral infection. *Genes Immun* **9**: 168–175.
543
544   Ank, N., Iversen, M.B., Bartholdy, C., Staeheli, P., Hartmann, R., Jensen, U.B., *et al.* (2008) An
545   Important Role for Type III Interferon (IFN- /IL-28) in TLR-Induced Antiviral Activity. *J*
546   *Immunol* **180**: 2474–2485 http://www.jimmunol.org/cgi/doi/10.4049/jimmunol.180.4.2474.
547
548   Baldridge, M.T., Lee, S., Brown, J.J., McAllister, N., Urbanek, K., Dermody, T.S., *et al.* (2017)
549   Expression of *Ifnlr1* on Intestinal Epithelial Cells Is Critical to the Antiviral Effects of Interferon
550   Lambda against Norovirus and Reovirus. *J Virol* **91**: e02079-16
551   http://jvi.asm.org/lookup/doi/10.1128/JVI.02079-16.
552
553   Baldridge, M.T., Nice, T.J., McCune, B.T., Yokoyama, C.C., Kambal, A., Wheadon, M., *et al.* (2014)
554   Commensal microbes and interferon-lambda determine persistence of enteric murine norovirus
555   infection_supplemental data. *Science* **3**: 1–9 http://www.ncbi.nlm.nih.gov/pubmed/25431490.
556
557   Baldridge, M.T., Nice, T.J., McCune, B.T., Yokoyama, C.C., Kambal, A., Wheadon, M., *et al.* (2015)
558   Commensal microbes and interferon-λ determine persistence of enteric murine norovirus
559   infection. *Science (80-)* **347**: 266–269.
560
561   Bartsch, S.M., Lopman, B.A., Ozawa, S., Hall, A.J., and Lee, B.Y. (2016) Global economic burden of
562   norovirus gastroenteritis. *PLoS One* **11**: e0151219
563   https://doi.org/10.1371/journal.pone.0151219.
564
565   Belliot, G., Lopman, B.A., Ambert-Balay, K., and Pothier, P. (2014) The burden of norovirus
566   gastroenteritis: An important foodborne and healthcare-related infection. *Clin Microbiol Infect*
567   **20**: 724–730.
568
569   Blazek, K., Eames, H.L., Weiss, M., Byrne, A.J., Perocheau, D., Pease, J.E., *et al.* (2015) IFN-λ
570   resolves inflammation via suppression of neutrophil infiltration and IL-1β production. *J Exp Med*
571   **212**: 845–853 http://www.jem.org/lookup/doi/10.1084/jem.20140995.
572
573   Chang, K.-O. (2009) Role of Cholesterol Pathways in Norovirus Replication. *J Virol* **83**: 8587–
574   8595 http://jvi.asm.org/cgi/doi/10.1128/JVI.00005-09.
575
576   Chang, K.-O., Sosnovtsev, S. V., Belliot, G., King, A.D., and Green, K.Y. (2006) Stable expression of a
577   Norwalk virus RNA replicon in a human hepatoma cell line. *Virology* **353**: 463–473
578   http://linkinghub.elsevier.com/retrieve/pii/S0042682206003965.
579
580   Chen, Q., Denard, B., Huang, H., and Ye, J. (2013) Epigenetic Silencing of Antiviral Genes Renders
581   Clones of Huh-7 Cells Permissive for Hepatitis C Virus Replication. *J Virol* **87**: 659–665
582   http://jvi.asm.org/cgi/doi/10.1128/JVI.01984-12.
583
584   Dang, W., Xu, L., Yin, Y., Chen, S., Wang, W., Hakim, M.S., *et al.* (2018) IRF-1, RIG-I and MDA5
585   display potent antiviral activities against norovirus coordinately induced by different types of
586   interferons. *Antiviral Res* **155**: 48–59.
587
588   Dickensheets, H., Sheikh, F., Park, O., Gao, B., and Donnelly, R.P. (2013) Interferon-lambda (IFN-)
589   induces signal transduction and gene expression in human hepatocytes, but not in lymphocytes
590   or monocytes. *J Leukoc Biol* **93**: 377–385
591   http://www.jleukbio.org/cgi/doi/10.1189/jlb.0812395.
592

593 Ding, S., Khoury-Hanold, W., Iwasaki, A., and Robek, M.D. (2014) Epigenetic Reprogramming of
594 the Type III Interferon Response Potentiates Antiviral Activity and Suppresses Tumor Growth.
595 *PLoS Biol* **12**.
596
597 Ettayebi, K., Crawford, S.E., Murakami, K., Broughman, J.R., Karandikar, U., Tenge, V.R., *et al.*
598 (2016) Replication of human noroviruses in stem cell-derived human enteroids. *Science (80-)*
599 **353**: 1387–1393.
600
601 Fankhauser, R.L., Noel, J.S., Monroe, S.S., Ando, T., and Glass, R.I. (1998) Molecular Epidemiology
602 of "Norwalk-like Viruses" in Outbreaks of Gastroenteritis in the United States. *J Infect Dis* **178**:
603 1571–1578 https://academic.oup.com/jid/article-lookup/doi/10.1086/314525.
604
605 Fernandez-vega, V., Sosnovtsev, S. V, King, A.D., Mitra, T., Gorbalenya, A., and Green, K.Y. (2004)
606 Norwalk Virus N-Terminal Nonstructural Protein Is Associated with Disassembly of the Golgi
607 Complex in Transfected Cells. *J Virol* **78**: 4827–4837.
608
609 Friedman, R.L., Manly, S.P., McMahon, M., Kerr, I.M., and Stark, G.R. (1984) Transcriptional and
610 posttranscriptional regulation of interferon-induced gene expression in human cells. *Cell* **38**:
611 745–755 http://www.ncbi.nlm.nih.gov/pubmed/6548414.
612
613 Grandvaux, N., Servant, M.J., tenOever, B., Sen, G.C., Balachandran, S., Barber, G.N., *et al.* (2002)
614 Transcriptional Profiling of Interferon Regulatory Factor 3 Target Genes: Direct Involvement in
615 the Regulation of Interferon-Stimulated Genes. *J Virol* **76**: 5532–5539
616 http://jvi.asm.org/cgi/doi/10.1128/JVI.76.11.5532-5539.2002.
617
618 Hall, A.J., Lopman, B.A., Payne, D.C., Patel, M.M., Gastañaduy, P.A., Vinjé, J., and Parashar, U.D.
619 (2013) Norovirus disease in the united states. *Emerg Infect Dis* **19**: 1198–1205.
620 Hemann, E.A., Gale, M., and Savan, R. (2017) Interferon lambda genetics and biology in
621 regulation of viral control. *Front Immunol* **8**.
622
623 Isaacs, A., and Lindenmann, J. (1988) Virus Interference: I. The Interferon. *CA Cancer J Clin* **38**:
624 280–290 http://doi.wiley.com/10.3322/canjclin.38.5.280.
625
626 Jones, M., Watanabe, M., Zhu, S., Graves, C., Keyes, L., Grau, K., *et al.* (2014) Enteric bacteria
627 promote human and mouse norovirus infection of B cells. *Science (80-)* **346**: 755–759.
628
629 Kotenko, S. V., Gallagher, G., Baurin, V. V., Lewis-Antes, A., Shen, M., Shah, N.K., *et al.* (2003) IFN-
630 λs mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat*
631 *Immunol* **4**: 69–77.
632
633 Lee, S., and Baldridge, M.T. (2017) Interferon-lambda: A potent regulator of intestinal viral
634 infections. *Front Immunol* **8**.
635
636 Lopman, B., Ahmed, S.M., Robinson, A., Verhoef, L., Koopmans, M., and Hall, A.J. (2013) The
637 global prevalence of norovirus among cases of gastroenteritis. *Bangkok, Thail Vaccines Enteritis*
638 *Dis* .
639
640 Ludwig, A., Adams, O., Laws, H.J., Schroten, H., and Tenenbaum, T. (2008) Quantitative detection
641 of norovirus excretion in pediatric patients with cancer and prolonged gastroenteritis and
642 shedding of norovirus. *J Med Virol* **80**: 1461–1467.
643
644 Mawatari, M., and Kato, Y. (2014) Norovirus Gastroenteritis. *Emerg Infect Dis Clin Case Stud* **361**:
645 203–212.
646

647 Mordstein, M., Neugebauer, E., Ditt, V., Jessen, B., Rieger, T., Falcone, V., *et al.* (2010) Lambda
648 Interferon Renders Epithelial Cells of the Respiratory and Gastrointestinal Tracts Resistant to
649 Viral Infections. *J Virol* **84**: 5670–5677 http://jvi.asm.org/cgi/doi/10.1128/JVI.00272-10.
650
651 Naka, K., Abe, K. ichi, Takemoto, K., Dansako, H., Ikeda, M., Shimotohno, K., and Kato, N. (2006)
652 Epigenetic silencing of interferon-inducible genes is implicated in interferon resistance of
653 hepatitis C virus replicon-harboring cells. *J Hepatol* **44**: 869–878.
654
655 Nice, T.J., Baldridge, M.T., McCune, B.T., Norman, J.M., Lazear, H.M., Artyomov, M., *et al.* (2015)
656 Interferon-λ cures persistent murine norovirus infection in the absence of adaptive immunity.
657 *Science (80-)* **347**: 269–273.
658
659 Odendall, C., Dixit, E., Stavru, F., Bierne, H., Franz, K.M., Durbin, A.F., *et al.* (2014) Diverse
660 intracellular pathogens activate type III interferon expression from peroxisomes. *Nat Immunol*
661 **15**: 717–726.
662
663 Patel, M.M., Widdowson, M.A., Glass, R.I., Akazawa, K., Vinjé, J., and Parashar, U.D. (2008)
664 Systematic literature review of role of noroviruses in sporadic gastroenteritis. *Emerg Infect Dis*
665 **14**: 1224–1231.
666
667 Payne, D.C., Vinjé, J., Szilagyi, P.G., Edwards, K.M., Staat, M.A., Weinberg, G.A., *et al.* (2013)
668 Norovirus and Medically Attended Gastroenteritis in U.S. Children. *N Engl J Med* **368**: 1121–
669 1130 http://www.nejm.org/doi/10.1056/NEJMsa1206589.
670
671 Pervolaraki, K., Stanifer, M.L., Münchau, S., Renn, L.A., Albrecht, D., Kurzhals, S., *et al.* (2017)
672 Type I and type III interferons display different dependency on mitogen-activated protein
673 kinases to mount an antiviral state in the human gut. *Front Immunol* **8**.
674
675 Phillips, G., Tam, C.C., Conti, S., Rodrigues, L.C., Brown, D., Iturriza-Gomara, M., *et al.* (2010)
676 Community incidence of norovirus-associated infectious intestinal disease in England:
677 Improved estimates using viral load for norovirus diagnosis. *Am J Epidemiol* **171**: 1014–1022.
678
679 Qu, L., Murakami, K., Broughman, J.R., Lay, M.K., Guix, S., Tenge, V.R., *et al.* (2016) Replication of
680 Human Norovirus RNA in Mammalian Cells Reveals Lack of Interferon Response. *J Virol* **90**:
681 8906–8923 http://jvi.asm.org/lookup/doi/10.1128/JVI.01425-16.
682
683 Reid, L.E., Brasnett, A.H., Gilbert, C.S., Porter, A.C., Gewert, D.R., Stark, G.R., and Kerr, I.M. (1989)
684 A single DNA response element can confer inducibility by both alpha- and gamma-interferons.
685 *Proc Natl Acad Sci U S A* **86**: 840–4
686 http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=286573&tool=pmcentrez&render
687 type=abstract.
688
689 Rocha-Pereira, J., Jacobs, S., Noppen, S., Verbeken, E., Michiels, T., and Neyts, J. (2018a)
690 Interferon lambda (IFN-λ) efficiently blocks norovirus transmission in a mouse model. *Antiviral*
691 *Res* **149**: 7–15 https://doi.org/10.1016/j.antiviral.2017.10.017.
692
693 Rocha-Pereira, J., Jacobs, S., Noppen, S., Verbeken, E., Michiels, T., and Neyts, J. (2018b)
694 Interferon lambda (IFN-λ) efficiently blocks norovirus transmission in a mouse model. *Antiviral*
695 *Res* **149**: 7–15.
696
697 Schmittgen, T.D., and Livak, K.J. (2008) Analyzing real-time PCR data by the comparative CT
698 method. *Nat Protoc* **3**: 1101–1108 http://www.nature.com/doifinder/10.1038/nprot.2008.73.
699
700 Sheppard, P., Kindsvogel, W., Xu, W., Henderson, K., Schlutsmeyer, S., Whitmore, T.E., *et al.*

701   (2003) IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* **4**: 63–68.
702
703   Sommereyns, C., Paul, S., Staeheli, P., and Michiels, T. (2008) IFN-lambda (IFN-λ) is expressed in
704   a tissue-dependent fashion and primarily acts on epithelial cells in vivo. *PLoS Pathog* **4**.
705
706   Son, K.-N., Liang, Z., and Lipton, H.L. (2015) Double-Stranded RNA Is Detected by
707   Immunofluorescence Analysis in RNA and DNA Virus Infections, Including Those by Negative-
708   Stranded RNA Viruses. *J Virol* **89**: 9383–9392
709   http://jvi.asm.org/lookup/doi/10.1128/JVI.01299-15.
710
711   Syedbasha, M., and Egli, A. (2017) Interferon Lambda: Modulating immunity in infectious
712   diseases. *Front Immunol* **8**.
713
714   Thorne, L.G., and Goodfellow, I.G. (2014) Norovirus gene expression and replication. *J Gen Virol*
715   **95**: 278–291.
716
717   Weber, F., Wagner, V., Rasmussen, S.B., Hartmann, R., and Paludan, S.R. (2006) Double-Stranded
718   RNA Is Produced by Positive-Strand RNA Viruses and DNA Viruses but Not in Detectable
719   Amounts by Negative-Strand RNA Viruses. *J Virol* **80**: 5059–5064
720   http://jvi.asm.org/cgi/doi/10.1128/JVI.80.10.5059-5064.2006.
721
722   Weerd, N.A. De, Samarajiwa, S.A., and Hertzog, P.J. (2007) Type I interferon receptors:
723   Biochemistry and biological functions. *J Biol Chem* **282**: 20053–20057.
724
725   Witte, K., Gruetz, G., Volk, H.D., Looman, A.C., Asadullah, K., Sterry, W., *et al.* (2009) Despite IFN-
726   receptor expression, blood immune cells, but not keratinocytes or melanocytes, have an
727   impaired response to type III interferons: Implications for therapeutic applications of these
728   cytokines. *Genes Immun* **10**: 702–714.
729
730   Wobus, C.E., Karst, S.M., Thackray, L.B., Chang, K.O., Sosnovtsev, S. V., Belliot, G., *et al.* (2004)
731   Replication of Norovirus in cell culture reveals a tropism for dendritic cells and macrophages.
732   *PLoS Biol* **2**: e432.
733
734

Fig. 1

A

B

C

HGT   HGT-NV   HGT-Cured

Fig. 2

Fig. 3

Fig. 4

Fig. 5

A

Viperin mRNA expression normalized to β-actin (ΔC_t)

878.0x ****

147.3x ****

256.7x ****

IFN-β

HGT | HGT NV | HGT Cured

B

Viperin mRNA expression normalized to β-actin (ΔC_t)

19.8x ***

0.9x ns

1.0x ns

IFN-λ2

HGT | HGT NV | HGT Cured

C

Mock | IFN-β (15 min) | IFN-β (30 min)

HGT HGT NV HGT Cured | HGT HGT NV HGT Cured | HGT HGT NV HGT Cured | kDa

STAT1 (pT701) — 75

STAT1 — 75

GAPDH — 37

D

Mock | IFN-λ2 (15 min) | IFN-λ2 (30 min)

HGT HGT NV HGT Cured | HGT HGT NV HGT Cured | HGT HGT NV HGT Cured | kDa

STAT1 (pT701) — 75

STAT1 — 75

GAPDH — 37

E

Mock | IFN-β (15 min)

HGT HGT-NV HGT Cured | HGT HGT-NV HGT Cured
C2 Pop | C2 Pop | kDa

STAT1 (pT701) — 75

STAT1 — 75

GAPDH — 37

F

Mock | IFN-λ2 (15 min)

HGT HGT-NV HGT Cured | HGT HGT-NV HGT Cured
C2 Pop | C2 Pop | kDa

STAT1 (pT701) — 75

STAT1 — 75

GAPDH — 37

Fig. 6

STAT1 (pT701)

STAT1

GAPDH

Empty vector — HGT, HGT NV, HGT Cured

IFNLR1 — HGT, HGT NV, HGT Cured

kDa

IFN-λ2

Fig. 7

Fig. 8

A. IFNLR1 — Fold change: HGT (ns), HGT NV (***), HGT Cured (**)

B. IL10RB — Fold change: HGT (ns), HGT NV (ns), HGT Cured (ns)