The symmetrical pattern of base-pair substitutions rates across the chromosome in

***Escherichia coli* has multiple causes**

Brittany A. Niccum[1], Heewook Lee[2], Wazim MohammedIsmail[3], Haixu Tang[3], and Patricia L. Foster[1]*

[1]Department of Biology, Indiana University, Bloomington, IN, USA, 47405

[2]School of Computer Science, Computational Biology Department, Carnegie-Mellon University

[3]School of Informatics and Computing, Indiana University, Bloomington, IN, USA, 47405

*Corresponding Author:

Patricia L. Foster

Department of Biology

Jordan Hall 447E

1001 East Third Street

Indiana University

Bloomington, IN 47405

812-855-4084

plfoster@indiana.edu

Author email addresses:

Patricia L. Foster, plfoster@indiana.edu

Brittany A. Niccum, brittanyniccum@gmail.com

Heewook Lee, heewookl@cs.cmu.edu

Wazim MohammedIsmail, wazimoha@indiana.edu

Haixu Tang, hatang@indiana.edu

Abstract

Background: Mutation accumulation experiments followed by whole-genome sequencing have

revealed that for several bacterial species, the rate of base-pair substitutions (BPSs) varies across the

chromosome in a wave-like pattern that is symmetrical about the origin of replication. The

experiments presented here investigate the causes of this pattern in *Escherichia coli*.

Results: The timing of replication initiation, the progression of the replication fork, and the structure of

the region where replication terminates all affect the mutational density pattern. As predicted

previously, the activities of the nucleoid binding proteins, HU and Fis, are also important, suggesting

that mutation rates increase when highly structured DNA regions are replicated. Biases in error-

correction by proofreading and mismatch repair are major factors in establishing the pattern. A

mutation accumulation experiment with mismatch-repair defective *Bacillus subtilis* revealed a similar

symmetrical pattern of BPS rates.

Conclusions: The symmetrical pattern of BPS rates does not have a single cause, but is, instead

determined by the initiation, progression, and termination of DNA replication, error-correction, and

chromosome structure. These factors should apply to most bacterial, and possibly eukaryotic,

genomes, and imply that different areas of the genome evolve at different rates.

Keywords *Escherichia coli*, mutation rate, mutation accumulation, whole genome sequencing,

mismatch repair, replication timing, nucleoid associated proteins, proofreading

Background

Recent studies of mutations accumulated non-selectively across bacterial chromosomes have revealed that rates of base-pair substitutions (BPSs) vary 2- to 4- fold in a wave like pattern that is mirrored in the two independently replicating halves of the chromosome. Such symmetrical patterns have been observed in mismatch-repair (MMR) defective strains of *Escherichia coli* (1), *Vibrio fischeri*, *V. cholerae* (2-4), *Pseudomonas fluorescens* (5), and *P. aeruginosa* (6). Such variation in mutation rates may affect the rate at which genes in different regions of the chromosome evolve, and may exert selective pressure on gene placement. Yet the causes of this variation are not known.

The fidelity of DNA replication, which, in *E. coli*, is about 1 mistake in 1000 generations (7), is determined by the intrinsic accuracy of the DNA polymerase and error correction by proofreading and mismatch repair (reviewed in (8, 9). In *E. coli*, proofreading is performed by epsilon, a subunit of the DNA polymerase III holoenzyme. If the polymerase inserts the incorrect base, epsilon's 5' to 3' exonuclease activity degrades a few bases of the new strand and polymerase then re-synthesizes it. The accuracy of DNA synthesis is improved about 4000-fold by proofreading (10). Mismatch repair is performed by three proteins, MutS, MutL, and MutH. MutS recognizes a mismatch and recruits MutL. Together they find a nearby GATC site in which, in *E. coli*, the A is methylated by the Dam methylase. Because methylation lags behind replication, unmethylated As identify the "new", and presumably error-containing, DNA strand. MutH is recruited by MutSL to the GATC site and activated to nick the unmethylated DNA strand, which is then degraded past the mismatch by the concerted activity of the UvrD helicase and one of four exonucleases. Pol III then re-synthesizes the strand. MMR improves the accuracy of DNA replication 100 to 200 fold (11).

In our previous study of the BPS density pattern in MMR-defective *E. coli* (1), we correlated

mutation rates to the chromosomal sites that are affected by two nucleoid-associated proteins (NAPs),

HU and Fis. We suggested that when the replication fork encounters regions of the chromosome with

high superhelical density due to the binding of these NAPs, the mutation rate increases. An alternative

explanation ties mutation rates to replication timing (2, 3). An intriguing hypothesis is that mutation

rates vary in concert with fluctuations in dNTP concentration when the replication origin fires

repeatedly during rapid growth (2).

In the work presented here, we investigate further the causes of the wave-like pattern of BPS

rates. MMR-defective strains of *E. coli* additionally defective for other activities were used in mutation

accumulation (MA) experiments and the mutations identified by whole genome sequencing (WGS). We

also investigated the effects of different growth conditions on BPS rates. We conclude that the BPS

density pattern does not have a single cause, but is the result of several factors affecting DNA

replication, repair, and chromosome structure. In addition, we report that a MMR-defective *Bacillus*

subtilis also has a wave-like BPS density pattern that is symmetrical about the origin of replication.

Results

The base-pair substitution (BPS) density pattern in mismatch repair-defective strains

In a previous paper (1) we demonstrated that the density of BPS that accumulated across the

chromosome in a MMR-defective strain during an MA experiment fell into a wave-like pattern that was

symmetrical about the origin. We have since performed nine additional MA experiments with MMR

mutant strains, each experiment resulting in the accumulation of over 1000 BPSs, for a total of 30,061

BPSs (11). As shown in Fig. 1A, the wave-like BPS density pattern was reproducible among the 10

experiments. Also as shown in Fig. 1A, the mutational density pattern does not exactly match the non-

interacting chromosomal macrodomains (MCs) as defined by Valens et al, 2004 (12). In particular, the

wave pattern of BPSs is symmetrical about the origin whereas the Ori MC is not.

Following the analysis of Dillon et al, 2018 (2), we computed the wavelet coherence of the

collected MMR⁻ data taken in the clockwise and the counterclockwise directions around the

chromosome; Additional file 1: Fig. S1A shows that, except for some asymmetry at the midpoint, the

wave is symmetrical across the chromosome. The coherence is greatest between 800 to 1,600 Kb per

cycle (= 8 to 16 bins per cycle), which is a similar result to that found by Dillon et al, 2018 (2). We also

computed the wavelet coherence of the collected MMR⁻ data against each of the experimental results

discussed below, and these graphs are also given in the Additional file material. In the analyses below,

the combined data from the 10 experiments with MMR-defective strains are used as the standard to

which the results obtained in other genetic backgrounds are compared.

The spectrum of BPSs in the MMR-defective strains is dominated by A:T transitions at

5'NAC3'/3'NTG5' sites (11). To test if the mutational density pattern was simply due to the distribution

of these sites, we removed all the A:T transitions at 5'NAC3'/3'NTG5' sites from the data set. As shown

in Additional file1: Figs. S5A, S6A, and Tables 1 and 2, although the mutation rate was reduced by 60%

when these BPSs were removed, the remaining 12,542 BPSs fell into the same wave pattern. Thus, the

variation in BPS rates must be reflective of regions of the chromosome and not the distribution of the

hotspot DNA sequences.

Transcription

One obvious hypothesis is that the mutational density pattern reflects transcriptional patterns.

Mutation rates have been reported to be both increased (13) and decreased (14) by high levels of

transcription. In a previous paper (7) we reported that highly expressed genes had normal mutation

rates, a finding that was confirmed in a recent study using deep sequencing to detect mutations (15).

To determine if, nonetheless, transcription influences the wave pattern, we used RNA-Seq to

quantitate the RNA levels in a Δ*mutL* mutant strain in the lag, exponential, and stationary phases of

growth. These results should be representative of the cells in colonies during our MA experiments. We

then compared these results to the BPS density pattern by binning the RNA-Seq results into the same

bins as used for mutational data. As shown in Fig. 1B and Additional file 1: Fig. S1B, there was no

similarity in the two data sets.

Because ribosomal operons are homologous, we could not call SNPs in these genes. The RNA

reads from the genes in ribosomal operons were also removed from the RNA-Seq data, but their

positions have been indicated in Fig. 1B. Interestingly, bin 35, which includes the ribosomal *rrnG*

operon, has a high number of RNA reads even in the absence of reads from the *rrnG* genes. This high

level of expression is almost exclusively due to the *ssrA* gene, which encodes transfer-messenger RNA

(16), that is highly expressed under all three conditions (the RNA-Seq data will be further analyzed in a

subsequent paper). The BPS density pattern of several of the strains that will be discussed below tends

to reach a minimum in this general area, but it is as often at bin 33 as at bin 35. Given that the bin size

is 100 Kb, it seems unlikely that the high level of transcription level of *ssrA*, located in the middle of bin

35, is causing these patterns.

Replication Initiation

The BPS density pattern is centered on *oriC*, the origin of replication, and over many experiments

in different genetic backgrounds the pattern around the origin has proved to be stable. In both

replichores BPS rates decline to a minimum about 300 Kb from the origin and then increase until a

peak is reached about 900 Kb from the origin. Mutation rates then fall again and reach a minimum

about 3/5th of the distance along each replichore.

We tested whether replication initiation was responsible for the maintenance of this pattern by

performing MA experiments with strains with errant replication start sites. The *rnhA* gene encodes

RNase H1, which degrades RNA-DNA hybrids; in the absence of RNase H1, persistent R-loops can

initiate aberrant DNA replication and disrupt normal fork migration (17, 18). But an MA experiment

with a Δ*rnhA* Δ*mutL* mutant strain showed no difference in the BPS density pattern from that of the

Δ*mutL* mutant strain (Fig. 1C, Additional file 1: Fig. S1C, Tables 1 and 2), indicating that aberrant

replication Initiation does not influence where mutations occur, at least not when a powerful *oriC* is

present.

To further test the influence of replication initiation on the mutational density pattern, we

performed MA experiments with strains that have a 5.1 Kb region containing *oriC* moved to the

midpoint of the right replichore, where it is called *oriZ* (19). These strains are derived from *E. coli* K12

strain AB1157, instead of MG1655 strain, the ancestor of our MA strains, and have a large inversion in

the right replichore that relieves the head-on collision between replication initiating at *oriZ* and

transcription of the *rrnCABE* operon (20). As a control we created an AB1157 Δ*mutL* mutant strain,

which had the similar wave-like BPS density pattern as our MMR⁻ strains, indicating that the pattern is

intrinsic to *E. coli* K12, not specific to sub-strain MG1655 (Additional file 1: Figs. S5B, S6B, Tables 1 and

2). We performed MA experiments on Δ*mutL* derivatives of two additional strains: one with both *oriZ*

and *oriC* (WX320 Δ*mutL*), and one with only *oriZ* (WX340 Δ*mutL*). The strain that contained two origins

had a similar BPS density pattern as the MMR⁻ strains (Fig. 1D, Additional file 1: Fig. S1D, Tables 1 and

2), suggesting that, under our conditions, firing of *oriZ* could not overcome the influence of *oriC*.

However, the strain containing only *oriZ* showed a decrease in BPS rate in the 200Kb-area surrounding

the new origin, similar to that normally observed about *oriC* (Fig. 1E, Additional file 1: Fig. S1E, Tables 1

and 2).

SeqA

As mentioned above, in *E. coli* the adenines in GATC sites are methylated by the Dam

methylase. The SeqA protein binds to hemimethylated GATC sites, many of which are clustered around

OriC, and, by so doing, SeqA occludes the replication initiation protein, DnaA, hindering origin firing.

Sequestering of the origin persists for about one third of a generation; however, the mechanism of

relief is not clear. In the absence of SeqA, unregulated initiation presumably results in over-replication,

at least when cells are rapidly growing in rich medium (21). Downstream events, such as replication

fork collapse, add to the phenotypes of *seqA* mutant cells (22).

Loss of SeqA affects chromosomal structure in areas distant from OriC. By binding to

hemimethylated DNA, SeqA forms complexes behind the replication fork as it progresses around the

chromosome (21). In addition, SeqA binds to areas of the chromosome with closely spaced GATC sites,

as well as to particular genes regulated by GATC methylation (23). In the absence of SeqA the

superhelicity of the chromosome increases, the nucleoid condenses (24), and transcription is altered

(25).

To test whether SeqA affects the BPS density pattern, we performed an MA experiment with a

Δ*mutL* Δ*seqA* mutant strain. As shown in Fig. 1F, Additional file 1: Fig. S1F, and Tables 1 and 2, loss of

SeqA somewhat amplified the BPS density pattern of the right replichore, but the pattern was still

highly correlated to that of right replichore of the MMR-defective strains. However, the pattern in the

left replichore was disrupted in the Δ*seqA* Δ*mutL* mutant strain. Based on chromatin

immunoprecipitation analysis, this area of the left replichore is not targeted by SeqA to a greater

extent than the same area of the right replichore (23, 26), suggesting that the disruption of the

mutational pattern is not due to loss of binding by SeqA. As shown in Fig. 1B, bins 44, 38, 37, and 35

contain a number of highly expressed genes; in addition, bins 42 and 35 contain highly transcribed

ribosomal RNA genes. Thus, we hypothesize that loss of SeqA makes the replication machinery

particularly susceptible to interference by transcription, disrupting the mutational pattern. If this

hypothesis is correct, the interference apparently makes replication more accurate, perhaps by slowing

the speed of DNA polymerase.

Replication Fork Progression

In a recent report, Dillon *et al*., 2018 (2), hypothesized that the periodic variation in mutation

rates across the chromosome was tied to the timing of DNA replication. Cellular levels of dNTPs are

controlled by ribonucleotide reductase (RNR), the expression of which increases when the origin fires

(27, 28) }. In fast-growing bacteria in which new rounds of replication are initiated before cell division,

the levels of dNTPs should be high when each origin fires, but then fall as progression of the multiple

forks dilute the dNTPs. High levels of dNTPs are predicted to increase the probability of

misincorporation, and thus the mutation rate, whereas low levels of dNTPs should slow replication and

improve fidelity. This systematic fluctuation in dNTP levels could account for the pattern of mutational

density across the chromosome (2).

This hypothesis can be tested by examining the wave pattern from several experiments that we

have already published (11). The number of replicating chromosomes is a positive function of the cell's

growth rate (29). When Δ*mutL* or Δ*mutS* mutant strains were grown on glucose minimal medium,

which reduces the growth rate about 2-fold relative to growth on LB, the BPS density pattern became

chaotic (Fig. 2A, Additional file 1: Figs. S2A, S5C, S6C, Tables 1 and 2). Supplementing the minimal

medium with just enough LB to increase the growth rate to normal restored the wave-like BPS density

pattern (Fig. 2B, Additional file 1: Fig. S2B, Tables 1 and 2). Growing the cells on diluted LB, on which

the growth rate was the same as on minimal glucose medium, also preserved the wave-like BPS density

pattern (Fig. 2C, Additional file 1: Fig. S2C, Tables 1 and 2). When the cells were grown at a lower

temperature, which also reduced the growth rate 2-fold, the overall shape of the BPS density pattern

was retained, but the increases in BPS rates that normally peak at about 900 Kb from each side of the

origin were shifted about 200 Kb further from OriC (Fig. 2D, Additional file 1: Fig. S2D, Tables 1 and 2).

In addition, the magnitude of the fluctuation of the BPS rate across the chromosome was doubled.

Thus, it appears that growth rate *per se* is not a major determinant of the BPS density pattern, but

other factors, such as the composition of the growth medium, are also important. For example, the

expressions and DNA binding characteristics of nucleoid associated proteins (NAPs) are different under

different growth conditions (30-32)

In *E. coli* growing aerobically dNTP levels are regulated by a Class Ia RNR encoded by the *nrdA*

and *nrdB* genes. Although most of this regulation appears to be due to DnaA interactions (28), the

NrdR repressor, which regulates a poorly expressed Class Ib RNR, also regulates *nrdAB* transcription;

loss of NrdR results in increased expression of RNR throughout the cell cycle (33). Increased RNR

should result in increased dNTP levels, and, indeed, the Δ*nrdR* Δ*mutL* mutant strain had twice the

mutation rate as the MMR⁻ strains, as expected when dNTP levels are high (Additional file 2: Table S1).

As shown in Fig. 2E, Additional file 1: Fig. S2E, Tables 1 and 2, loss of NrdR did not change the basic BPS

density pattern, but the peak rate was shifted away from the origin about 200 Kb on each side. This

pattern was similar to that observed when cells were grown at low temperature, as described above.

Replication fork progression is aided by the accessory replication helicase, Rep, and, in its

absence, the time required for chromosome duplication is doubled (34). Rep removes proteins bound

to the DNA in front of the fork (34, 35). While these nucleoproteins are primarily transcription

complexes (36, 37), Rep could also free the DNA of blocking NAPs. In addition, Rep aids in restarting

replication forks after they stall or collapse (38, 39). As shown in Fig. 2F, Additional file 1: Fig. S2F,

Tables 1 and 2, with the exception of a region close to the origin, loss of Rep disrupted the BPS density

pattern across the chromosome, suggesting that slowing or stalling the fork results in a distribution of

BPS that is essentially random.

Replication Termination

The results from almost all the strains tested show an increase in the BPS rate in the region

where replication terminates. The pattern of this increase varies somewhat among experiments.

Usually there are two unequal peaks, as shown in Fig.1A, but in some experiments these peaks are

better defined and of equal heights and occasionally there is just one peak. We do not know the source

of this variation, but suspect it is simply random noise.

Replication terminates approximately 180° from the origin in a 1200 Kb region bounded by

replication pause (Ter) sites; this region extends from bin 18 to bin 31 in our Figures. The anti-helicase

Tus protein binds to the Ter sites and allows each replication fork to enter but not to exit, creating a

replication fork "trap", within which the two forks fuse and the chromosome dimer is resolved (40). To

determine if the interaction of replication forks with Tus contributes to the increased mutation rate

within this region, we performed an MA experiment on a Δ*tus* Δ*mutL* mutant strain. As shown in Fig.

2G, Additional file 1: Fig. S2G, Tables 1 and 2, loss of Tus did not affect the BPS density pattern.

The Ter macrodomain (MD) extends from 1200 Kb to 2200 Kb (12), which is roughly from bin 20

to bin 28 in our Figures. The structure of the Ter MD is maintained by the MatP protein, which binds to

23 *matS* sites within this region (41). In the absence of MatP, the Ter MD is disorganized, the DNA is

less compact, and the Ter MD segregates too early in the cell cycle and fails to localize properly at

midcell (41, 42). Because the mobility of the Ter MD is increased in the absence of MatP, DNA

interactions across MD barriers can occur in Δ*matP* mutant cells (41).

As shown in Fig. 2H, Additional file 1: Fig. S2H, Tables 1 and 2, loss of MatP caused a severe

disruption of the BPS density pattern. The mutation rates in the Ter MD were depressed whereas new

peaks appeared on either side of the Ter MD in the Right and Left MDs. Interestingly, the BPS pattern

near the origin was maintained in the right but not in the left replichore.

Recombination and the SOS response

Homologous recombination is intimately connected to replication. As replication proceeds,

various blocks, such as DNA lesions, transcription complexes, and DNA secondary structures, can cause

the replisome to pause and to eventually disassemble. This potentially lethal event is prevented by

recombination, which can repair and restart the replisome (43). In addition, the termination region is

subject to hyperrecombination (44, 45) particularly in the region bounded by TerA and TerB (our bins

21 to 24), named the terminal recombination zone (TRZ) (46, 47).

Elimination of *E. coli*'s major recombinase, RecA, had a modest effect on the mutational density

pattern. As shown in Fig. 3A, Additional file 1: Figs. S3A, S5D, S6D, Tables 1 and 2, in the *recA mutL* and

recA mutS mutant strains, BPS rates declined in bins 21 to 25, corresponding fairly well to the TRZ

described above. However, the hyperrecombination within the TRZ is dependent on the recombination

pathway defined by participation of the RecBCD complex. When we eliminated RecB, the mutational

density pattern did not phenocopy that seen when RecA was absent (Additional file 1: Figs. S5E, S6E,

Tables 1 and 2). Either our protocol is not sensitive enough to detect an effect of loss of RecB, or

another recombination pathway, *e.g.* RecFOR (48), is sufficient to maintain the mutation rate in the

region.

In addition to its role in recombination, RecA is also a master regulator of the SOS response to

DNA damage, which includes the induction of two error-prone DNA polymerases, DNA Pol IV and V. To

test whether these polymerases are involved in determining the BPS density pattern, we performed an

MA experiment on a strain deleted for the genes that encode Pol IV, *dinB*, and PolV, *umuDC*. As shown

in Fig. 3B, Additional file 1: Fig. S3B, Tables 1 and 2, the BPS density pattern in the *mutL dinB umuDC*

mutant strain was not significantly different than the MMR⁻ pattern. The genes of the SOS response

are repressed by the LexA protein; the *lexA3* allele encodes a super-repressor LexA protein that

prevents the SOS genes from being induced (49). When this allele was present the BPS density pattern

was also unaffected (Fig. 3C, Additional file 1: Fig. S3C, Tables 1 and 2). Thus, the SOS response appears

to play no role in determining the pattern of BPSs across the chromosome.

Nucleoid Associated Proteins

In a previous report (1), we found that the BPS density pattern of a Δ*mutL* strain was correlated

with the density of genes activated by the HU protein and repressed by the Fis protein. Combining

these two factors in a linear correlation model accounted for 33% of the variation in the mutational

data. HU constrains supercoils and compacts the DNA into nucleosome-like particles; Fis also

constrains supercoils but, in addition, bends the DNA (31). While both of these NAPs affect

transcription, the general lack of correlation of the BPS rate with transcriptional levels (7); also see

above) led us to hypothesize that mutation rates across the chromosome were correlated not with

transcription *per se*, but with areas of high DNA structure (1). To further test this hypothesis we

preformed MA experiments with MMR⁻ mutant strains also defective for each of a number of NAPs.

HU exists as a dimer of its two subunits, HUα and HUβ, encoded by the paralogous genes *hupA*

and *hupB*, respectively, in the three possible configurations. While loss of both subunits confers a

severe growth defect, loss of only one has little consequence during a normal growth cycle, suggesting

they can substitute for each other. HU$\alpha\beta$ is the dominant form over most of the cell cycle, but

significant amounts of HUα_2 are found during lag phase and early exponential phase, and HUβ_2 is

prominent in stationary phase (30). Chromatin immunoprecipitation sequencing (ChIP-Seq) results

revealed that HU binds non-specifically to the chromosome and the DNA binding patterns of the three

dimers appear to be identical (50).

We performed MA experiments with both $\Delta hupA$ $\Delta mutL$ and $\Delta hupB$ $\Delta mutL$ mutant strains. As

shown in Figs. 3D, 3E, Additional file 1: Figs. S3D, S3E, Tables 1 and 2, although loss of *hupB* appeared

to affect the BPS density pattern, the differences from the MMR⁻ pattern were not significant.

However, loss of *hupA* depressed mutation rates across the chromosome, particularly in the terminus

region, while creating new peaks on either side of the Ter MD.

Fis has many binding sites across the chromosome, but, based on data at RegulonDB, Sᴏʙᴇᴛᴢᴋᴏ *et*

al. (2012) (51) reported more Fis binding sites in the origin region. However both Chromatin

immunoprecipitation plus microarray analysis (ChIP-chip) and ChIP-Seq studies found the density of Fis

binding to be more-or-less constant across the chromosome (32, 52). Loss of Fis in both Δfis $\Delta mutL$ and

Δfis $\Delta mutS$ mutant strains tended to flatten the BPS density pattern across the chromosome except in

the region around the origin (Fig. 3F, Additional file 1: Figs. S3F, S5F, S6F, Tables 1 and 2).

The NAP HNS binds to DNA at its high-affinity binding sites and then spreads by oligomerization

along A:T rich regions of DNA. Bridging between HNS-DNA complexes condenses the DNA into a few

clusters per chromosome (53-55). However, as shown in Fig. 3G, Additional file 1: Fig. S3G, Tables 1

and 2), loss of HNS had little effect on the BPS density pattern, and, thus, the long-range structures

produced by HNS appear not to affect BPS rates.

The DPS protein accumulates in stationary phase cells, condenses the nucleoid into a crystalline-

like state, and protects the DNA from oxidative and other damage (56). Despite this radical physical

change, loss of DPS had little effect on the BPS density pattern Fig. 3H, Additional file 1: Fig. S3H, Tables

1 and 2). Of course, we do not know the degree to which cells in stationary phase contribute to the BPS

rates under our experimental conditions.

Proofreading

As mentioned above, epsilon is the proofreading subunit of the DNA polymerase III holoenzyme.

The *mutD5* allele encodes an epsilon protein that is inactive for proofreading, and strains carrying this

allele have a mutation rate 4000-fold greater than that of wild-type strains, and 35-fold greater than

that of MMR-defective strains (10). As shown in Fig. 4A, Additional file 1: Fig. S4A, Tables 1 and 2, when

proofreading was inactive but MMR was active, the BPS density pattern was less dramatic than when

MMR was inactive and proofreading was active; but, nonetheless, the wave pattern was basically the

same. When both MMR and proofreading were inactive, which reveals the mutations solely due to

replication errors, the BPS density pattern was nearly flat but around the origin it retained significant

correlations to the patterns of both the MMR-defective and *mutD5* mutant strains (Fig. 4B, Additional

file 1: Fig. S4B, Tables 1 and 2). Thus, the pattern of BPS on both sides of the origin appears to be

established by replication errors and then elsewhere across the chromosome the density pattern is

largely due to differential error-correction by both proofreading and MMR.

Wild-type

The mutational density wave patterns evident in our data, and in data from other bacteria (2, 3),

were obtained when MMR was inactive. Thus, the question arises: does the pattern appear in wild-

type strains? It is difficult to answer this question because mutation rates in wild-type strains are so

low (in *E. coli*, 120-fold lower than that of MMR-defective strains (7, 11) that enormous experiments

would have to be conducted in order to accumulate enough mutations to approach statistical

confidence. In a recent study we compared the mutation rates and spectra of a number of *E. coli*

strains defective in various DNA repair activities; of these, the results from seven strains were

indistinguishable from those of the wild-type parent (57). By combining the mutations from these

strains, we achieved 1933 BPSs (11), enough to expect to see a wave pattern if it existed. As is evident

in Figs. 4C, 4D, Additional file 1: Figs. S4C, S4D, Tables 1 and 2, these BPSs did not create fall into a

recognizable pattern. Indeed, the pattern from the wild-type strains appears to be random; the

variance to mean ratio of the binned mutations is 1.4, indicating the values are not disperse, and the

MatLab "runstest", a test for runs, returns a P value of 0.58, also indicating that the bin values are

random. In the wild-type strain both MMR and proofreading are active, and, while these two activities

have similar correction biases (Fig. 4A), proofreading is much more powerful, producing the pattern

seen in the MMR-defective strains. Although we cannot conclude that the mutational density pattern

in the wild-type strain is other than random, it does have similarity to both the patterns seen in the

MMR-defective strains and in the *mutD5* mutant strain, particularly around the terminus (Figs. 4C, 4D,

Additional file 1: Figs. S4C, S4D and Tables 1 and 2).

Bacillus subtilis

In additional to *E. coli* strains, symmetrical mutational density patterns have been demonstrated

in MMR- derivatives of *Vibrio fischeri*, *V. cholera* (2-4), *Pseudomonas fluorescens* (5), and *P. aeruginosa*

(6). Here we add *Bacillus subtilis* to this list. As shown in Fig. 4E, Additional file 1: Fig. S4E, Tables 1 and

2, the BPS mutation rates in a *B. subtilis mutS::*Tn*10* mutant strain fell into a wave like pattern that was

symmetrical about the origin. Although similar in shape, the pattern was significantly different from

that of *E. coli* (Fig. 4E, Additional file 1: Fig. S4F, Tables 1 and 2). However, as in *E. coli*, the BPS rate

appeared to increase in the terminus region, which in *B. subtilis* is not 180° from the origin and

corresponds to bins 20-24 in Fig. 4E.

Discussion

In this report we have examined a number of factors that could be responsible for establishment

and maintenance of the symmetrical wave-like BPS density pattern across the chromosome. In broad

terms these factors were: transcription; DNA replication initiation, progression, and termination;

recombination and the SOS response to DNA damage; the binding of nucleoid-associated proteins;

and, error-correction by MMR and proofreading. As discussed above, we found that transcription and

the SOS response had little effect, and the effect of recombination was modest and confined to the

terminal region. We discuss the more significant, factors in greater detail here.

DNA replication initiation, progression, and termination

Providing additional replication origins, either by eliminating RNase H1 or by inserting an ectopic

oriC (*oriZ*), did not disrupt the wave (Fig. 1C and 1D). However, when *oriZ* was the only origin of

replication, the region of depressed BPS rate that surrounds *oriC* when it is in the normal position was

re-established about the new origin (Fig. 1E). Note that only 5.1 Kb of DNA containing *oriC* was

relocated (19), whereas the region of reduced mutation rate is about 200 Kb; thus, the mutation rate is

not determined just by the DNA sequence surrounding the origin. We hypothesize that the process of

replication initiation protects the DNA from damage and/or newly established replication forks have a

low error-rate. In addition, the BPS mutation rate was increased for about 1000 Kb (10 bins) on either

side the new origin so that it resembled the same region about the normal origin. The size of this area

is close to the same size detected as "interacting zones" around *oriZ* (58). However, the mutational

density pattern across the rest of the chromosome did not re-establish itself to be symmetrical about

oriZ, but remained symmetrical about the absent *oriC*. Thus, other factors must be important at distant

regions. We can also conclude that the overall structure of the BPS pattern across the chromosome is

not determined by active replication initiation *per se*, but may have evolved in response to replication

initiation.

Both *V. fischeri* and *V. cholera* have multiple circular chromosomes of different sizes; by

comparing the mutational density patterns of these chromosomes, Dillon et al, 2018 (2) identified the

timing of replication as a significant determinant of the mutational density patterns. As mentioned

above, they suggested that the pattern could be the result of variations in the levels of dNTPs as origins

fire during rapid growth. Our results provide partial support for this hypothesis, but also indicate that

growth on rich medium, not growth rate *per se*, is a significant determinant of the mutational density

pattern, possibly because of effects on the expression and DNA binding of HU and Fis (see below). Our

results with a Δ*nrdR* mutant strain (Fig. 2E) also show that dNTP levels are important, but, again, other

factors are driving the overall wave pattern. In addition, the results with the Δ*rep* mutant strain (Fig.

2F) indicate that radically interrupting the progression of the replication fork disrupts the mutational

density pattern.

After declining to a local minimum about 3/5th of the distance along each replichore, mutation

rates rise in the terminus region (Fig. 1A). We originally hypothesized that this increase was due to

collisions of the replication complexes with the Tus anti-helicase, most of which would take place in

the region between bins 18 and 31 (1). However, elimination of Tus had no effect on the wave pattern

(Fig. 2G, Additional file 1: Fig. S2G), refuting this hypothesis. But, eliminating MatP, the protein that

maintains the structure of the Ter MD, reduced mutation rates in the Ter MD, suggesting that

replication of highly structured DNA is error-prone (Fig. 2H, Additional file 1: Fig. S2H). Interestingly, in

Δ*matP* mutant cells mutation rates increased in the Right and Left MCs, suggesting that in the absence

of MatP, these adjacent regions of DNA gained structure. The mechanism by which MatP structures

the Ter MC is not clear (59, 60), but it probably involves supercoils which, when unconstrained, could

migrate into adjacent areas. However, two studies have found that loss of MatP increases the mobility

and long-range interactions of adjacent DNA (41, 61). Resolution of these conflicts will require further

experimentation.

Nucleoid-associated proteins

Our previous results predicted that the NAPs HU and Fis should play a role in establishing or

maintaining the wave-pattern of BPS, but HNS should not (1). The results presented here confirmed

that prediction. In addition, we also found that Dps had no effect on the BPS density pattern. Because

the NAPs affect gene expression in various ways, we cannot conclude that DNA binding by the NAPs

themselves is responsible for the mutational pattern. And, indeed, we found no significant positive

correlations between the BPS pattern and published binding sites of the NAPs, although the location of

the binding sites themselves vary widely among published results (e.g. see (32, 50) and the data in

RegulonDB (62). Nonetheless, we favor the hypothesis that structuring of the DNA by the NAPs,

directly or indirectly, contributes to the BPS pattern.

The local effect of HU binding is to bend the DNA, but dimer-dimer interactions produce higher-

order HU-DNA complexes that can constrain negative supercoils (63, 64). The analysis of the effect of

HU on the mutational density pattern is also complicated by the variation in the cellular concentrations

of the three forms with growth cycle (30). Although the two subunits can compensated for each other

for viability, the affinities of three forms of HU for various DNA structures (linear, nicked, and gapped)

differ (65). Both $HU\alpha_2$ and $HU\alpha\beta$ can constrain supercoils, but $HU\beta_2$ apparently cannot, at least *in vitro*

(30). Since $HU\beta$ is not stable (66), the phenotype of a $\Delta hupA$ mutant strain may simply be that of low

cellular HU concentration.

While loss of $HU\beta$ had little effect on the mutational density pattern (Fig. 3D, Additional file 1:

Fig. S3D), loss of $HU\alpha$ reduced the overall BPS rate by 33% and also changed the pattern (Fig. 3E,

Additional file 1: Fig. S3E). In the $\Delta hupA$ $\Delta mutL$ mutant strain, mutation rates were low in the terminus

region but elevated in the Right and Left MCs, very similar to the pattern seen in the $\Delta matP$ $\Delta mutL$

mutant strain (Fig. 2H). The low mutation rate in the Ter MD corresponds well with the density of

genes activated in wild-type cells but not in HU-deleted cells (67). HU regulates transcription by

modifying the DNA superhelical density, suggesting that, like MatP, HU increases the structure of the

DNA in the terminal region, causing DNA replication to become error-prone.

The mutational pattern in the $\Delta hupA$ $\Delta mutL$ mutant strain is also similar to that of the $\Delta recA$

$\Delta mutL$ strain (Fig. 3A). An mutation accumulation experiment with a $\Delta hupA$ $\Delta recA$ $\Delta mutL$ mutant strain

resulted in a pattern similar to both single mutant strains, and so was not informative (data not

shown).

Fis is a major transcriptional regulator, either activating or repressing, directly or indirectly,

nearly a thousand genes (52). At some promotors Fis acts as a classic transcriptional regulator by

interacting with RNA polymerase, but at other promotors Fis affects gene expression by altering the

DNA superhelical density (68). The Fis DNA binding sequence is degenerate and the estimated number

of binding sites identified by sequence analysis across the genome varies widely among studies. The

results of ChIP analyses also vary: the number of DNA regions associated with Fis in different studies

ranges from 200 to 1500, which may reflect the effects of different growth conditions (32, 52, 69).

The loss of Fis nearly doubled the overall BPS rate, but flattened the wave pattern outside of the

origin region (Fig. 3F, Additional file 1: Figs. S5F, S6F). As mentioned above, the mutational density

pattern MMR⁻ strains is correlated to the density of genes activated in a Δ*fis* mutant strain as reported

by Blot *et al*, 2006 (70). This correlation is particularly strong in bins 14 to 33 ($\rho = 0.62$, $P = 0.003$),

which corresponds to the area flattened in the Δ*fis* Δ*mutL* and Δ*fis* Δ*mutS* mutant strains. However, no

correlation exists between our mutational data and genes found to be responsive to Fis in a recent

study (32). Clearly more studies are needed to resolve these conflicts.

Error-correction

Assuming that the BPS recovered from the *mutD5* Δ*mutL* mutant strain are due to intrinsic errors

made by DNA polymerase, we conclude that the polymerase is accurate close to the origin, then

becomes increasingly less accurate as replication proceeds to about 1/3 of the replichore, at which

point accuracy increases again (Fig. 4B). The mutational pattern in the MMR-defective strains must

reflect the biased ability of proofreader to correct these polymerase errors. Thus, proofreading is

effective close to the origin, but then declines for about 1/3 of the replichore, increases in the Right

and Left MCs, but declines again in the terminus region (Figs. 1A, 4B). Mutations in the *mutD5* mutant

strain are due to the failure of MMR to correct the polymerase errors that survive in the absence of

proofreading. Since the mutational pattern in the *mutD5* mutant strain mimics that of the MMR-

defective strains (Fig. 4A, Additional file 1: Fig. S4A, Tables 1 and 2), error correction by MMR, although

much less powerful, apparently has the same biases as error correction by proofreader.

Given the above considerations, we would expect the mutational density pattern in the wild-type

parental strain to mimic that of the MMR-defective strains. However, the nearly 2000 BPS accumulated

in eight strains with wild-type mutational phenotypes appear randomly distributed across the

chromosome (Figs. 4C, 4D). One explanation for this discrepancy is that when both proofreading and

MMR are active, the mutation rate is reduced to the point that other, weaker DNA repair and

mutagenic activities obscure the underlying pattern. Although not significantly correlated, the wild-

type BPS pattern is similar to the MMR⁻ pattern, particularly around the terminus, supporting this

hypothesis. Alternatively, although the eight strains that we combined to produce 2000 BPS have the

same mutation rates, they may have different wave patterns that tend to negate each other and

obscure the underlying wild-type pattern.

Conclusions

By applying mutation accumulation followed by whole-genome sequencing to bacterial strains mutant

in various activities we have determined that the most important factors determining the symmetrical

pattern of BPS rates across the chromosome are: the initiation, progression, and termination of DNA

replication, replication error-correction, and chromosome structure. Because of the conservation of

these factors, our results should apply to most bacteria, and possibly eukaryotes, and imply that

different regions of the genome evolve at different rates.

Methods

479 **Bacterial Strains and Media**

480 The strains used in this study are listed in Additional file 2: Table S1 and the methods of their

481 construction are in Additional file Materials and Methods. Standard media and antibiotics were used

482 (see Additional file 3: Methods).

483 **Mutation accumulation experiments**

484   The complete MA procedure has been extensively described (7, 11, 57). Details are given in the

485 Additional file 3: Methods

486 **Genomic DNA Preparation, Library construction, Sequencing, and Sequence Analysis**

487   Details are given in the Additional file 3: Methods.

488 **RNA sequencing**

489   The *E. coli* strain PFM144, which is PFM2 Δ*mutL* (11), was grown in LB and aliquots collected

490 during lag (OD = 0.022), log (OD = 0.3), and stationary (OD = 1.5) phase. The number of cells collected

491 was kept constant for each growth phase. Total RNA was extracted, DNA and rRNA were removed, and

492 libraries constructed and sequenced as detailed in the Additional file 3: Methods. Three biological

493 triplicates were prepared for each growth phase.

494 A complete analysis of the RNA-Seq results will be the subject of a subsequent report. For this report

495 the numbers of RNA-Seq reads for each condition were first normalized to the number of reads

496 mapped to the gene *holD*, which was determined by rtPCR to be expressed at the same level in all

497 phases of growth. The means of the normalized RNA reads from the triplicates were then binned into

498 the same bins used for the mutational analysis. A fourth-order Daubechies wavelet transform was

499 performed on the binned RNA-Seq reads as described for the mutational data (1).

500 **Statistical Analysis**

To obtain the BPS density patterns, the numbers of BPSs were binned into 46 bins, each 100 Kb long, as

described (1). A fourth-order Daubechies wavelet transform was performed on the binned mutation

data as described (1). For presentation in the Figures, these results were converted into rates by

dividing the number of BPS by the appropriate number of generations. Pearson's product-moment

correlation coefficient, ρ, was used to evaluate the correlations between the binned BPS data (Table

1). Spearman's nonparametric correlation coefficient was also computed for a few data sets, but gave

similar results. To account for multiple comparisons, p values were adjusted using the Benjamini-

Hochberg method (71) with the false discovery rate set at 25%, implemented with the MatLab R2018a

"mafdr" command. Because comparisons using the data from the same strain are not independent,

this adjustment was made separately for each column in Tables 1 and 2.

To further compare the BPS density patterns between two data sets, wavelet coherence was calculated

and plotted using the MatLab R2018a "wcoherence" command. While the Daubechies wavelet

provides a good visual representation of the binned data, it is not continuous and thus not easily

adapted for wavelet coherence analysis. The MatLab program first converts the binned data to Morlet

wavelets and then computes the coherence between two of these wavelets. We chose to analyze the

data with wavelet coherence because it gives a measure of the correlation between the signals

(displayed as colors in the Figures) (72, 73). In addition, the MatLab wavelet coherence plot indicates,

as a dashed curve, the 'cone of influence' within which results are free of artifactual edge effects (72).

The relative phase-lag between the two signals is indicated by small arrows: arrows pointing right

indicate in-phase, arrows pointing left indicate 180° out-of-phase, and arrows pointing in other

directions indicate the various degrees in between. Because the MatLab program assumes a

frequency-time series, the X axis of the plot is cycles/sample and the Y axis is time; we converted these

to bins/cycle (on an inverted scale) and bins, respectively.

Declarations

525   **Ethics approval and consent to participate**: Not applicable

526   **Consent for publication**: Not applicable

527   **Availability of data and material**

Competing interests

The authors declare no competing interests

Funding

This research was supported by US Army Research Office Multidisciplinary University Research

Initiative (MURI) Award W911NF-09-1-0444 to P.L.F. and H.T., the National Institutes of Health T32

GM007757 to B.A.N., and the US Army Undergraduate Research Apprenticeship Program to J. H. and S.

R

Authors' contributions

B.A.N. and P.L.F. designed the research, analyzed the data, and wrote the paper. B.A.N executed the

experiments. W.M., H.L., and H.T. performed the bioinformatic analyses. P.L.F. and H.T. supplied

resources. All authors read and approved the final manuscript.

Acknowledgements

We thank H. Bedwell-Ivers, C. Coplen, M. Durham, J. Eagan, N. Gruenhagen, J. A. Healy, N. Ivers,

C. Klineman., H. Lee, E. Popodi, I. Rameses, S. Riffert, H. Rivera, D. Simon, K. Smith, J. Townes, L. Tran,

and L. Whitson for technical help. Bacterial strains were kindly provided by R. Schaaper, R. Reyes-

Lamothe, D. Kearns, M. Konkol, and The National BioResource Project at the (Japanese) National

Institute of Genetics. We also thank S. E.Bell and X. Wang for technical help and discussions.

References

1.   Foster PL, Hanson AJ, Lee H, Popodi EM, Tang H. On the mutational topology of the bacterial

genome. G3 (Bethesda). 2013;3(3):399-407.

2.   Dillon MM, Sung W, Lynch M, Cooper VS. Periodic variation of mutation rates in bacterial

genomes associated with replication timing. MBio. 2018;9(4).

3.   Dillon MM, Sung W, Sebra R, Lynch M, Cooper VS. Genome-wide biases in the rate and molecular

spectrum of spontaneous mutations in *Vibrio cholerae* and *Vibrio fischeri*. Mol Biol Evol. 2017;34(1):93-

109.

4.   Wei W, Xiong L, Ye YN, Du MZ, Gao YZ, Zhang KY, et al. Mutation landscape of base substitutions,

duplications, and deletions in the representative current cholera pandemic strain. Genome Biol Evol.

2018;10(8):2072-85.

5.   Long H, Sung W, Miller SF, Ackerman MS, Doak TG, Lynch M. Mutation rate, spectrum, topology,

and context-dependency in the DNA mismatch repair-deficient Pseudomonas fluorescens ATCC948.

Genome Biol Evol. 2014;7(1):262-71.

6.   Dettman JR, Sztepanacz JL, Kassen R. The properties of spontaneous mutations in the

opportunistic pathogen Pseudomonas aeruginosa. BMC Genomics. 2016;17:27.

7.   Lee H, Popodi E, Tang H, Foster PL. Rate and molecular spectrum of spontaneous mutations in

the bacterium *Escherichia coli* as determined by whole-genome sequencing. Proceedings of the

National Academy of Sciences of the United States of America. 2012;109(41):E2774-E83.

8.   Marinus MG. DNA Mismatch Repair. EcoSal Plus. 2012; doi:10.1128/ecosalplus.7.2.5

9.   Ganai RA, Johansson E. DNA replication-a matter of fidelity. Mol Cell. 2016;62(5):745-55.

10.   Niccum BA, Lee H, MohammedIsmail W, Tang H, Foster PL. The spectrum of replication errors in

the absence of error correction assayed across the whole genome of *Escherichia coli*. Genetics.

2018;209(4):1043-54.

11.   Foster PL, Niccum BA, Popodi E, Townes JP, Lee H, MohammedIsmail W, et al. Determinants of

base-pair substitution patterns revealed by whole-genome sequencing of DNA mismatch repair

defective *Escherichia coli*. Genetics. 2018;209(4):1029-42.

12.   Valens M, Penaud S, Rossignol M, Cornet F, Boccard F. Macrodomain organization of the

Escherichia coli chromosome. EMBO J. 2004;23(21):4330-41.

13.   Jinks-Robertson S, Bhagwat AS. Transcription-associated mutagenesis. Annu Rev Genet.

2014;48:341-59.

14.   Rasouly A, Pani B, Nudler E. A magic spot in genome maintenance. Trends Genet. 2017;33(1):58-

67.

15.   Zhang X, Zhang X, Zhang X, Liao Y, Song L, Zhang Q, et al. Spatial vulnerabilities of the *Escherichia*

coli genome to spontaneous mutations revealed with improved duplex sequencing. Genetics.

2018;210(2):547-58.

16.   Withey JH, Friedman DI. A salvage pathway for protein structures: tmRNA and trans-translation.

Annu Rev Microbiol. 2003;57:101-23.

17.   Kogoma T. Stable DNA replication: interplay between DNA replication, homologous

recombination, and transcription. Microbioland MolecBiolRev. 1997;61:212-38.

18.   Maduike NZ, Tehranchi AK, Wang JD, Kreuzer KN. Replication of the *Escherichia coli* chromosome

in RNase HI-deficient cells: multiple initiation regions and fork dynamics. Mol Microbiol. 2014;91(1):39-

56.

19.   Wang X, Lesterlin C, Reyes-Lamothe R, Ball G, Sherratt DJ. Replication and segregation of an

Escherichia coli chromosome with two replication origins. Proceedings of the National Academy of

Sciences of the United States of America. 2011;108(26):E243-50.

20.   Ivanova D, Taylor T, Smith SL, Dimude JU, Upton AL, Mehrjouy MM, et al. Shaping the landscape

of the *Escherichia coli* chromosome: replication-transcription encounters in cells with an ectopic

replication origin. Nucleic Acids Res. 2015;43(16):7865-77.

21.   Waldminghaus T, Skarstad K. The *Escherichia coli* SeqA protein. Plasmid. 2009;61(3):141-50.

22.   Sutera VA, Jr., Lovett ST. The role of replication initiation control in promoting survival of

replication fork damage. MolMicrobiol. 2006;60(1):229-39.

23.   Sanchez-Romero MA, Busby SJ, Dyer NP, Ott S, Millard AD, Grainger DC. Dynamic distribution of

SeqA protein across the chromosome of *Escherichia coli* K-12. MBio. 2010;1(1):e00012-10.

24.   Weitao T, Nordstrom K, Dasgupta S. *Escherichia coli* cell cycle control genes affect chromosome

superhelicity. EMBO Rep. 2000;1(6):494-9.

25.   Lobner-Olesen A, Marinus MG, Hansen FG. Role of SeqA and Dam in *Escherichia coli* gene

expression: a global/microarray analysis. Proceedings of the National Academy of Sciences of the

United States of America. 2003;100(8):4672-7.

26.   Waldminghaus T, Skarstad K. ChIP on Chip: surprising results are often artifacts. BMC Genomics.

2010;11:414.

27.   Sun L, Fuchs JA. *Escherichia coli* ribonucleotide reductase expression is cell cycle regulated. Mol

Biol Cell. 1992;3(10):1095-105.

28.   Gon S, Camara JE, Klungsoyr HK, Crooke E, Skarstad K, Beckwith J. A novel regulatory mechanism

couples deoxyribonucleotide synthesis and DNA replication in *Escherichia coli*. EMBO J.

2006;25(5):1137-47.

29.   Cooper S, Helmstetter CE. Chromosome replication and the division cycle of *Escherichia coli* B/r. J

Mol Biol. 1968;31(3):519-40.

30.   Claret L, Rouviere-Yaniv J. Variation in HU composition during growth of *Escherichia coli*: the

heterodimer is required for long term survival. J Mol Biol. 1997;273(1):93-104.

31.   Dorman CJ. Function of nucleoid-associated proteins in chromosome structuring and

transcriptional regulation. J Mol Microbiol Biotechnol. 2014;24(5-6):316-31.

32.   Kahramanoglou C, Seshasayee AS, Prieto AI, Ibberson D, Schmidt S, Zimmermann J, et al. Direct

and indirect effects of H-NS and Fis on global gene expression control in *Escherichia coli*. Nucleic Acids

Res. 2011;39(6):2073-91.

33.   Torrents E, Grinberg I, Gorovitz-Harris B, Lundstrom H, Borovok I, Aharonowitz Y, et al. NrdR

controls differential expression of the *Escherichia coli* ribonucleotide reductase genes. J Bacteriol.

2007;189(14):5012-21.

34.   Lane HE, Denhardt DT. The rep mutation. IV. Slower movement of replication forks in *Escherichia*

coli rep strains. J Mol Biol. 1975;97(1):99-112.

35.   Guy CP, Atkinson J, Gupta MK, Mahdi AA, Gwynn EJ, Rudolph CJ, et al. Rep provides a second

motor at the replisome to promote duplication of protein-bound DNA. Mol Cell. 2009;36(4):654-66.

36.   Boubakri H, de Septenville AL, Viguera E, Michel B. The helicases DinG, Rep and UvrD cooperate

to promote replication across transcription units in vivo. EMBO J. 2010;29(1):145-57.

37.   Myka KK, Hawkins M, Syeda AH, Gupta MK, Meharg C, Dillingham MS, et al. Inhibiting translation

elongation can aid genome duplication in *Escherichia coli*. Nucleic Acids Res. 2017;45(5):2571-84.

38.   Heller RC, Marians KJ. Non-replicative helicases at the replication fork. DNA Repair (Amst).

2007;6(7):945-52.

39.   Sandler SJ, Marians KJ. Role of PriA in replication fork reativation in *Escherichia coli*. J Bact.

2000;182:9-13.

40.   Duggin IG, Wake RG, Bell SD, Hill TM. The replication fork trap and termination of chromosome

replication. MolMicrobiol. 2008;70(6):1323-33.

41.   Mercier R, Petit MA, Schbath S, Robin S, El Karoui M, Boccard F, et al. The MatP/matS site-specific

system organizes the terminus region of the *E. coli* chromosome into a macrodomain. Cell.

2008;135(3):475-85.

42.   Espeli O, Borne R, Dupaigne P, Thiel A, Gigant E, Mercier R, et al. A MatP-divisome interaction

coordinates chromosome segregation with cell division in *E. coli*. EMBO J. 2012;31(14):3198-211.

43.   Syeda AH, Hawkins M, McGlynn P. Recombination and replication. Cold Spring Harb Perspect

Biol. 2014;6(11):a016550.

44.   Louarn JM, Louarn J, Francois V, Patte J. Analysis and possible role of hyperrecombination in the

termination region of the *Escherichia coli* chromosome. J Bacteriol. 1991;173(16):5097-104.

45.   Bierne H, Ehrlich SD, Michel B. The replication termination signal terB of the *Escherichia coli*

chromosome is a deletion hot spot. EMBO J. 1991;10(9):2699-705.

46.   Louarn J, Cornet F, Francois V, Patte J, Louarn JM. Hyperrecombination in the terminus region of

the *Escherichia coli* chromosome: possible relation to nucleoid organization. J Bacteriol.

1994;176(24):7524-31.

47.   Corre J, Cornet F, Patte J, Louarn JM. Unraveling a region-specific hyper-recombination

phenomenon: genetic control and modalities of terminal recombination in *Escherichia coli*. Genetics.

1997;147(3):979-89.

48.   Cox MM. Regulation of bacterial RecA protein function. Crit Rev Biochem Mol Biol.

2007;42(1):41-63.

49.   Little JW, Harper JE. Identification of the lexA gene product of *Escherichia coli* K-12. Proceedings

of the National Academy of Sciences of the United States of America. 1979;76(12):6147-51.

50.   Prieto AI, Kahramanoglou C, Ali RM, Fraser GM, Seshasayee AS, Luscombe NM. Genomic analysis

of DNA binding and gene regulation by homologous nucleoid-associated proteins IHF and HU in

Escherichia coli K12. Nucleic Acids Res. 2012;40(8):3524-37.

51.   Sobetzko P, Travers A, Muskhelishvili G. Gene order and chromosome dynamics coordinate

spatiotemporal gene expression during the bacterial growth cycle. ProcNatlAcadSciUSA.

2012;109(2):E42-E50.

52.   Cho BK, Knight EM, Barrett CL, Palsson BO. Genome-wide analysis of Fis binding in *Escherichia*

coli indicates a causative role for A-/AT-tracts. Genome Res. 2008;18(6):900-10.

53.   Wang W, Li GW, Chen C, Xie XS, Zhuang X. Chromosome organization by a nucleoid-associated

protein in live bacteria. Science. 2011;333(6048):1445-9.

54.   Lang B, Blot N, Bouffartigues E, Buckle M, Geertz M, Gualerzi CO, et al. High-affinity DNA binding

sites for H-NS provide a molecular basis for selective silencing within proteobacterial genomes. Nucleic

Acids Res. 2007;35(18):6330-7.

55.   Bouffartigues E, Buckle M, Badaut C, Travers A, Rimsky S. H-NS cooperative binding to high-

affinity sites in a regulatory element results in transcriptional silencing. Nat Struct Mol Biol.

2007;14(5):441-8.

56.   Wolf SG, Frenkiel D, Arad T, Finkel SE, Kolter R, Minsky A. DNA protection by stress-induced

biocrystallization. Nature. 1999;400(6739):83-5.

57.   Foster PL, Lee H, Popodi E, Townes JP, Tang H. Determinants of spontaneous mutation in the

bacterium *Escherichia coli* as revealed by whole-genome sequencing. Proceedings of the National

Academy of Sciences of the United States of America. 2015;112(44):E5990-E599.

58.   Duigou S, Boccard F. Long range chromosome organization in *Escherichia coli*: The position of the

replication origin defines the non-structured regions and the Right and Left macrodomains. PLoS

Genet. 2017;13(5):e1006758.

59.   Dupaigne P, Tonthat NK, Espeli O, Whitfill T, Boccard F, Schumacher MA. Molecular basis for a

protein-mediated DNA-bridging mechanism that functions in condensation of the *E. coli* chromosome.

Mol Cell. 2012;48(4):560-71.

60.   Lioy VS, Cournac A, Marbouty M, Duigou S, Mozziconacci J, Espeli O, et al. Multiscale structuring

of the *E. coli* chromosome by nucleoid-associated and condensin proteins. Cell. 2018;172(4):771-83

e18.

61.   Thiel A, Valens M, Vallet-Gely I, Espeli O, Boccard F. Long-range chromosome organization in *E.*

coli: a site-specific system isolates the Ter macrodomain. PLoSGenet. 2012;8(4):e1002672.

62.   Salgado H, Peralta-Gil M, Gama-Castro S, Santos-Zavaleta A, Muniz-Rascado L, Garcia-Sotelo JS, et

al. RegulonDB v8.0: omics data sets, evolutionary conservation, regulatory phrases, cross-validated

gold standards and more. Nucleic Acids Res. 2013;41(Database issue):D203-13.

63.   Guo F, Adhya S. Spiral structure of *Escherichia coli* HUalphabeta provides foundation for DNA

supercoiling. Proceedings of the National Academy of Sciences of the United States of America.

2007;104(11):4309-14.

64.   van Noort J, Verbrugge S, Goosen N, Dekker C, Dame RT. Dual architectural roles of HU:

formation of flexible hinges and rigid filaments. Proceedings of the National Academy of Sciences of

the United States of America. 2004;101(18):6969-74.

65.   Pinson V, Takahashi M, Rouviere-Yaniv J. Differential binding of the *Escherichia coli* HU,

homodimeric forms and heterodimeric form to linear, gapped and cruciform DNA. J Mol Biol.

1999;287(3):485-97.

66.   Bonnefoy E, Almeida A, Rouviere-Yaniv J. Lon-dependent regulation of the DNA binding protein

HU in *Escherichia coli*. Proceedings of the National Academy of Sciences of the United States of

America. 1989;86(20):7691-5.

67.   Berger M, Farcas A, Geertz M, Zhelyazkova P, Brix K, Travers A, et al. Coordination of genomic

structure and transcription by the main bacterial nucleoid-associated protein HU. EMBO Rep.

2010;11(1):59-64.

68.   Browning DF, Grainger DC, Busby SJ. Effects of nucleoid-associated proteins on bacterial

chromosome structure and gene expression. CurrOpinMicrobiol. 2010;13(6):773-80.

69.   Grainger DC, Hurd D, Goldberg MD, Busby SJ. Association of nucleoid proteins with coding and

non-coding segments of the *Escherichia coli* genome. Nucleic Acids Res. 2006;34(16):4642-52.

70.   Blot N, Mavathur R, Geertz M, Travers A, Muskhelishvili G. Homeostatic regulation of

supercoiling sensitivity coordinates transcription of the bacterial genome. EMBO Rep. 2006;7(7):710-5.

71.   Benjamini Y, Hochberg Y. Controlling the false discovery rate - a practical and powerful approach

to multiple testing. J Roy Stat Soc Ser B (Stat Method). 1995;57(1):289-300.

72.   Grinsted A, Moore JC, Jevrejeva S. Application of the cross wavelet transform and wavelet

coherence to geophysical time series. Nonlinear Processes in Geophysics. 2004;11:561-6.

73.   Maraun D, Kurths J. Cross wavelet analysis: significance testing and pitfalls. Nonlinear Processes

in Geophysics. 2004;11:505-14.

Table 1. Pearson's correlation, ρ, of each strain versus MMR⁻

Strain	Description	Whole Chromosome Bins 1 to 46		Right Replichore Bins 1 to 23		Left Replichore Bins 46 to 24		Origin Bins 1 to 13, 46 to 34		Terminus Bins 14 to 33	
		ρ	P^a	ρ	P^a	ρ	P^a	ρ	P^a	ρ	P^a
Collective[b]	MMR⁻	—	—	—	—	—	—	—	—	—	—
Collective	MMR⁻ minus A:T ts at NAC/NTG	0.94	<0.001	0.93	<0.001	0.96	<0.001	0.97	<0.001	0.71	0.004
PFM421	Δ*rnhA* Δ*mutL*	0.74	<0.001	0.63	0.002	0.84	<0.001	0.81	<0.001	0.47	0.075
PFM669	Δ*mutL* (AB1157)	0.59	<0.001	0.40	0.064	0.75	<0.001	0.71	<0.001	0.45	0.079
PFM430/ 431	*oriC⁺ oriZ⁺* Δ*mutL* (AB1157)	0.75	<0.001	0.71	<0.001	0.81	<0.001	0.83	<0.001	0.46	0.075
PFM426	Δ*oriC oriZ⁺* Δ*mutL* (AB1157)	0.39	0.008	0.21	0.34	0.62	0.003	0.40	0.044	0.41	0.11
PFM533/ 534	Δ*seqA* Δ*mutL*	0.49	0.001	0.71	<0.001	0.27	0.23	0.53	0.006	0.38	0.14
PFM5[b]	Δ*mutL* on minimal medium	0.50	0.001	0.40	0.069	0.61	0.003	0.57	0.003	0.58	0.029

PFM343[b]	Δ*mutS* on minimal medium	0.18	0.22	0.18	0.41	0.18	0.41	0.21	0.31	0.22	0.38
PFM343[b]	Δ*mutS* on diluted LB	0.84	<0.001	0.84	<0.001	0.85	<0.001	0.87	<0.001	0.71	0.004
PFM343[b]	Δ*mutS* on supplemented minimal medium	0.81	<0.001	0.82	<0.001	0.81	<0.001	0.83	<0.001	0.65	0.008
PFM342[b]	Δ*mutS* at low temperature	0.57	<0.001	0.60	0.003	0.53	0.013	0.69	<0.001	-0.02	0.95
PFM799	Δ*nrdR* Δ*mutL*	0.49	0.001	0.58	0.005	0.39	0.080	0.69	<0.001	-0.31	0.24
PFM677	Δ*rep* Δ*mutL*	0.48	0.001	0.49	0.022	0.48	0.027	0.62	0.001	-0.12	0.67
PFM256	Δ*tus* Δ*mutL*	0.82	<0.001	0.79	<0.001	0.85	<0.001	0.87	<0.001	0.70	0.004
PFM257	Δ*matP* Δ*mutL*	0.40	0.007	0.60	0.004	0.21	0.34	0.70	<0.001	-0.47	0.075
PFM422	Δ*recA* Δ*mutL*	0.71	<0.001	0.70	<0.001	0.73	<0.001	0.85	<0.001	0.06	0.84
PFM424	Δ*recA* Δ*mutS*	0.80	<0.001	0.78	<0.001	0.83	<0.001	0.91	<0.001	0.44	0.082
PFM456	Δ*recB* Δ*mutL*	0.82	<0.001	0.87	<0.001	0.79	<0.001	0.89	<0.001	0.39	0.12
PFM118	Δ*umuDC* Δd*inB* Δ*mutL*	0.59	<0.001	0.62	0.003	0.55	0.010	0.61	0.001	0.69	0.004
PFM120	*lexA3* Δ*sulA* Δ*mutL*	0.82	<0.001	0.82	<0.001	0.83	<0.001	0.83	<0.001	0.80	0.001
PFM259	Δ*hupB* Δ*mutL*	0.73	<0.001	0.70	<0.001	0.77	<0.001	0.81	<0.001	0.57	0.031
PFM258	Δ*hupA* Δ*mutL*	0.49	0.001	0.43	0.053	0.58	0.005	0.67	<0.001	-0.23	0.38
PFM317/ 318	Δ*fis* Δ*mutL*	0.51	<0.001	0.61	0.003	0.42	0.060	0.67	<0.001	-0.30	0.25
PFM482	Δ*fis* Δ*mutS*	0.49	0.001	0.31	0.15	0.62	0.003	0.65	<0.001	-0.28	0.27

PFM741	Δhns ΔmutL	0.81	<0.001	0.81	<0.001	0.81	<0.001	0.88	<0.001	0.46	0.075
PFM713	Δdps ΔmutL	0.80	<0.001	0.84	<0.001	0.78	<0.001	0.89	<0.001	0.47	0.075
PFM163[c]	mutD5	0.76	<0.001	0.79	<0.001	0.74	<0.001	0.82	<0.001	0.78	0.001
PFM165/397/399[c]	mutD5 ΔmutL	0.64	<0.001	0.76	<0.001	0.52	0.014	0.86	<0.001	-0.46	0.075
Collective[b]	Wild type	0.36	0.014	0.41	0.061	0.31	0.17	0.37	0.065	0.51	0.060
Collective[d]	Bacillus subtilis mutS::Tn10	0.52	<0.001	0.63	0.003	0.40	0.068	0.74	<0.001	-0.52	0.055

[a] P values were adjusted for multiple comparisons using the Benjamini-Hochberg method (71) with the false discovery rate set at 25%

[b] Data are from (11)

[c] Data are from (10)

[d] Data are combined from four MA experiments with DK2140 (NCIB3610 mutS::Tn10), DK2141 (NCIB3610 mutS::Tn10), DK2142 (NCIB3610 mutS::Tn10 ΔcomI), and DK2143 (NCIB3610 mutS::Tn10 ΔcomI). There were no differences in mutation rates or spectra among these strain

Table 2. Pearson's correlation, ρ, of right versus left replichore for each strain

Strain	Description	Whole Chromosome Bins 1 to 23, 46 to 24		Origin Bins 1 to 13, 46 to 34		Terminus Bins 14 to 23, 33 to 24	
		ρ	P[a]	ρ	P[a]	ρ	P[a]
Collective[b]	MMR⁻	0.91	<0.001	0.95	<0.001	0.66	0.23
Collective	MMR⁻ minus A:T ts at NAC/NTG	0.89	<0.001	0.94	<0.001	0.42	0.60
PFM421	ΔrnhA ΔmutL	0.49	0.026	0.58	0.051	0.24	0.82
PFM669	ΔmutL (AB1157)	0.52	0.018	0.64	0.026	0.29	0.77
PFM430/ 431	oriC⁺ oriZ⁺ ΔmutL (AB1157)	0.60	0.005	0.80	0.003	0.19	0.84
PFM426	ΔoriC oriZ⁺ ΔmutL (AB1157)	0.28	0.22	0.33	0.31	-0.06	0.91
PFM533/ 534	ΔseqA ΔmutL	0.41	0.062	0.53	0.079	0.32	0.71
PFM5[b]	ΔmutL on minimal medium	0.27	0.22	0.19	0.53	0.05	0.91
PFM343[b]	ΔmutS on minimal medium	-0.02	0.93	0.32	0.31	-0.53	0.39
PFM343[b]	ΔmutS on diluted LB	0.87	<0.001	0.93	<0.001	0.66	0.23
PFM343[b]	ΔmutS on supplemented minimal medium	0.74	<0.001	0.79	0.003	0.54	0.39
PFM342[b]	ΔmutS at low temperature	0.84	<0.001	0.89	<0.001	0.73	0.19
PFM799	ΔnrdR ΔmutL	0.79	<0.001	0.78	0.003	0.84	0.07
PFM677	Δrep ΔmutL	0.43	0.053	0.58	0.051	-0.16	0.84
PFM256	Δtus ΔmutL	0.70	0.001	0.88	<0.001	0.26	0.81
PFM257	ΔmatP ΔmutL	0.45	0.042	0.40	0.20	0.72	0.19

PFM422	*ΔrecA ΔmutL*	0.66	0.001	0.81	0.003	0.11	0.88
PFM424	*ΔrecA ΔmutS*	0.64	0.002	0.80	0.003	0.07	0.91
PFM456	*ΔrecB ΔmutL*	0.67	0.001	0.78	0.003	0.03	0.94
PFM118	*ΔumuDC ΔdinB ΔmutL*	0.39	0.075	0.68	0.018	-0.13	0.87
PFM120	*lexA3 ΔsulA ΔmutL*	0.51	0.019	0.57	0.055	0.42	0.60
PFM259	*ΔhupB ΔmutL*	0.64	0.002	0.78	0.003	0.35	0.65
PFM258	*ΔhupA ΔmutL*	0.54	0.014	0.69	0.015	0.15	0.84
PFM317/ 318	*Δfis ΔmutL*	0.61	0.004	0.67	0.019	0.58	0.34
PFM482	*Δfis ΔmutS*	0.34	0.13	0.49	0.108	0.19	0.84
PFM741	*Δhns ΔmutL*	0.81	<0.001	0.85	0.001	0.60	0.33
PFM713	*Δdps ΔmutL*	0.52	0.017	0.77	0.004	-0.20	0.84
PFM163[c]	*mutD5*	0.75	<0.001	0.77	0.004	0.52	0.39
PFM165/397/399[c]	*mutD5 ΔmutL*	0.78	<0.001	0.88	<0.001	0.39	0.62
Collective[b]	Wild type	0.17	0.46	0.24	0.45	0.17	0.84
Collective[d]	*Bacillus subtilis mutS::*Tn*10*	0.73	<0.001	0.87	0.000	0.37	0.65

[a] P values were adjusted for multiple comparisons using the Benjamini-Hochberg method (71) with the false discovery rate set at 25%

[b] Data are from (11)

[c] Data are from (10)

Figure Legends

Fig. 1. The BPS density pattern is not due to transcription but is affected by moving the origin of replication. For each plot the data were collected into 100 Kb bins starting at the origin of replication on the left and continuing clockwise around the chromosome back to the origin on the right. **Fig. 1A**. The BPS density pattern of MMR-defective strains. Bars represent the mean mutation rate in each bin calculated from the BPSs collected from 10 experiments with MMR-defective strains. Error bars are the 95% CLs of the means. The grey line represents the Daubechies wavelet transform of the binned data. The chromosomal macrodomains (MCs) are indicated at the top of the plot: green, Ori MC; blue, Right and Left MC; red, Ter MC. **Fig. 1B**. The Daubechies wavelet transform of the binned reads from RNA-Seq samples taken during lag, log, and stationary growth phases of the Δ*mutL* mutant strain, PFM144. The data used for the wavelet transforms were the means of three biological replicates. Arrows indicate the positions of the rRNA operons, but the reads from the ribosomal genes in those operons were not included in the plot. **Fig. 1C, D, E, and F**. The BPS density patterns (bars) and Daubechies wavelet transforms (pink line) of the indicated mutant strains compared to the Daubechies wavelet transform of the combined MMR⁻ strains (grey line). See text for a description of the strains. In each plot the left-hand scale has been adjusted to bring the wavelets close together for comparison. The red arrow in 1E indicates the position of *oriZ*. Strains: 1C, PFM421; 1D, PFM430/431; 1E, PFM426; 1F, PFM533/534.

Fig. 2. Altering the progression of DNA replication affects the BPS density pattern. For each plot the data were collected into 100 Kb bins starting at the origin of replication on the left and continuing clockwise around the chromosome back to the origin on the right. The BPS density

patterns (bars) and Daubechies wavelet transforms (pink lines) of the indicated strains are

compared to the Daubechies wavelet transform of the combined MMR⁻ strains (grey line). In

each plot the left-hand scale has been adjusted to bring the wavelets close together for

comparison. **Fig. 2A, B, C, and D**. Altering the cellular growth rate by growing a Δ*mutS* mutant

strain on different medium and at different temperatures differentially affects the BPS density

pattern. In the experiments shown in Plot A, C, and D, the cells grew at about half the normal

rate, whereas in the experiment shown in plot B they grew at the normal rate. Strains: 2A, 2B,

and 2C, PFM343; 2D, PFM342. **Fig. 2E and F**. Dysregulation of dNTP levels and loss of Rep, the

auxiliary replication helicase, affect the BPS density pattern. 2E, PFM799; 2F, PFM677.

Fig. 2G and H. Loss of the Tus antihelicase has no effect, but loss of the terminus organizing

protein, MatP, changes the BPS density pattern across the chromosome. The arrows in Fig. 2G

mark the major Ter sites where Tus binds. The bar in Fig. 2H shows the region in which MatP

binds. Strains: 2G, PFM256; 2H, PFM257.

Fig. 3. Loss of RecA, HUα, or Fis changes the BPS pattern, but the SOS response and loss of

other NAPs do not. For a description of the plots, see the legend to Fig. 2. **Fig. 3A, B, and C.** Loss

of RecA alters the BPS density pattern in the terminus region, but not through its role as a

master regulator of the SOS response. RecA (Fig. 3A) is *E. coli*'s major recombinase. The *dinB*

and *umuDC* genes (Fig. 3B) encode the error-prone DNA polymerases, Pol IV and Pol V, that are

induced as part of the SOS response. The *lexA3* allele encodes a non-inducible repressor of the

SOS genes. Deletion of *sulA* prevents lethal filamentation after induction of the SOS response

(not relevant to this study). Strains: 3A, PFM422; 3B, PFM118; 3C, PFM120. **Fig. 3D, E, F, G, and**

H. Loss of the HUα subunit of HU or of Fis changes the BPS density pattern, but loss of the HUβ

subunit of HU, HNS, or DPS has only minor effects. Strains: 3D, PFM259; 3E, PFM258; 3F,

PFM317/318; 3G, PFM741/742; 3H, PFM713.

Fig. 4. The BPS density pattern is a result of biased correction by replication proofreading and

MMR. For a description of the plots, see the legend to Fig. 2. **Fig. 4A and B**. A strain with

deficient proofreading but active MMR yields a BPS pattern similar to a MMR-defective strain,

but a strain with neither proofreading nor MMR does not. The *mutD5* allele encodes an

exonuclease deficient proofreader. Strains: 4A, PFM163; 4B, PFM165/397/399. **Fig. 4C and D.**

The BPS density pattern in the wild-type strain does not match the pattern of either the MMR-

defective strain, or the *mutD5* mutant strain. The green line in Fig. 1D is the Daubechies

wavelet transform of the *mutD5* mutant strain (pink line in Fig. 4A). Strains: 4C and D, eight

strains with wild-type mutational phenotypes (see text). **Fig. 4E**. MMR-defective *B. subtilis* also

has a symmetrical BPS density pattern, but it is different than *E. coli*'s pattern. Strains: three *B.*

subtilis mutant strains with the same mutational phenotype (see Additional file 2: Tables S1 and

S3).

Figure 1

Figure 2

Figure 3

Figure 4