Mosquito and primate ecology predict human risk of yellow fever virus spillover in Brazil

Marissa L. Childs[1]*, Nicole Nova[2], Justine Colvin[2], Erin A. Mordecai[2]

[1]Emmett Interdisciplinary Program in Environment and Resources, Stanford University, Stanford, CA 94305, USA
[2]Biology Department, Stanford University, Stanford, CA 94305, USA
*To whom correspondence should be addressed: marissac@stanford.edu, ORCID ID 0000-0002-8597-2161

Keywords: Brazil, disease ecology, mosquito, pathogen spillover, vector-borne disease, yellow fever

Abstract

Many (re)emerging infectious diseases in humans arise from pathogen spillover from wildlife or

livestock, and accurately predicting pathogen spillover is an important public health goal. In the

Americas, yellow fever in humans primarily occurs following spillover from non-human

primates via mosquitoes. Predicting yellow fever spillover can improve public health responses

through vector control and mass vaccination. Here, we develop and test a mechanistic model of

pathogen spillover to predict human risk for yellow fever in Brazil. This environmental risk

model, based on the ecology of mosquito vectors and non-human primate hosts, distinguished

municipality-months with yellow fever spillover from 2001 to 2016 with high accuracy (AUC =

0.72). Incorporating hypothesized cyclical dynamics of infected primates improved accuracy

(AUC = 0.79). Using boosted regression trees to identify gaps in the mechanistic model, we

found that important predictors include current and one-month lagged environmental risk,

vaccine coverage, population density, temperature, and precipitation. More broadly, we show

that for a widespread human viral pathogen, the ecological interactions between environment,

vectors, reservoir hosts, and humans can predict spillover with surprising accuracy, suggesting

the potential to improve preventative action to reduce yellow fever spillover and prevent onward

epidemics in humans.

Introduction

Many important (re)emerging infectious diseases in humans—including Ebola, sudden acute respiratory syndrome (SARS), influenza, *Plasmodium knowlesi* and other primate malarias, yellow fever, and leptospirosis—arise from spillover of pathogens from wildlife or livestock into human populations (1,2). While spillover is an important mechanism of human disease emergence, the drivers and dynamics of spillover are poorly understood and difficult to predict (3). Pathogen spillover requires favorable conditions to align in the reservoir (non-human animal), human, and pathogen populations and in the environment (3–5). Because these conditions interact, nonlinear relationships among the environment, host populations, and spillover probability are likely to emerge. Moreover, spillover is a probabilistic process that does not always occur, even when suitable conditions align. Despite these challenges, it is critical to predict pathogen spillover to enhance public health preparedness. Predicting spillover also provides an opportunity to test ecological approaches to solving globally important human health problems.

Most previous attempts to predict pathogen spillover have used statistical models (6–8). These models may be locally accurate for within-sample prediction, but may struggle to detect multidimensional, nonlinear, and stochastic relationships among host populations, pathogens, the environment, and spillover. In contrast, mechanistic models can test our understanding of transmission ecology, reproduce the complex, nonlinear interactions emerging in disease systems, and potentially improve our ability to predict spillover. In particular, Plowright *et al.* (3) recently proposed a mechanistic model, which remains untested, that integrates multiple ecological requirements to identify when conditions will align for pathogen spillover. Yellow fever in Brazil presents an ideal opportunity to test this model because the ecology of the

pathogen has been studied for nearly 120 years (9), providing a wealth of mechanistic

information and data, and because almost all recent cases in South America have occurred via

spillover from the sylvatic cycle (10,11)

Yellow fever virus is a mosquito-borne *Flavivirus* that mainly persists in a sylvatic

transmission cycle between forest mosquitoes (primarily *Haemagogus janthinomys, Hg.*

leucocelaenus, and *Sabethes chloropterus* in South America) and non-human primates, and

occasionally spills over into human populations (12). In some settings, these spillover events

lead to onward human epidemics in an urban transmission cycle between humans and *Aedes*

aegypti mosquitoes (9). Spillover of yellow fever requires the virus to be transmitted locally,

mosquito vectors to acquire the virus from infected non-human vertebrate hosts, survive the

extrinsic incubation period, and feed on human hosts, and human hosts to be susceptible to

infection following exposure. These events require distributions of reservoirs, vectors, and

humans, their interactions, and immune dynamics to align in space and time. In humans, yellow

fever is the most severe vector-borne virus circulating in the Americas (10) with an estimated

fatality rate for severe cases of 47% (13). While no urban transmission of yellow fever has

occurred in the Americas since 1997 (14) and in Brazil since 1942 (15), a large epidemic began

in December 2016 in Minas Gerais and by June 2018 had caused 2,154 confirmed cases and 745

deaths (16). Despite these large case numbers, molecular and epidemiological evidence suggests

that human cases were caused by spillover from the sylvatic cycle, rather than urban

transmission (11), most recently in areas previously believed to be free of yellow fever.

Prior statistical models have found climate and weather (including precipitation,

temperature, and normalized difference vegetation index), non-human primate richness, land use

intensiveness, and a latitudinal gradient to be predictive of the spatial and spatio-temporal

distribution of yellow fever (6,8). We build on previous efforts by incorporating a mechanistic

understanding of how ecological and human population factors affect yellow fever transmission

and spillover. A mechanistic model allows for known relationships between the environment and

transmission mechanisms, estimated from empirical data, to be included to test our

understanding of the disease ecology. Additionally, mechanistic models allow for extrapolation

beyond known regions to identify other regions where conditions are also suitable for yellow

fever spillover. We use a mechanistic model encapsulating sylvatic yellow fever ecology to

predict the spatial and temporal distribution of yellow fever spillover in Brazil, and we test the

model on human yellow fever case data using a receiver operating characteristic curve and

logistic regression. Here, we use "predict" to refer to independently estimating spillover risk

mechanistically from simultaneous covariates and "forward prediction" to refer to estimating

future spillover. We contrast this mechanistic prediction with statistical models that are fit to the

spillover data, and therefore not able to make independent, out-of-sample predictions. We then

incorporate the mechanistic model into further statistical analyses with boosted regression trees

to understand what mechanisms our model does not capture.

Specifically, we ask: (1) Does the environmental suitability for sylvatic vectors, reservoir

hosts, vector-human contact, and vector transmission—together termed environmental risk—

predict geographic, seasonal, and interannual variation in yellow fever virus spillover into

humans? (2) Are human population size and vaccine coverage, above and beyond environmental

risk, critical for predicting spillover? (3) What additional environmental and population drivers

might improve predictions of spillover? (4) Do the ecological processes that predict spillover in

other parts of Brazil predict the recent yellow fever outbreak in the Southeast region of Brazil in

2016–2018, and if so, was risk elevated above historical baseline levels?

Methods

Our goals were (1) to construct mechanistic estimates of yellow fever spillover risk over space and time, (2) to test these mechanistic risk models against observed cases of yellow fever spillover to humans, and (3) to statistically test for associations between observed spillover occurrence, mechanistically predicted risk, and environmental covariates to identify potential gaps in the mechanistic models. We constructed mechanistic risk estimates by modeling the ecological processes expected to drive transmission within reservoir hosts—vector distribution and seasonal abundance, vector dispersal, vector infectiousness, vector survival, vector–reservoir contact, and reservoir host distributions—and the risk of spillover to humans—human population density, vector–human contact rates, and human susceptibility (Fig. 1, Mechanistic model). For each of these ecological or human population factors, we parameterized a submodel using data from the literature and remotely-sensed covariates (Fig. 1 lists data sources and Fig. 2 shows the data and/or fitted submodels). We modeled several different risk metrics, as described below (see Methods: Spillover model). We then predict monthly risk of yellow fever spillover from the component submodels for each 1 km × 1 km pixel from December 2000 to December 2016 (Fig. 1; Supplementary Materials, Section 1.1). The risk estimates from January 2001 to December 2016 were aggregated to a municipality-level estimate to compare to available reports of human cases. Next, to test for relationships that were absent or mis-specified in our mechanistic model, we used both current and lagged aggregated municipality-wide environmental risk from December 2000 to December 2016 as covariates in a statistical model (a boosted regression tree) along with other environmental and demographic covariates to identify the traits of municipalities and months where yellow fever spillover occurred during the available human case data from 2001 to 2016 (Fig. 1, Statistical model). Finally, we sought to identify whether

the mechanistic models predicted high suitability for spillover during the recent outbreaks

(December 2016 – April 2018) (16). Given the limited time range of some covariates, we

extrapolate model covariates for 2017 and 2018 by assuming that they were identical to 2016 or

followed the same linear trend as was observed from 2015 to 2016. We then calculate the

environmental risk metric for January 2017 to June 2018 in the region where the large outbreak

occurred.

<u>Spillover model</u>

Yellow fever spillover risk is first estimated monthly from December 2000 to December

2016 using an adapted version of the model from Plowright *et al.* (3). We then estimate monthly

spillover risk using extrapolated covariates (Supplementary Materials, Table S1) for the duration

of the 2016 – 2018 outbreaks. We define environmental risk at a location $\vec{y}$ and time t —

proportional to the number of infectious mosquito bites—as:

$b(\vec{y},t)\beta_h(\vec{y},t)\int_{\tau=0}^{\tau=t}\int_{\vec{x}}\rho_v(\vec{x},\tau)b(\vec{x},\tau)\beta_p(\vec{x},\tau)\kappa(\vec{x},\tau)EIP(T(\vec{x},\tau),t-\tau)s(T(\vec{x},\tau),t-\tau)d(\|\vec{y}-\vec{x}\|)dxd\tau$ (1)

as a function of sylvatic vector density ($\rho_v(\vec{x},t)$, Fig. 2b and 2e), probability of biting non-human

primates ($\beta_p(\vec{x},t)$, Fig. 2c) contingent on primate presence (Fig. 2a), probability of biting humans

($\beta_h(\vec{y},t)$, Fig. 2d) which depends on human presence (Fig. 2j), non-human primate infection

prevalence ($\kappa(\vec{x},\tau)$, Fig. 2k), vector biting rate ($b(\vec{y},t)$), vector probability of becoming infectious

($EIP(T(\vec{x},t),t-\tau)$, Fig. 2h), vector survival ($s(T(\vec{x},\tau),t-\tau)$, Fig. 2g), and vector dispersal

($d(\|\vec{y}-\vec{x}\|)$, Fig. 2f), as described in Table 1. This model is a case study of a more general family

of percolation models of pathogen spillover with alternative pathogen sources in space and time

(Washburne *et al.*, this issue).

We hypothesized that yellow fever spillover could be limited by environmental

conditions, human susceptibility, human population distribution, and primate infection dynamics.

To compare their relative importance, we define four metrics of model-predicted yellow fever

spillover risk. First, we approximate *environmental risk* (Eq. 1, Fig. 2l), assuming that biting rate

($b(\vec{y}, t)$ in Eq. 1,) and reservoir infection prevalence ($\kappa(\vec{x}, \tau)$ in Eq. 1) are constant over space and

time in the absence of empirical data on these parameters, as described in Table 1. Since this

metric ignores variation in human susceptibility, we then calculate *immunological risk* (Fig. 2m)

as environmental risk multiplied by the estimated proportion of the human population that is

susceptible to yellow fever (Fig. 2i), using previously estimated vaccine coverage rates (17). We

then consider the influence of human population size on spillover risk by calculating *population-*

scaled risk (Fig. 2n) as the immunological risk scaled by the number of people in a given

location (Fig. 2j). Finally, we incorporate the effects of cycles of reservoir susceptibility and

infection dynamics, for which data are not available, by calculating *periodic risk* (Fig. 2o), which

uses a phenomenological periodic curve (Fig. 2k) for primate infection prevalence ($\kappa(\vec{x}, \tau)$ in Eq.

1). This periodic curve is designed to represent cycles of reservoir infection prevalence, driven

by the demography of primate populations as naïve individuals are born, susceptible individuals

accumulate, and epizootics become more likely (18). The full spillover model was run in Google

Earth Engine (19). We estimate risk metrics monthly for 1 km × 1 km pixels using built-in

functionality of Google Earth Engine that allows for calculations across differing scales by

performing calculations for a specified output pixel scale.

<u>Mechanistic submodels</u>

We fit mechanistic submodels from data for all key components of spillover (Fig. 1). For

primate distribution (Fig. 2a), human susceptibility (Fig. 2i), and human population distribution

(Fig. 2j), we used previously published estimates (17,20,21). All other mechanistic models

(terms in Eq. 1) were fit with the R programming language, version 3.5.1 (22), with additional

packages used for data processing, manipulation, and visualization (23–31).

Given limited information on the vector species, we use data for *Hg. janthinomys, Hg.*

leucocelaenus, and *Sa. chloropterus* to fit models for the sylvatic vectors collectively for all

mechanistic vector trait models. All data used were publicly available or results from previously

published papers, as described in the Supplementary Materials (Table S1 and Mechanistic

submodel details). Additional details on mechanistic model methods and data are available in the

Supplementary Materials.

Vector distribution and seasonal density

To estimate the geographic distribution of sylvatic vector species (Fig. 2b), we fit a

species distribution model (32,33) to *Hg. janthinomys, Hg. leucocelaenus,* and *Sa. chloropterus*

occurrence data identified from the Global Biodiversity Information Facility (GBIF) (34,35) and

a review of the literature (36–90), using the maxnet package in R (91). We included maximum,

median, and minimum annual land surface temperature, total annual precipitation, precipitation

in the driest month, precipitation in the wettest month, elevation, forest cover (%), land cover

category, median annual enhanced vegetation index, and absolute latitude as predictors in the

model (Supplementary Materials, Table S2). To account for uneven sampling effort across the

geographic range, we corrected the background (pseudo-absence) points by subsampling from

occurrence data of other mosquito species from GBIF (92). We calculated vector density as

$\log\left(\frac{1}{1-p}\right)$, where p is the probability of occurrence estimated from the species distribution model

(93). To estimate seasonal variation in vector abundance (Fig. 2e) due to rainfall seasonality

(94), we fit a logistic regression of relative monthly vector abundance on current and one-month

lagged relative monthly rainfall using field data (80,95–99) with glm in R.

Vector survival

To capture effects of temperature on vector survival (Fig. 2f), we used empirical data

(100–102) and Bayesian inference to fit a quadratic function to the relationship between lifespan

and temperature using rstan in R (103). Assuming constant vector mortality at a given

temperature, we calculated daily survival probability as $p = e^{-1/L}$, where L is vector lifespan

(104).

Vector infectiousness

Virus infection, dissemination, and infectiousness in the vector are temperature-

dependent (Fig. 2h) (105). We assume that vector competence—the probability that a vector

exposed to an infectious blood meal becomes infectious with virus in its salivary glands—is a

quadratic function of temperature, as shown for other flaviviruses (106). Additionally, we

assume that at a given temperature, the extrinsic incubation period—the length of time required

for an exposed vector to become infectious—is log-normally distributed across individuals

(107,108). We fit a Bayesian model using experimental data (109–116) with the package rstan

(103).

Vector dispersal

To estimate the range on which sylvatic mosquitoes disperse (Fig. 2f), we fit a negative

binomial dispersal kernel (117) to mark-recapture data (118) using a Bayesian framework with

the package rstan (103).

Vector contacts

We approximated reservoir-vector contact (Fig. 2c) as percent forest cover (119)

contingent on the presence of at least one reservoir species (Fig. 2a). Similarly, we approximated

human-vector contact (Fig. 2d) as percent non-forest cover (119) contingent presence of human

population (Fig. 2j).

Phenomenological primate dynamics

Primate population dynamics and susceptibility have been suggested as important

constraints on yellow fever spillover (18), which remain poorly characterized. In the absence of

primate infection data, we assumed that human spillover events are a proxy for infection

prevalence during reservoir epizootics. This is the only mechanistic submodel that uses the

human yellow fever spillover data directly—all other submodels are independent of human

infection data. For this submodel, we used human cases of yellow fever reported by month of

first symptoms and municipality of infection (2001–2016) from the Brazilian Ministry of Health

(120). We define a spillover municipality-month as one in which at least one human case of

yellow fever occurred. As an estimate of reservoir infection dynamics, we fit a

phenomenological sine curve with a seven year period (121) to the yearly number of

municipality-months with spillover (Fig. 2k) and then transformed the curve to be positive and

less than one. The resulting curve is used as a spatially constant estimate of primate reservoir

infection prevalence. Phenomenological primate dynamics are used in the periodic risk estimate

(Fig. 2o) to account for a missing ecological process but are not used in any other risk metric, so

all other risk metrics are parameterized independent of human spillover data.

Model-data comparison

We compared spatially- and temporally-explicit mechanistic model predictions for

spillover risk to observed human cases of yellow fever spillover using a statistical model. We

limit the comparison to 2001–2016 based on the availability of human case data. We considered

four modeled risk metrics (defined above): environmental risk, immunological risk, population-

scaled risk, and periodic risk. Because risk was modeled by pixel, to compare the model output

with municipality-month observations of human cases, we calculate both mean risk and

maximum risk in each municipality and month. While mean risk may be more representative of

the entire municipality, we hypothesized that maximum risk in the municipality-month might

better predict the small-scale processes that drive spillover. The use of maximum risk also may

also help to avoid spatial aggregation that can lead to bias or mask the relationships, for example

the modifiable areal unit problem (122).

We compared municipality means and maxima for all four risk metrics to human yellow

fever data for model evaluation in the following three ways. First, for each modeled risk metric

and each municipality summary statistic (mean and maximum), we fit a logistic regression of

spillover probability as a function of model-predicted risk (Supplementary Materials, Table S4)

using glm in R (22). Second, we calculated a receiver operating characteristic curve to calculate

the area under the curve (AUC), a measure of goodness of fit, for each modeled risk metric and

municipality summary statistic (Supplementary Materials, Table S4). As this analysis focuses on

prediction of spillover as a way to compare hypothesized mechanisms, comparison of AUC

values to a null model is beyond the scope of this paper. Finally, for all eight mechanistic

predictions and estimated vaccine coverage, we regressed the number of reported yellow fever

cases given that spillover occurred and calculated Spearman's rank correlation coefficient with

number of reported cases to consider nonlinear but monotonic associations (Supplementary

Materials, Table S5).

Statistical model

We used a boosted regression tree (123,124) to understand any potential gaps in the

mechanistic model and its relationship to environmental and human population covariates. As

predictors of yellow fever spillover in the boosted regression tree, we included the following

covariates for each municipality-month: current and one-month lagged maximum predicted

environmental risk, current and one-month lagged fire area, average and maximum number of

primate species, estimated municipality vaccine coverage, average human population density,

average monthly air temperature, average monthly precipitation, phenomenological primate

dynamics, region, and month (Supplementary Materials, Table S6). Each observation is a

municipality-month and the response variable is the binary indicator of whether or not yellow

fever spillover occurred in a municipality-month (see Methods: Model-data comparison). While

some of predictor covariates contribute to the environmental risk metric (i.e., air temperature,

rainfall, and primate reservoir ranges), we also include them in the boosted regression tree

analysis to identify whether the environmental covariates have any predictive power beyond their

role in the mechanistic model, which could indicate that the mechanistic model does not fully

capture their influence on spillover. We included fire area as a proxy for land conversion (125),

which has previously been shown to be predictive of yellow fever spillover (8). We also included

vaccine coverage and human population density despite their poor predictive performance in the

mechanistic model to identify whether these human population factors are predictive of spillover

in ways not previously hypothesized, and therefore not captured in the mechanistic model.

Boosted regression trees repeatedly fit regression trees, which create multiple binary splits in the

dataset based on predictor variables. Each successive tree is fit to the residuals of the previous

best model. The model is then updated to include the next tree (123). Variable importance is

calculated as a weighted sum of the number of times a variable is used for splitting, with weights

determined by the squared improvement due to the split (123).

We fit the boosted regression tree to data from 2001 to 2016, given this is the range of the

available human case data for inferring spillover. We partition the dataset into spatially- and

temporally-balanced training (80%) and test (20%) sets prior to the analysis. Optimal learning

rate, tree complexity, and number of trees were selected as the set of parameters that minimized

cross-validation predictive deviance (Supplementary Materials, Table S7; 63). The dataset was

split in R using the BalancedSampling package (126), models were fit in R using the gbm and

dismo (127,128) packages, and variable effects were calculated with the pdp package (129).

Additional details can be found in the Supplementary Materials.

Results

Primate species distribution (Fig. 2a), vector distribution (Fig. 2b; Supplementary

Materials, Fig. S1 and S2), reservoir-vector contact (Fig. 2c), human-vector contact (Fig. 2d),

and human susceptibility (Fig. 2i) varied over space and time based on estimates and models fit

to empirical data. In addition, vector survival (Fig. 2g) and infectiousness (Fig. 2h,

Supplementary Materials, Fig. S4 and S5) varied with temperature, vector abundance varied

seasonally with rainfall (Fig. 2e; Supplementary Materials, Fig. S3 and Table S3), and vector

dispersal declined exponentially with distance (Fig. 2f; Supplementary Materials, Fig. S6).

Together, these empirical relationships between environment and host, vector, and virus ecology

compose an estimate of environmental risk of yellow fever spillover (Supplementary File 2).

The environmental risk model strongly predicted episodes of yellow fever spillover into

humans (AUC = 0.72) and adding phenomenological reservoir infection dynamics in periodic

risk further improved the model (AUC = 0.79; Fig. 3). Surprisingly, models that included human

vaccination coverage and human population size performed worse than the environment-driven

models (AUC = 0.64 and 0.64; Fig. 3). For all risk metrics, maximum value in the municipality-

month was a better predictor of spillover than mean value (Fig. 3). Logistic regressions of

spillover probability as a function of model-predicted risk showed similar patterns in Akaike

Information Criterion (AIC) values (Supplementary Materials, Table S4). Model-predicted

environmental (mean and maximum), periodic (mean and maximum), and immunological

(maximum) risk metrics were statistically significant predictors of spillover probability at the 5%

level after correcting for multiple hypothesis testing (Supplementary Materials, Table S4; 71).

By contrast, given that spillover occurred, none of the eight mechanistic model risk summaries

were statistically significant predictors of number of cases, nor was estimated vaccine coverage

(Supplementary Materials, Table S5). In Spearman's rank correlations, we find that of the

independent predictors, vaccine coverage is most correlated with risk, followed by maximum

environmental risk, although these correlations are weak (Supplementary Materials, Table S5).

Mechanistic model estimates matched seasonal variation in spillover, and accurately

captured differences in seasonality by region (Fig. 4a). Risk peaked in April in the North and

Northeast regions and in February in Central-West, South, and Southeast regions. The seasonal

regional correlation between number of municipality-months with spillover and average

environmental risk was highest in the Southeast (0.77), followed by the South (0.61), Central

West (0.58), and North (0.42) regions. The periodic risk matched interannual variation in

spillover (Fig. 4c), an unsurprising finding given periodic risk incorporated phenomenological

primate dynamics derived from human cases of spillover. Interannual regional correlations were

weaker than seasonal correlations but similarly highest in the Southeast (0.54), followed by the

Central-West (0.45), North (0.21), and South (0.13) regions.

The boosted regression tree found one-month lagged environmental risk and current

environmental risk to be the second and fifth most important predictors of spillover, respectively

(Fig. 5). Not surprisingly, the boosted regression tree significantly improved predictive

performance from the mechanistic model because it was trained on the human spillover data

(training AUC > 0.99, test AUC = 0.95). Vaccine coverage, temperature, population density and

precipitation were also among the six most important predictors in the boosted regression tree.

As expected, municipality-months with spillover had higher current and one-month lagged

environmental risk (Fig. 5b and e), as well as high (phenomenologically) estimated primate

infection prevalence and high primate species richness (Supplementary Materials, Fig. S8). We

find that municipality-months with spillover have low monthly average temperatures (Fig. 5a),

which may in part be due to poorly captured effects of temperature in the mechanistic model

from averaging temperature before calculating mosquito trait values for survival and

infectiousness (131). We also find that municipality-months with spillover have low rates of

precipitation (Fig. 5d), which may correspond to settings with increased human activity in the

forest, and therefore increased chance of spillover. However, current and lagged fire area,

hypothesized indicators of deforestation activity, were not significant predictors of spillover in

the boosted regression tree models (Supplementary Materials, Fig. S8).

Unexpectedly, municipality-months with spillover tended to have vaccine coverage

above 90%, suggesting that high rates of vaccine coverage do not prevent spillover from

occurring. While estimated vaccine coverage was included as a measure of human susceptibility,

it is likely capturing other patterns in the spatial distribution of spillover; regions known to

experience yellow fever spillover are likely to have high vaccination rates, while those where

spillover is rare or nonexistent are likely to have low vaccination rates. Accordingly, estimated

vaccine coverage is bimodal, potentially due to a group of lower risk municipalities and a group

of higher risk municipalities. The partial dependence plot also displays two plateaus in the

marginal effect in the vaccine coverage on model estimates, which roughly correspond to the two

vaccine coverage groups.

The recent outbreaks in Brazil in the 2016–2017 and 2017–2018 transmission seasons

have been the largest in over 50 years (16). The environmental risk model predicts persistent,

low environmental risk of spillover in the affected states (Minas Gerais, Espírito Santo, Sao

Paulo, and Rio de Janeiro) and does not predict any increase in spillover risk during the recent

transmission seasons (Fig. 6). The date ranges of confirmed human cases during the 2016–2017

and 2017–2018 outbreaks are shown in pink bands (Fig. 6) based on a World Health

Organization epidemiological update (16). The mechanistic model predicts spillover risk in

Espírito Santo and Rio de Janeiro, where no spillover occurred from 2001 to 2016, at levels

similar to those in Minas Gerais and Sao Paulo, where spillover had previously occurred. As in

other regions, the model accurately captures the seasonality of spillover risk in this region (Fig.

6), which is distinct from that of other regions (Fig. 4a).

Discussion

Our mechanistic understanding of environmental risk of spillover—which combines

reservoir host and sylvatic vector distributions, vector contact with reservoirs and humans, and

vector dispersal, survival, infectiousness, and seasonal abundance—predicts yellow fever

spillover into humans with high accuracy (AUC = 0.72; Fig. 3). Within each municipality and

month, the maximum risk, rather than the mean risk, was the best predictor of spillover

occurrence, suggesting that local heterogeneity in risk within municipalities is important for

determining spillover probability. Rainfall-driven seasonality in the vector populations and

temperature-driven seasonality in vector survival and infectiousness accurately predicted

seasonal variation in spillover (Fig. 4a). While interannual variation in risk was not well-

predicted in the environmental risk model based on climate and land cover information alone,

including phenomenologically modeled variation in primate yellow fever infection prevalence

improved predictions of year-to-year variation in spillover (AUC = 0.79).

Although we hypothesized that low vaccination coverage and high human population

density would each increase spillover risk, neither improved model accuracy for predicting

spillover in the mechanistic model (Fig. 3). However, we found that vaccine coverage was the

third most important predictor of municipality-months with spillover when allowing for a

nonlinear but generally positive relationship between coverage and spillover probability (Fig. 5).

The recent outbreak is also consistent with the ecological processes driving past spillover

in this region (Fig. 6). While environmental risk in 2016–2018 was not elevated above historical

levels (2001–2015) and spillover had not occurred in in the states of Espírito Santo or Rio de

Janeiro during the previous 15 years, it has previously occurred in Minas Gerais and Sao Paulo

states in 2001–2003 and 2008–2009. Data from the recent 2016–2018 outbreak past December

2016 are not included in the statistical models because consistent monthly municipality-scale

spillover data across the country are not available for that period.

The boosted regression tree analysis, which aimed to detect candidate drivers of spillover

that might be missing from our mechanistic model, identified vaccine coverage, current and

lagged environmental risk, temperature, population density, and precipitation as important

predictors, which together improved upon mechanistic model predictive performance of

pathogen spillover (out-of-sample AUC = 0.95). The relative importance of lagged and current

environmental risk provides evidence that the mechanistic model captures the potentially

nonlinear and interactive relationship between environmental variables that drive spillover in

mosquitoes, reservoir hosts, and humans better than the environmental variables alone. One-

month lagged environmental risk may be more important than current environmental risk for

predicting spillover because of a lag between cases and reporting. Additionally, environmental

suitability for reservoir and vectors may drive reservoir infection dynamics, causing a lag

between conditions suitable for virus amplification in the primate reservoir and vector

populations, and spillover into humans. Moreover, the relative importance of one-month lagged

environmental risk creates the potential for forward prediction of spillover. The boosted

regression tree also identified municipality-months with spillover to have low temperatures. As

mosquito thermal performance traits often have steep drop-offs at high temperatures, temperature

variation affects mosquito traits (132). Our mechanistic model using monthly average

temperature may overestimate the suitability in warm temperatures and underestimate the

suitability in cool temperatures (131), resulting in the decreasing relationship observed between

average monthly temperature and spillover in the boosted regression tree.

In a recent publication, Kaul et al. (8) also used a machine learning approach to predict

municipality-months with spillover in Brazil and similarly found rainfall and temperature to be

important predictors. However, their model also identified primate richness and fire density as

important predictors, while our boosted regression tree analysis ranked municipality average

primate richness tenth, municipality maximum primate richness fourteenth, one-month lagged

fire area ninth, and current fire area twelfth for variable importance out of fourteen variables.

Our covariates add to those used by Kaul *et al.* (8) by including vaccine coverage and our

mechanistic environmental risk estimate (current and lagged), which boosted regression trees

found to be three of the five most important predictors. We expect that our mechanistic

environmental risk estimates capture much of the variation attributed to other environmental

variables in the Kaul *et al.* model. Despite the differing relative importance of variables for

predicting spillover in the two models, they both predict that seasonal patterns vary by regions of

Brazil and find Southeast Brazil seasonally suitable for yellow fever spillover. Our mechanistic

model further illustrates that this differing seasonality can be explained by seasonal variation in

vector survival and infectiousness driven by temperature and vector abundance driven by

rainfall.

Given the importance of vaccination campaigns in limiting yellow fever outbreaks, we

expected that the number of susceptible (unvaccinated) people would be an important positive

predictor of yellow fever spillover occurrence, yet mechanistic population-scaled risk performed

worse at predicting spillover than environmental risk alone (Fig. 3). For example, scaling by

population size predicts areas of very high risk along the coast of Brazil, where environmental

risk is low, but population sizes are high. Additionally, we expected that vaccination coverage

and human density might be more predictive of the number of cases in spillover events (for

example, the recent outbreak in Southeast Brazil) than the probability of spillover occurring,

given that very low environmental suitability will be amplified in large, unvaccinated

populations. However, vaccine coverage was not a significant predictor of the number of human

cases of yellow fever given that spillover occurred (Supplementary Materials, Table S5).

Anecdotally, it is worth noting that prior to the recent large outbreak in Southeastern Brazil in

2016–2018, vaccination rates in the region were low, potentially allowing that outbreak to reach

an unusually high magnitude.

The substantial improvement in model prediction from environmental to periodic risk

(AUC = 0.72 vs. 0.79) suggests that primate population dynamics, immunity, and infection

prevalence may be a key missing component of this mechanistic model. Ongoing surveillance

efforts in Brazil are used to detect non-human primate cases of yellow fever as an advanced

warning system (133). While this advanced warning system can make a critical difference, the

recent outbreaks in Southeast Brazil displayed that in some cases this surveillance may not

provide sufficient time to respond to prevent spillover, especially in areas with high populations

and low vaccine coverage rates as were found in the Southeast. Incorporating a mechanistic

model of non-human primate infection prevalence, driven by local primate surveillance data,

could help to indicate when primate cases of yellow fever are likely, to provide additional time

for public health officials to respond. This remains a significant and potentially very fruitful gap

in our understanding of yellow fever transmission and spillover.

Vector-human contact rates are another important empirical gap in the mechanistic

model, which could further refine the relationships between land use, human occupations and

behavior, and spillover risk. We approximate human contact rates with sylvatic vectors with

percent forest cover, but the relationship is likely much more complex. The surprising decreasing

relationship between precipitation and spillover probability in the boosted regression tree (Fig.

5d) may be due to the influence of precipitation on human activities in and around forests, and

therefore its influence on human–vector contact (94). Additionally, while vector contacts depend

on biting rate of the vector and mosquito biting rates are known to depend on temperature for

other species (106,134), we assume constant biting rate in the mechanistic model due to a lack of

empirical evidence.

While it was beyond the scope of this paper, the most influential mechanisms in the

model could be further identified through sensitivity analyses of specific submodel components.

Additionally, associations between model components and spillover probability could be

estimated using the framework of percolation models (Washburne *et al.*, this issue). Finally, a

thorough uncertainty analysis could highlight the model components most in need of further

study to improve prediction of spillover.

Yellow fever is an ancient, historically important human disease that played a central role

in the discovery of mosquito transmission of pathogens and the subsequent development of

vector control as a public health measure (135). The wealth of existing knowledge about the

ecology of yellow fever virus and its sylvatic reservoir hosts and vectors allowed us to synthesize

data from 73 published papers to mathematically formalize our ecological understanding of

sylvatic transmission and spillover. Although spillover is a stochastic process that is expected to

be difficult to predict, the mechanistic model that integrates vector, human host, non-human

reservoir, and virus ecology allowed us to predict spillover with surprising accuracy. Historically

in the Americas and presently in other regions such as sub-Saharan Africa, yellow fever

regularly has entered urban transmission cycles that lead to major human epidemics. The model

framework presented here could be extended to include the ecology of different vectors, hosts,

and environments, including urban *Ae. aegypti* and more human immune interactions with other

flaviviruses, to ask intriguing questions such as: What prevents yellow fever from entering urban

transmission cycles in the Americas, where other flavivirus epidemics regularly occur? Why has

urban transmission occurred recently in Africa and not in South America? What prevents yellow

fever circulation and spillover in Southeast Asia, where sylvatic vectors and non-human primate

hosts are present and the climate is suitable? Answers to these questions would further our

understanding of the ecology of (re)emerging diseases in different parts of the world. More

fundamentally, this work provides clear evidence for the predictive power of mechanistic,

ecological models—even for rare events like pathogen spillover—and can provide useful

information to enhance public health interventions of zoonotic diseases.

Additional Information

Acknowledgements

Freya Shearer kindly provided data on vaccine coverage estimates. We thank the Defense

Advanced Research Projects Agency whose funding provided the opportunity to discuss and

develop this project at a workshop on pathogen spillover. We would also like to thank the many

previous studies who have made their data available as well as publicly available remote sensing

data that made this project possible. The manuscript was improved by comments from

anonymous reviewers.

Data Accessibility

All data and code are available on GitHub

(https://github.com/marissachilds/YellowFeverSpillover).

Authors' Contributions

MLC and EAM conceived of the project and designed the analyses. MLC, NN, and JC collected

the data and performed the analyses. NN created the artwork in Fig. 1 and 2. MLC drafted the

manuscript. EAM, NN, and JC revised the manuscript and all authors read and approved the final

manuscript.

Competing Interests

We have no competing interests.

Funding

MLC was funded by the Lindsay Family E-IPER Fellowship. NN was funded by The Bing

Fellowship in Honor of Paul Ehrlich. JC was funded by the Stanford University Biology Summer

Undergraduate Research Program. EAM was funded by the National Science Foundation

512   Ecology and Evolution of Infectious Diseases program (DEB-1518681), the Stanford University

513   Woods Institute for the Environment – Environmental Ventures Program, and the Hellman

514   Faculty Fellowship.

References

1. Woolhouse MEJ, Haydon DT, Antia R. Emerging pathogens: The epidemiology and evolution of species jumps. Trends Ecol Evol. 2005;20(5):238–44. https://doi.org/10.1016/j.tree.2005.02.009

2. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. Nature. 2008;451(7181):990–3. https://doi.org/10.1038/nature06536

3. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic spillover. Nat Rev Microbiol [Internet]. 2017 May 30;15(8):502–10. https://doi.org/10.1038/nrmicro.2017.45

4. Plowright RK, Eby P, Hudson PJ, Smith IL, Westcott D, Bryden WL, et al. Ecological dynamics of emerging bat virus spillover. Proc R Soc B Biol Sci [Internet]. 2014;282(1798):20142124–20142124. https://doi.org/10.1098/rspb.2014.2124

5. Lloyd-Smith JO, George D, Pepin KM, Pitzer VE, Pulliam JRC, Dobson AP, et al. Epidemic Dynamics at the Human-Animal Interface. Science (80-) [Internet]. 2009;326(5958):1362–7. https://doi.org/10.1126/science.1177345

6. Hamrick PN, Aldighieri S, Machado G, Leonel DG, Vilca LM, Uriona S, et al. Geographic patterns and environmental factors associated with human yellow fever presence in the Americas. PLoS Negl Trop Dis [Internet]. 2017;11(9). https://doi.org/10.1371/journal.pntd.0005897

7. Schmidt JP, Park AW, Kramer AM, Han BA, Alexander LW, Drake JM. Spatiotemporal fluctuations and triggers of ebola virus spillover. Emerg Infect Dis. 2017;23(3):415–22. https://doi.org/10.3201/eid2303.160101

8. Kaul RB, Evans M V., Murdock CC, Drake JM. Spatio-temporal spillover risk of yellow fever in Brazil. Parasit Vectors [Internet]. 2018;11(1):488. https://doi.org/10.1186/s13071-018-3063-6

9. Monath TP, Vasconcelos PFC. Yellow fever. J Clin Virol [Internet]. 2015;64:160–73. https://doi.org/10.1016/j.jcv.2014.08.030

10. Paules CI, Fauci AS. Yellow Fever — Once Again on the Radar Screen in the Americas. N Engl J Med [Internet]. 2017;376(15):1397–9. https://doi.org/10.1056/NEJMp1002530

11. Faria NR, Kraemer MUG, Hill SC, Goes de Jesus J, Aguiar RS, Iani FCM, et al. Genomic and epidemiological monitoring of yellow fever virus transmission potential. Science (80-) [Internet]. 2018 Aug 31;361(6405):894–9. https://doi.org/10.1126/science.aat7115

12. Hanley KA, Monath TP, Weaver SC, Rossi SL, Richman RL, Vasilakis N. Fever versus fever: The role of host and vector susceptibility and interspecific competition in shaping the current and future distributions of the sylvatic cycles of dengue virus and yellow fever virus. Infect Genet Evol [Internet]. 2013 Oct;19:292–311. https://doi.org/10.1016/j.meegid.2013.03.008

13. Johansson MA, Vasconcelos PFC, Staples JE. The whole iceberg: Estimating the incidence of yellow fever virus infection from the number of severe cases. Trans R Soc Trop Med Hyg. 2014;108(8):482–7. https://doi.org/10.1093/trstmh/tru092

14. Valderrama A, Díaz Y, López-Vergès S. Interaction of Flavivirus with their mosquito vectors and their impact on the human health in the Americas. Biochem Biophys Res Commun. 2017;492:541–7. https://doi.org/10.1016/j.bbrc.2017.05.050

15. de Souza RP, Foster PG, Sallum MAM, Coimbra TLM, Maeda AY, Silveira VR, et al. Detection of a new yellow fever virus lineage within the South American genotype I in

Brazil. J Med Virol [Internet]. 2010 Jan;82(1):175–85. https://doi.org/10.1002/jmv.21606

16.   Pan American Health Organization / World Health Organization. Epidemiological Update:
Yellow Fever (7 December 2018). 2018.

17.   Shearer FM, Moyes CL, Pigott DM, Brady OJ, Marinho F, Deshpande A, et al. Global
yellow fever vaccination coverage from 1970 to 2016: An adjusted retrospective analysis.
Lancet Infect Dis. 2017;3099(17):1–9. https://doi.org/10.1016/S1473-3099(17)30419-X

18.   Vasconcelos PF da C. Yellow fever in Brazil: thoughts and hypotheses on the emergence
in previously free areas. Rev Saude Publica [Internet]. 2010;44(6):1144–9.
https://doi.org/10.1590/S0034-89102010005000046

19.   Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D, Moore R. Google Earth
Engine: Planetary-scale geospatial analysis for everyone. Remote Sens Environ [Internet].
2017; https://doi.org/10.1016/j.rse.2017.06.031

20.   IUCN. The IUCN Red List of Threatened Species. Version 2018-2. 2018.

21.   Center for International Earth Science Information Network - CIESIN - Columbia
University. Gridded Population of the World, Version 4 (GPWv4): Population Count
Adjusted to Match 2015 Revision of UN WPP Country Totals. 2016.
https://doi.org/10.7927/H4SF2T42

22.   R Core Team. R: A Language and Environment for Statistical Computing [Internet].
Vienna, Austria; 2018. Retrieved from https://www.r-project.org/

23.   Wickham H. The Split-Apply-Combine Strategy for Data Analysis. J Stat Softw
[Internet]. 2011;40(1):1–29. Retrieved from http://www.jstatsoft.org/v40/i01/

24.   Wickham H, François R, Henry L, Müller K. dplyr: A Grammar of Data Manipulation
[Internet]. 2018. Retrieved from https://cran.r-project.org/package=dplyr

25.   Bache SM, Wickham H. magrittr: A Forward-Pipe Operator for R [Internet]. 2014.
Retrieved from https://cran.r-project.org/package=magrittr

26.   Bivand R, Keitt T, Rowlingson B. rgdal: Bindings for the "Geospatial" Data Abstraction
Library [Internet]. 2018. Retrieved from https://cran.r-project.org/package=rgdal

27.   Wickham H. stringr: Simple, Consistent Wrappers for Common String Operations
[Internet]. 2018. Retrieved from https://cran.r-project.org/package=stringr

28.   Hijmans RJ. raster: Geographic Data Analysis and Modeling [Internet]. 2018. Retrieved
from https://cran.r-project.org/package=raster

29.   Wickham H. ggplot2: Elegant Graphics for Data Analysis [Internet]. Springer-Verlag New
York; 2016. Retrieved from http://ggplot2org

30.   Borchers HW. pracma: Practical Numerical Math Functions [Internet]. 2018. Retrieved
from https://cran.r-project.org/package=pracma

31.   Bivand R, Rundel C. rgeos: Interface to Geometry Engine - Open Source ('GEOS')
[Internet]. 2018. Retrieved from https://cran.r-project.org/package=rgeos

32.   Phillips SJ, Anderson RP, Schapire RE. Maximum entropy modeling of species
geographic distributions. Ecol Modell [Internet]. 2006 Jan;190(3–4):231–59.
https://doi.org/10.1016/j.ecolmodel.2005.03.026

33.   Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ. A statistical explanation of
MaxEnt for ecologists. Divers Distrib. 2011;17(1):43–57. https://doi.org/10.1111/j.1472-
4642.2010.00725.x

34.   GBIF.org. GBIF Occurrence Download [Internet]. [cited 2018 Jul 13].
https://doi.org/10.15468/dl.ozsvnj

35.   GBIF.org. GBIF Occurrence Download [Internet]. [cited 2018 Jul 13].

https://doi.org/10.15468/dl.gxbxtq

36. Alencar J, de Mello CF, Gil-Santana HR, Guimarï¿½es AE, de Almeida SAS, Gleiser
RM. Vertical oviposition activity of mosquitoes in the Atlantic Forest of Brazil with
emphasis on the sylvan vector, Haemagogus leucocelaenus (Diptera: Culicidae). J Vector
Ecol [Internet]. 2016;41(1):18–26. https://doi.org/10.1111/jvec.12189

37. Alencar J, De Mello VS, Serra-Freire NM, Silva JDS, Morone F, Guimarães AÉ.
Evaluation of mosquito (Diptera: Culicidae) species richness using two sampling methods
in the hydroelectric reservoir of Simplício, Minas Gerais, Brazil. Zoolog Sci [Internet].
2012;29(4):218–22. https://doi.org/10.2108/zsj.29.218

38. Barbosa AA, Navarro-Silva MA, Calado D. Culicidae activity in a restrict forest inside
Curitiba urban area (Parana, Brazil). Rev Bras Zool. 2003;20(1):59–63.
https://doi.org/10.1590/S0101-81752003000100007

39. Cardoso JC, de Almeida MAB, dos Santos E, da Fonseca DF, Sallum MAM, Noll CA, et
al. Yellow fever virus in Haemagogus leucocelaenus and Aedes serratus mosquitoes,
Southern Brazil, 2008. Emerg Infect Dis [Internet]. 2010;16(12):1918–24.
https://doi.org/10.3201/eid1612.100608

40. Cardoso JDC, de Paula MB, Fernandes A, dos Santos E, de Almeida MAB, da Fonseca
DF, et al. Ecological aspects of mosquitoes (Diptera: Culicidae) in an Atlantic forest area
on the north coast of Rio Grande do Sul State, Brazil. J Vector Ecol [Internet].
2011;36(1):175–86. https://doi.org/10.1111/j.1948-7134.2011.00155.x

41. Chadee DD, Beier JC. Natural variation in blood-feeding kinetics of four mosquito
vectors. J Vector Ecol [Internet]. 1996;21(2):150–5. Retrieved from
https://www.scopus.com/inward/record.uri?eid=2-s2.0-
2842603690&partnerID=40&md5=dc7235ba2320e841aa683b49f301029f

42. Correa FF, Gleiser RM, Leite PJ, Fagundes E, Gil-Santana HR, Mello CF, et al. Mosquito
communities in Nova Iguaçu Natural Park, Rio de Janeiro, Brazil. J Am Mosq Control
Assoc [Internet]. 2014;30(2):83–90. https://doi.org/10.2987/13-6372.1

43. D'Oria JM, Martí DA, Rossi GC. Culicidae, province of Misiones, northeastern
Argentina. Check List [Internet]. 2010;6(1):176–9. https://doi.org/10.15560/6.1.176

44. das Virgens TM, Rezende HR, Pinto IS, Falqueto A. Fauna of mosquitoes (Diptera:
Culicidae) in Goytacazes national forest and surrounding area, State of Espírito Santo,
southeastern Brazil . Biota Neotrop [Internet]. 2018;18(1). https://doi.org/10.1590/1676-
0611-bn-2016-0250

45. De Figueiredo ML, De C Gomes A, Amarilla AA, De S Leandro A, De S Orrico A, De
Araujo RF, et al. Mosquitoes infected with dengue viruses in Brazil. Virol J [Internet].
2010;7. https://doi.org/10.1186/1743-422X-7-152

46. de Souza RP, Petrella S, Coimbra TLM, Maeda AY, Rocco IM, Bisordi I, et al. Isolation
of yellow fever virus (YFV) from naturally infected haemagogus (conopostegus)
leucocelaenus (diptera, culicidae) in São Paulo state, Brazil, 2009 . Rev Inst Med Trop
Sao Paulo [Internet]. 2011;53(3):133–9. https://doi.org/10.1590/S0036-
46652011000300004

47. Dos Santos EB, Orlandin E, Piovesan M, Favretto MA. Short communication on the
mosquitoes of a forested urban area at the municipality of Joaçaba, Santa Catarina, Brazil .
Entomotropica [Internet]. 2016;31(2016):91–4. Retrieved from
https://www.scopus.com/inward/record.uri?eid=2-s2.0-
84980004951&partnerID=40&md5=17adfc0672bed30a457cba1884c38bd7

48.   Alencar J, Gil-Santana HR, Oliveira RDFND, Dgallier N, Guimares AR. Natural breeding
654   sites for haemagogus mosquitoes (Diptera, Culicidae) in Brazil. Entomol News [Internet].
655   2010;121(4):393–6. https://doi.org/10.3157/021.121.0414
656   49.   Fernandez Z, Richartz R, Da Rosa AT, Soccol VT. Identification of the encephalitis
657   equine virus, Brazil. Rev Saude Publica. 2000;34(3):232–5.
658   https://doi.org/10.1590/S0034-89102000000300004
659   50.   Forattini OP, Gomes AD. Biting Activity of Aedes-Scapularis (Rondani) and Haemagogus
660   Mosquitos in Southern Brazil (Diptera, Culicidae). Rev Saude Publica. 1988;22(2):84–93.
661   https://doi.org/10.1590/S0034-89101988000200003
662   51.   Freitas Silva SO, de Mello CF, Figueiro R, Maia D de A, Alencar J. Distribution of the
663   Mosquito Communities (Diptera: Culicidae) in Oviposition Traps Introduced into the
664   Atlantic Forest in the State of Rio de Janeiro, Brazil. Vector-Borne Zoonotic Dis.
665   2018;18(4):214–21. https://doi.org/10.1089/vbz.2017.2222
666   52.   Inácio CLS, Da Silva JHT, De Melo Freire RC, Gama RA, Marcondes CB, De Melo
667   Ximenes MDFF, et al. Checklist of mosquito species (Diptera: Culicidae) in the Rio
668   Grande do Norte State, Brazil-Contribution of Entomological Surveillance. J Med
669   Entomol [Internet]. 2017;54(3):763–73. https://doi.org/10.1093/jme/tjw236
670   53.   Linares MA, Laurito M, Visintin AM, Rossi GC, Stein M, Almirón WR. New mosquito
671   records (Diptera: Culicidae) from northwestern Argentina. Check List [Internet].
672   2016;12(4). https://doi.org/10.15560/12.4.1944
673   54.   Lira-Vieira AR, Gurgel-Goncalves R, Moreira IM, Cavalcanti Yoshizawa MA, Coutinho
674   ML, Prado PS, et al. Ecological aspects of mosquitoes (Diptera: Culicidae) in the gallery
675   forest of Brasilia National Park, Brazil, with an emphasis on potential vectors of yellow
676   fever. Rev Soc Bras Med Trop. 2013;46(5):566–74. https://doi.org/10.1590/0037-8682-
677   0136-2013
678   55.   Mangudo C, Aparicio JP, Rossi GC, Gleiser RM. Tree hole mosquito species composition
679   and relative abundances differ between urban and adjacent forest habitats in northwestern
680   Argentina. Bull Entomol Res [Internet]. 2018;108(2):203–12.
681   https://doi.org/10.1017/S0007485317000700
682   56.   Marassá AM, Paula MB, Gomes AC, Consales CA. Biotin-avidin sandwich ELISA with
683   specific human isotypes IgG1 and IgG4 for culicidae mosquito blood meal identification
684   from an epizootic yellow fever area in Brazil. J Venom Anim Toxins Incl Trop Dis
685   [Internet]. 2009;15(4):696–706. https://doi.org/10.1590/S1678-91992009000400008
686   57.   Medeiros-Sousa AR, Fernandes A, Ceretti-Junior W, Wilke ABB, Marrelli MT.
687   Mosquitoes in urban green spaces: Using an island biogeographic approach to identify
688   drivers of species richness and composition. Sci Rep [Internet]. 2017;7(1).
689   https://doi.org/10.1038/s41598-017-18208-x
690   58.   Medeiros AS, Marcondes CB, De Azevedo PRM, Jerônimo SMR, Silva VPME, De De
691   Ximenes MFFM. Seasonal variation of potential flavivirus vectors in an urban biological
692   reserve in Northeastern Brazil. J Med Entomol [Internet]. 2009;46(6):1450–7.
693   https://doi.org/10.1603/033.046.0630
694   59.   Alencar J, Gleiser RM, Morone F, de Mello CF, dos Silva JS, Serra-Freire NM, et al. A
695   comparative study of the effect of multiple immersions on Aedini (Diptera: Culicidae)
696   mosquito eggs with emphasis on sylvan vectors of yellow fever virus. Mem Inst Oswaldo
697   Cruz [Internet]. 2014;109(1):114–7. https://doi.org/10.1590/0074-0276130168
698   60.   Méndez-López MR, Attoui H, Florin D, Calisher CH, Florian-Carrillo JC, Montero S.

Association of vectors and environmental conditions during the emergence of Peruvian horse sickness orbivirus and Yunnan orbivirus in northern Peru. J Vector Ecol [Internet]. 2015;40(2):355–63. https://doi.org/10.1111/jvec.12174

61. Montes J. Culicidae fauna of Serra da Cantareira, Sao Paulo, Brazil. Rev Saude Publica. 2005;39(4):578–84. https://doi.org/10.1590/S0034-89102005000400010

62. Mucci LF, Medeiros-Sousa AR, Ceretti-Júnior W, Fernandes A, Camargo AA, Evangelista E, et al. Haemagogus leucocelaenus and Other Mosquitoes Potentially Associated with Sylvatic Yellow Fever in Cantareira State Park in the São Paulo Metropolitan Area, Brazil. J Am Mosq Control Assoc [Internet]. 2016;32(4):329–32. https://doi.org/10.2987/16-6587.1

63. Muller GA, Dalavequia MA, Wagner G, Marcondes CB. Blood sucking Diptera (Culicidae, Psychodidae, Simuliidae) in forest fragment under impact of dam in the borderland of Rio Grande do sul and Santa Catarina states, Brazil . Cienc Rural [Internet]. 2014;44(7):1194–6. https://doi.org/10.1590/0103-8478cr20131656

64. Müller GA, Kuwabara EF, Duque JE, Navarro-Silva MA, Marcondes CB. New records of mosquito species (Diptera: Culicidae) for Santa Catarina and Paraná (Brazil) . Biota Neotrop [Internet]. 2008;8(4):211–8. https://doi.org/10.1590/S1676-06032008000400021

65. Orlandin E, Santos EB, Piovesan M, Favretto MA, Schneeberger AH, Souza VO, et al. Mosquitoes (Diptera: Culicidae) from crepuscular period in an Atlantic Forest area in Southern Brazil . Brazilian J Biol [Internet]. 2017;77(1):60–7. https://doi.org/10.1590/1519-6984.09815

66. Pauvolid-Corrêa A, Tavares FN, Alencar J, Silva JDS, Murta M, Serra-Freire NM, et al. Preliminary investigation of culicidae species in South Pantanal, Brazil and their potential importance in arbovirus transmission. Rev Inst Med Trop Sao Paulo [Internet]. 2010;52(1):17–23. https://doi.org/10.1590/S0036-46652010000100004

67. Pecor JE, Jones J, Turell MJ, Fernandez R, Carbajal F, O'Guinn M, et al. Annotated checklist of the mosquito species encountered during arboviral studies in Iquitos, Peru (Diptera: Culicidae). J Am Mosq Control Assoc [Internet]. 2000;16(3):210–8. Retrieved from https://www.scopus.com/inward/record.uri?eid=2-s2.0-0034263438&partnerID=40&md5=475b03ef4a586aada5abff1d0a150829

68. Pinto CS, Confalonieri UEC, Mascarenhas BM. Ecology of Haemagogus sp. and Sabethes sp. (Diptera: Culicidae) in relation to the microclimates of the Caxiuanã National Forest, Pará, Brazil. Mem Inst Oswaldo Cruz. 2009;104(4):592–8. https://doi.org/10.1590/S0074-02762009000400010

69. Rossi GC. Annotated checklist, distribution, and taxonomic bibliography of the mosquitoes (Insecta: Diptera: Culicidae) of Argentina. Check List [Internet]. 2015;11(4). https://doi.org/10.15560/11.4.1712

70. Alencar J, Morone F, Mello CFD, Dégallier N, Lucio PS, Serra-Freire NMD, et al. Flight height preference for oviposition of mosquito (diptera: Culicidae) vectors of sylvatic yellow fever virus near the hydroelectric reservoir of simplicío, minas Gerais, Brazil. J Med Entomol [Internet]. 2013;50(4):791–5. https://doi.org/10.1603/ME12120

71. Rubio-Palis Y, Moreno JE, Bevilacqua M, Medina D, Martínez Á, Cardenas L, et al. Ecological characterization of anophelines and culicines in the indigenous territory of the Lower Caura River, Bolívar State, Venezuela . Bol Malariol y Salud Ambient [Internet]. 2010;50(1):95–107. Retrieved from https://www.scopus.com/inward/record.uri?eid=2-s2.0-80051939902&partnerID=40&md5=6996d3e624d3f07bb245aec69e24ff03

72.   Santos CF, Borges M. Impact of livestock on a mosquito community (Diptera: Culicidae)
in a Brazilian tropical dry forest. Rev Soc Bras Med Trop [Internet]. 2015;48(4):474–8.
https://doi.org/10.1590/0037-8682-0022-2015

73.   Santos CF, Silva AC, Rodrigues RA, Jesus JSR, Borges MAZ. Inventory of mosquitoes
(Diptera: Culicidae) in conservation units in Brazilian tropical dry forests . Rev Inst Med
Trop Sao Paulo [Internet]. 2015;57(3):227–32. https://doi.org/10.1590/S0036-
46652015000300008

74.   Serra OP, Cardoso BF, Maria Ribeiro AL, Leal dos Santos FA, Slhessarenko RD. Mayaro
virus and dengue virus 1 and 4 natural infection in culicids from Cuiaba, state of Mato
Grosso, Brazil. Mem Inst Oswaldo Cruz. 2016;111(1):20–9. https://doi.org/10.1590/0074-
02760150270

75.   Silva JDS, Pacheco JB, Alencar J, Guimarães AÉ. Biodiversity and influence of climatic
factors on mosquitoes (Diptera: Culicidae) around the Peixe Angical hydroelectric scheme
in the state of Tocantins, Brazil. Mem Inst Oswaldo Cruz [Internet]. 2010;105(2):155–62.
https://doi.org/10.1590/S0074-02762010000200008

76.   Talaga S, Dejean A, Carinci R, Gaborit P, Dusfour I, Girod R. Updated Checklist of the
Mosquitoes (Diptera: Culicidae) of French Guiana. J Med Entomol [Internet].
2015;52(5):770–82. https://doi.org/10.1093/jme/tjv109

77.   Talaga S, Murienne J, Dejean A, Leroy C. Online database for mosquito (Diptera,
Culicidae) occurrence records in French guiana. Zookeys [Internet]. 2015;2015(532):107–
15. https://doi.org/10.3897/zookeys.532.6176

78.   Tatila-Ferreira A, Maia D de A, Alencar J. Development of Preimaginal Stages of
Haemagogus Leucocelaenus (Diptera: Culicidae) in Laboratory Conditions. Entomol
News. 2017;127(2):142–50. https://doi.org/10.3157/021.127.0209

79.   Tatila-Ferreira A, Maia D de A, Santos de Abreu FV, Rodrigues WC, Alencar J.
Oviposition behavior of Haemagogus leucocelaenus (Diptera: culicidae), a vector of wild
yellow fever in Brazil. Rev Inst Med Trop Sao Paulo. 2017;59:UNSP e60-UNSP e60.
https://doi.org/10.1590/S1678-9946201759060

80.   Tubaki RM, Menezes RMT De, Vesgueiro FT, Cardoso RP. Observations on
Haemagogus janthinomys Dyar (Diptera: Culicidae) and other mosquito populations
within tree holes in a gallery forest in the northwestern region of Sao Paulo state, Brazil.
Neotrop Entomol. 2010;39(4):664–70. https://doi.org/Doi 10.1590/S1519-
566x2010000400030

81.   Alencar J, Serra-Friere NM, Marcondes CB, Silva JDS, Correa FF, Guimarães AE.
Influence of climatic factors on the population dynamics of haemogogus janthinomys
(Diptera: Culicidae), a vector of sylvatic yellow fever. Entomol News [Internet].
2010;121(1):45–52. https://doi.org/10.3157/021.121.0109

82.   Vale Barbosa M das G, Fe NF, Ribera Marciao AH, Thome da Silva AP, Monteiro WM,
de Farias Guerra MV, et al. Record of epidemiologically important Culicidae in the rural
area of Manaus, Amazonas. Rev Soc Bras Med Trop. 2008;41(6):658–63.

83.   Vasconcelos PFC, Rosa APAT, Rodrigues SG, Rosa EST, Monteiro HAO, Cruz ACR, et
al. Yellow fever in Para State, Amazon region of Brazil, 1998-1999: Entomologic and
epidemiologic findings. Emerg Infect Dis. 2001;7(3):565–9.
https://doi.org/10.3201/eid0703.010338

84.   Vasconcelos PFC, Sperb AF, Monteiro HAO, Torres MAN, Sousa MRS, Vasconcelos
HB, et al. Isolations of yellow fever virus from Haemagogus leucocelaenus in Rio Grande

do Sul State, Brazil. Trans R Soc Trop Med Hyg. 2003;97(1):60–2.
https://doi.org/10.1016/S0035-9203(03)90023-X

85. Yanoviak SP, Lounibos LP, Weaver SC. Land use affects macroinvertebrate community
composition in phytotelmata in the Peruvian Amazon. Ann Entomol Soc Am [Internet].
2006;99(6):1172–81. https://doi.org/10.1603/0013-
8746(2006)99[1172:LUAMCC]2.0.CO;2

86. Zequi JAC, Lopes J, Medri IM. Immature specimens of Culicidae (Diptera) found in
installed recipients in forest fragments in the Londrina, Parana, Brazil. Rev Bras Zool.
2005;22(3):656–61. https://doi.org/10.1590/S0101-81752005000300021

87. Alencar J, de Mello CF, Barbosa LS, Gil-Santana HR, Maia D de A, Marcondes CB, et al.
Diversity of yellow fever mosquito vectors in the Atlantic forest of Rio de Janeiro, Brazil.
Rev Soc Bras Med Trop. 2016;49(3):351–6. https://doi.org/10.1590/0037-8682-0438-
2015

88. Alencar J, Marcondes CB, Serra-Freire NM, Lorosa ES, Pacheco JB, Guimarães AÉ.
Feeding Patterns of Haemagogus capricornii and Haemagogus leucocelaenus (Diptera:
Culicidae) in Two Brazilian States (Rio de Janeiro and Goiás). J Med Entomol [Internet].
2008 Sep 1;45(5):873–6. https://doi.org/10.1093/jmedent/45.5.873

89. Aragão A de O, Nunes Neto JP, Cruz ACR, Casseb SMM, Cardoso JF, da Silva SP, et al.
Description and phylogeny of the mitochondrial genome of Sabethes chloropterus ,
Sabethes glaucodaemon and Sabethes belisarioi (Diptera: Culicidae). Genomics [Internet].
2018 Mar; https://doi.org/10.1016/j.ygeno.2018.03.016

90. Aragão NC, Müller GA, Balbino VQ, Costa Junior CRL, Figueirêdo Júnior CS, Alencar J,
et al. A list of mosquito species of the Brazilian State of Pernambuco, including the first
report of Haemagogus janthinomys (Diptera: Culicidae), yellow fever vector and 14 other
species (Diptera: Culicidae). Rev Soc Bras Med Trop [Internet]. 2010 Aug;43(4):458–9.
https://doi.org/10.1590/S0037-86822010000400024

91. Phillips S. maxnet: Fitting "Maxent" Species Distribution Models with "glmnet"
[Internet]. 2017. Retrieved from https://cran.r-project.org/package=maxnet

92. GBIF.org. GBIF Occurrence Download [Internet]. [cited 2018 Jul 19].
https://doi.org/10.15468/dl.wvs9g2

93. Fox SJ, Bellan SE, Perkins TA, Johansson MA, Meyers LA. Downgrading disease
transmission risk estimates using terminal importations. bioRxiv [Internet]. 2018;
https://doi.org/10.1101/265942

94. Kumm HW. Seasonal variations in rainfall: Prevalence of Haemagogus and incidence of
jungle yellow fever in Brazil and Colombia. Trans R Soc Trop Med Hyg. 1950;43(6):673–
82. https://doi.org/10.1016/0035-9203(50)90009-5

95. Causey OR, Dos Santos G V. Diurnal Mosquitoes in an Area of Small Residual Forests in
Brazil1. Ann Entomol Soc Am [Internet]. 1949 Dec 1;42(4):471–82.
https://doi.org/10.1093/aesa/42.4.471

96. Chadee DD. Seasonal abundance and diel landing periodicity of Sabethes chloropterus
(Diptera: Culicidae) in Trinidad, West Indies. J Med Entomol. 1990;27(6):1041–4.
https://doi.org/10.1093/jmedent/27.6.1041

97. Chadee DD, Tikasingh ES, Ganesh R. Seasonality, biting cycle and parity of the yellow
fever vector mosquito Haemagogus janthinomys in Trinidad. Med Vet Entomol.
1992;6(2):143–8. https://doi.org/10.1111/j.1365-2915.1992.tb00592.x

98. Chadee DD, Ganesh R, Hingwan JO, Tikasingh ES. Seasonal abundance, biting cycle and

parity of the mosquito Haemagogus leucocelaenus in Trinidad, West Indies. Med Vet
Entomol [Internet]. 1995 Oct;9(4):372–6. https://doi.org/10.1111/j.1365-
2915.1995.tb00006.x

99.   Medeiros AS, Marcondes CB, De Azevedo PRM, Jerônimo SMB, e Silva VPM, Ximenes
MDFFDM. Seasonal variation of potential flavivirus vectors in an urban biological
reserve in northeastern Brazil. J Med Entomol [Internet]. 2009;46(6):1450–7.
https://doi.org/10.1603/033.046.0630

100. Bates M. The Development and Longevity of Haemagogus Mosquitoes under Laboratory
Conditions1. Ann Entomol Soc Am [Internet]. 1947 Mar 1;40(1):1–12.
https://doi.org/10.1093/aesa/40.1.1

101. Galindo P. Bionomics of Sabethes Chloropterus Humboldt, a Vector of Sylvan Yellow
Fever in Middle America 1. Am J Trop Med Hyg [Internet]. 1958 Jul 1;7(4):429–40.
https://doi.org/10.4269/ajtmh.1958.7.429

102. Dégallier N, SÁ Filho GC, Monteiro HAO, Castro FC, Vaz Da Silva O, Brandão RCF, et
al. Release–Recapture Experiments with Canopy Mosquitoes in the Genera Haemagogus
and Sabeihes (Diptera: Culicidae) in Brazilian Amazonia. J Med Entomol [Internet]. 1998
Nov 1;35(6):931–6. https://doi.org/10.1093/jmedent/35.6.931

103. Team SD. RStan: the R interface to Stan [Internet]. 2018. Retrieved from http://mc-
stan.org/

104. Clements AN, Paterson GD. The Analysis of Mortality and Survival Rates in Wild
Populations of Mosquitoes. J Appl Ecol [Internet]. 1981 Aug;18(2):373.
https://doi.org/10.2307/2402401

105. Tesla B, Demakovsky LR, Mordecai EA, Ryan SJ, Bonds MH, Ngonghala CN, et al.
Temperature drives Zika virus transmission: Evidence from empirical and mathematical
models. Proc R Soc B Biol Sci. 2018;285(1884). https://doi.org/10.1098/rspb.2018.0795

106. Mordecai EA, Cohen JM, Evans M V, Gudapati P, Johnson LR, Lippi CA, et al. Detecting
the impact of temperature on transmission of Zika, dengue, and chikungunya using
mechanistic models. Althouse B, editor. PLoS Negl Trop Dis [Internet]. 2017 Apr
27;11(4):e0005568. https://doi.org/10.1371/journal.pntd.0005568

107. Johansson MA, Arana-Vizcarrondo N, Biggerstaff BJ, Staples JE. Incubation periods of
yellow fever virus. Am J Trop Med Hyg. 2010;83(1):183–8.
https://doi.org/10.4269/ajtmh.2010.09-0782

108. Chan M, Johansson MA. The Incubation Periods of Dengue Viruses. PLoS One.
2012;7(11):1–7. https://doi.org/10.1371/journal.pone.0050972

109. Kumm HW, Waddell MB. Haemagogus Capricornii Lutz as a Laboratory Vector of
Yellow Fever 1. Am J Trop Med Hyg [Internet]. 1948 Mar 1;s1-28(2):247–52.
https://doi.org/10.4269/ajtmh.1948.s1-28.247

110. Roca-García M, Bates M. The Development of the Virus of Yellow Fever in Haemagogus
Mosquitoes. Am J Trop Med Hyg [Internet]. 1946 Sep 1;s1-26(5):585–605.
https://doi.org/10.4269/ajtmh.1946.s1-26.585

111. ANDERSON CR, OSORNO-MESA E. The laboratory transmission of yellow fever virus
by Haemagogus splendens. Am J Trop Med Hyg [Internet]. 1946 Sep;26(5):613–8.
Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/21003269

112. Waddell MB, Taylor RM. Studies on Cyclic Passage of Yellow Fever Virus in South
American Mammals and Mosquitoes: Marmosets (Callithrix aurita) and Cebus Monkeys
(Cebus versutus) in Combination with Aedes aegypti and Haemagogus equinus. Am J

883   Trop Med Hyg [Internet]. 1945 May 1;s1-25(3):225–30.
884   https://doi.org/10.4269/ajtmh.1945.s1-25.225
885 113. Roca-Garcia M, Bates M. Laboratory Studies of the Saimiri-Haemagogus Cycle of Jungle
886   Yellow Fever. Am J Trop Med Hyg [Internet]. 1945 May 1;s1-25(3):203–16.
887   https://doi.org/10.4269/ajtmh.1945.s1-25.203
888 114. Galindo P, Trapido H, de Rodaniche E. Experimental Transmission of Yellow Fever by
889   Central American Species of Haemagogus and Sabethes Chloropterus. Am J Trop Med
890   Hyg [Internet]. 1956 Nov 1;5(6):1022–31. https://doi.org/10.4269/ajtmh.1956.5.1022
891 115. WADDELL MB. Comparative efficacy of certain South American Aëdes and
892   Haemagogus mosquitoes as laboratory vectors of yellow fever. Am J Trop Med Hyg
893   [Internet]. 1949 Jul;29(4):567–75. Retrieved from
894   http://www.ncbi.nlm.nih.gov/pubmed/18153054
895 116. Waddell MB, Taylor RM. Studies on the Cyclic Passage of Yellow Fever Virus in South
896   American Mammals and MosquitoesI III. Further Observations on Haemagogus equinus
897   as a vector of the virus. Am J Trop Med Hyg [Internet]. 1947 Jul 1;s1-27(4):471–6.
898   https://doi.org/10.4269/ajtmh.1947.s1-27.471
899 117. Estep LK, Burkett-Cadena ND, Hill GE, Unnasch RS, Unnasch TR. Estimation of
900   Dispersal Distances of Culex erraticus in a Focus of Eastern Equine Encephalitis Virus in
901   the Southeastern United States. J Med Entomol [Internet]. 2010;47(6):977–86.
902   https://doi.org/10.1603/ME10056
903 118. CAUSEY OR, KUMM HW. Dispersion of forest mosquitoes in Brazil; preliminary
904   studies. Am J Trop Med Hyg [Internet]. 1948 May;28(3):469–80. Retrieved from
905   http://www.ncbi.nlm.nih.gov/pubmed/18859835
906 119. Dimiceli C, Carroll M, Sohlberg R, Kim DH, Kelly M, Townshend JRG. MOD44B
907   MODIS/Terra Vegetation Continuous Fields Yearly L3 Global 250m SIN Grid V006
908   [Data set]. NASA EOSDIS Land Processes DAAC. 2015.
909   https://doi.org/10.5067/MODIS/MOD44B.006
910 120. Ministério da Saúde do Brasil. Epidemiológicas e Morbidade. 2017.
911 121. Camara FP, Gomes ALBB, Carvalho LMF de, Castello LGV. Dynamic behavior of
912   sylvatic yellow fever in Brazil (1954-2008). Rev Soc Bras Med Trop [Internet]. 2011
913   Jun;44(3):297–9. https://doi.org/10.1590/S0037-86822011005000024
914 122. Dark SJ, Bram D. The modifi able areal unit problem (MAUP) in physical geography.
915   2007;31(5):471–9. https://doi.org/10.1177/0309133307083294
916 123. Elith J, Leathwick JR, Hastie T. A working guide to boosted regression trees. J Anim
917   Ecol. 2008;77(4):802–13. https://doi.org/10.1111/j.1365-2656.2008.01390.x
918 124. Han BA, Schmidt JP, Alexander LW, Bowden SE, Hayman DTS, Drake JM.
919   Undiscovered Bat Hosts of Filoviruses. PLoS Negl Trop Dis. 2016;10(7):1–10.
920   https://doi.org/10.1371/journal.pntd.0004815
921 125. Millones M, Rogan J, II BL, Parmentier B, Harris R, Griffith D. Fire Data as Proxy for
922   Anthropogenic Landscape Change in the Yucatán. Land [Internet]. 2017;6(3):61.
923   https://doi.org/10.3390/land6030061
924 126. Grafström A, Lisic J. BalancedSampling: Balanced and Spatially Balanced Sampling
925   [Internet]. 2018. Retrieved from https://cran.r-project.org/package=BalancedSampling
926 127. Greenwell B, Boehmke B, Cunningham J, Developers GBM. gbm: Generalized Boosted
927   Regression Models [Internet]. 2018. Retrieved from https://cran.r-
928   project.org/package=gbm

128. Hijmans RJ, Phillips S, Leathwick J, Elith J. dismo: Species Distribution Modeling
[Internet]. 2017. Retrieved from https://cran.r-project.org/package=dismo
129. Greenwell BM. pdp: An R Package for Constructing Partial Dependence Plots. R J
[Internet]. 2017;9(1):421–36. Retrieved from https://journal.r-
project.org/archive/2017/RJ-2017-016/index.html
130. Goeman JJ, Solari A. Multiple hypothesis testing in genomics. Stat Med.
2014;33(11):1946–78. https://doi.org/10.1002/sim.6082
131. Bernhardt JR, Sunday JM, Thompson PL, O'Connor MI. Nonlinear averaging of thermal
experience predicts population growth rates in a thermally variable environment. Proc R
Soc B Biol Sci [Internet]. 2018 Sep 12;285(1886):20181076.
https://doi.org/10.1098/rspb.2018.1076
132. Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB. Influence of
climate on malaria transmission depends on daily temperature variation. Proc Natl Acad
Sci [Internet]. 2010;107(34):15135–9. https://doi.org/10.1073/pnas.1006422107
133. Almeida MAB, da Cardoso JC, dos Santos E, da Fonseca DF, Cruz LL, Faraco FJC, et al.
Surveillance for Yellow Fever Virus in Non-Human Primates in Southern Brazil, 2001-
2011: A Tool for Prioritizing Human Populations for Vaccination. PLoS Negl Trop Dis.
2014;8(3). https://doi.org/10.1371/journal.pntd.0002741
134. Shapiro LLM, Whitehead SA, Thomas MB. Quantifying the effects of temperature on
mosquito and parasite traits that determine the transmission potential of human malaria.
PLoS Biol. 2017;15(10):1–21. https://doi.org/10.1371/journal.pbio.2003489
135. Gorgas WC. Sanitation in Panama. Appleton; 1915.
951
952

953 **Tables**
954
955 **Table 1. Spillover model variables and definitions.**

Variable	Definition	Model
$\beta_h(\vec{x}, t)$	Proportion of mosquito bites from sylvatic vectors on humans at location $\vec{x}$ at time t	Approximated as $1 - f(\vec{x}, t)$ in locations with non-zero human population density, where $f(\vec{x}, t)$ is the percent forest cover at location $\vec{x}$ at time t
$\beta_p(\vec{x}, t)$	Proportion of mosquito bites from sylvatic vectors on non-human primates at location $\vec{x}$ at time t	Approximated as $f(\vec{x}, t)$ in locations within at least one non-human primate range, where $f(\vec{x}, t)$ is the percent forest cover at location $\vec{x}$ at time t
$\rho_v(\vec{x}, t)$	Sylvatic vector density at location $\vec{x}$ at time t	Approximated as location maximum mosquito density multiplied by relative seasonal abundance, where maximum mosquito density is determined by a species distribution model and seasonal abundance is modeled from field capture data (see Methods: Mechanism models – Mosquito distribution and seasonality)
$EIP(T(\vec{x}, t), \Delta t)$	Probability a mosquito that took an infectious blood meal becomes infectious with yellow fever virus given a temperature $T(\vec{x}, t)$ and Δt days elapsing	See Methods: Mechanistic submodels – Mosquito infectiousness
$s(T(\vec{x}, t), \Delta t)$	Probability that a mosquito survives Δt days given a temperature $T(\vec{x}, t)$	See Methods: Mechanistic submodels – Mosquito survival
$d(\|\vec{x} - \vec{y}\|)$	Probability that a mosquito disperses from $\vec{x}$ to $\vec{y}$	See Methods: Mechanistic models – Mosquito dispersal
$b(\vec{x}, t)$	Biting rate of sylvatic vectors at location $\vec{x}$ at time t	Assumed constant given limited information on determinants of vector biting rates
$\kappa(\vec{x}, t)$	Infection prevalence in nonhuman primate reservoir at location $\vec{x}$ at time t	For environmental risk metric, assumed constant given limited information of nonhuman primate infection prevalence. For periodic risk metric, used periodic curve fit to yearly case data (see Methods: Phenomenological primate dynamics).

956
957

Figures

Figure 1. Mechanistic and statistical model schematic. Submodels of components in the mechanistic model are parameterized using independent data on reservoir species, vector species occurrences, seasonal abundances, vector mark-recapture studies, vector survival, transmission experiments, forest cover, estimated vaccine coverage, and human population estimates. Reservoir disease prevalence is estimated from annual number of municipality-months with spillover. The output from the submodels are used in a mechanistic spillover model to predict four metrics risk of yellow fever in humans: periodic disease risk, environmental disease risk, immunological disease risk, and population-scaled disease risk. Environmental disease risk metric is then used as a covariate in a boosted regression tree to predict the municipality-months with spillover and identify covariates important for predicting spillover. Other environmental covariates are also included in the boosted regression tree. Details on data used in the mechanistic model can be found in the Supplementary Materials (S). Specific locations within the Supplementary Materials are noted parenthetically by either the section or table in which details can be found. Data used in the boosted regression tree are described in the Supplementary Materials (Table S6). Layers shown on the left correspond to mechanistic model components in Fig. 2a–k.

Figure 2. Data used to estimate ecological and human population components of spillover (a–k) and estimates of overall spillover risk (l–o). Number of primate reservoir species (a), vector species probability of occurrence (b), reservoir-vector contact probability (c), human-vector contact probability (d), human susceptibility approximated by one minus estimated vaccine coverage (i), and human distribution (j) vary spatially. Vector seasonal abundance is modeled as a function of rainfall using mosquito capture data (e). Vector dispersal depends on distance and is estimated from mark-recapture studies (f). Vector survival has been measured at different temperatures in laboratory (open circles) and field (closed circles) settings and was used to estimate temperature-dependent vector lifespan (g). Transmission studies at different temperatures inform modeled probability of vector infectiousness as a function of days since infecting bite and temperature (h). Phenomenologically modeled reservoir disease prevalence

(light blue line, k) is approximated from human case data (blue dots, k). All mechanistic model
components (a–k) are derived from empirical data in previously published studies. Components
a–h are used to predict environmental risk of disease spillover (l), components a–i are used for
immunological risk (m), components a–j are used for population-scaled risk (n) and components
a–h and k are used for periodic risk (o). The four disease risk metrics presented here for
illustrative purposes were estimated for January 2001 (l–o).
996

Figure 3. Municipality maximum periodic risk best predicts spillover. Each point is the calculated area under the curve (AUC) from spillover predicted by modeled risk, where higher AUC represents a model better able to distinguish between spillover and non-spillover observations. The risk models (from left to right on the *x*-axis) are environmental risk, periodic risk, immunological risk, and population-scaled risk. Municipality-wide maxima (red dashed lines and circles) and means (blue dotted lines and triangles) are shown for each metric.

1005
Figure 4. Modeled environmental risk captures seasonal variation and periodic risk
captures interannual variation in spillover. Each colored line is the seasonal average of
modeled maximum environmental risk in a municipality (a) and the yearly average of modeled
maximum periodic risk in a municipality (c). White lines are the regional average over the
municipal curves. Black points represent the total number of municipality-months with spillover
in that region per month (a) and per year (c), or the municipalities with at least one month with
spillover (b). Correlations between regional average environmental risk (white lines) and
regional number of municipality-months with spillover (black points) shown in parentheses (a, b)
for regions where spillover has occurred (all except the Northeast). Regions of Brazil are shown
with corresponding colors (c). The Southeast (shown in blue) was the region with the majority of
cases during the large outbreaks in 2016–2018.
1017

1018
Figure 5. Partial dependence plots of top six predictors of spillover in a municipality-month
from boosted regression tree analysis. Plots are listed in order of predictive importance with
relative influence (%) listed. In order, the variables identified as most important predictors were
average temperature in the municipality-month (a), one-month lagged maximum environmental
risk (b), estimated vaccine coverage (c), average rate of precipitation in the municipality-month
(d), current month maximum environmental risk (e), and municipality population density (log-
scaled for visibility, f). Histograms show the distribution of observed municipality-months at
each covariate value (left *y*-axis) and solid lines show the marginal effects of covariate on model
prediction (right *y*-axis). Marginal effects highlight the characteristics of municipality-months
that experienced spillover in Brazil 2001–2016.
1029

1030
Figure 6: Mechanistic model predicts consistent low, seasonal risk across states in
Southeast region of Brazil, where a large outbreak occurred in 2016–2018. Data are only
available until the end of 2016 (blue dashed line), so do not include the duration of the 2016–
2018 outbreaks (pink boxes). Only 2001–2016 spillovers are shown, defined as municipality-
months with human yellow fever cases (red points). Grey lines are municipality estimates of
maximum environmental risk and the black line is the environmental risk averaged over all
municipalities in the state. Prior to the large outbreak in 2017–2018, spillover had occurred in
Minas Gerais and São Paulo (right panels) but not in Espírito Santo or Rio de Janeiro (left
panels).