Comprehensive genome-wide identification of angiosperm upstream ORFs with peptide sequences conserved in

various taxonomic ranges using a novel pipeline, ESUCA

Hiro Takahashi[1,2#]*, Noriya Hayashi[3#], Yui Yamashita[3], Satoshi Naito[3,4], Anna Takahashi[5], Kazuyuki Fuse[6], Kenji

Satou[7], Toshinori Endo[8], Shoko Kojima[9], and Hitoshi Onouchi[3]*

[1]Graduate School of Medical Sciences, Kanazawa University, Kanazawa 920-1192, Japan

[2]Graduate School of Horticulture, Chiba University, Matsudo 271-8510, Japan

[3]Graduate School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

[4]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan

[5]Faculty of Information Technologies and Control, Belarusian State University of Informatics and Radio

Electronics, Minsk 220013, Belarus

[6]New Business Development Office, Churitsu Electric Corporation, Toyoake 470-1112, Japan

[7]Faculty of Biological Science and Technology, Institute of Science and Engineering, Kanazawa University,

920-1192, Kanazawa, Japan

[8] Graduate School of Information Science and Technology, Hokkaido University, Sapporo 060-0814, Japan

[9] Graduate School of Bioscience and Biotechnology, Chubu University, Kasugai 487-8501 Japan

*To whom correspondence should be addressed. Tel: +81-11-706-3887; Fax: +81-11-706-4932; Email:

onouchi@abs.agr.hokudai.ac.jp

Correspondence may also be addressed to Hiro Takahashi. Email: takahasi@p.kanazawa-u.ac.jp

[#]The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First

Authors.

[Key words: upstream ORF; translational regulation; bioinformatics; nascent peptide]

Abstract

Background: Upstream open reading frames (uORFs) in the 5′-untranslated regions (5′-UTRs) of certain

eukaryotic mRNAs encode evolutionarily conserved functional peptides, such as cis-acting regulatory peptides

that control translation of downstream main ORFs (mORFs). For genome-wide searches for uORFs with

conserved peptide sequences (CPuORFs), comparative genomic studies have been conducted, in which uORF

sequences were compared between selected species. To increase chances of identifying CPuORFs, we previously

developed an approach in which uORF sequences were compared using BLAST between *Arabidopsis* and any

other plant species with available transcript sequence databases. If this approach is applied to multiple plant

species belonging to phylogenetically distant clades, it is expected to further comprehensively identify CPuORFs

conserved in various plant lineages, including those conserved among relatively small taxonomic groups.

Results: To efficiently compare uORF sequences among many species and efficiently identify CPuORFs

conserved in various taxonomic lineages, we developed a novel pipeline, ESUCA. We applied ESUCA to the

genomes of five angiosperm species, which belong to phylogenetically distant clades, and selected CPuORFs

conserved among at least three different orders. Through these analyses, we identified 88 novel CPuORF families.

As expected, ESUCA analysis of each of the five angiosperm genomes identified many CPuORFs that were not

identified from ESUCA analyses of the other four species. However, unexpectedly, these CPuORFs include

those conserved in wide taxonomic ranges, indicating that the approach used here is useful not only for

comprehensive identification of narrowly conserved CPuORFs but also for that of widely conserved CPuORFs.

Examination of the effects of 11 selected CPuORFs on mORF translation revealed that CPuORFs conserved

only in relatively narrow taxonomic ranges can have sequence-dependent regulatory effects, suggesting that most

of the identified CPuORFs are conserved because of functional constraints of their encoded peptides.

Conclusions: This study demonstrates that ESUCA is capable of efficiently identifying CPuORFs likely to be

conserved because of the functional importance of their encoded peptides. Furthermore, our data show that the

approach in which uORF sequences from multiple species are compared with those of many other species, using

ESUCA, is highly effective in comprehensively identifying CPuORFs conserved in various taxonomic ranges.

Background

The 5′-untranslated regions (5′-UTRs) of many eukaryotic mRNAs contain upstream open reading frames (uORFs) [1-4]. Although most uORFs are not thought to encode functional proteins or peptides, certain uORFs encode regulatory peptides that have roles in post-transcriptional regulation of gene expression [5-8]. During translation of some of these regulatory uORFs, nascent peptides act inside the ribosomal exit tunnel to cause ribosome stalling [9]. Ribosome stalling on a uORF results in translational repression of the downstream main ORF (mORF) because stalled ribosomes block the access of subsequently loaded ribosomes to the mORF start codon [10]. Additionally, if ribosome stalling occurs at the stop codon of a uORF, nonsense-mediated mRNA decay may be induced [11, 12]. In some genes, uORF-encoded nascent peptides cause ribosome stalling in response to metabolites to down-regulate mORF translation under specific cellular conditions [10, 12-17]. In contrast to the uORFs encoding cis-acting regulatory nascent peptides, a uORF in the *Medicago truncatula MtHAP2-1* gene encodes a trans-acting regulatory peptide, which binds to the 5′-UTR of *MtHAP2-1* mRNA and causes mRNA degradation [18].

To comprehensively identify uORFs that encode functional peptides, genome-wide searches for uORFs with conserved peptide sequences (CPuORFs) have been conducted using comparative genomic approaches in various organisms [19-23]. In plants, approximately 40 CPuORF families have been identified by comparing the uORF-encoded amino acid sequences of orthologous genes in some of *Arabidopsis*, rice, cotton, orange, soybean, grape and tobacco, or those of paralogous genes in *Arabidopsis* [20, 22, 23]. Recently, 29 additional CPuORF families, which include CPuORFs with non-canonical initiation codons, have been identified by comparing 5′-UTR sequences between *Arabidopsis* and 31 other plant species [24].

In conventional comparative genomic approaches, uORF sequences are compared among selected species. Therefore, homology detection depends on the selection of species for comparison. In searches using this approach, if a uORF amino acid sequence is not conserved among the selected species, this uORF is not identified as a CPuORF, even if it is evolutionarily conserved between one of the selected species and other unselected species. To overcome this problem, we previously developed the BAIUCAS (for BLAST-based algorithm for identification of uORFs with conserved amino acid sequences) pipeline [25]. In BAIUCAS, homology searches of uORF amino acid sequences are performed using BLAST between a certain species and any other

species for which expressed sequence tag (EST) databases are available, and uORFs conserved beyond a certain

taxonomic range are selected. Using BAIUCAS, we searched for *Arabidopsis* CPuORFs conserved beyond the

order Brassicales, which *Arabidopsis* belongs to, and identified 13 novel CPuORF families [25]. We examined

the sequence-dependent effects of the CPuORFs identified by BAIUCAS on mORF translation using a transient

expression assay, and identified six regulatory CPuORFs that repress mORF translation in an amino acid

sequence-dependent manner [26, 27]. These sequence-dependent regulatory CPuORFs include ones conserved

only among relatively small taxonomic groups, such as a part of eudicots. Therefore, it is expected that

sequence-dependent regulatory CPuORFs conserved in various plant lineages, including narrowly conserved

ones, will be more comprehensively identified if BAIUCAS is applied to many plant species.

Before applying BAIUCAS to many species, improvement of BAIUCAS was desired to more efficiently

identify CPuORFs that were conserved because of the functional importance of their encoded peptides. One

major problem with identifying CPuORFs is that there are cases where a uORF found in the 5′-UTR of a

transcript is fused to the mORF in an isoform of the transcript, and in some of these cases, such uORF sequences

are conserved because they actually encode parts of mORF-encoded protein sequences. In other words, there are

cases where the protein-coding mORF is split into multiple ORFs in a splice variant and the ORF coding for the

N-terminal region of the protein appears like a uORF. Such an ORF can be extracted as a CPuORF if the amino

acid sequence in the N-terminal region of the protein is evolutionarily conserved. It is difficult to distinguish

between this type of 'spurious' CPuORFs and 'true' CPuORFs because even 'true' CPuORF-containing genes

produce splice variants in which a CPuORF is fused to the mORF, as seen in the At2g31280, At5g01710, and

At5g03190 genes [20, 25, 28]. Another major point to be improved is the method of calculating nonsynonymous

to synonymous nucleotide substitution (K_a/K_s) ratios for CPuORF sequences. These K_a/K_s ratios are used to

evaluate whether uORF sequences are conserved at the amino acid level or at the nucleotide level. However,

K_a/K_s ratios largely depend on the selection of uORF sequences used for their calculation. If uORF sequences

used for the calculation of a K_a/K_s ratio include many sequences from closely related species, the K_a/K_s ratio tends

to be high. For appropriate calculation of K_a/K_s ratios, uORF sequences need to be selected using proper criteria.

Here, we present an improved BAIUCAS version ESUCA (for evolutionary search for uORFs with

conserved amino acid sequences) and genome-wide identification of CPuORFs from five angiosperm genomes

using ESUCA. To distinguish between 'spurious' CPuORFs conserved because they code for parts of

mORF-encoded proteins and 'true' CPuORFs conserved because of functional constraints of their encoded small

peptides, ESUCA includes an algorithm to assess whether, for each uORF, transcripts bearing a uORF-mORF

fusion are minor or major forms among orthologous transcripts. Another new function of ESUCA is systematic

calculation of K_a/K_s ratios for CPuORF sequences. ESUCA includes an algorithm to select a uORF sequence

from each order for calculation of the K_a/K_s ratio of each CPuORF. Additionally, ESUCA is capable of

determining the taxonomic range within which each CPuORF is conserved. Although ESUCA can identify

CPuORFs conserved only among a small taxonomic group because ESUCA compares uORF sequences

between a certain species and any other species with available transcript databases, CPuORFs conserved among a

small taxonomic group may be less likely to encode functional peptides than those conserved across a wide

taxonomic range. The automatic determination of the taxonomic range of CPuORF conservation provides useful

information for the selection of CPuORFs likely to encode functional peptides. The current study demonstrates

that ESUCA efficiently identifies CPuORFs likely to be conserved because of functional constraints of their

encoded peptides. Furthermore, the data presented here show that the approach in which uORF sequences from

multiple species are compared with those of many other species, using ESUCA, is highly effective in

comprehensively identifying CPuORFs conserved in various taxonomic lineages.

Results

The ESUCA pipeline

In this study, to efficiently identify CPuORFs likely to be conserved because of functional importance of their

encoded peptides, we developed a novel pipeline, ESUCA, which consists of a six-step procedure (Fig. 1). The

first step is extraction of uORF sequences from a transcript sequence dataset. The uORFs are extracted by

searching the 5′-UTR sequence of each transcript for an ATG codon and its nearest downstream in-frame stop

codon. Although uORFs overlapping their downstream mORFs are also usually considered uORFs, we focus on

the type of uORFs that has both the start and stop codons within the 5′-UTR to avoid including uORFs whose

sequences are conserved because of functional constraints of mORF-encoded proteins. When there are splice

variants of a gene, uORFs in all splice variants are extracted. The second step assesses whether, for each uORF,

uORF-mORF fusion type transcripts are minor or major forms among orthologous transcripts. If transcripts with

a uORF-mORF fusion are found as a major form in a majority of species with their orthologs, the uORF

sequence is likely to code for a part of the mORF-encoded protein. Therefore, such a uORF should be discarded

as a 'spurious' uORF. In contrast, if transcripts with a uORF-mORF fusion are found in only a small proportion

of species with their orthologs, the uORF-mORF fusion type transcripts are considered minor form transcripts

and therefore can be ignored. For this assessment, the NCBI reference sequence (RefSeq) database is used, which

provides curated non-redundant transcript sequences [29]. For each uORF, the ratio of RefSeq RNAs with a

uORF-mORF fusion to all RefSeq RNAs with both sequences similar to the uORF and its downstream mORF is

calculated (Fig. 2). We define this ratio as the uORF-mORF fusion ratio. If the uORF-mORF fusion ratio of a

uORF is equal to or greater than 0.3, then the uORF is discarded. The third step is uORF amino acid sequence

homology searches. In this step, tBLASTn searches are performed against a transcript sequence database, using

the amino acid sequences of the uORFs as queries (uORF-tBLASTn analysis). The uORFs with tBLASTn hits

from other species are selected. The fourth step is selection of uORFs conserved among homologous genes. To

confirm whether the uORF-tBLASTn hits are derived from homologs of the original uORF-containing gene, the

downstream sequences of putative uORFs in the uORF-tBLASTn hits are subjected to another tBLASTn

analysis, which uses the mORF amino acid sequence of the original uORF-containing transcript as a query

(mORF-tBLASTn analysis) (Fig. 3). If a uORF-tBLASTn hit has a partial or intact ORF that contains a sequence

similar to the mORF amino acid sequence downstream of the putative uORF, it is considered to be derived from

a homolog of the original uORF-containing gene. If uORF-tBLASTn and mORF-tBLASTn hits are found in at

least two orders other than that of the original uORF, then the uORF is selected as a candidate CPuORF. This is

because at least three uORF sequences from different orders are necessary to confirm at the later manual

validation step that the same region is conserved among homologous uORF sequences. The fifth step is K_a/K_s

analysis. In this step, K_a/K_s ratios for the selected candidate CPuORFs is calculated to assess whether the

candidate CPuORF sequences are conserved at the nucleotide or amino acid level. A K_a/K_s ratio close to 1

indicates neutral evolution, whereas a K_a/K_s ratio close to 0 suggests that purifying selection acted on the amino

acid sequences. For each candidate CPuORF, a representative uORF-tBLASTn and mORF-tBLASTn hit is

selected from each order, and the putative uORF sequences in the representative uORF-tBLASTn and

mORF-tBLASTn hits are used for the calculation of the K_a/K_s ratio. If the K_a/K_s ratio of a candidate CPuORF is

less than 0.5 and significantly different from that of the negative control with q less than 0.05, then the candidate

CPuORF is selected for further analysis. The final step is to determine the taxonomic range of uORF sequence

conservation. In this step, the representative uORF-tBLASTn and mORF-tBLASTn hits selected in the fifth step

are classified into taxonomic categories (Fig. 4). On the basis of the presence of the uORF-tBLASTn and

mORF-tBLASTn hits in each taxonomic category, the taxonomic range of sequence conservation is determined

for each CPuORF.

Identification of angiosperm CPuORFs using ESUCA

We applied ESUCA to five angiosperm species, *Arabidopsis*, rice, tomato, poplar and grape, which belong to

phylogenetically distant clades of angiosperm, and for which entire genomic DNA and transcript sequence

datasets were available. Rice is a monocot, whereas the others are eudicots. *Arabidopsis* and poplar belong to two

different groups of rosids (marvids and fabids), whereas tomato belongs to asterids. Grape belongs to neither

rosids nor asterids. In the first step of ESUCA, we extracted uORF sequences from the 5′-UTR sequence of each

transcript of these species, using the transcript sequence datasets described in the Materials and Methods. In these

datasets, different transcript IDs are assigned to each splice variant from the same gene. To extract sequences of

uORFs and their downstream mORFs from all splice variants, we extracted uORF and mORF sequences from

each of the transcripts with different transcript IDs. In the second step, we calculated the uORF-mORF fusion

ratio of each uORF-containing transcript, using the extracted uORF and mORF sequences, and removed uORFs

with uORF-mORF fusion ratios equal to or greater than 0.3 (Supplementary Table S1). We also discarded

uORFs whose numbers of RefSeq RNAs containing both sequences similar to the uORF and its downstream

mORF were less than 10. This was done because appropriate evaluations of uORF-mORF fusion ratios were

difficult with a few related RefSeq RNAs and such uORFs are unlikely to be evolutionarily conserved. In the

third step, using the amino acid sequences of the remaining uORFs as queries, we performed uORF-tBLASTn

searches against a plant transcript sequence database that contained contigs of assembled EST and transcriptome

shotgun assembly (TSA), singleton EST/TSA sequences, and RefSeq RNAs (See Materials and Methods for

details). In the fourth step, the uORF-tBLASTn hits were subjected to mORF-tBLASTn analysis, and

uORF-tBLASTn and mORF-tBLASTn hits were extracted. Plant EST and TSA databases can include

contaminant sequences from other organisms, such as parasites, plant-feeding insects and infectious

microorganisms. We checked the possibility that the extracted uORF-tBLASTn and mORF-tBLASTn hits

included contaminant EST/TSA sequences, using BLASTn searches. The BLASTn searches were performed

using each uORF-tBLASTn and mORF-tBLASTn hit EST/TSA sequence as a query against EST/TSA and

RefSeq RNA sequences from all organisms, with an E-value cutoff of 10^{-100} and an identity threshold of 95%.

Contaminant EST/TSA sequences were identified by this analysis, as described in Materials and Methods, and

were removed from the uORF-tBLASTn and mORF-tBLASTn hits. We selected uORFs whose remaining

uORF-tBLASTn and mORF-tBLASTn hits were found in homologs from at least two orders other than that of

the original uORF. Thereafter, we generated amino acid sequence alignments of each selected uORF and its

homolog, using a putative homologous uORF sequence from each order in which uORF-tBLASTn and

mORF-tBLASTn hits were found. When multiple original uORFs derived from splice variants of the same gene

partially or completely shared amino acid sequences, the one with the longest conserved region was manually

selected on the basis of the uORF amino acid sequence alignments. In the fifth step, the remaining uORFs were

subjected to K_a/K_s analysis. The uORFs with K_a/K_s ratios less than 0.5 showing significant differences from those

of negative controls ($q < 0.05$) were selected as candidate CPuORFs (Supplementary Table S1). Through

ESUCA analyses of *Arabidopsis*, rice, tomato, poplar, and grape genomes, 104, 59, 42, 148, and 79 candidate

CPuORFs were extracted, respectively. Of these, 87 *Arabidopsis*, 53 rice, 30 tomato, 77 poplar, and 44 grape

uORFs belong to the previously identified CPuORF families, homology groups (HGs) 1 to 53 [20, 22-25]

(Supplementary Table S1). The amino acid sequences of the remaining candidate CPuORFs are not similar to

those of the known CPuORFs. Therefore, 17, 6, 12, 71, and 35 novel candidate CPuORFs were extracted from

Arabidopsis, rice, tomato, poplar, and grape genomes, respectively.

Validation of candidate CPuORFs

If the amino acid sequence of a uORF is evolutionarily conserved because of functional constraints of the

uORF-encoded peptide, it is expected that the amino acid sequence in the functionally important region of the

peptide is conserved among the uORF and its orthologous uORFs. Therefore, we manually checked whether the

amino acid sequences in the same region are conserved among uORF sequences in the alignment of each novel

candidate CPuORF. We found that the alignments of 13 novel candidate CPuORFs contain sequences that do not

share the consensus amino acid sequence in the conserved region, and removed these sequences from the

alignments. We also removed sequences derived from genes not related to the corresponding original

uORF-containing gene from the alignments of five novel candidate CPuORFs. When these changes resulted in

the number of orders with the uORF-tBLASTn and mORF-tBLASTn hits becoming less than two, the candidate

CPuORFs were discarded. Eight novel candidate CPuORFs were discarded for this reason. Supplementary

Figure S1 shows the uORF amino acid sequence alignments without the removed sequences. The K_a/K_s ratios

were recalculated after the manual removal of the sequences (Supplementary Table S1), and six additional novel

candidate CPuORFs were discarded because their K_a/K_s ratios were greater than 0.5.

Using genomic position information from Ensembl Plants (http://plants.ensembl.org/index.html) [30] and

Phytozome v12.1 (https://phytozome.jgi.doe.gov/pz/portal.html) [31], we manually checked whether the

positions of the remaining novel candidate CPuORFs overlap with those of the mORFs of other genes or the

mORFs of splice variants of the same genes. We found that the genomic position of the candidate CPuORF of the

Arabidopsis ROA1 (AT1G60200) gene overlaps with that of an intron in the mORF region of a splice variant.

Protein sequences with an N-terminal region similar to the amino acid sequence encoded by the 5′-extended region

of the mORF in this splice variant are found in most orders from which the uORF-tBALASTn and

mORF-tBLASTn hits of this candidate CPuORF were extracted, suggesting that the splice variant with the

5′-extended mORF is not a minor form among orthologous transcripts. Therefore, this candidate CPuORF was

discarded.

In the second step of ESUCA, we excluded uORF sequences likely to encode parts of the mORF-encoded

proteins, by removing uORFs with high uORF-mORF fusion ratios. To confirm that the novel candidate

CPuORFs do not code for parts of the mORF-encoded proteins, each of the putative uORF sequences used for

the alignment and K_a/K_s analysis was queried against the UniProt protein database (https://www.uniprot.org/),

using BLASTx. When putative uORF sequences matched protein sequences with low *E*-values, we manually

checked whether amino acid sequences similar to those encoded by the putative uORFs were contained within

mORF-encoded protein sequences. In this analysis, mORF-encoded proteins with N-terminal sequences similar

to the amino acid sequences encoded by the candidate CPuORFs of the rice *OsUAM2* gene and its poplar

ortholog, POPTR_0019s07850, were identified in many orders. This suggests that the sequences encoded by

these candidate CPuORFs are likely to function as parts of the mORF-encoded proteins. Therefore, we discarded

these candidate CPuORFs. For some other novel candidate CPuORFs, mORF-encoded proteins with sequences

similar to those encoded by the candidate CPuORF and/or its homologous putative uORFs were also found.

However, we did not exclude these candidate CPuORFs, because such uORF-mORF fusion type proteins were

found in only a few species for each candidate.

After manual validation, 12, 4, 11, 70, and 34 uORFs were identified as novel CPuORFs in *Arabidopsis*, rice,

tomato, poplar and grape, respectively. Among these novel CPuORFs, those with both similar uORF and mORF

amino acid sequences were classified into the same HGs. Also, using OrthoFinder ver. 1.1.4 [32], an algorithm

for ortholog group inference, we classified the genes with novel CPuORFs and those with previously identified

CPuORFs into ortholog groups. The same HG number with a different sub-number was assigned to CPuORFs of

genes in the same ortholog group with dissimilar uORF sequences (e.g. HG56.1 and HG56.2). Of the newly

identified CPuORF genes, six were classified into the same ortholog groups as previously identified CPuORF

genes, but the amino acid sequences of these six CPuORFs are dissimilar to those of the known CPuORFs.

Including this type of CPuORFs, we identified 131 novel CPuORFs that belong to 88 novel HGs (HG2.2, HG9.2,

HG16.2, HG43.2, HG50.2, HG52.2 and HG54-HG130) (Supplementary Table S1).

Determination of the taxonomic range of CPuORF sequence conservation

As the final step of ESUCA, we determined the taxonomic range of the sequence conservation of each CPuORF

identified, including previously identified CPuORFs. For this purpose, the uORF-tBLASTn and

mORF-tBLASTn hits selected for the K_a/K_s analysis and retained after manual validation were classified into 13

plant taxonomic categories (See Materials and Methods for details.), on the basis of taxonomic lineage

information of EST, TSA, and RefSeq RNA sequences (Fig. 4). Fig. 5 and Supplementary Table S2 show the

taxonomic range of sequence conservation for each HG and each CPuORF, respectively. In general, CPuORFs

belonging to previously identified HGs tend to be conserved in a wider range of taxonomic categories than those

belonging to the newly identified HGs. For 19 of the novel HGs, CPuORF sequences are conserved both in

eudicots and monocots or in wider taxonomic ranges. In contrast, for 69 of the novel HGs, CPuORF sequences

are conserved only among eudicots. For 12 of these, CPuORF sequences are conserved in narrower taxonomic

ranges, only among rosids or asterids. These results indicate that the taxonomic range of CPuORF sequence

conservation varies, and that ESUCA can identify CPuORFs conserved in a relatively narrow taxonomic range.

Sequence-dependent effects of CPuORFs on mORF translation

To address the relationship between the taxonomic range of CPuORF sequence conservation and the

sequence-dependent effects of CPuORFs on mORF translation, we selected 11 poplar CPuORFs and examined

their sequence-dependent effects on expression of the downstream reporter gene using a transient expression

assay. Of the selected CPuORFs, those belonging to HG46, HG55, HG57, HG66 and HG103 are conserved in

diverse angiosperms or in wider taxonomic ranges (Fig. 5, Supplementary Table S2). The CPuORFs belonging

to HG65, HG80, HG81 and HG87 are conserved in a wide range of eudicots, whereas the CPuORFs belonging

to HG88 and HG107 are conserved only among rosids (Fig. 5, Supplementary Table S2). In the 5′-UTR of the

poplar gene with the HG107 CPuORF, there is another uORF immediately upstream of the CPuORF

(Supplementary Table S2K). To focus on the sequence-dependent effect of the CPuORF on mORF translation,

the upstream uORF was eliminated by mutating its initiation codon because the presence of the immediate

upstream uORF may reduce the translation efficiency of the CPuORF and therefore potentially make the effect of

the CPuORF ambiguous. The 5′-UTR sequences containing the selected CPuORFs were fused to the firefly

luciferase (Fluc) coding sequence and were placed under the control of the 35S promoter to generate the

wild-type (WT) reporter constructs (Fig. 6a, Supplementary Figure S2). To assess the importance of the amino

acid sequences for the effects of these CPuORFs on mORF translation, frameshift mutations were introduced into

the CPuORFs so that the amino acid sequences of their conserved regions could be altered. A + 1 or − 1

frameshift was introduced upstream or within the conserved region of each CPuORF, and another frameshift was

introduced before the stop codon to shift the reading frame back to the original frame (Supplementary Figure S2).

These reporter constructs were each transfected into protoplasts from *Arabidopsis thaliana* MM2d

suspension-cultured cells. After 24 h of incubation, cells were harvested and disrupted to analyze luciferase

activity. In five of the 11 CPuORFs, the introduced frameshift mutations significantly upregulated the expression

of the reporter gene, indicating that these CPuORFs repress mORF translation in a sequence-dependent manner

(Fig. 6b). These five novel sequence-dependent regulatory CPuORF include the HG107 CPuORF, which is one

of the CPuORFs conserved only among rosids. Therefore, this result suggests that CPuORFs conserved only

among rosids can have sequence-dependent regulatory effects.

Discussion

Comprehensive identification of angiosperm CPuORFs by ESUCA analyses of multiple species' genomes

In this study, we developed ESUCA, a pipeline for efficient genome-wide identification of CPuORFs. By

applying ESUCA to five angiosperm genomes, we identified 131 CPuORFs that belong to 88 novel HGs. Of

these HGs, 70 were identified through ESUCA analysis of only one of the *Arabidopsis*, rice, tomato, poplar or

grape genomes (Fig. 5). This means that CPuORFs belonging to these HGs cannot be identified by comparing

uORF sequences of orthologous genes between these five species. Therefore, this result demonstrates that the

approach used in this study, in which uORF sequences from multiple species are compared with those of many

other species, is highly effective in comprehensively identifying CPuORFs. We expected this approach to be

particularly useful for comprehensive identification of CPuORFs conserved in relatively narrow taxonomic

ranges. Therefore, we used the five angiosperm species belonging to relatively distant lineages in order to identify

CPuORFs conserved only among the taxonomic categories to which each of the five species belong. However,

unexpectedly, in 41 of 70 HGs identified through ESUCA analysis of only one of the five plant genomes,

CPuORF sequences are conserved in both rosids and asterids, two major groups of eudicots, indicating that these

CPuORFs are conserved across diverse eudicots (Fig. 5). One possible explanation of this observation is that

sequence conservation of uORFs is often lost in small taxonomic groups, such as orders or families, during

evolution. For example, while the CPuORF of the tomato LOC101264451 gene, which belongs to HG43.1,

exerts a sequence-dependent repressive effect on mORF translation, the CPuORF of its *Arabidopsis* ortholog,

ANAC096, lacks the C-terminal half of the amino acid sequence in the highly conserved region and does not have

a sequence-dependent regulatory effect [25-27]. In contrast to the *Arabidopsis ANAC096* CPuORF, all the critical

amino acid residues in the highly conserved region are retained in the HG43.1 CPuORF of *Tarenaya hassleriana*,

which belongs to the same order as *Arabidopsis*, Brassicales, but a different family. CPuORFs involved in

preferable but not essential post-transcriptional regulations may be lost in some taxonomic clades during

evolution. Such CPuORFs would be difficult to be identified by comparing uORF sequences between a few

selected species if a selected species is included in clades where the CPuORF sequences were lost. Therefore, the

ESUCA method is advantageous for comprehensive identification of CPuORFs compared with conventional

comparative genomic approaches. Our results reveal that the approach used here (*i.e.* ESUCA analyses of

multiple species' genomes) is further advantageous and is highly useful not only for comprehensive identification

of narrowly conserved CPuORFs but also for that of widely conserved CPuORFs.

The transient expression assays in the current study identified five novel poplar regulatory CPuORFs that

exert sequence-dependent repressive effects on mORF translation. One of the identified regulatory CPuORFs is

conserved only among rosids (*i.e.* only among fabids and malvids). This result suggests that at least, CPuORFs

conserved only among fabids and malvids can have sequence-dependent regulatory effects, although we cannot

rule out the possibility that CPuORFs conserved in narrower taxonomic ranges can have sequence-dependent

regulatory effects. Of the 92 CPuORF HGs identified through ESUCA analysis of the poplar genome, HG101.1

is conserved only among fabids, whereas the others are conserved in at least one other taxonomic category in

addition to fabids (Fig. 5). Likewise, all the HGs identified through ESUCA analysis of the other four plant

genomes are conserved in multiple taxonomic categories. Altogether, these results suggest that most the

CPuORFs identified in this study, in which CPuORFs conserved in more than two orders were extracted, are

likely to be conserved because of functional constraints of their encoded peptides. Of the 11 poplar CPuORFs

analyzed by the transient expression assays, five are conserved beyond eudicots, and three of them exhibited

sequence-dependent repressive effects (Figs. 5 and 6b). In addition, we previously examined the effects of 16

CPuORFs, which belong to HG27-HG29, HG33-HG43.1, HG44, and HG45, on mORF translation and

identified six sequence-dependent regulatory CPuORFs. In this analysis, five of 11 CPuORFs conserved beyond

eudicots showed sequence-dependent repressive effects, whereas one of five CPuORFs conserved only among

eudicots showed a sequence-dependent repressive effect [26, 27]. These results may suggest that CPuORFs

conserved beyond eudicots are more likely to encode functional peptides than CPuORFs conserved only among

eudicots. ESUCA is capable of selecting CPuORFs conserved in certain taxonomic ranges on the basis of two

criteria, the numbers of orders in which CPuORFs are conserved and/or taxonomic categories in which

CPuORFs are conserved. Our results demonstrate that this function of ESUCA is highly useful for the efficient

selection of CPuORFs likely to encode functional peptides.

Of the CPuORFs analyzed for their sequence-dependent regulatory effects in the present study, the CPuORFs

belonging to HG55 and HG66 showed no significant sequence-dependent effect on mORF translation, despite

their widespread sequence conservation beyond eudicots (Figs. 5 and 6b). These CPuORFs might encode

peptides that have functions other than the control of mORF translation, or they might exert sequence-dependent

regulatory effects only under certain conditions. In fact, many known sequence-dependent regulatory uORFs

repress mORF translation in response to metabolites, such as polyamine, arginine, and sucrose [10, 14, 15, 17].

Likewise, the other CPuORFs that exhibited no significant sequence-dependent regulatory effect might encode

peptides that have other functions or exert regulatory effects only under specific conditions.

Filtering using the uORF-mORF fusion ratio

To distinguish between 'spurious' CPuORFs conserved because they code for parts of mORF-encoded proteins

and 'true' CPuORFs conserved because of functional constraints of their encoded small peptides, we employed

the criterion of the uORF-mORF fusion ratio and discarded uORFs with uORF-mORF fusion ratios equal to or

greater than 0.3. We checked how effectively the 'spurious' CPuORFs were removed with this criterion, using a

protein sequence database. Although uORF-mORF fusion type protein sequences were found in OsUAM2

homologs from widespread angiosperm species, no other candidate CPuORFs were extracted in which

uORF-mORF fusion type protein sequences were found in many species that have sequences similar to the

candidate CPuORFs. This indicates that the uORF-mORF fusion ratio filtering worked effectively to exclude

'spurious' CPuORFs that code for parts of the mORF-encoded proteins.

ESUCA extracted all known plant cis-acting regulatory CPuORFs that control mORF translation in a

sequence-dependent manner (*i.e.* HG1, HG3, HG6, HG12, HG13, HG14, HG15.3, HG18, HG19, HG27, HG34,

HG36, HG41, HG42, and HG43.1 CPuORFs [15, 17, 26, 27, 33-38]) (Supplementary Table S1). Of these, the

HG3 CPuORFs, associated with an mORF coding for an *S*-adenosylmethionine decarboxylase (AdoMetDC; EC

4.1.1.50) [20, 39], showed relatively high uORF-mORF fusion ratios, although still below 0.3 (Supplementary

Table S1). In this study, 11 CPuORFs belonging to this HG were extracted using ESUCA and the uORF-mORF

fusion ratios of these CPuORFs were in the range of 0.17 to 0.27, with a median value of 0.26 (Supplementary

Table S1). Consistent with these relatively high uORF-mORF fusion ratios, uORF-mORF fusion type protein

sequences that contain amino acid sequences resembling HG3 CPuORFs are found in widespread angiosperm

species. The uORF-mORF fusion ratios of the candidate CPuORFs of rice *OsUAM2* and its poplar ortholog were

0.28 (Supplementary Table S1). The rice *OsUAM2* gene codes for a protein similar to UDP-arabinopyranose

mutases (EC 5.4.99.30) [40]. Two protein isoforms, a uORF-mORF fusion type and one lacking the

uORF-encoded region, produced from splice variants of an *OsUAM2* orthologous gene, are found in diverse

angiosperm species. The functions of both of isoforms are not yet known. Considering the example of HG3

CPuORFs, we cannot rule out the possibility that a candidate CPuORF functions as a regulatory uORF even if

uORF-mORF fusion type protein sequences are found in widespread species. However, to avoid including

potential 'spurious' CPuORFs whose amino acid sequences are likely to be evolutionarily conserved because of

their function as N-terminal regions of the mORF-encoded proteins, we excluded the candidate CPuORFs of the

rice *OsUAM2* gene and its poplar ortholog. In conclusion, the criterion of the uORF-mORF fusion ratio used in

this study appears appropriate because all known cis-acting sequence-dependent regulatory CPuORFs were

extracted and most 'spurious' CPuORFs were removed.

Conclusion

The present study demonstrates that the approach in which uORF sequences from multiple species are compared

with those of many other species, using ESUCA, is highly effective in comprehensive identification of CPuORFs.

Using this approach, we identified many novel angiosperm CPuORFs, which include CPuORFs conserved

among limited clades and widely conserved CPuORFs. Our results also showed that ESUCA is capable of

efficiently selecting CPuORFs likely to be conserved because of functional importance of their encoded peptides.

The approach used here can be applied to any eukaryotic organism with available genome and transcript

sequence databases and therefore is expected to contribute to the comprehensive identification of CPuORFs

encoding functional peptides in various organisms. Furthermore, besides CPuORFs, the algorithms developed for

ESUCA and the approach used here can be applied to the identification of other sequences conserved in various

taxonomic ranges.

Materials and Methods

Extraction of uORF sequences

We used genome sequence files in FASTA format and genomic coordinate files in GFF3 format obtained from

Ensemble Plants Release 33 (https://plants.ensembl.org/index.html) [30] to extract *Arabidopsis* (*Arabidopsis*

thaliana), tomato (*Solanum lycopersicum*), poplar (*Populus trichocarpa*), and grape (*Vitis vinifera*) uORF

sequences. We used a genome sequence file in FASTA format and a genomic coordinate files in GFF3 format

obtained from Phytozome v11 (https://phytozome.jgi.doe.gov/pz/portal.html) [31] for rice (*Oryza sativa*). We

extracted exon sequences from genome sequences, on the basis of genomic coordinate information, and

constructed transcript sequence datasets by combining exon sequences. On the basis of the transcription start site

and the translation initiation codon of each transcript in the genomic coordinate files, we extracted 5´-UTR

sequences from the transcript sequence datasets. Then, we searched the 5´-UTR sequences for an ATG codon and

its nearest downstream in-frame stop codon. Sequences starting with an ATG codon and ending with the nearest

in-frame stop codon were extracted as uORF sequences. When multiple uORFs from a gene shared the same

stop codon, only the longest uORF sequence was used for further analyses.

Assembly of EST and TSA sequences

EST, TSA, and RefSeq RNA sequence datasets were obtained from the International Nucleotide Sequence

Database Collaboration databases (NCBI release of 2016-12-03 and DDBJ release 106.0 for EST and TSA

sequences, NCBI release 79 for RefSeq RNA sequences). On the basis of taxonomic lineage information

provided by NCBI Taxonomy (https://www.ncbi.nlm.nih.gov/taxonomy), EST and TSA sequences derived from

Viridiplantae were extracted from the databases. EST and TSA sequences from the same species were assembled

using Velvet ver. 1.2.10 [41] with k-mer length of 99. The k-mer length was optimized using *A. thaliana* EST and

TSA sequences to minimize the total numbers of assembled contigs and unclustered singleton sequences.

Unclustered singleton EST/TSA sequences, derived from species for which RefSeq RNA sequences were

available, were mapped to the RefSeq RNA sequences using Bowtie2 ver. 2.2.9 [42] and the default parameters.

We discarded singleton EST/TSA sequences that matched any RefSeq RNA sequences from the same species.

We created a plant transcript sequence database for BLAST searches in our local computers by using the

remaining singleton EST/TSA sequences, assembled contigs, Viridiplantae RefSeq RNA sequences, and

makeblastdb, a program contained in the NCBI-BLAST package.

Calculation of the uORF-mORF fusion ratio

The uORF-mORF fusion ratio for each of the extracted uORFs was assessed as follows. We performed tBLASTx using each uORF sequence as a query against plant (Viridiplantae) RefSeq RNA sequences with an E-value cutoff of 2000 (uORF-tBLASTx). We used standalone NCBI-BLAST + ver. 2.6.0 [43] for all BLAST analyses. We next performed tBLASTx with an E-value threshold of 10^{-1} using the mORF sequence associated with each uORF as a query against the uORF-tBLASTx hit sequences (mORF-tBLASTx). Using these two-step tBLASTx searches, we selected RefSeq RNAs that contain both sequences similar to the original uORF and its downstream mORF. Then, we examined whether the largest ORF of each of the selected RefSeq RNAs included the region that matched the original mORF in the same reading frame (Fig. 2). We also examined whether the largest ORF included the region that matched the original uORF, irrespective of the reading frame. The RefSeq RNA was considered to have a uORF-mORF fusion if the largest ORF contained both regions that matched the original uORF and mORF (Fig. 2). The RefSeq RNA was considered to have a uORF separated from the downstream mORF if the largest ORF contained the region that matched the original mORF but not the region that matched the original uORF (Fig. 2). RefSeq RNA numbers of the former and latter types were defined as X and Y, respectively. We calculated a uORF-mORF fusion ratio as $X / (X + Y)$ for each of the original uORF-containing transcript sequences.

BLAST-based search for uORFs conserved between homologous genes

To search for uORFs with amino acid sequences conserved between homologous genes, we first performed tBLASTn searches against the assembled plant transcript sequence database, using the amino acid sequences of the uORFs as queries. In these uORF-tBLASTn searches, we extracted transcript sequences that matched a uORF with an E-value less than 2000 and derived from species other than that of the original uORF. The downstream in-frame stop codon closest to the 5′-end of the matching region of each uORF-tBLASTn hit was selected (Fig. 3). Then, we looked for an in-frame ATG codon upstream of the selected stop codon, without any other in-frame stop codon between them. uORF-tBLASTn hits without such an ATG codon were discarded. If one or more in-frame ATG codons were identified, the 5′-most ATG codon was selected. The ORF beginning with the selected ATG codon and ending with the selected stop codon was extracted as a putative uORF (Fig. 3). The downstream sequences of putative uORFs were subjected to another tBLASTn analysis to examine whether

the transcripts were derived from homologs of the original uORF-containing gene. In this analysis, the amino

acid sequence of the mORF associated with the original uORF was used as a query sequence, and transcript

sequences matching the mORF with an E-value less than 10^{-1} were extracted. For each of the uORF-tBLASTn

and mORF-tBLASTn hits, the upstream in-frame stop codon closest to the 5′-end of the region matching the

original mORF was selected, and the 5′-most in-frame ATG codon located downstream of the selected stop

codon was identified as the putative mORF initiation codon (Fig. 3). uORF-tBLASTn and mORF-tBLASTn hits

were discarded as uORF-mORF fusion type sequences if the putative mORF overlapped with the putative uORF.

The original uORF was selected as a candidate CPuORF if the remaining uORF-tBLASTn and

mORF-tBLASTn hits belonged to at least two orders other than that from which the original uORF was derived.

Identification of contaminant ESTs and TSAs

In the uORF-tBLASTn and mORF-tBLASTn analyses described above, we excluded tBLASTn hit ESTs and

TSAs derived from contaminated organisms. To examine whether each uORF-tBLASTn and mORF-tBLASTn

hit sequence is derived from contaminated organisms, we performed BLASTn searches against EST, TSA, and

RefSeq RNA sequences from all organisms except for those of metagenomes, using each uORF-tBLASTn and

mORF-tBLASTn hit EST/TSA sequence as a query. If a uORF-tBLASTn hit EST/TSA sequence matched an

EST, TSA, or RefSeq RNA sequence of a different order from the species of the uORF-tBLASTn and

mORF-tBLASTn hit, with an E-value less than 10^{-100} and an identity equal to or greater than 95%, it was

considered a candidate contaminant sequence. In this case, either the uORF-tBLASTn and mORF-tBLASTn hit

or the BLASTn hit may be a contaminant sequence. To distinguish these possibilities, we compared the ratio of

the BLASTn hit number to the total EST/TSA and RefSeq RNA sequence number between the species of each

uORF-tBLASTn and mORF-tBLASTn hit and the species of its BLASTn hits. Appropriate comparisons are

difficult unless species used for this comparison have enough number of EST/TSA and RefSeq RNA sequences.

Therefore, if the total EST/TSA and RefSeq RNA sequence number of a species is less than 5000, BLASTn hits

derived from the species were not used for this analysis. If the ratio of the BLASTn hit number to the total

EST/TSA and RefSeq RNA sequence number of a uORF-tBLASTn and mORF-tBLASTn hit species is less

than that of any other BLASTn hit species, the uORF-tBLASTn and mORF-tBLASTn hit sequence was

identified as a contaminant sequence.

K_a/K_s analysis

For K_a/K_s analysis of each candidate CPuORF, a putative uORF sequence was selected from each order in which

uORF-tBLASTn and mORF-tBLASTn hits were found, using the following criteria. First, we selected transcript

sequences that matched the mORF associated with the candidate CPuORF with an *E*-value less than 10^{-20} in

mORF-tBLASTn analysis. Of these sequences, we then selected the one with the smallest geometric means of

mORF-tBLASTn and uORF-tBLASTn *E*-values. When mORF-tBLASTn *E*-values of all uORF-tBLASTn and

mORF-tBLASTn hits in an order were equal to or greater than 10^{-20}, we selected the transcript sequence with the

smallest geometric means of mORF-tBLASTn and uORF-tBLASTn *E*-values in the order. Putative uORF

sequences in the selected transcript sequences were used for generation of uORF amino acid sequence

alignments and for K_a/K_s analysis. Multiple alignments were generated by using standalone Clustal Omega

(ClustalO) ver. 1.2.2 [44]. For each candidate CPuORF, a median K_a/K_s ratio for all pairwise combinations of the

original uORF and its homologous putative uORFs was calculated using the LWL85 algorithm [45] in the seqinR

package [46].

For statistical tests of K_a/K_s ratios, we calculated the distribution of mutation rates between the original uORF

and its homologous putative uORFs and those between the original uORF and its artificially generated mutants,

using the observed mutation rate distribution. Then, observed empirical K_a/K_s ratio distributions were compared

with null distributions (negative controls) using the Mann-Whitney *U* test to validate statistical significance. The

one-sided *U* test was used to investigate whether the observed distributions were significantly lower than the null

distributions. Adjustment for multiple comparisons was achieved by controlling the false discovery rate using the

Benjamini and Hochberg procedure [47].

Determination of the taxonomic range of uORF sequence conservation

To automatically determine the taxonomic range of the sequence conservation of each CPuORF, we first defined

13 plant taxonomic categories. The 13 defined taxonomic categories are lamiids, asterids other than lamiids,

mavids, fabids, eudicots other than rosids and asterids, commelinids, monocots other than commelinids,

Angiospermae other than eudicots and monocots, Gymnospermae, Polypodiopsida, Embryophyta other than

Euphyllophyta, Streptophyta other than Embryophyta, and Viridiplantae other than Streptophyta. On the basis of

taxonomic lineage information of EST, TSA, and RefSeq RNA sequences, which were provided by NCBI

Taxonomy, the uORF-tBLASTn and mORF-tBLASTn hit sequences selected for K_a/K_s analysis were classified

into the 13 taxonomic categories (Fig. 4). It should be noted that, in NCBI taxonomy, eudicots, Angiospermae,

and Ggymnospermae are referred to as eudicotyledons, Magnoliophyta, and Acrogymnospermae, respectively.

For each CPuORF, the numbers of transcript sequences classified into each category were counted and shown in

Supplementary Table S2. These numbers represent the numbers of orders in which the amino acid sequence of

each CPuORF is conserved.

Statistical and informatic analyses

All programs, except for existing stand-alone programs, such as BLAST [43], ClustalO [44] and Jalview [48],

were written in R (www.r-project.org). We also used R libraries, GenomicRanges [49], exactRankTests,

Biostrings and seqinr [46].

Plasmid construction

Plasmid pNH006 harbors the cauliflower mosaic virus 35S RNA (35S) promoter, the Fluc coding sequence, and

the polyadenylation signal of the *A. thaliana HSP18.2* gene in pUC19. To construct this plasmid, pMT61 [26]

was digested with *Sal*I and *Sac*I, and the *Sal*I-*Sac*I fragment containing the Fluc coding sequence was ligated into

the *Sal*I and *Sac*I sites between the 35S promoter and the *HSP18.2* polyadenylation signal of plasmid pKM56

[38]. To generate reporter plasmids pNH92-pNH101 (Supplementary Table S3) for transient expression assays,

the 5´-UTR sequences of ten poplar genes were amplified by PCR from poplar (*Populus nigra*) full-length

cDNA clones pds25559, pds10965, pds14390, pds12940, pds13862, pds15817, pds28294, pds26157, pds14623

and pds23234 (Supplementary Table S4), obtained from RIKEN [50]. Primer sets used are shown in

Supplementary Tables S3 and S5. To construct pNH92, pNH94-pNH98 and pNH100-pNH101, amplified

fragments containing the 5´-UTR sequences were digested with *Xba*I and *Sal*I and ligated between the *Xba*I and

*Sal*I sites of pNH006. To create pNH93 and pNH99, amplified fragments containing the 5´-UTR sequences were

inserted between the *Xba*I and *Sal*I sites of pNH006 using the SLiCE method [51]. To make pHN102, the

5´-UTR sequence of the *Populus trichocarpa* POPTR_0013s08000 gene with a mutation at the initiation codon

of the uORF located immediately upstream of the HG107 CPuORF was synthesized by Fasmac (Atsugi, Japan)

on the basis of NCBI RefSeq accession no. XM_002319213.3. The synthesized 5´-UTR sequence was amplified

by PCR using primers 35S_XbaI_SLiCE-F and FLUC_SalI_SLiCE-R (Supplementary Table S5) and were

inserted between the *Xba*I and *Sal*I sites of pNH006 using the SLiCE method [51]. Frameshift mutations were

introduced into each CPuORF using overlap extension PCR [52], with primers listed in Supplementary Tables S4

and S5, to yield pNH103-pNH112. Sequence analysis confirmed the integrity of the PCR-amplified regions of all

constructs.

Transient expression assay

Transient expression assays were performed as described in Hayashi *et al*. 2017 [38]. Protoplasts from *A.*

thaliana MM2d suspension cells [53] were used, as were the reporter plasmids described above and the pKM5

[38] internal control plasmid. pKM5 contains the 35S promoter, the *Renilla* luciferase (Rluc) coding sequence,

and the *NOS* polyadenylation signal in pUC19. For each experiment, 5 µg each of a reporter plasmid and pKM5

were transfected into protoplasts.

Declarations

Funding

This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI [Grant Nos.

JP16H05063 to S.N., JP16K07387 to H.O., JP18H03330 to H.T and H.O., JP18K06297 to H.T, JP19H02917 to

H.O., JP19K22892 to H.T., JP19K22299 to H.O.]; the Ministry of Education, Culture, Sports, Science and

Technology (MEXT) KAKENHI [Grant Nos. JP26113519 to S.K, JP16H01246 to S.K, JP17H05658 to S.N.,

JP26114703 to H.T, JP17H05659 to H.T]; the Research Foundation for the Electrotechnology of Chubu to H.T;

and the Naito Foundation to H.O.

Authors' contributions

HT, HO, SK, KS, and SN conceived and designed the experiments. NH and YY performed the experiments. HT,

HO, TE, and AT analyzed the data. HT, NH, YY, AT, and KF contributed to reagents/materials/analysis tools.

HO and HT wrote the paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable.

Acknowledgements

We thank Ms. Kazuko Harada for general assistance.

List of abbreviations

35S: cauliflower mosaic virus 35S RNA; 5′-UTR: 5′-untranslated region; CPuORF: conserved peptide upstream

open reading frame; EST: expressed sequence tag; Fluc: firefly luciferase; HG: homology group; mORF: main

open reading frame; RefSeq: NCBI reference sequence; Rluc: Renilla luciferase; TSA: transcriptome shotgun

assembly; uORF: upstream open reading frame

References

1.   Churbanov A, Rogozin IB, Babenko VN, Ali H, Koonin EV. Evolutionary conservation suggests a regulatory function of AUG triplets in 5'-UTRs of eukaryotic genes. Nucleic Acids Res. 2005; 33:5512-5520.

2.   Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, Batzoglou S, et al. Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. Nature 2005; 438:1105-1115.

3.   Kawaguchi R, Bailey-Serres J. mRNA sequence features that contribute to translational regulation in *Arabidopsis*. Nucleic Acids Res. 2005; 33:955-965.

4.   Rogozin IB, Kochetov AV, Kondrashov FA, Koonin EV, Milanesi L. Presence of ATG triplets in 5' untranslated regions of eukaryotic cDNAs correlates with a 'weak' context of the start codon. Bioinformatics 2001; 17:890-900.

5.   Morris DR, Geballe AP. Upstream open reading frames as regulators of mRNA translation. Mol. Cell Biol. 2000; 20:8635-8642.

6.   Cruz-Vera LR, Sachs MS, Squires CL, Yanofsky C. Nascent polypeptide sequences that influence ribosome function. Curr. Opin. Microbiol. 2011; 14:160-166.

7.   Ito K, Chiba S. Arrest peptides: cis-acting modulators of translation. Annu. Rev. Biochem. 2013; 82:171-202.

8.   Somers J, Poyry T, Willis AE. A perspective on mammalian upstream open reading frame function. Int. J. Biochem. Cell Biol. 2013; 45:1690-1700.

9.   Bhushan S, Meyer H, Starosta AL, Becker T, Mielke T, Berninghausen O, et al. Structural basis for translational stalling by human cytomegalovirus and fungal arginine attenuator peptide. Mol. Cell 2010; 40:138-146.

10. Wang Z, Sachs MS. Ribosome stalling is responsible for arginine-specific translational attenuation in *Neurospora crassa*. Mol. Cell Biol. 1997; 17:4904-4913.

11.   Gaba A, Jacobson A, Sachs MS. Ribosome occupancy of the yeast *CPA1* upstream open reading frame

termination codon modulates nonsense-mediated mRNA decay. Mol. Cell 2005; 20:449-460.

12.   Uchiyama-Kadokura N, Murakami K, Takemoto M, Koyanagi N, Murota K, Naito S, et al.

Polyamine-responsive ribosomal arrest at the stop codon of an upstream open reading frame of the

AdoMetDC1 gene triggers nonsense-mediated mRNA decay in *Arabidopsis thaliana*. Plant Cell

Physiol. 2014; 55:1556-1567.

13.   Wang Z, Gaba A, Sachs MS. A highly conserved mechanism of regulated ribosome stalling mediated

by fungal arginine attenuator peptides that appears independent of the charging status of arginyl-tRNAs.

J. Biol. Chem. 1999; 274:37565-37574.

14.   Law GL, Raney A, Heusner C, Morris DR. Polyamine regulation of ribosome pausing at the upstream

open reading frame of S-adenosylmethionine decarboxylase. J. Biol. Chem. 2001; 276:38036-38043.

15.   Hanfrey C, Elliott KA, Franceschetti M, Mayer MJ, Illingworth C, Michael AJ. A dual upstream open

reading frame-based autoregulatory circuit controlling polyamine-responsive translation. J. Biol. Chem.

2005; 280:39229-39237.

16.   Yamashita Y, Takamatsu S, Glasbrenner M, Becker T, Naito S, Beckmann R. Sucrose sensing through

nascent peptide-meditated ribosome stalling at the stop codon of Arabidopsis *bZIP11* uORF2. FEBS

Lett. 2017; 591:1266-1277.

17.   Rahmani F, Hummel M, Schuurmans J, Wiese-Klinkenberg A, Smeekens S, Hanson J. Sucrose control

of translation mediated by an upstream open reading frame-encoded peptide. Plant Physiol. 2009;

150:1356-1367.

18.   Combier JP, de Billy F, Gamas P, Niebel A, Rivas S. Trans-regulation of the expression of the

transcription factor MtHAP2-1 by a uORF controls root nodule development. Genes Dev. 2008;

22:1549-1559.

19.   Crowe ML, Wang XQ, Rothnagel JA. Evidence for conservation and selection of upstream open

reading frames suggests probable encoding of bioactive peptides. BMC genomics 2006; 7:16.

20.   Hayden CA, Jorgensen RA. Identification of novel conserved peptide uORF homology groups in

Arabidopsis and rice reveals ancient eukaryotic origin of select groups and preferential association with

transcription factor-encoding genes. BMC Biol. 2007; 5:32.

21.   Hayden CA, Bosco G. Comparative genomic analysis of novel conserved peptide upstream open

reading frames in *Drosophila melanogaster* and other dipteran species. BMC genomics 2008; 9:61.

22.   Tran MK, Schultz CJ, Baumann U. Conserved upstream open reading frames in higher plants. BMC

genomics 2008; 9:361.

23.   Vaughn JN, Ellingson SR, Mignone F, Arnim A. Known and novel post-transcriptional regulatory

sequences are conserved across plant families. Rna 2012; 18:368-384.

24.   van der Horst S, Snel B, Hanson J, Smeekens S. Novel pipeline identifies new upstream ORFs and

non-AUG initiating main ORFs with conserved amino acid sequences in the 5' leader of mRNAs in

Arabidopsis thaliana. Rna 2018; 25:292-304.

25.   Takahashi H, Takahashi A, Naito S, Onouchi H. BAIUCAS: a novel BLAST-based algorithm for the

identification of upstream open reading frames with conserved amino acid sequences and its application

to the *Arabidopsis thaliana* genome. Bioinformatics 2012; 28:2231-2241.

26.   Ebina I, Takemoto-Tsutsumi M, Watanabe S, Koyama H, Endo Y, Kimata K, et al. Identification of

novel *Arabidopsis thaliana* upstream open reading frames that control expression of the main coding

sequences in a peptide sequence-dependent manner. Nucleic Acids Res. 2015; 43:1562-1576.

27.   Noh AL, Watanabe S, Takahashi H, Naito S, Onouchi H. An upstream open reading frame represses

expression of a tomato homologue of Arabidopsis *ANAC096*, a NAC domain transcription factor gene,

in a peptide sequence-dependent manner. Plant Biotechnol. 2015; 32:157-163.

28.   Jorgensen RA, Dorantes-Acosta AE. Conserved Peptide Upstream Open Reading Frames are

Associated with Regulatory Genes in Angiosperms. Front. Plant Sci. 2012; 3:191.

29.   Pruitt KD, Tatusova T, Maglott DR. NCBI reference sequences (RefSeq): a curated non-redundant

sequence database of genomes, transcripts and proteins. Nucleic Acids Res. 2007; 35:D61-D65.

30.   Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, et al. Ensembl 2018. Nucleic

677   Acids Res. 2018; 46:D754-D761.

678   31.   Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a comparative

679   platform for green plant genomics. Nucleic Acids Res. 2012; 40:D1178-D1186.

680   32.   Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons

681   dramatically improves orthogroup inference accuracy. Genome Biol. 2015; 16:157.

682   33.   Alatorre-Cobos F, Cruz-Ramirez A, Hayden CA, Perez-Torres CA, Chauvin AL, Ibarra-Laclette E, et al.

683   Translational regulation of Arabidopsis XIPOTL1 is modulated by phosphocholine levels via the

684   phylogenetically conserved upstream open reading frame 30. J. Exp. Bot. 2012; 63:5203-5221.

685   34.   Guerrero-Gonzalez ML, Ortega-Amaro MA, Juarez-Montiel M, Jimenez-Bremont JF. Arabidopsis

686   Polyamine oxidase-2 uORF is required for downstream translational regulation. Plant Physiol. Biochem.

687   2016; 108:381-390.

688   35.   Imai A, Hanzawa Y, Komura M, Yamamoto KT, Komeda Y, Takahashi T. The dwarf phenotype of the

689   Arabidopsis acl5 mutant is suppressed by a mutation in an upstream ORF of a bHLH gene.

690   Development 2006; 133:3575-3585.

691   36.   Ribone PA, Capella M, Arce AL, Chan RL. A uORF Represses the Transcription Factor AtHB1 in

692   Aerial Tissues to Avoid a Deleterious Phenotype. Plant Physiol. 2017; 175:1238-1253.

693   37.   Tabuchi T, Okada T, Azuma T, Nanmori T, Yasuda T. Posttranscriptional regulation by the upstream

694   open reading frame of the phosphoethanolamine N-methyltransferase gene. Biosci. Biotechnol.

695   Biochem. 2006; 70:2330-2334.

696   38.   Hayashi N, Sasaki S, Takahashi H, Yamashita Y, Naito S, Onouchi H. Identification of *Arabidopsis

697   thaliana* upstream open reading frames encoding peptide sequences that cause ribosomal arrest. Nucleic

698   Acids Res. 2017; 45:8844-8858.

699   39.   Franceschetti M, Hanfrey C, Scaramagli S, Torrigiani P, Bagni N, Burtin D, et al. Characterization of

700   monocot and dicot plant S-adenosyl-l-methionine decarboxylase gene families including identification

in the mRNA of a highly conserved pair of upstream overlapping open reading frames. Biochem. J.

2001; 353:403-409.

40.   Konishi T, Takeda T, Miyazaki Y, Ohnishi-Kameyama M, Hayashi T, O'Neill MA, et al. A plant

mutase that interconverts UDP-arabinofuranose and UDP-arabinopyranose. Glycobiology 2007;

17:345-354.

41.   Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs.

Genome Res. 2008; 18:821-829.

42.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat. Methods 2012; 9:357-359.

43.   Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and

PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 1997;

25:3389-3402.

44.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of

high-quality protein multiple sequence alignments using Clustal Omega. Mol. Syst. Biol. 2011; 7:539.

45.   Li WH. Unbiased estimation of the rates of synonymous and nonsynonymous substitution. J. Mol. Evol.

1993; 36:96-99.

46.   Charif D, Lobry JR: SeqinR 1.0-2: a contributed package to the R project for statistical computing

devoted to biological sequences retrieval and analysis. In *Structural Approaches to Sequence Evolution:*

Molecules, Networks, Populations. Edited by Bastolla U, Porto M, Roman HE, Vendruscolo M. New

York: Springer Verlag; 2007: 207-232

47.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to

multiple testing. J. R. Statist. Soc. ser.B 1995; 57:298-300.

48.   Clamp M, Cuff J, Searle SM, Barton GJ. The Jalview Java alignment editor. Bioinformatics 2004;

20:426-427.

49.   Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, et al. Software for computing

and annotating genomic ranges. PLoS Comput. Biol. 2013; 9:e1003118.

50.   Nanjo T, Futamura N, Nishiguchi M, Igasaki T, Shinozaki K, Shinohara K. Characterization of

727   full-length enriched expressed sequence tags of stress-treated poplar leaves. Plant Cell Physiol. 2004;

728   45:1738-1748.

729   51.   Motohashi K. Seamless Ligation Cloning Extract (SLiCE) Method Using Cell Lysates from Laboratory

730   Escherichia coli Strains and its Application to SLiP Site-Directed Mutagenesis. Methods Mol. Biol.

731   2017; 1498:349-357.

732   52.   Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. Site-directed mutagenesis by overlap extension

733   using the polymerase chain reaction. Gene 1989; 77:51-59.

734   53.   Menges M, Murray JA. Synchronous *Arabidopsis* suspension cultures for analysis of cell-cycle gene

735   activity. Plant J. 2002; 30:203-212.

Figure legends

Figure 1. Outline of the ESUCA pipeline.

Figure 2. Schematic representation of the algorithm to calculate uORF-mORF fusion ratios. For each original uORF-containing transcript sequence, RefSeq RNAs are selected that match an original uORF sequence, irrespective of the reading frame, and the original mORF sequence in the same reading frame as the largest ORF of the RefSeq RNA, using tBLASTx. The shaded regions in the open boxes represent the tBLASTx-matching regions. If the uORF-tBLASTx-matching region is within the largest ORF, the RefSeq RNA is considered a uORF-mORF fusion type. The number of this type of RefSeq RNA is defined as 'X'. If the uORF-tBLASTx-matching region is not within the largest ORF, the RefSeq RNA is considered a uORF-mORF separate type. The number of this type of RefSeq RNA is defined as 'Y'. For each of the original uORF-containing transcripts, the uORF-mORF fusion ratio is calculated as $X / (X + Y)$.

Figure 3. Schematic representation of BLAST-based search for uORFs conserved between homologous genes. In the third step of ESUCA, tBLASTn searches are conducted against a transcript sequence database that consists of assembled EST/TSA contigs, unclustered singleton EST/TSA sequences and RefSeq RNAs, using original uORF sequences as queries (uORF-tBLASTn). The shaded regions in the open boxes show the tBLASTn-matching regions. Asterisks represent stop codons. (i) The downstream in-frame stop codon closest to the 5′-end of the matching region of each uORF-tBLASTn hit is selected. (ii) The 5′-most in-frame ATG codon located upstream of the stop codon is selected. The ORF beginning with the selected ATG codon and ending with the selected stop codon is extracted as a putative uORF. In the fourth step of ESUCA, the downstream sequences of putative uORFs in the transcript sequences are subjected to mORF-tBLASTn analysis. Transcript sequences matching the original mORF with an E-value less than 10^{-1} are extracted. (iii) For each of the uORF-tBLASTn and mORF-tBLASTn hits, the upstream in-frame stop codon closest to the 5′-end of the matching region is selected. (iv) The 5′-most in-frame ATG codon located downstream of the selected stop codon is identified as the

initiation codon of the putative partial or intact mORF. If the putative mORF overlaps with the putative uORF, the

uORF-tBLASTn and mORF-tBLASTn hit is discarded as a uORF-mORF fusion type.

Figure 4. Schematic representation of the algorithm to determine the taxonomic range of uORF sequence

conservation. In the fifth step of ESUCA, for K_a/K_s analysis of each candidate CPuORF, a uORF-tBLASTn and

mORF-tBLASTn hit sequence is selected from each order that has uORF-tBLASTn and mORF-tBLASTn hits

(See Materials and Methods for the criteria for the selection). In the sixth step of ESUCA, the selected transcript

sequences are classified into the 13 plant taxonomic categories to determine the taxonomic range of uORF

sequence conservation, using taxonomic lineage information of EST, TSA, and RefSeq RNA sequences.

Figure 5. Taxonomic range of the sequence conservation of the CPuORF families. The blue, yellow, red, green,

and purple lines show the conservation range of CPuORF HGs determined by applying ESUCA to *Arabidopsis*

(*Arabidopsis thaliana*), rice (*Oryza sativa*), tomato (*Solanum lycopersicum*), poplar (*Populus trichocarpa*), and

grape (*Vitis vinifera*) genomes, respectively. The presence of a line within a cell in each taxonomic category

indicates the presence of uORF-tBLASTn and mORF-tBLASTn hits for any of the CPuORFs that belong to

each HG. In taxonomic categories with a category name with an asterisk, uORF-tBLASTn and

mORF-tBLASTn hits found in lower taxonomic categories were excluded. In the case where no

uORF-tBLASTn and mORF-tBLASTn hit was found in the taxonomic category that contain a species from

which the original uORF was derived, the line showing the species was still drawn in the cell of the taxonomic

category because this category contained the species with the original uORF. *Arabidopsis*, rice, tomato, poplar

and grape belong to malvids, commelinids, lamiids, fabids and eudicots*, respectively. HG1-HG53 are

previously identified HGs, except for HG2.2, HG9.2, HG16.2, HG43.2, HG50.2 and HG52.2, whereas HG2.2,

HG9.2, HG16.2, HG43.2, HG50.2, HG52.2 and HG54-HG130 are newly identified HGs.

Figure 6. Sequence-dependent effects of novel CPuORFs on main ORF translation. (A) Schematic

representation of the WT (*35S::UTR(WT):Fluc*) and frameshift mutant (*35S::UTR(fs):Fluc*) reporter constructs.

The 5′-UTR sequence containing each of the 11 poplar CPuORFs was inserted between the 35S promoter

(*35Spro*) and the *Fluc* coding sequence. The hatched box in the frameshift (fs) mutant CPuORF indicates the

frame-shifted region. The dotted boxes represent the first five nucleotides of the mORF associated with each of

the 11 poplar CPuORFs. See Supplementary Figure S2 for the exact position and length of each CPuORF and the

exact frame-shifted region. The polyadenylation signal of the *A. thaliana HSP18.2* gene is designated as 'ter'. (B)

Transient expression assay. Each reporter plasmid containing a WT or fs mutant CPuORF, belonging to an

indicated HG, was co-transfected into MM2d protoplasts with the *35S::Rluc* internal control plasmid by PEG

treatment. After incubation for 24 h, cells were collected and disrupted for dual luciferase assay. Fluc activity was

normalized to Rluc activity, and the activity relative to that of the corresponding WT reporter construct was

calculated. Means ± SD of three biological replicates are shown. Single and double asterisks indicate significant

differences between the WT and fs constructs at $p < 0.05$ and $p < 0.01$, respectively, as determined by Student's

t-test. Representative results of two experiments using independently prepared protoplasts are shown.

Figure 1

Step 1: Extraction of uORF sequences from a transcript sequence dataset

Step 2: Calculation of uORF-mORF fusion ratios

uORF-mORF fusion ratio < 0.3

uORF-mORF fusion ratio ≥ 0.3

Discarded

Step 3: Homology searches of the uORF amino acid sequences against a transcript sequence database (uORF-tBLASTn)

uORFs with tBLASTn hits from the other orders

uORFs with no tBLASTn hits from the other orders

Discarded

Step 4: Selection of uORFs conserved among homologous genes (mORF-tBLASTn)

uORFs conserved among homologs from more than two orders

uORFs not conserved among homologs from more than two orders

Discarded

Step 5: Calculation of K_a/K_s ratios

K_a/K_s ratio < 0.5 and q < 0.05

K_a/K_s ratio ≥ 0.5 or q ≥ 0.05

Discarded

Step 6: Determination of the taxonomic range of uORF sequence conservation

Figure 2

Original uORF-containing transcript sequence

Original uORF-containing transcript sequence

$$\text{uORF-mORF fusion ratio} = \frac{X}{X + Y}$$

Figure 3

Original uORF-containing
transcript sequence

uORF-tBLASTn and mORF-tBLASTn
hit sequence (Assembled EST/TSA
and RefSeq RNA)

Figure 4

uORF-tBLASTn and mORF-tBLASTn hits
selected for K_a/K_s analysis

Figure 5

Figure 5 (continued)

Figure 6