Protein features for assembly of the RNA editing helicase 2 subcomplex (REH2C) in

Trypanosome holo-editosomes

Vikas Kumar[1#], Pawan K. Doharey[1#], Shelly Gulati[2#], Joshua Meehan[1], Mary G. Martinez[1],

Karrisa Hughes[2], Blaine H.M. Mooers[2*] and Jorge Cruz-Reyes[1*]

[1] Department of Biochemistry and Biophysics, Texas A&M University, College Station, Texas,

United States of America.

[2] Department of Biochemistry & Molecular Biology, University of Oklahoma Health Sciences

Center, Oklahoma City, Oklahoma, United States of America.

* Corresponding author:

e-mail: cruzrey@tamu.edu

* Correspondence may also be addressed to:

e-mail: blaine-mooers@ouhsc.edu

# These authors contributed equally to this work.

Abstract

Uridylate insertion/deletion RNA editing in *Trypanosoma brucei* is a complex system that is not found in humans, so there is interest in targeting this system for drug development. This systeme uses hundreds of small non-coding guide RNAs (gRNAs) to modify the mitochondrial mRNA transcriptome. This process occurs in holo-editosomes that assemble several macromolecular *trans* factors around mRNA: the RNA-free RNA editing core complex (RECC) and auxiliary RNPs. Yet, the regulatory mechanisms of editing remain obscure. The enzymatic accessory RNP complex, termed the REH2C, includes mRNA substrates and products, the multi-domain 240 kDa RNA Editing Helicase 2 (REH2) and an intriguing 8-zinc finger protein termed REH2-Associated Factor 1 ([H2]F1). Both proteins are essential in editing. REH2 is a member of the DExH/RHA subfamily of RNA helicases with a conserved C-terminus that includes a regulatory OB-fold domain. In trypanosomes, [H2]F1 recruits REH2 to the editing apparatus, and [H2]F1 downregulation causes REH2 fragmentation. Our systematic mutagenesis dissected determinants in REH2 and [H2]F1 for the assembly of REH2C, the stability of REH2, and the RNA-mediated association of REH2C with other editing *trans* factors. We identified functional OB-fold amino acids in eukaryotic DExH/RHA helicases that are conserved in REH2 and impact the assembly and interactions of REH2C. [H2]F1 upregulation stabilized the large REH2 polypeptide *in vivo*. Mutation of the core cysteines or basic amino acids in individual zinc fingers affected the stabilizing property of [H2]F1 but not its interactions with other examined editing components. Thus, most if not all fingers may contribute to REH2 stabilization. Finally, a recombinant REH2 (240 kDa) established that the full-length protein is a *bona fide* RNA helicase with ATP-dependent unwinding activity. REH2 is the only DExH/RHA-type helicase in kinetoplastid holo-editosomes.

Introduction

Kinetoplastid protozoa including *Trypanosoma brucei* are early-branching eukaryotes that require extensive insertion and deletion of uridylates in most mitochondrial mRNAs (reviewed in 1). This RNA editing is protein-catalyzed and directed by small guide RNAs (gRNAs) that exhibit complementarity to fully edited mRNA via Watson-Crick and wobble G•U base pairs. Most editing progresses in blocks that typically overlap and each block is directed by a single gRNA (2). The editing machinery includes multiple subcomplexes: the ~20S catalytic RNA editing core complex (RECC, also called the 20S editosome) and auxiliary editing RNPs. The ~15S REH2-associated complex (REH2C) characterized here includes RNA editing helicase 2 (REH2), REH2-associated protein factor 1 ([H2]F1), [H2]F2, and mRNA substrates and products (3, 4). However, only REH2 and [H2]F1 participate in editing. REH2C interacts physically and functionally with a larger accessory editing RNP complex termed the RNA editing substrate binding complex (RESC), which also contains mRNA substrates and products (5, 6). The presence of mRNA in RESC was confirmed by others (7, 8). RESC contains two modules that are intimately associated with each other: the gRNA-binding complex (GRBC) typified by the GAP1/GAP2 (alias GRBC2/GRBC1) heterotetramer that binds and stabilizes gRNA, and the more loosely defined RNA editing mediator complex (REMC) typified by RGG2 (6). REMC includes proteins that promote editing progression (8-10). Other mRNA *trans* factors include a subcomplex containing MRB6070 and MRB1590, and RNA helicase REH1 that associate with holo-editosomes via stable or transient RNA contacts, respectively (11-13). The editing apparatus is further complicated by the occurrence of variants of GRBC, REMC and RECC (reviewed in 1). The dynamic nature of the holo-editosome may be relevant in editing control and is reminiscent of the intricate molecular dynamics of the RNP complexes involved in mRNA splicing and transcription.

The finding that the editing RNPs described above carry mRNA and gRNA implied that

mRNA-gRNA hybrids form on multi-RNP platforms, and that transient addition of the RECC

enzyme to the substrate-loaded platforms completes the assembly of the holo-editosome (3-8).

The precise control mechanisms in editing remain unclear. However, the helicase REH2C

subcomplex may play an important regulatory role. REH2 is the largest characterized member

of the DExH/RNA helicase A (RHA) subfamily of proteins (14). These proteins are monomeric

and contain a conserved C-terminal domain cluster that includes the characteristic auxiliary

oligonucleotide-binding (OB fold) domain (3, 14-16). The OB fold in some RNA helicases is

known to bind protein regulators or mediate RNA-dependent activation of these enzymes (17).

DExH/RHA proteins have a N-terminus with a variable domain organization of less clear

function. The N-terminal half of the REH2 carries two predicted double-stranded RNA binding

domains: dsRBD1 and dsRBD2 (14). Other characterized DExH/RHA-type proteins that carry

two N-terminal dsRBDs are the RNA helicase RHA (aka DHX9) in vertebrates and its

orthologous RNA helicase MLE (*maleless*) in flies (18). Recombinant versions of the REH2 N-

terminal half and full-size octa-zinc finger ^{H2}F1 formed a stable complex *in vitro* (4). However,

^{H2}F1 is an unusual binding partner of DExH/RHA proteins that typically associate with G-patch

proteins (17). A genetic knockdown of ^{H2}F1 prevented RNA-mediated association of REH2 with

other editing components and caused fragmentation of REH2 *in vivo*, so ^{H2}F1 serves as an

adaptor protein and may stabilize REH2 (4).

The current study identified features in REH2 and ^{H2}F1 that affect the binding of these

proteins with each other, the stability of REH2, and RNA-mediated stable or transient

association of the REH2C subcomplex with other editing components *in vivo*. About 20 different

protein variants were examined *in vivo*. Comparisons of the REH2 with other eukaryotic

DExH/RHA helicases identified conserved OB fold residues that affect the assembly and

interactions of REH2C and are also important in distant processes, namely mRNA splicing in

yeast, and the assembly of the dosage compensation complex in fly. ^{H2}F1 upregulation

stabilized the large REH2 polypeptide *in vivo*. However, mutation of individual zinc fingers

compromised the stabilizing property of ^{H2}F1 but not its association with REH2 or other

examined components of the editing apparatus. Thus, most if not all fingers may contribute to

REH2 stabilization. Finally, we used a recombinant protein construct to establish that the

isolated full-length REH2 (240 kDa) is a *bona fide* RNA helicase enzyme with ATP-dependent

unwinding activity. REH2C is most likely an enzymatic editing RNP in kinetoplastid holo-

editosomes.

Results

Previous studies revealed that the native REH2C exhibits RNA-mediated interactions

with at least two variants of RESC (3-5). These variants contain gRNA and comparable levels of

GAP1 and RGG2, that typify GRBC and REMC, respectively. However, these variants differ

minimally in their relative content of the canonical MRB3010 in GRBC. Additional variations in

protein composition have been reported in purifications of RESC components by different labs

(reviewed in 1). Interestingly, native and genetically-induced changes in MRB3010 stoichiometry

may be common. While the precise reasons of these changes are unclear, we reasoned that

they reflect relevant dynamic changes in the assembly and function of the editing apparatus. In

this study, we examined protein features that affect the formation of REH2C, its stable

association with RESC components, and transient contacts with the RECC enzyme. Because

the examined helicase-associated RESC variant exhibits substoichiometric MRB3010, we refer

to its variant GRBC module as GRBC* (4). We have detected other proteins in this RESC

variant using antibodies made available to us, including: MRB8170 and MRB6070 (data not

shown). However, further analysis of relative composition between variants is needed.

Analysis of REH2 mutants *in vivo*
124
We analyzed the importance of specific features of REH2, both in the formation of REH2C

and in the association of this subcomplex with other components in holo-editosomes (constructs

listed in Fig. 1A). To this end, we expressed a tetracycline-inducible tagged REH2 wild-type (WT)

construct and examined its co-purification with endogenous [H2]F1 and GAP1 proteins, in the REH2C

and GRBC, respectively. We found that ectopic REH2 was able to reconstitute interactions of the

native REH2 protein in trypanosomes (Fig. 1B-C, lanes 1-2, respectively). With this system in hand,

we began testing the assembly of REH2 variants bearing truncations or point mutations in an

approach that has been used in other eukaryotic DExH/RHA helicases (16, 19). Neither

overexpression or depletion of REH2 affects the steady-state level of its binding partner [H2]F1 or

the examined GRBC proteins (3, 4).

Fig. 1. *In vivo* analyses of REH2 variants. (A) Scheme of REH2 WT (2167 residues) and

variants with the examined truncations or point mutations in the N- and C-terminal regions (NTR

and CTR). The mutations, affected domains, and predicted molecular weight of the constructs

are listed. Identified domains: dsRBD1 and dsRBD2 (double-stranded RNA binding domains 1

and 2), CORE (catalytic core RecA1 and RecA2 domains), HA2 (helicase associated helical

bundle domain), and OB (OB fold-like domain). **(B-C)** Western blots of IgG pulldowns from

extracts to examine the association of tagged REH2 WT or deletion mutants with endogenous

[H2]F1 and GAP1. An uninduced (-Tet) control lane is included. The GAP1 panels derive from

equivalent pulldowns with the same recovery of tagged protein (data not shown). The data are

representative of at least three biological replicates for each mutant. A few non-specific species

cross-react with the antibodies in some assays. The position of IgG is marked. Sizing markers

(M) are in kDa. **(D)** Charts of the relative level of ^{H2}F1 and GAP1 in the pulldowns.

Representative assays from panels A-B and other pulldowns in this study were used to plot +/-1

SD, *n*=3 for each mutant. The ^{H2}F1/REH2 and GAP1/REH2 ratios in each REH2 mutant were

normalized to the ratios in REH2 WT (= 1, in the plot). Assays with signals that too low were not

measured (not determined, n.d.). REH2 was often fragmented, so only the full-length

polypeptide was scored. Some short variants were present at higher level than REH2 WT in the

induced extracts.

N- and C-truncations. We examined N- and C-terminal regions (NTR and CTR) flanking the

catalytic and helicase associated domains RecA1/RecA2 and HA2, respectively, in REH2. We

removed sequences in the unique NTR (>1000 residues) in the REH2 variants ΔN and ΔNds, or in

the CTR after the OB fold (123 residues) or from the OB fold (260 residues) in the ΔAOB and ΔOB

variants, respectively. REH2 and its orthologs in kinetoplastids are the largest known DExH/RHA-

type helicases and carry the longest helicase-specific N terminus (~1300 residues in REH2 from *T.*

brucei) (14). Direct comparisons with the REH2 WT construct showed that the examined N- and C-

resections reduced the association of the shortened REH2 polypeptides with ^{H2}F1 and GAP1 proteins

(Figs. 1B-D). This suggested that both termini of REH2 may contribute to the normal assembly of

REH2C and its RNA-mediated interactions with other editing components *in vivo*. Motifs in the

dsRBDs, the OB fold, and undefined features in the N and C termini may be involved. It is also

conceivable that the examined sequence resections impact the global conformation of REH2, thereby

indirectly affecting its protein interactions. Thus, the normal REH2 interactions in the editing

apparatus may be sensitive to overall changes in the integrity or conformation of REH2. Because the

CTR includes a potentially regulatory OB fold in DExH/RHA-type RNA helicases (16), we focused on

this domain in the subsequent studies of REH2.

Point mutations. Our structural searches of the OB fold-like domain in REH2 using Phyre2

(20) found the best agreements with reported structures of yeast ADP-bound Prp43p helicase and

Drosophila RNA helicase MLE RNA ADP Alf4 complex (4, 16, 18)(and data not shown). Prp43p is a

bi-functional helicase in mRNA splicing and ribosome biogenesis (15, 21). The helicase MLE

remodels roX RNA substrates to promote assembly of the dosage compensation complex in

Drosophila (18). We examined specific residues in the predicted OB fold of REH2 that seem to be

conserved in reported DExH/RHA helicase structures (Fig. 2 A-D). OB point mutations were

compared to two controls: REH2 WT and a reported mutation in dsRBD2 (dsR) that reduces the

association of REH2 with GAP1, gRNA and mRNA (Fig. 2A, lanes 1 and 2, respectively) (3). The

$\beta1-\beta2$ and $\beta4-\beta5$ loops in the OB fold may include determinants in nucleic acid recognition (Fig. 2D)

(16, 18). In helicase Prp43p, mutation of R664 or K704, which are exposed in the Prp43p structure,

decreased the RNA affinity and ATPase activity of the helicase *in vitro* (16). Sequence alignments

and a homology model using Prp43p as the template matched R664 and K704 in Prp43p with R1979

and R2023 in REH2, respectively (data not shown). Pulldowns of the R1979A and R2023A mutants

suggest that neither mutation alters the association of REH2 with [H2]F1 (Figs. 2A-C; and data not

shown). R1979A did not appear to affect the REH2 association with GAP1. However, R2023A

caused a moderate decrease in the association with GAP1, implying that at least R2023A could be

disruptive in trypanosomes. In helicase MLE, a few identified residues in the OB fold bind specific

uridylates in U-rich sequences in roX transcripts, and mutation of some of these residues impaired

RNA binding by MLE *in vitro* or the proper localization of MLE in chromosomes (18). Sequence

alignments and homology models indicated that the residues H1032 and K1033 in MLE likely

represent residues H1998 and R1999 in the $\beta3-\beta4$ loop in the OB fold in REH2 (Fig. 2D; S1 Fig; and

data not shown). The H1998E mutant appeared to retain a normal association with [H2]F1 but exhibited

a moderate decrease in association with GAP1. In contrast, R1999E either alone or together with

H1998E (98/99) exhibited a more robust decrease (over 60%) in association with both ^{H2}F1 and

GAP1 (Fig. 2A-C). This indicated that H1998 and R1999 contribute to the assembly of the REH2

helicase with other editing components. However, R1999 has a higher impact on such interactions

than the adjoining H1998. As expected, the dsR mutant used as a control exhibited a strong

decrease in association with GAP1. However, the dsR mutation did not seem to affect the

association of REH2 with ^{H2}F1 (Figs. 2A, 2C). Pulldowns of the mutants were normalized to the

REH2 WT control.

Fig. 2. *In vivo* analyses of REH2 variants. (A) Western blots of IgG pulldowns as in Fig. 1 to

examine the association of endogenous ^{H2}F1 and GAP1 with tagged REH2 WT or variants with

point mutations in the OB fold domain: single at R1979A (1979), H1998E (1998) and R1999E

(1999), or double at 1998/1999 (98/99). The previously characterized dsR mutant with an

inactivating double mutation in the dsRBD2 was included for comparison (3). The GAP1 panels

in Figs. 2A-B derived from identical pulldowns with the same recovery of tagged protein (data

not shown). A few non-specific species crossreact with the antibodies in some assays. **(C)**

Charts of the relative levels of ^{H2}F1 and GAP1 in the pulldowns. Assays from panels A-B and

biological replicate pulldowns in this study were used to plot +/-1 SD, *n*=2 or *n*=3 for each

mutant. The ^{H2}F1/REH2 and GAP1/REH2 ratios in each REH2 mutant were normalized to the

ratios in REH2 WT (= 1, in the plot). **(D)** Secondary structure prediction of the OB fold with α-

helix (cylinders) and β-strand (arrows) elements, and the position of point mutations examined

indicated by arrows.

Together, our systematic mutagenesis of REH2 suggests an intricate interaction network

including features across both N- and C- terminal regions of REH2 that affect the assembly of

REH2C, and the association of this subcomplex with other components of the editing apparatus *in*

vivo. Remarkably, these observations also indicate that equivalent residues in the auxiliary

conserved OB fold domain in DExH/RHA helicases function in disparate RNA processes in

taxonomically distant species of trypanosomes, yeast, and fly. While large protein truncations in

REH2 may alter the global conformation of the polypeptide, relevant point mutations in REH2 are

more likely to cause specific functional effects. Table 1 summarizes observed effects of the examined

mutations in REH2 on its association with ^{H2}F1 and GAP1.

Table 1. Association of REH2 constructs with ^{H2}F1 and GAP1

REH2-TAP variant	Association in pulldown		Reference
	^{H2}F1	GAP1	Construct described
WT	control*	control*	Hernandez et al. 2010
ΔN	decreased	decreased	This study
ΔNds	decreased	decreased	This study
ΔOB	decreased	decreased	This study
ΔAOB	decreased	decreased	This study
dsR	control-like †	decreased	Madina et al. 2016
1979	control-like	control-like	This study
1998	control-like	decreased	This study
1999	decreased	decreased	This study
98/99	decreased	decreased	This study
2023	control-like	decreased	This study

  * The WT construct served as positive control to assess relative association of mutant constructs.

228   † Comparable to the level observed with the WT construct.

Analysis of ^{H2}F1 mutants *in vivo*

To analyze specific features of the ^{H2}F1 zinc finger protein in trypanosomes, we expressed a

tagged ^{H2}F1 WT construct and examined its association with endogenous REH2, GAP1 and REL1

proteins. We found that the tagged ^{H2}F1 WT was able to reconstitute interactions of the native ^{H2}F1

protein in trypanosomes (Fig. 3). Notably, expression of the ectopic ^{H2}F1 WT increased the steady-

state level of endogenous REH2 compared to a control lane without tetracycline (Fig. 3A, lanes 1-2;

Fig. 3B). As a control to mark the position of REH2 we used a reported RNAi-based knockdown of

^{H2}F1 that decreases the level of endogenous REH2 (Fig. 3A, lanes 5-7; Fig. 3B)(4). The contrasting

effects of the upregulation and downregulation of ^{H2}F1 on the endogenous REH2 are consistent with

the idea that [H2]F1 stabilizes REH2 *in vivo*. These effects by [H2]F1 also imply that the interaction

between REH2 and [H2]F1 may vary *in vivo* or that these proteins are not always associated. We

examined canonical proteins in REH2C, RESC and RECC in sedimentation analyses of

mitochondrial extract (S2 Fig). REH2 is heterodispersed as shown before (22) and overlaps with [H2]F1

in the gradient. However, the relative abundance of these proteins at low and high densities differs

substantially. It is indeed possible that the association between REH2 and [H2]F1 or their stoichiometry

in REH2C varies *in vivo*. Additional studies will be necessary to examine these possibilities.

Fig 3. Effect of expressing [H2]F1 WT or a variant lacking Znf5 (core residues C2>A2) on

the REH2 steady state. (A) Western blots of extracts with tagged [H2]F1 WT (lanes 1-2) or Znf5

mutant (Z5; lanes 3-4). Control assays include lanes without Tet-induction. Also, control lanes

with a reported [H2]F1-RNAi construct mark the position of the endogenous REH2 and [H2]F1. The

RNAi lanes show the induced decrease of [H2]F1 and concurrent fragmentation of REH2 full-

length (4). The level of endogenous REH2 was examined. GAP1 was used as a loading control

in all assays in this panel. Molecular size markers (M). A typical C2H2 zinc finger domain fold is

depicted with core cysteine substitutions C2>A2 (arrows), other conserved residues, and

variable residues (black circles). **(B)** Charts of the relative levels of endogenous REH2 full-

length in the extracts +/-1 SD, *n*=3. Induced/uninduced ratios of REH2 for each construct were

further normalized to GAP1. **(C-D)** IgG pulldowns of WT and Z5 constructs showing association

with endogenous REH2, GAP1, REL1, and endogenous [H2]F1. Non-specific species (*) may

represent fragments of the tagged protein that react with the antibodies. **(E)** Charts of the

relative levels of REH2, GAP1 and REL1 in the pulldowns +/-1 SD, *n*=3. The ratios

(protein/tagged-bait) in the Z5 mutant were normalized to ratios in the [H2]F1 WT construct.

263
To determine if the stabilizing effect of [H2]F1 requires one or more zinc finger (Znf) domains,

we introduced amino acid substitutions that either disrupt the canonical fold of the domain (by

replacing the canonical residues) or eliminate the positive charge (by replacing various basic

residues with neutral residues) in individual fingers of the ectopic construct. The examined ^{H2}F1

mutants are summarized in Table 2.

Table 2. ^{H2}F1 mutants with reduced ability to stabilize REH2 *in vivo*

^{H2}F1-TAP variant	Endogenous REH2	Reference
	Steady-state level	Construct described
WT	Increased*	This study
ZnF1 +>A	no change†	This study
ZnF2 +>A	no change	This study
ZnF3 +>A	no change	This study
ZnF4 +>A	no change	This study
ZnF5 +>A	no change	This study
ZnF5 C2>A2	no change	This study
ZnF1-8 C2>A2&	no change	This study
ZnF6-8 C2>A2&	no change	This study
ZnF1-4 C2>A2&	no change	This study
ΔN	no change	This study
ΔC	no change	This study

* ^{H2}F1-WT ectopic expression increased the level of endogenous REH2

†   No change (i.e., level within stardard deviation of at least two biological replicates)

&   Not shown. Ectopic expression was unstable or eventually lost in culture.

Canonical cysteine residues. We changed two cysteines to two alanines (C2>A2) to disrupt

the native conformation of the predicted zinc fingers (Fig. 3A). We began by disrupting Znf5 because

of its significant homology to a reported dsRNA-bound C2H2 zinc finger domain structure (4, 14, 23).

The resulting mutant protein (^{H2}F1 minus Znf5) did not induce an evident increase in the steady-state

level of endogenous REH2 as the WT ^{H2}F1. This was evident in direct comparisons of extracts that

contain similar levels of ectopic ^{H2}F1 protein, WT or Znf5-less (Fig. 3A, lanes 1-4; Fig. 3B). This

differential effect between the ^{H2}F1 WT and Znf5-less variants was similarly observed in cultures 1 or

3 days after induction (data not shown). So, the stabilizing property of ^{H2}F1 is independent of the

period of induction *in vivo*. To examine whether the lack of Znf5 affected the association of ^{H2}F1 with

other editing components, we performed IgG pulldown assays of tagged ^{H2}F1. These assays showed

that the ^{H2}F1 WT and Znf5-less variants similarly associated with endogenous helicase REH2, GAP1

and REL1 ligase (Figs. 3C, 3E). Notably, pulldowns of both tagged [H2]F1 WT and Znf5-less, contain

endogenous [H2]F1 (Fig. 3D). Thus, Znf5 is not essential for the normal association of [H2]F1 with other

editing components.

286   We also tested a [H2]F1 variant bearing a triple C2>A2 substitution in Znf6, Znf7 and Znf8. The

287   [H2]F1 triple mutant failed to stabilize endogenous REH2 but it was able to bind REH2 and GAP1 as

288   we observed with both [H2]F1 WT and Znf5-less constructs (data not shown). Expression of this

289   construct was unstable and eventually lost in culture (data not shown). These observations suggest

290   that [H2]F1 may require a full complement of zinc fingers in their native conformation to stabilize REH2

291   but not to associate with the examined editing components.

293   *Variable basic residues*. Because the native conformation of the zinc fingers seems to be

required for [H2]F1 to stabilize REH2 we introduced mutations to decrease the positive charge in the

zinc fingers. Yet these mutated fingers are likely to retain their native conformation (24). We

expressed [H2]F1 variants to test the role of several basic residues around the core residues that are

mostly located between the second cysteine and the first histidine (Fig. 4A) (25). Specifically, we

introduced R/K>A substitutions to decrease the positive charge in the zinc fingers and to remove

potential charge-charge interactions with REH2 or associated RNAs. Our functional analyses of Znf1,

Znf2, Znf3 or Zn5 showed that the loss of positive charge in any of these zinc fingers reduced the

ability of [H2]F1 to induce the accumulation of intact endogenous REH2 in extracts (Figs. 4B-C).

However, these mutations did not prevent the association of [H2]F1 with REH2, GAP1 and REL1 (Figs.

4D, 4F). These expressed [H2]F1 variants also co-purified with endogenous [H2]F1 at a level comparable

to the WT construct (Figs. 4D-F). So, the wildtype and R/K>A variant proteins associated similarly

with the examined editing components. Overall, [H2]F1 may require that most zinc fingers retain their

native conformation and positive charge to efficiently increase the stability of REH2. However, the

loss of the native conformation or positive charge in individual fingers does not preclude the

assembly of [H2]F1 in the REH2C, or the association of this subcomplex with the examined editing

components. Detection of endogenous [H2]F1 in our pulldowns suggest the presence of [H2]F1

multimers. Pulldowns treated with an RNaseA/T1 mixture suggest that the association between

tagged [H2]F1 and endogenous [H2]F1 is RNase-resistant (S3 Fig.). However, further studies using

isolated recombinant proteins are needed to establish direct protein contacts between [H2]F1

monomers as it was shown between REH2 and [H2]F1 (4).

Fig. 4. *In vivo* analyses of [H2]F1 variants with reduced positive charge in Znf1, Znf2, Znf3

or Znf5 (variable residues R/K>A). (A) Cartoon of [H2]F1 (524 residues). Multi-sequence

alignment of the eight C2H2 zinc fingers, indicating the core cysteines and histidines (boxes)

and variable basic residues in each finger (open circles). Depiction of a C2H2 zinc finger

domain fold, and variable residue substitutions R/K>A. **(B)** Extracts as in Fig. 3A of induced

tagged [H2]F1 WT and Z1, Z2, Z3 and Z5 mutants or uninduced control. GAP1 was used as a

loading control (lanes 1-7). **(C)** Chart of steady-state level of endogenous REH2 full-length in

extracts with each construct as in Fig. 3B. **(D-E)** IgG pulldowns of [H2]F1 WT and Z1, Z2, Z3 and

Z5 mutants showing association with endogenous REH2, [H2]F1, REL1 and GAP1. Non-specific

species in pulldowns (*) are indicated. (**F)** Charts of the relative levels of associated

endogenous proteins in the pulldowns +/-1 SD, *n=2* or *n*=3, as in Fig. 3.

N- and C-truncations. Because mutagenesis of the tested Znf domains did not prevent the

association of [H2]F1 with other editing components *in vivo*, we asked if fragments of [H2]F1 retain these

interactions. To test this question, we expressed N- or C-terminal fragments including Znf1-to-Znf3

(termed ΔC) or Znf4-to-Znf8 (termed ΔN) that span 49% and 51% of the [H2]F1 polypeptide chain,

respectively (Fig. 5A, bottom). These two fragments exhibited minimal or no association with the

examined endogenous REH2, [H2]F1, or REL1 ligase (Figs. 5A). Independent pulldowns of the [H2]F1

truncated constructs confirmed these defects in association (data not shown). These mutants were

directly compared with the [H2]F1 WT and Znf4 (R/K>Ala) constructs. The Znf4 like other Znf mutations

in the current study retained all tested interactions that are observed with the [H2]F1 WT control. Thus,

the ΔN and ΔC constructs failed to establish basic interactions with the tested editing components.

Because the native REH2C subcomplex carries mRNA (4), we wondered if the [H2]F1 ΔN and ΔC

constructs also fail to co-purify with mRNA. To this end, we quantitated the relative content of

mitochondrial transcripts in the pulldowns of the WT, ΔN, and ΔC constructs in biological replicate

experiments that were performed in triplicate (Fig. 5B). The examined mitochondrial transcripts

include examples of unedited, edited, and never-edited mRNA, the never-edited mRNA ND4, and

ribosomal RNA 9S. We also examined cytosolic transcripts (Tubulin, 1400 and 1390) to score non-

specific RNA interactions of editing subcomplexes as in previous pulldown studies (5). Relative to the

[H2]F1 WT control used for normalization, both ΔN and ΔC pulldowns exhibited a dramatic loss in

bound mitochondrial mRNAs (Fig. 5B, upper and middle panels). Interestingly, the levels of cytosolic

mRNAs were similar in the ΔN and WT proteins but dramatically less in the ΔC variant. We also

compared the relative impact of the ΔN and ΔC deletions with a more discrete alteration in the Znf4

construct (R/K>Ala); Fig. 5B, lower panel). The level of mitochondrial RNAs in the Znf4 pulldown was

similar to that of the [H2]F1 WT control. However, the level of cytosolic transcripts decreased ~70 fold

in Znf4. The independent pulldowns and RNA quantitation in three equivalent experiments showed

consistent results in the protein interactions and RNA content of these [H2]F1 constructs. (Fig. 5A).

Thus, the tested [H2]F1 fragments appear to have lost the normal direct interaction of [H2]F1 WT with

REH2 in REH2C and the RNA-dependent association of [H2]F1 WT with other examined proteins in the

editing apparatus. Quantitation of the examined transcripts in the input extracts (including those used

in the pulldowns in Figs. 5A-B) confirmed that the observed dramatic loss in RNA association with ΔN

and ΔC constructs was not due to large changes in the transcripts levels at steady state (S4 Fig.).

The above observations indicated that features in both termini of [H2]F1 are required for efficient

association of mitochondrial transcripts in the pulldowns of [H2]F1. Also, C-terminal features in [H2]F1

potentially including Znf4-to-Znf8, may be largely responsible for the observed non-specific

association of cytosolic transcripts in the pulldowns of ^{H2}F1 (Fig. 5B, compare the top and middle

panels). Finally, the more discrete mutation in Znf4 may have caused a relatively mild decrease in

RNA affinity that mostly affected weak, non-specific interactions, e.g., by cytosolic RNA in the

pulldowns of ^{H2}F1.

Figure 5. *In vivo* analyses of ^{H2}F1 terminal deletion and Znf4 mutants. (A) IgG pulldowns of

N and C truncation mutants (ΔN and ΔC) that removed Znf1-3 and Znf4-8, respectively, and the

Znf4 R/K>A substituted (Z4) variant. Association of the constructs with endogenous REH2, ^{H2}F1

and REL1 was examined. Cartoon of ^{H2}F1 with zinc fingers and a line marking the N and C

truncations. **(B)** qRT-PCR quantitation of RNA transcripts in the pulldowns in panel A. dCq of

IgG pulldowns of tagged protein (induced extract) normalized to a mock pulldown (uninduced

extract) [dCq = $2^{(test\ Cq\ -\ mock\ Cq)}$] for each construct. The mutant proteins were then normalized to

the tagged WT protein. All values took into account the recovery of tagged protein in each

pulldown. Three independent biological replicates with Cq average values and one standard

deviation (+/-1 SD, *n*=3) were plotted. The replicates suggest a similar quality and loading of

samples in the pulldowns. Non-specifically associated cytosolic transcripts differed substantially

in pulldowns of different mutants, so these transcripts were not used as reference. Mitochondrial

unedited (UE) and edited (ED) and cytosolic RNA transcripts are indicated. 1390 and 1400 are

abbreviations for the cytosolic HGPRT isoforms Tb927.10.1390 and Tb927.10.1400,

respectively.

Analysis of recombinant REH2 *in vitro*

The conserved predicted catalytic domains in REH2, the impact of mutating catalytic amino

acids *in vivo*, and the observed unwinding activity of pulldowns of REH2 and ^{H2}F1 from extracts

indicate that REH2 in the context of its native subcomplex in trypanosomes functions as an RNA

helicase (3, 4, 22). However, the unwinding activity of the isolated REH2 has not been formally

established. We have now generated a recombinant full-length rREH2 polypeptide (240 kDa) that

supports robust ATP-dependent unwinding of a synthetic dsRNA substrate *in vitro.* This dsRNA

substrate is a mimic of the native A6 editing system formed by pre-annealing of a 3' fragment of

the pre-mRNA A6 with a cognate gRNA. The synthetic A6 mRNA/gRNA pair supports a full

round of U-deletion editing in an *in vitro* assay with cell extracts and purified RECC enzyme (26,

27). The unwinding activity of the recombinant enzyme increased in a titration with increasing

concentrations of rREH2, and also in a time course up to 30 min when most of the annealed

radiolabeled RNA was unwound (Figs. 6A-B). An upshift of radiolabeled RNA in a ribonucleoprotein

complex at the highest tested concentrations of rREH2 in the titration may be due to the known RNA

binding activity of the native REH2 established by UV crosslinking (5, 22). The shifted species is

sensitive to proteinase K (data now shown). The radiolabeled strand alone (gRNA) and in heat-

treated controls of the dsRNA substrate exhibit the same gel mobility as the helicase-unwound gRNA

(Figs. 6B-C). Interestingly, a N-terminal truncated rREH2 polypeptide that lacked both dsRBDs was

inactive *in vitro* (data not shown). In summary, the isolated 240 kDa recombinant REH2 is a *bona fide*

RNA helicase enzyme *in vitro*, and further supports the idea that REH2 may remodel RNA or its

RNPs in editing.

Figure 6. Recombinant rREH2 is a catalytically active RNA helicase. (A) Unwinding assays

in 10 μL reaction mixtures with increasing concentrations of rREH2[30-2167] +/- ATP; **(B)**

Unwinding assays at increasing incubation times with rREH2[30-2167] +/- ATP using the

standard reaction mixture described in the material and methods section. The standard reaction

used a molar excess of enzyme over substrate. The dsRNA (ds) was assembled by pre-

annealing of synthetic transcript that mimics a 3' fragment of the *T. brucei* A6 pre-edited mRNA

and a radiolabeled cognate guide RNA (26). Radiolabeled unwound ssRNA (ss) is indicated.

Input dsRNA was heat denatured (∆) as control. **(C)** Additional controls showing that the starting

radiolabeled ssRNA used to generate the dsRNA substrate in the assay and the unwound

radiolabeled ssRNA in the heat-denatured (Δ) dsRNA (in panel B) have the same gel mobility.

Discussion

417   In the current study, we generated and characterized over 20 constructs of REH2 and [H2]F1

418   (Tables 1 and 2) to identify features in these editing proteins that impact the stability of REH2, the

419   assembly of the REH2C RNP, and its RNA-mediated coupling with other components of the RNA

420   editing holoenzyme (or RNA holo-editosome). We also formally established that the isolated REH2 is

421   a *bona fide* RNA helicase enzyme.

423   The core RECC enzyme and its accessory *trans* factors RESC and REH2C that assemble

around mRNA exhibit transient and stable contacts with each other via RNA. Also, variants of RECC

or the GRBC and REMC modules in RESC are emerging in studies by different labs (reviewed in 1).

These variants imply a dynamic nature of holo-editosomes that may be important in editing control

and needs further characterization. The helicase complex REH2C exhibits transient and stable

contacts with variants of RESC that minimally differ in the relative content of MRB3010 (in the GRBC

module). In this study, we focused on stable interactions by REH2C that can be readily examined.

Prior studies *in vivo* showed that inactivating mutations in the helicase ATP binding site or the

dsRBD2 domain dissociate REH2 from GAP1, gRNA and mRNA, reflecting a general helicase

detachment from the editing apparatus (3). [H2]F1 RNAi dissociates the helicase from the canonical

RESC markers GAP1 and RGG2. The RNAi knockdown of either [H2]F1 or REH2 inhibits editing

associated with a RESC variant that is particularly active and transiently interacts with the helicase

REH2 (3, 4). These and other observations indicate that transient and stable contacts in dynamic

holo-editosomes including those involving REH2C are important.

The section below discusses examined mutations in REH2 and [H2]F1 in the current study that

identified features affecting the proper assembly of the REH2C RNP, or its association with other

components of holo-editosomes *in vivo.*

REH2 mutants *in vivo*

The NTR truncations examined here (>1000 residues each), ∆N and ∆Nds, exhibited a

reduced binding with [H2]F1 and reduced RNA-mediated association with GAP1. The only difference

between these constructs is that ∆N carries the ~70 amino acid dsRBD2 domain but ∆Nds does not.

The dsRBD2 is necessary for the proper assembly of REH2 in holo-editosomes (3). Also, the

dsRBD2 in the related helicase MLE in flies seems to provide a major RNA-binding surface,

presumably due to its potential to bind A-form double helical RNA (28, 29). The tested truncations

suggests that other N-terminal features besides dsRBD2, e.g., dsRBD1 or unidentified motifs

contribute to normal interactions by REH2 *in vivo*. These truncations may also introduce changes in

topology that alter the normal REH2 interactions. Conformational changes are known to modulate

functional interactions by other eukayotic DExH/RHA helicases (18, 30). Thus, the tested NTR

truncations may indicate the presence of unidentified protein features or global changes in

conformation that affect the REH2 interactions. Further studies will be needed to resolve these

possibilities. However, relatively short CTR truncations and specific point mutations discussed below

also affected the REH2 interactions with other editing components. These short truncations and

particularly point mutations in the C-terminus are less likely to cause major conformation changes in

REH2.

We have speculated that the highly conserved OB fold in DExH/RHA subfamily helicases

may share functional residues in kinetoplastid RNA editing and other RNA processes (14). Our

examined C-terminal truncations included most of the CTR (260 residues in ∆OB) or elements

beyond the OB fold that may be specific of REH2 (123 residues in ∆AOB). The conserved and

specific C-terminal features of REH2 could work in synergy. We compared specific conserved

residues in REH2 and in helicase Prp43p in yeast spliceosomes. R1979 and R2023 in REH2

(R664 and R708 in the RNA helicase Prp43p) (16) should be located in two critical loops, β1-β2

and β4-β5, respectively. In yeast, R664 and R708 are poised to interact with the RNA substrate

entering the helicase cavity, and mutation of either residue inhibits the RNA binding and RNA-

mediated activation of Prp43p (16). In trypanosomes, the R2023A substitution had a moderate

effect on the stable association of REH2C with GAP1. However, R1979A did not evidently affect

this interaction. Neither R1979A or R2023A seem to significant effect on the REH2 interaction

with [H2]F1.

Similarly, H1998 and R1999 in REH2 were compared to the equivalent H1032 and

K1033 positions in the RNA helicase MLE in flies. The later residues in MLE bind specific

uridylates in U-rich roX boxes of the cognate roX RNA. H1032 and K1033 are also required for

the proper function and chromosomal localization of MLE in the *Drosophila* dosage

compensation complex (18). The H1998E substitution had a moderate effect on the REH2

association with GAP1 but had little or no effect on the interaction with [H2]F1. In contrast, the

adjacent substitution R1999E affected both the direct REH2 interaction with [H2]F1 and the RNA-

based REH2 association with GAP1. These results are consistent with a propose role of [H2]F1 as

an adaptor protein of REH2 (4). So, structure-function correlations may be tractable to specific

conserved helicase amino acids in protozoa and vertebrate species. This is in line with the idea

that the OB fold in DExH/RHA helicases, including REH2, is a conserved regulatory domain.

[H2]F1 mutants *in vivo*
486
[H2]F1 has eight predicted C2H2 Znf domains that are spread out and cover over ~45% of the

polypeptide sequence. Apart from the Znf domains, [H2]F1 has no evident sequence or structural

conservation outside of the kinetoplastids. ^{H2}F1 is as an adaptor protein that brings the REH2 to the

editing machinery (4) and it is also a stabilizing factor of REH2. That is, the induced downregulation

of ^{H2}F1 caused fragmentation of REH2 (4), and we showed here that the converse upregulation of

^{H2}F1 increased the steady-state level of REH2 *in vivo*. ^{H2}F1 may require most of its zinc fingers to

stabilize REH2 because substitution of core residues or variable basic residues in individual fingers

compromised this function. Thus, ^{H2}F1 binding to REH2 may induce substantial changes in the

helicase conformation and stability. The stabilizing role of ^{H2}F1 may involve the coordinated topology

of its zinc fingers. Notably, all examined ^{H2}F1 substitution-mutants were able to associate with REH2,

GAP1, and REL1. So, the basic fold or positive charge of the fingers is not essential for the

association of ^{H2}F1 with other editing components. However, N- or C-terminal truncation of nearly half

the size of ^{H2}F1 hampered its normal interactions with the examined editing components. Additional

studies are needed to pinpoint specific determinants for ^{H2}F1 direct binding to REH2 or its RNA-

mediated contacts with other editing components.

Besides the REH2•^{H2}F1 system, other DExH/RHA helicase•Znf protein pairs have been

characterized (14): the MOP•MEP-1 system in worm embryogenesis, and the DXH30•ZAP system in

antivirus protection in humans (31, 32). The antiviral protein ZAP has four Znf motifs that participate

in viral RNA recognition (32). All four fingers are on a positively charged surface in a crystal structure

of ZAP (24). and substitution of the basic residues in the fingers inhibited the antiviral functions of

ZAP (32). Insights from the ZAP structure imply that stabilization of REH2 by ^{H2}F1 involves a precise

topology of its zinc fingers and a role of its fingers in RNA binding. Notably, the Znf proteins ZAP and

MEP-1 multimerize. If ^{H2}F1 dimerizes as it was shown for recombinant ZAP (24) a single REH2C

RNP would include 16 zinc fingers. The co-purification of tagged and endogenous ^{H2}F1 from RNase-

treated extracts suggest the presence of direct protein contacts between the ^{H2}F1 copies (S3 Fig.).

However, analyses of recombinant ^{H2}F1 are necessary to establish the potential of ^{H2}F1 to

multimerize or bind RNA. The RNA content in pulldowns of a ^{H2}F1 variant with four R/K>Ala

substitutions in Znf4 suggested minimal changes in association with target RNAs in mitochondria but

a significance loss in non-specific contacts with unrelated transcripts. So, subtle changes in RNA

affinity are more likely to affect weak non-specific RNA contacts. The dramatic loss of all transcripts

examined in the pulldowns of the tested ^{H2}F1 fragments suggest that efficient RNA association by the

REH2C subcomplex *in vivo* requires the full array of zinc fingers in ^{H2}F1. Unfortunately, we were

unable to maintain stable expression an ^{H2}F1 construct with mutations in all fingers or subsets of

fingers (data not shown). We note that the transcripts examined may associate with either ^{H2}F1,

REH2 or both proteins in the native REH2C RNP.

REH2 may function as an RNA helicase in kinetoplastid RNA editing
524
Prior studies support the idea that REH2 is a functional RNA helicase in the context of the

editing apparatus. That is, REH2 pulldowns from GAP1-depleted extracts exhibit RNA unwinding

activity. Also, mutation of conserved catalytic residues in the predicted RecA1 domain inhibits the

unwinding activity of REH2 pulldowns from mitochondrial extracts (22). We have now established

that the isolated full-length recombinant REH2 (240 kda) is a *bona fide* RNA helicase enzyme.

Additional studies are needed to determine if ^{H2}F1 affects the unwinding activity of REH2. Thus,

REH2, the only DExH/RHA-type RNA helicase in kinetoplastid holo-editosomes most likely provides

an ATP-dependent unwinding activity in RNA editing. Besides REH2, the editing apparatus also

requires the DEAD-box RNA helicase REH1 (12, 13). This helicase is not considered to be a subunit

of the RNPs studied here and no potential binding partner of REH1 has been identified. A transient

RNA-mediate interaction of REH1 with other editing proteins has been detected (22, 33).

Important functional differences are known between the two ATP-dependent RNA helicases.

REH2 is needed for editing within single blocks, including the first "initiating" block (3). In

contrast, REH1 is needed for editing of two or more blocks (i.e., by overlapping gRNAs) but has

no effect on the first block (13). Both RNA helicases may be involved in critical remodeling of

RNA or RNP structure. REH1 may promote the relay of gRNAs as editing progresses from one

block to the next. Our ongoing RNA-seq analyses suggest that REH2 may impact the editing

efficiency at the individual sites and may provide additional functional insights concerning

REH2C (Kumar et al., unpublished).

Possible roles of editing complex variants

545
546
The editing apparatus much like the splicing and transcription molecular machines may

exhibit dynamic phases of assembly and function. The native RECC enzyme has multiple

isoforms, and several RNase III-like protein subunits in RECC predict a binary combinatorial

potential that may add fine-tuning in editing function (34-36). Also, natural GRBC and REMC

isoforms and genetically-induced changes in the organization of their protein and RNA

components are emerging. A recurring theme in the known examples is the variable interaction

between the GAP1/2 heterotetramer that is required for all editing involving GRBC-bound

gRNAs, and MRB3010 (GRBC6) that participates in early editing (1, 6, 37). A controlled

interaction of the GAP1/2 tetramer with MRB3010 may be used to modulate editing. For

example, GRBC variants in direct pulldowns of MRB3010 or REH2 pulldowns differ in their

relative content of MRB3010 and editing level in examined associated mRNAs (3, 5). However,

RNAi of REH2 or [H2]F1 reduced the editing level in mRNAs in the MRB3010 pulldowns (3, 4).

REH2 and [H2]F1 features, including those examined in the current studies, are expected to affect

specific steps in the editing mechanism and current studies in our lab are addressing this.

Variations in MRB3010 or other critical RESC proteins may distinguish pre-editosomes from

fully-active editosomes. Dynamics in complex assembly could also pause editing facilitating

helicase remodeling of RNA or RNPs. Additional studies are needed to examine these

possibilities in editing control.

Materials and methods

Cell Culture

T. brucei Lister strain 427 29-13 procyclic was grown axenically in log phase in SDM79

medium (38) and harvested at a cell density of $1–3 \times 10^7$ cells/mL. All transgenic cell lines were

induced with tetracycline at 1μg/mL. The TbRGG2 (alias RGG2) RNAi cell line was provided by

the Read lab (39). The GAP1 and [H2]F1 RNAi cell lines that were used as controls in some

analyses were generated in our lab (3, 4). We have seen that the TAP tag in REH2 or [H2]F1,

including WT constructs, seems to slow the growth of trypanosomes after several days of

culture (22)(data not shown). This and presumably incomplete cloning after transfection may

account for the uneven expression levels in extracts from independent subcultures. To partially

control for this issue, most plasmid constructs in the current study were re-made to use

puromycin selection. Transfections were also repeated with tetracycline-screened HyClone

Fetal Bovine Serum (FBS) (SH30070.03T; GE Healthcare. This enabled a faster cell recovery

after transfection than with constructs using phleomycin selection that were originally

characterized (data not shown). Regardless of this issue, independent analyses of protein

(complex) interactions were consistent with either puromycin or phleomycin constructs.

DNA constructs

All overexpression and RNAi studies used inducible plasmid constructs. The constructs

for [H2]F1 RNAi and the REH2 dsRBD2 mutant K1078A/A1086D were previously reported (3, 4).

N-terminal deletion constructs for REH2 (ΔN and ΔNds) and [H2]F1 (ΔN) were prepared using the

In-Fusion® HD Cloning Kit (639648; Clontech). Two PCR amplicons were prepared in each

case using CloneAmp™ HiFi PCR Premix (638500; Clontech) and were then gel-eluted using

NucleoSpin® Gel and PCR Clean-up (740609.10; Clontech). One amplicon encodes the first 34

amino acids of REH2 (primers: F-1503, R-1522), including the predicted REH2 mitochondrial

leader sequence (REH2-MLS) (22), that was used for all N-deletion constructs. The other

amplicon included the remaining relevant gene fragment sequence: 3070-6501 bases (REH2

ΔN), 3280-6501 bases (REH2 ΔNds), or 766-1572 bases (^{H2}F1 ΔN). Three-piece in-fusion

reactions were performed to join the two amplicons and pLEW79-ada-TAP plasmid linearized at

XhoI and *BamHI* (40). The PCR primers (S1 Table) included 5' and 3' terminal homology (15

bases) to enable the recombination-based fusion between the amplicons and the plasmid DNA.

The REH2 (ΔAOB and ΔOB) and ^{H2}F1 ΔC deletion constructs are also recombination-based

fusions of the amplified relevant gene fragment with linearized plasmid DNA. The REH2

constructs R2023A, H1998E, R1999E and H1998E/R1999E, and the ^{H2}F1 constructs Z5 C>A

and Z5 R/K>A, were created by site-directed mutagenesis in inverse-PCR reactions of the

pLEW79-ada-TAP WT plasmid as the template. pLEW79-ada-TAP was modified from the

original pLEW79-TAP (40, 41). The resulting linearized mutant constructs were circularized

using the In-Fusion HD Cloning Kit. Other ^{H2}F1 constructs used synthetic gBlock gene

fragments (IDT) with mutations in single or multiple zinc-fingers. The gBlock fragment

sequences were fused as follows. A full-length ^{H2}F1 sequence with all fingers mutated (Z1-8

C>A) was cloned into *XhoI/BamHI* sites of the plasmid vector. N-terminal fragments of ^{H2}F1 (Z1,

Z2 and Z3 R/K>A single finger mutants) replaced the corresponding sequence in the ^{H2}F1 WT

construct after digestion with *XhoI* and the internal *ApaI* site in ^{H2}F1. C-terminal fragments of

^{H2}F1 (Z4 and the multi-finger mutant Z6-Z8) replaced corresponding sequence in the ^{H2}F1 WT

construct after digestion with *ApaI* and *BamH1*. All constructs were confirmed by DNA

sequencing, linearized with *NotI*, and transfected in trypanosomes (38).

Western Blots and Radioactivity Assays
617
Western blots of REH2, ^{H2}F1 and ^{H2}F2 (subunits of REH2C), GAP1 (alias GRBC2;

subunit of GRBC) and A2 (alias MP42; subunit of RECC) were performed as reported (4, 22).

Western blots of the GAP1/GAP2 (alias GRBC2/GRBC1) paralogs simultaneously, and of

MRB8170 and MRB6070 (subunits of REMC and MRB6070/1590 assemblies, respectively),

were performed as previously described (42, 43). Peroxidase Anti-peroxidase antibody (P1291;

Sigma) diluted to 1:200 (v/v) was used to detect the TAP-tag. Protein quantitative analyses

using Amersham ECL^{TM} western blot detection reagents were performed in an Amersham

Imager 600. RNA ligases in the RECC enzyme were radiolabeled by self-adenylation directly on

the beads in IgG pulldowns (44). Unwinding assays of REH2 used reported conditions with a

few modifications (4) and full-length recombinant His-REH2. The dsRNA substrate in these

assays used an A6 mRNA/gRNA pair (22). The A6 pair mimics the endogenous ATPase subunit

6 substrate. The gRNA was transcribed with RNA T7 polymerase from an amplicon (S1 Table)

after *DraI* restriction digestion and gel purification of the product. The 61-nt gRNA was 5'-

labelled with γ^{32}P-ATP using T4 Polynucleotide kinase. The 62-nt mRNA was synthetic (IDT). A

mix with 3 pmoles of labelled gRNA and 15 pmoles of mRNA was heated at 95° C for 3 min,

and then immediately incubated for 30 min at room temperature in the annealing buffer (10 mM

MOPS pH 6.5, 1 mM EDTA, and 50 mM KCl). The annealed RNA hybrids were isolated from

native 12% PAGE run at 4 °C at 50 V for 120 min in 0.5x Tris-borate-EDTA buffer. Standard 10

μL reaction mixtures in unwinding buffer (40 mM Tris-Cl pH 8.0, 0.5 mM MgCl^2, 0.01% Nonidet

P-40, and 2 mM DTT) were incubated for 30 min at 19 °C with 20 cps of hybrid (~20 fmoles)

and 3 μg of recombinant REH2 (~12 pmoles). Variations to the standard assay including in the

concentration of REH2 or the reaction time are indicated in the text. All assays contained 100

fmoles of unlabeled gRNA as competitor to prevent re-annealing of unwound γ^{32}P-gRNA. The

assays were stopped with an equal volume of 2x helicase reaction stop buffer (50 mM EDTA,

1% SDS, 0.1% bromophenol blue, 0.1% xylene cyanol, and 20% glycerol) and kept on ice for 5

min. The assays in Fig. 6B were also treated with 0.4U of proteinase K (P8107S; NEB), 25 mM

EDTA and 0.5% SDS at 22° C for 30 min. The entire reaction was loaded onto a native 12%

PAGE and resolved at 25 V for 120 min at 4°C.

Cell extracts and purification of protein and RNA from IgG pulldowns

648
Whole cell extracts as well as mitochondria-enriched extracts were used in this study.

Whole-cell extracts were generated by incubation of cell pellets in MRB-Triton buffer (25 mM

Tris-Cl pH 8.0, 10 mM MgOAC, 1 mM EDTA, 10 mM KCl, 5% glycerol, 0.5% Triton X-100) at 4

°C for 45 minutes followed by centrifugation at 15,000g for 30 minutes. Mitochondria-enriched

extracts were prepared using a simplified protocol similar to others reported earlier (2, 45).

Namely, cell pellets were resuspended into hypotonic DTE buffer (1 mM Tris-HCl pH 8.0, 1 mM

EDTA pH 8.0) supplemented with protease inhibitors and disrupted in a Dounce homogenizer.

The suspension was passed through a 26-gauge needle as originally described (46). The

enriched mitochondrial vesicles were spun down and the cytosolic content in the supernatant

discarded. This step improved the recovery of intact REH2 compared to whole-cell extracts.

REH2 is more prone to fragmentation than other proteins examined in this study. The following

steps were also as in (46). The pellet was resuspended into STM buffer (0.25M Sucrose, 20 mM

Tris-HCl pH 8.0, 2 mM $MgCl_2$) supplemented with DNase I, RNase-free (EN0523; Fisher

Scientific). The reaction was stopped with one volume of STE buffer (0.25 M Sucrose, 20 mM

Tris-HCl pH 8.0, 2 mM EDTA pH 8.0). The enriched mitochondria were spun down and washed

in STE buffer. The pellet was resuspended into MRB-Triton buffer and incubated in ice for 15

min. The suspension was spun down at 15,000g at 4 °C for 30 min. The supernatant was

collected and stored at -80 °C. In IgG pulldowns of the extracts, ectopically expressed TAP-

REH2 and TAP-[H2]F1 were recovered as reported (4, 22) with some modifications.

Approximately 2 mg of protein was mixed with 1x SUPERase·In[TM] RNase inhibitor (Invitrogen[TM])

and incubated with Dynabeads IgG (11203D; Invitrogen[TM]). The beads were washed five times

with 1ml of wash buffer (150 mM NaCl, 1 mM EDTA, 10 mM MgAOc, 0.1% NP-40, and 25 mM

Tris pH 8.0). Protein was eluted from the beads with 15 μL of 1x SDS loading buffer at 95 °C for

2 min. RNA was extracted by treating the beads with 4U proteinase K (NEB) for 2 hrs at 55 °C,

followed by phenol extraction and ethanol precipitation. For some pulldowns, RNase A/T1 mix

(EN0551; Fisher Scientific) was applied at 20 U/mg of protein both in the input extract and again

while bound to the beads.

Quantitative RT-PCR

677
678
Isolated RNA from the pulldowns or directly from input extracts was treated with DNase

I, RNase-free (EN0523; Thermo Scientific^{TM}) prior to cDNA synthesis with the iScript™ Reverse

Transcription Supermix (Bio-Rad) as described elsewhere (5). The amplifications mixtures (10

µl) used SsoAdvanced™ Universal SYBR$^®$ Green Supermix (Bio-Rad) and reported

oligonucleotides for unedited mRNAs, fully edited mRNA, and reference transcripts (47). Diluted

samples of the examined cDNA produced a single amplicon during linear amplification under

our conditions. The end-point amplicons described here have been gel-isolated, cloned and

confirmed by sequencing (5) (data not shown). Fold-enrichment of mitochondrial and cytosolic

transcripts in IgG pulldowns of tagged proteins (induced extracts), relative to a mock pulldown

(uninduced extract), was calculated as follows: Fold = $2^{[-ddCq]}$, where ddCq = Cq test IP – Cq

mock IP, as in (5). The relative value for each mutant construct was adjusted to the amount of

recovered tagged protein, and subsequently normalized to the tagged WT construct (= 1). The

plot in Fig. 5B was constructed using data from 3 independent biological replicate experiments,

including a total of six amplicons per data point (i.e., two technical replicate amplicons per

assay). Two of the independent experiments (including cultures +/- Tet for each mutant) were

performed concurrently. The third independent experiment (one set of cultures per mutant +/-

Tet) was conducted on a subsequent date. The replicate experiments showed a consistent

sample loading in the assays used in our plots. The examined mutant constructs differed in the

amount of non-specific association with Tubulin and other cytosolic transcripts that are normally

used as internal reference in antibody pulldowns using a common protein bait. The raw Cq

values of examined editing transcripts in the pulldowns and the plotted values after

normalization indicated a substantial loss of these transcripts in the truncated mutants.

qRT-PCR assays of input extract in S4 Fig. were normalized using the following equation: Ratio

(reference/target) = $2^{[Cq(ref) - Cq(target)]}$. This scores the relative difference between the reference

and target Cq values. The current plots of input extract used 18S rRNA as reference (S4 Fig.).

Similar results were obtained with any of these cytosolic transcripts used as a reference: tubulin,

1400 or 1390 (Tb927.10.1400 or Tb927.10.1390, respectively) (data not shown).

Homology modeling and bioinformatic analysis

709   The multiple-sequence alignment of REH2 and MLE was done with Clustal Omega (48),

710   and that of the ^{H2}F1 zinc finger domains was done with BOXSHADE version 3.21

711   (http:/www.ch.embnet.org/software/BOX_form.html). The homology models of the helicase

712   region of *T. brucei* REH2 were built using the program Phyre2 (20).

Recombinant protein

6xHis-REH2 (amino acid residues 30-2167) was amplified by PCR (S1 Table). The gel-

purified product was cloned into the NcoI and BamHI sites in the expression vector pET15b by

using the NEBuilder® HiFi DNA Assembly Master Mix (E262, NEB) and the appropriate primers.

The ligation reaction mixtures were transformed into chemically competent Omnimax cells from

ThermoFisher Scientific. Colonies with the correct insert were identified by single-colony PCR

and then amplified for plasmid purification. The sequence of the flanking vector region and the

insert was checked by DNA sequencing. The plasmid DNA was transformed into Rosetta2 DE3

cells (Novagen Inc.) and overexpressed in 4 L of Terrific Broth media by supplementing 2% (v/v)

ethanol and 2 mM MgCl$_2$ at 37 °C. Protein induction with 0.5 mM IPTG was done at a OD$_{600}$ of

0.8, and the expression was continued for 22 hours at 16 °C. The cell pellet was stored at -80

°C overnight and was resuspended in lysis buffer (5 mL/g of cell pellet) of 50 mM Tris-HCl pH

8.0, 250 mM NaCl, 1 mM EDTA, and 1 mM DTT along with 10 µg of RNase-free DNase and 15

mg of hen egg white lysozyme. The cells were lysed in an Emulsiflex hydraulic press. The cell

debris was pelleted at 39,200 x g for 30 minutes at 4 °C. The supernatant with 6xHis-REH2 was

batch attached to equilibrated Qiagen Ni-NTA Agarose at 4 °C for 16 hrs. The complex was

eluted with 50 mM Tris-HCl pH 8.0, 300 mM NaCl, 1 mM DDT, and 250 mM imidazole. The first

elution of the purification was dialyzed into unwinding assay buffer. This material was tested for

the presence of the recombinant protein in western blots and antibody pulldowns, and for

activity in dsRNA unwinding activity assays *in vitro*.

Acknowledgements
737

We thank the Read lab for their gift of antibodies against GAP1, TbRGG2 (alias RGG2)

and [H2]F2. The Afasizhev lab kindly provided antibodies that react against the GAP1/GAP2

paralogs (alias GRBC2/GRBC1) simultaneously. We also thank the Stuart lab for providing the

antibodies against A2 (alias MP42). Amna Mushtaq provided detailed comments on the

manuscript. Jana Gomez, Elizabeth Alvarez and Melissa Yang helped with the preparation of

figures.

References

1.   Cruz-Reyes J, Mooers BHM, Doharey PK, Meehan J, Gulati S. Dynamic RNA holoeditosomes with subcomplex variants: Insights into the control of trypanosome editing. WIREs RNA. 2018;e1502.

2.   Koslowsky DJ, Sun Y, Hindenach J, Theisen T, Lucas J. The insect-phase gRNA transcriptome in Trypanosoma brucei. Nucleic acids research. 2013.

3.   Madina BR, Kumar V, Mooers BH, Cruz-Reyes J. Native Variants of the MRB1 Complex Exhibit Specialized Functions in Kinetoplastid RNA Editing. PloS one. 2015;10(4):e0123441.

4.   Kumar V, Madina BR, Gulati S, Vashisht AA, Kanyumbu C, Pieters B, et al. REH2C Helicase and GRBC Subcomplexes May Base Pair through mRNA and Small Guide RNA in Kinetoplastid Editosomes. J Biol Chem. 2016;291(11):5753-64.

5.   Madina BR, Kumar V, Metz R, Mooers BH, Bundschuh R, Cruz-Reyes J. Native mitochondrial RNA-binding complexes in kinetoplastid RNA editing differ in guide RNA composition. RNA. 2014;20(7):1142-52.

6.   Aphasizheva I, Zhang L, Wang X, Kaake RM, Huang L, Monti S, et al. RNA binding and core complexes constitute the U-insertion/deletion editosome. Mol Cell Biol. 2014;34(23):4329-42.

7.   Huang Z, Faktorova D, Krizova A, Kafkova L, Read LK, Lukes J, et al. Integrity of the core mitochondrial RNA-binding complex 1 is vital for trypanosome RNA editing. RNA. 2015.

8.   McAdams NM, Simpson RM, Chen R, Sun Y, Read LK. MRB7260 is essential for productive protein-RNA interactions within the RNA editing substrate binding complex during trypanosome RNA editing. RNA. 2018;24(4):540-56.

9.   Simpson RM, Bruno AE, Bard JE, Buck MJ, Read LK. High-throughput sequencing of partially edited trypanosome mRNAs reveals barriers to editing progression and evidence for alternative editing. RNA. 2016;22(5):677-95.

10.   Simpson RM, Bruno AE, Chen R, Lott K, Tylec BL, Bard JE, et al. Trypanosome RNA

Editing Mediator Complex proteins have distinct functions in gRNA utilization. Nucleic acids

research. 2017;45(13):7965-83.

11.   Shaw PL, McAdams NM, Hast MA, Ammerman ML, Read LK, Schumacher MA.

Structures of the T. brucei kRNA editing factor MRB1590 reveal unique RNA-binding pore motif

contained within an ABC-ATPase fold. Nucleic acids research. 2015;43(14):7096-109.

12.   Missel A, Souza AE, Norskau G, Goringer HU. Disruption of a gene encoding a novel

mitochondrial DEAD-box protein in Trypanosoma brucei affects edited mRNAs. Mol Cell Biol.

1997;17(9):4895-903.

13.   Li F, Herrera J, Zhou S, Maslov DA, Simpson L. Trypanosome REH1 is an RNA helicase

involved with the 3'-5' polarity of multiple gRNA-guided uridine insertion/deletion RNA editing.

Proc Natl Acad Sci U S A. 2011;108(9):3542-7.

14.   Cruz-Reyes J, Mooers BH, Abu-Adas Z, Kumar V, Gulati S. DEAH-RHA helicase•Znf

cofactor systems in kinetoplastid RNA editing and evolutionarily distant RNA processes. RNA

Dis. 2016;3(2):e1336.

15.   Jarmoskaite I, Russell R. RNA helicase proteins as chaperones and remodelers. Annu

Rev Biochem. 2014;83:697-725.

16.   Walbott H, Mouffok S, Capeyrou R, Lebaron S, Humbert O, van Tilbeurgh H, et al.

Prp43p contains a processive helicase structural architecture with a specific regulatory domain.

EMBO J. 2010;29(13):2194-204.

17.   Robert-Paganin J, Rety S, Leulliot N. Regulation of DEAH/RHA helicases by G-patch

proteins. Biomed Res Int. 2015;2015:931857.

18.   Prabu JR, Muller M, Thomae AW, Schussler S, Bonneau F, Becker PB, et al. Structure

of the RNA Helicase MLE Reveals the Molecular Mechanisms for Uridine Specificity and RNA-

ATP Coupling. Mol Cell. 2015;60(3):487-99.

19.   Dhote V, Sweeney TR, Kim N, Hellen CU, Pestova TV. Roles of individual domains in

the function of DHX29, an essential factor required for translation of structured mammalian

mRNAs. Proc Natl Acad Sci U S A. 2012;109(46):E3150-9.

20.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for

protein modeling, prediction and analysis. Nat Protoc. 2015;10(6):845-58.

21.   Jankowsky E. RNA helicases at work: binding and rearranging. Trends in biochemical

sciences. 2011;36(1):19-29.

22.   Hernandez A, Madina BR, Ro K, Wohlschlegel JA, Willard B, Kinter MT, et al. REH2

RNA helicase in kinetoplastid mitochondria: ribonucleoprotein complexes and essential motifs

for unwinding and guide RNA (gRNA) binding. J Biol Chem. 2010;285(2):1220-8.

23.   Segal DJ, Crotty JW, Bhakta MS, Barbas CF, 3rd, Horton NC. Structure of Aart, a

designed six-finger zinc finger peptide, bound to DNA. J Mol Biol. 2006;363(2):405-21.

24.   Chen S, Xu Y, Zhang K, Wang X, Sun J, Gao G, et al. Structure of N-terminal domain of

ZAP indicates how a zinc-finger protein recognizes complex RNA. Nat Struct Mol Biol.

2012;19(4):430-5.

25.   Klug A. The discovery of zinc fingers and their development for practical applications in

gene regulation and genome manipulation. Q Rev Biophys. 2010;43(1):1-21.

26.   Seiwert SD, Heidmann S, Stuart K. Direct visualization of uridylate deletion in vitro

suggests a mechanism for kinetoplastid RNA editing. Cell. 1996;84(6):831-41.

27.   Rusche LN, Cruz-Reyes J, Piller KJ, Sollner-Webb B. Purification of a functional

enzymatic editing complex from Trypanosoma brucei mitochondria. EMBO J. 1997;16(13):4069-

81.

28.   Izzo A, Regnard C, Morales V, Kremmer E, Becker PB. Structure-function analysis of the

RNA helicase maleless. Nucleic acids research. 2008;36(3):950-62.

29.   Masliah G, Barraud P, Allain FH. RNA recognition by double-stranded RNA binding

domains: a matter of shape and sequence. Cell Mol Life Sci. 2013;70(11):1875-95.

30.   Christian H, Hofele RV, Urlaub H, Ficner R. Insights into the activation of the helicase

Prp43 by biochemical studies and structural mass spectrometry. Nucleic acids research.

2014;42(2):1162-79.

31.   Ye P, Liu S, Zhu Y, Chen G, Gao G. DEXH-Box protein DHX30 is required for optimal

function of the zinc-finger antiviral protein. Protein & cell. 2010;1(10):956-64.

32.   Guo X, Carroll JW, Macdonald MR, Goff SP, Gao G. The zinc finger antiviral protein

directly binds to specific viral mRNAs through the CCCH zinc finger motifs. J Virol.

2004;78(23):12781-7.

33.   Kruse E, Voigt C, Leeder WM, Goringer HU. RNA helicases involved in U-

insertion/deletion-type RNA editing. Biochimica et biophysica acta. 2013;1829(8):835-41.

34.   Carnes J, Soares CZ, Wickham C, Stuart K. Endonuclease associations with three

distinct editosomes in Trypanosoma brucei. J Biol Chem. 2011;286(22):19320-30.

35.   McDermott SM, Luo J, Carnes J, Ranish JA, Stuart K. The Architecture of Trypanosoma

brucei editosomes. Proc Natl Acad Sci U S A. 2016;113(42):E6476-E85.

36.   Carnes J, McDermott S, Anupama A, Oliver BG, Sather DN, Stuart K. In vivo cleavage

specificity of Trypanosoma brucei editosome endonucleases. Nucleic acids research.

2017;45(8):4667-86.

37.   Ammerman ML, Hashimi H, Novotna L, Cicova Z, McEvoy SM, Lukes J, et al. MRB3010

is a core component of the MRB1 complex that facilitates an early step of the kinetoplastid RNA

editing process. RNA. 2011;17(5):865-77.

38.   Wirtz E, Leal S, Ochatt C, Cross GA. A tightly regulated inducible expression system for

conditional gene knock-outs and dominant-negative genetics in Trypanosoma brucei. Molecular

and biochemical parasitology. 1999;99(1):89-101.

39.   Fisk JC, Ammerman ML, Presnyak V, Read LK. TbRGG2, an essential RNA editing

accessory factor in two Trypanosoma brucei life cycle stages. J Biol Chem.

2008;283(34):23016-25.

40.   Hernandez A, Panigrahi A, Cifuentes-Rojas C, Sacharidou A, Stuart K, Cruz-Reyes J.

Determinants for association and guide RNA-directed endonuclease cleavage by purified RNA

editing complexes from Trypanosoma brucei. J Mol Biol. 2008;381(1):35-48.

41.   Panigrahi AK, Schnaufer A, Ernst NL, Wang B, Carmean N, Salavati R, et al.

Identification of novel components of Trypanosoma brucei editosomes. RNA. 2003;9(4):484-92.

42.   Ammerman ML, Downey KM, Hashimi H, Fisk JC, Tomasello DL, Faktorova D, et al.

Architecture of the trypanosome RNA editing accessory complex, MRB1. Nucleic acids

research. 2012;40(12):5637-50.

43.   Weng J, Aphasizheva I, Etheridge RD, Huang L, Wang X, Falick AM, et al. Guide RNA-

binding complex from mitochondria of trypanosomatids. Mol Cell. 2008;32(2):198-209.

44.   Sabatini R, Hajduk SL. RNA ligase and its involvement in guide RNA/mRNA chimera

formation. Evidence for a cleavage-ligation mechanism of Trypanosoma brucei mRNA editing. J

Biol Chem. 1995;270(13):7233-40.

45.   Hashimi H, Zikova A, Panigrahi AK, Stuart KD, Lukes J. TbRGG1, an essential protein

involved in kinetoplastid RNA metabolism that is associated with a novel multiprotein complex.

RNA. 2008;14(5):970-80.

46.   Harris ME, Hajduk SL. Kinetoplastid RNA editing: in vitro formation of cytochrome b

gRNA-mRNA chimeras from synthetic substrate RNAs. Cell. 1992;68(6):1091-9.

47.   Carnes J, Stuart KD. Uridine insertion/deletion editing activities. Methods in enzymology.

2007;424:25-54.

48.   Sievers F, Higgins DG. Clustal Omega, accurate alignment of very large numbers of

sequences. Methods Mol Biol. 2014;1079:105-16.

Supporting information

S1 Fig. Multi-sequence alignment of a C-terminal segment in *T. brucei* REH2 and

***Drosophila* MLE DExH/RHA RNA helicases**. The alignment was generated with Clustal

Omega (48). Boxes were inserted manually to improve the match between the MLE and REH2

residues. Note that H1032, K1033 and T1034 (in red), which make U-specific contacts in MLE,

are aligned with H1998, R1999 and T2000 (in red) in REH2.

S2 Fig. Sedimentation analysis of endogenous editing proteins. (A) 10-30% glycerol

gradients of freshly-made mitochondria-enriched extract from 29:13 procyclic trypanosomes.

Catalase and RECC complex were used as 11 S and 20 S markers, respectively (22).

Endogenous REH2, ^{H2}F1 and ^{H2}F2, GAP1 (GRBC2), GAP2 (GRBC1), and A2 (MP42) were

examined in western blots. All panels in this figure derived from the same extract fractions. The

data shown is representative of at least two panels for each protein in biological replicate

gradients.

S3 Fig. RNase-resistant co-purification of tagged-^{H2}F1 and endogenous ^{H2}F1. Western

blots of IgG pulldowns from extracts with or without an RNaseA/T1 mix. All panels in this figure

derive from the same blot. The upper blot with the tagged-^{H2}F1 bait was cut below the 75 kDa

marker. The middle and lower panels were divided between the 50 kDa and 37 kDa marker.

The 34.4 kDa RGG2, a typical subunit of the REMC module in the RESC complex. As expected,

the RNA-mediated association of RGG2 decreased with the RNase treatment.

S4 Fig. Quantitation of steady-state RNA transcripts in the input mitochondrial extracts

used in the IgG pulldowns. Independent biological replicates (two independent cultures used

in **Fig. 5**) with Cq average values and one standard deviation (+/-1SD, *n*=2) were plotted. dCq

of steady-state RNA transcripts in lysates relative to background 18s rRNA used as reference

[dCq = $2^{(target\ Cq\ -\ ref\ Cq)}$]. Shorter bars indicate a smaller differential versus 18s rRNA in the

sample. For example, ND7 is relatively abundant compared to other transcripts in the sample

(i.e., it has a lower Cq). The WT construct is induced or not (+/-). All mutants are induced. All

end-point amplicons were examined in gels to confirm that they were single products during

linear amplification.

S1 Table. DNA oligonucleotides, gBlocks (IDT) and synthetic RNA.

Figure 2

Figure 3

Figure 4

Figure 5

A

NAME	MUTATION	DOMAINS AFFECTED	MW
dsR	K1078A/A1086D	dsRBD2	242
ΔOB	Δ2045-2167		228
ΔAOB	Δ1908-2167	OB fold	213
ΔN	Δ34-1023	dsRBD1	132
ΔNds	Δ34-1093	dsRBD1/dsRBD2	124
1979	R1979A	OB fold	242
1998	H1998E	OB fold	242
1999	R1999E	OB fold	242
98/99	H1998/R1999	OB fold	242
2023	R2023	OB fold	242

Figure 1

Figure 1

Figure 6

Figure 6