1

## Modeling enamel matrix secretion in mammalian teeth

**Teemu J. Häkkinen**[1,2,*]**, S. Susanna Sova**[1,3]**, Ian J. Corfe**[1]**, Leo Tjäderhane**[4]**, Antti Hannukainen**[2] **& Jukka Jernvall**[1,*]

**1** Developmental Biology Program, Institute of Biotechnology, University of Helsinki, Helsinki, Finland, **2** Department of Mathematics and Systems Analysis, Aalto University, Aalto, Finland, **3** Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland, **4** Department of Oral and Maxillofacial Diseases, Helsinki University Hospital, University of Helsinki, Helsinki, Finland.

[*] hakkinen@fastmail.com (TJH); jernvall@fastmail.fm (JJ)

## Abstract

The most mineralized tissue of the mammalian body is tooth enamel. Especially in species with thick enamel, three-dimensional (3D) tomography data has shown that the distribution of enamel varies across the occlusal surface of the tooth crown. Differences in enamel thickness among species and within the tooth crown have been used to examine taxonomic affiliations, life history, and functional properties of teeth. Before becoming fully mineralized, enamel matrix is secreted on the top of a dentine template, and it remains to be explored how matrix thickness is spatially regulated. To provide a

predictive framework to examine enamel distribution, we introduce a computational

model of enamel matrix secretion that maps the dentine topography to the enamel

surface topography. Starting from empirical enamel-dentine junctions, enamel matrix

deposition is modeled as a diffusion-limited free boundary problem. Using laboratory

microCT and synchrotron tomographic data of pig molars that have markedly different

dentine and enamel surface topographies, we show how diffusion-limited matrix

deposition accounts for both the process of matrix secretion and the final enamel

distribution. Simulations reveal how concave and convex dentine features have distinct

effects on enamel surface, thereby explaining why the enamel surface is not a

straightforward extrapolation of the dentine template. Human and orangutan molar

simulations show that even subtle variation in dentine topography can be mapped to

the enamel surface features. Mechanistic models of extracellular matrix deposition can

be used to predict occlusal morphologies of teeth.

## Author summary

Teeth of most mammals are covered by a layer of highly mineralized enamel that cannot be

replaced or repaired. The enamel layer is not uniform over the underlying dentine, and spatial

regulation of enamel formation is critical for making a functional tooth. To explore which kind

of mechanisms could underlie the complex patterns of enamel distribution, we present a

computational model. Starting from tomography-imaged teeth from which enamel has been

digitally removed, enamel is restored on dentine surfaces by simulating diffusion-limited

secretion of enamel matrix. Our simulations show how the combination of subtle features of

dentine and diffusion-limited secretion of the enamel matrix can substantially increase the

complexity of the enamel surface. We propose that the strength of the diffusion-limited process

is a key factor in determining enamel distribution among mammalian species.

**Introduction**

Most mammalian species have their teeth covered by a layer of highly mineralized enamel. The

thickness of the enamel layer relative to the tooth size ranges from thin to very thick. These

differences among species, and also increasingly within the tooth crown, have been informative

in studies focused on functional properties of teeth, taxonomy, and life history [1–9]. Even

though mutations in genes required for enamel matrix secretion and maturation are known to

affect the enamel thickness in mammals [10], relatively little is known about the regulatory

changes that might underlie the variation in enamel thickness among populations or species [1,

11, 12]. Even less is known about the regulation of enamel thickness variation within the tooth

crown, which contrasts with the increasing availability of 3D tomography data on various

species. Analyses of such tomography data show that even though the enamel surface

topography reflects the enamel-dentine junction (EDJ) topography, the enamel surface is not a

simple extrapolation of the EDJ shape [13-15]. Because enamel distribution is not

developmentally remodeled after formation, and because the internal structure of mineralized

enamel retains developmental information, tomography data of fully formed teeth can be used to

examine mechanisms underlying variation in enamel thickness. To provide mechanistic insights

into the regulation of enamel thickness, here we combine tomography data on enamel

distribution with a computational approach and introduce a model to simulate enamel matrix

secretion.

**Results**

**The model principles and simulation of artificial shapes**

The enamel matrix is secreted by specialized epithelial cells, the ameloblasts. When the matrix

secretion begins, ameloblasts detach from the EDJ and advance as a front secreting the enamel

matrix that will later mineralize into the enamel (Fig 1A). The EDJ is defined by the

mesenchymal dentine matrix, whose secretion begins first (Fig 1A). For empirical tests, we used

EDJs of real teeth as the starting point to simulate matrix secretion. Matrix deposition is

modeled as a diffusion-limited free boundary problem, motivated by the classical Stefan

problem that models phase transition of undercooled liquid by assuming that the rate of phase

transition from liquid to solid is limited by a diffusion process (see Methods and S1 Appendix)

[16, 17]. Here we similarly assume that the growth of the matrix front is a diffusion-limited

process: The advancement of the ameloblast layer is assumed to be limited by the diffusion of

nutrients, by which we refer collectively to all the factors that ameloblasts require for the

secretion of the matrix (Fig 1B). Biologically, both the extracellular milieu and capillary

networks present in the enamel organ during matrix secretion [18, 19] can be hypothesized to

limit the supply of nutrients in a diffusion-limited fashion. The model parameters adjust the

nutrient diffusion rate, the amount of nutrients required for growth, and the interfacial tension or

stiffness of the advancing ameloblast layer (Methods). Model equations are solved using the

finite element method, and the matrix interface (the ameloblast layer) is tracked using the level

set method (S1 Appendix). The source code of the Matlab implementation of the model is freely

available (Methods). For computational efficiency, the model is implemented in 2D and 3D

reconstructions are obtained by simulating multiple sections that are combined into volumes.

**Fig 1. Modeling tooth enamel matrix secretion.**

(A) A schematic illustration of a cross section of enamel matrix being secreted on the top of dentine. In

real teeth (a pig molar on the right), the enamel surface is not a linear representation of the dentine

template (EDJ). (B) In a diffusion-limited model, differences in surface topography lead to ridges (black

circle) receiving more nutrients (dashed arrows) than valleys (white circle). (C) Starting from a synthetic

EDJ shape, diffusion-limited matrix deposition advances faster in convex than concave features. (D)

Excess production of nutrients overcomes diffusion-limited effects and produces a uniform distribution

of matrix. (E) Reducing interfacial tension of the simulation in (C) results in a crenulated matrix surface.

For details of the model, see Methods, and parameters used in simulation are in S1 Table.

The fundamental component of the model is the assumption that the growth of the matrix

requires a net influx of a diffusing nutrient substance. At the initial stage nutrients are present

exterior to the dentine, which in real teeth, acts as an internal nutrient barrier [18]. The nutrients

are further replenished over time by a constant background source exterior to the dentine

(Methods). By controlling the relative amount of background production, we examine two

hypothetical matrix secretion processes. The primary process tested is a diffusion-limited

secretion in which concave surfaces are progressively exaggerated as the features protruding

into the nutrient-rich domain receive more nutrients than the concavities (Fig 1C). An

alternative process assumes excess availability of nutrients through strong background

production, leading to a moving boundary of uniform thickness (Fig 1D). This latter process in

fact closely approximates a simple geometric extrapolation of matrix thickness from the EDJ,

which we use as a null hypothesis to demonstrate the non-linearity of the matrix deposition.

Simulations of matrix secretion using a synthetic EDJ shape show that whereas convex EDJ

surfaces result in relatively linear extrapolation of the enamel surface in both simulations (Fig

1C and 1D), concave surfaces of diffusion-limited simulations behave nonlinearly (Fig 1C).

Additionally, reducing interfacial tension in the simulations increases small undulations in the

moving front (Fig 1E), suggesting that lowered stiffness of the ameloblast layer may underlie

crenulated enamel found in taxa such as *Chiropotes* (saki monkeys) with relatively smooth EDJ

[14]. Next we applied the model to real teeth with convex and concave features (Fig 2A).

**Fig 2. Diffusion-limited simulations approximate complex patterns of enamel thickness in the pig**

**molar.**

(A) A heat map and a horizontal section of 3D tomography reconstruction of a pig molar shows the

variable enamel thickness. (B) Using a horizontal EDJ section of a pig molar as an empirical template

(A), diffusion-limited simulations of matrix secretion produce deep fissures present in concave surfaces.

(C) Geometric extrapolation shows how the fissures are filled-in. In contrast, convex slopes are relatively

similar between the simulations. Note how the enamel is redirected and remains separated between the

buccal and lingual cusps in the real tooth (A) and diffusion-limited simulation (B). All images show

occlusal views of the left lower first molar. Simulations run until the lateral matrix thicknesses

approximate the empirical enamel thicknesses, see Methods, and for parameters S1 Table. Scale bar, 5

mm.

**Diffusion-limited simulations predict enamel distribution on pig molar teeth**

To simulate enamel matrix secretion in real teeth, first we focused on domestic pig molar teeth

which exhibit substantial variation in enamel thickness and EDJ topography (Fig 1A and 2A)

[20]. EDJ and enamel surface shapes were reconstructed from microCT scans of first lower

molars (Fig 2A, Methods). From the data, horizontal slices of cusps were extracted (Fig 2A) and

the EDJs were used as the starting point for the simulations. The horizontal plane represents a

relatively synchronous front of enamel matrix secretion and captures the complex EDJ

morphology of the pig molars [20]. The simulations show that the diffusion-limited process

reproduces the deep narrow furrows or fissures present on the concave sides of the real cusps

(Fig 2A and 2B). In contrast, these features are lost when the enamel matrix is geometrically

extrapolated from the EDJ, or when modeling with excess background nutrients (Fig 2C). These

results support the role of a diffusion-limited-like process in the regulation of enamel matrix

secretion, and underscore the distinct effects that the convex and concave EDJ features impose

on the enamel distribution. Indeed, whereas the overall distribution of convex and concave

features is conserved among molars of individual pigs, small differences in EDJ shape

correspond to large differences in enamel features (Fig 3).

**Fig 3. Small differences in EDJ shape produce large differences in diffusion-limited simulations.**

Horizontal slices of three different entoconid cusps of pig first lower molars and their diffusion-limited simulations using the same parameter values. Small differences in the size of EDJ ridges (arrowheads) correspond to large differences in enamel surface shape both in empirical and simulated enamel surfaces. The cusp in the center is 3D-simulated with the same parameters in Fig 4. Scale bar, 1 mm.

Next we simulated the matrix secretion in a whole cusp with both convex and concave features (Methods). A 3D reconstruction of these simulations show that the diffusion-limited model captures the overall enamel thickness patterns in which concavities show reduced enamel thickness whereas ridges show increased enamel thickness (Fig 4 and S1 Fig). A small ridge present in the middle of an EDJ concavity results in a local thickening of the enamel within an otherwise deep fissure (arrow heads in Fig 4), a feature completely lost in geometrically extrapolated surfaces (Fig 4D).

**Fig 4. 3D distribution of enamel is predicted by the diffusion-limited model.**

(A) The EDJ of the pig entoconid cusp (arrowhead in Fig 2B) has four ridges and only the buccal slope is convex whereas the other slopes are concave. (B) The enamel surface is thickest around the ridges and the EDJ concavities correspond to deep fissures with thinner enamel. The enamel cap, which has no EDJ, has been removed from the 3D data. (C) The diffusion-limited matrix simulation, matched to have the enamel thickness of the convex lateral slope, captures the enamel distribution patterns of the pig cusp (mean distance to the real surface is 61 μm, maximum difference = 362 μm, s.d. = 66.4). The small ridge present in the mesial slope of the cusp (arrowhead in the oblique views) corresponds to the narrow ridge of the empirical enamel surface (4B) and small ridge in an EDJ concavity (4A). (D) Geometric extrapolation from the EDJ results in relatively uniform 3D matrix thickness (mean distance to the real surface is 80 μm, maximum difference = 467 μm, s.d. = 92.0). The diffusion-limited simulations were done from 51 individual EDJ slices using the same parameter values (S1 Table). The thinner enamel in the lower parts of the lingual cusp ridge of the real cusp (towards the bottom in occlusal view) is due to the vicinity of the adjacent hypoconid cusp. Scale bar, 1 mm.

**Diffusion-limited simulations reproduce the progression of matrix secretion**

In addition to the distribution of enamel in fully formed teeth, the diffusion-limited simulations can be used to examine the progression of the matrix secretion process itself. The successive positions of the matrix-secreting front during development is recorded in teeth by incremental lines (laminations or striae of Retzius, [9]) that are broadly analogous to growth rings in trees [21]. These are preserved in mature enamel and can be observed from thin sections or through

phase contrast synchrotron imaging [22]. We obtained synchrotron data from a pig molar and

compared the positions of individual incremental lines in convex cusp ridges with the lines in

cusp concavities (Fig 5A). Both the virtual incremental lines of diffusion-limited simulations

and empirical incremental lines show initially relatively uniform distances from the EDJ, but

this uniformity disappears and the differences between ridges and valleys become progressively

larger as the secretion accelerates in the ridges and slows down in the valleys (Fig 5). These

results suggest that in addition to the final patterns of enamel distribution, the diffusion-limited

model captures aspects of the actual secretion process.

**Fig 5. Incremental lines in pig molar tooth show comparable patterns to diffusion-limited matrix lines.**

(A) Incremental lines (arrowheads) are visible in a synchrotron imaged fully formed but unerupted cusp

of the second lower molar. (B) Diffusion-limited simulation and (C) geometric extrapolation of matrix

show contrasting patterns. The different shades of grey correspond to progressive steps in simulations

(every fourth) and geometric extrapolation. (D) The arrows in Fig 5A, 5B, and 5C) show the lines that

are used to measure the progression of matrix secretion in the valleys (concave EDJ regions) and ridges

(convex EDJ regions). Both the empirical and diffusion-limited simulations produce progressively

thicker matrix in the ridges relative to the valleys (mean lines shown, error bars denote s.d.). The border

size (0.79) was set to produce the empirical enamel thickness using the same number of iterations as in

Fig 3 and 4. The target enamel thickness for the simulations was measured from the left side of the cusp.

Because the cusp was physically trimmed for synchrotron imaging, the section is missing the lower side.

Scale bar, 1 mm.

**Subtle EDJ concavities are sufficient to produce complex enamel surface features**

Domestic pigs are an example of species with relatively pointed molar cusps, allowing the

simulation of matrix secretion in the horizontal plane. In contrast, molars of primates with thick

enamel, including humans, typically have relatively low cusp relief. Therefore, to capture matrix

secretion of human molar morphology, we run the diffusion-limited simulations vertically.

Because the enamel secretion period is shorter towards the base of the tooth and the enamel is

therefore thinner, we implemented a nutrient sink at the base to simulate the shorter secretion

period (Methods). The sink decreases the rate of matrix formation towards the base of the crown

(Fig 6A), thereby approximating crown formation and the apical decline in matrix secretion

before the initiation of root development [9]. Biologically, the sink implementation can be

considered a simplification of the crown-tip-to-base growth of the capillaries providing nutrients

[18]. The simulations show the subtle waviness of the human EDJ, with mainly concave ripple-

like features, is enough to produce the characteristic undulations of the enamel surface (Fig 6B

and 6C). These features are further refined by simulations using lateral braces mimicking the

presence of adjacent teeth and alveolar bone (Fig 6B), in agreement with the suggested role of

the surrounding tissues in the regulation of tooth shape and mineralization [23, 24]. Taken

together, these simulations indicate that even subtle EDJ features present in many hominids are

important for the functional surface morphology of the tooth.

**Fig 6. The subtle EDJ topography of human molars is sufficient to produce enamel surface features in diffusion-limited simulations.**

(A) To simulate the thick enamel of human molars that also have a low cusp relief, we combined vertically oriented simulations with a basal nutrient sink. The sink is used to simulate the shorter time of matrix secretion in the lower parts of the crown. (B) Diffusion-limited simulations reproduce the surface features that are lost in excess nutrient simulations and geometric extrapolations. Braces mimicking adjacent teeth and bone constrain lateral expansion of enamel matrix. (C) Obliquely lingual views of empirical and simulated enamel surfaces with a half of the EDJ (yellow) visible. The diffusion-limited simulations were done from 27 individual EDJ slices using the same parameter values (S1 Table). The tooth shown is a human third lower molar. Scale bar, 5 mm.

**Diffusion-limited matrix secretion increases surface complexity**

The hallmark of diffusion-limited aggregation, which is closely related to the present model, is the fractal geometry, or self-similarity of the outcome [e.g. 25]. Although both the interfacial tension approximating the stiffness of the ameloblast layer and the background nutrient production weaken the fractal-like outcomes in our model, it is still interesting to explore the fractal nature of the enamel distribution, as it provides a quantitative measure of surface complexity. We contrasted complexities of the contours [26] of human and orangutan molars,

the latter being a species with highly complex enamel distribution (Fig 7A) [14]. Box-counting

dimensions ($D_B$) of the human and orangutan molars show that although the enamel surfaces do

not exhibit self-similarity per se, simulations with low background nutrient levels show stronger

diffusion-limited effects (Fig 7B, S2 Table). The main factor determining enamel surface

complexity of the orangutan molar appears to be the diffusion-limited effect of matrix secretion

(Fig 7B). Swapping the parameter values of the human and orangutan simulations (S1 Table)

produces orangutan-like enamel on human EDJ and human-like enamel on orangutan EDJ (Fig

7B, S2 Table). In contrast, excess background nutrient production results in no increase in

enamel complexity relative to the EDJ in either of the species (S2 Table), indicating that

diffusion-limited matrix secretion is required for the increase in surface complexity.

**Fig 7. Strong diffusion-limited effect generates complex enamel surfaces.**

(A) Compared to human molars, orangutan molars have a large number of small surface features. (B)

Vertically oriented simulations of human and orangutan molars with corresponding box counting

dimensions (mean values, see S2 Table) show that lower background nutrient production, resulting in a

stronger diffusion-limited effect, is enough to produce the relatively complex enamel surface of the

orangutan, and increase the complexity of the simulated human enamel surface. In contrast, using human

parameter values produces human-like enamel on orangutan EDJ. The human parameters are the same as

in Fig 6 (S1 Table) and the longitudinal sections simulated are from approximately corresponding

positions between the species. Teeth in (A) are not to scale. Scale bar, 1 mm in (B).

**Discussion**

Our diffusion-limited model suggests an objective basis for mapping the EDJ morphology to

enamel surface morphology. The simulations point to the critical role of the diffusion-limited

effect of matrix secretion in increasing tooth surface complexity and, together with convex and

concave EDJ features, in fine-tuning the shape of the enamel surface. Previously the rate of

matrix secretion has been reported to be highest in the outer parts of the enamel [9, 27]. The

result is supported by our simulations and synchrotron data, but only in the convex parts of the

crown (Fig 4). Nevertheless, complex patterns of enamel thickness on a tooth crown can result

from a single developmental process, without the need to evoke specific control or explanations

for the thickness of individual features. Combined with in-depth analyses of enamel formation

rates [4, 9, 27] and isotope compositions [28, 29] obtainable from sections, our model should

help to move studies using enamel thickness towards more mechanistic and predictive science.

The model also points to the need to investigate the role of extracellular space and capillaries in

regulating diffusion, and to the potential effects of cell-cell tension in the ameloblast layer [30].

Our approach should also be applicable to other systems with extracellular matrix secretion, or

organs in which directional tissue growth may be limited by a diffusion process. Diffusion-

limited free boundary problems have a long history in mathematics [16, 17, 31], and as shown

here, they can contribute to solving biological problems.

## Materials and methods

### Diffusion-limited model

We solve the model using a finite element method algorithm presented in [32]. The model is

motivated by the classical Stefan problem [16, 17], and our formulation follows the main

principles of the problem with some modifications. The model does not consider individual cells

in the system, rather it treats both the enamel matrix and the dentin as continuums.

Fundamentally, we are solely focused on how a 1-manifold interface representing the

ameloblast front on a cross-sectional plane evolves over time subject to interfacial tension and

restricted flow of nutrients. For mathematical and implementation details, see S1 Appendix. The

Matlab source code is available at https://github.com/tjhakkin/biomatrix.

**Dental data and processing**

Pig and human molar samples are described in [20] and the orangutan sample is from the

Finnish Museum of Natural History (UN2720). The microCT voxel resolutions were 10 to 24

µm (pig), 17 µm (human), and 32 µm (orangutan). For the analyses, the scans were

downsampled to 44 µm (pig molars) and 33 µm (human and orangutan molars). Synchrotron

data was collected at beamline ID19 of the European Synchrotron Radiation Facility, with voxel

resolution 2.24 µm, keV = 91, and 6000 projections in 4x accumulation mode. Synchrotron data

reconstruction used Paganin style single propagation distance phase contrast. All the image

processing steps after primary tomography reconstructions were carried out with Fiji 2.0 [33].

To digitize EDJs for simulations, EDJs in each individual section were traced with a freehand

selection or wand tool, and the area was converted to a line (Edit/Selection/Area to Line). The

line was then interpolated with an interval of 10 pixels (Edit/Selection/Interpolate), fitted to a

spline (Edit/Selection/Fit spline), and adjusted manually to follow the EDJ if needed. The

splines were saved in ROI Manager (Analyze/Tools/ROI manager/more/save). The splines were

saved as XY coordinates (File/Save As/XY Coordinates) that were converted to level sets using

a Python script, included with the source code. During the conversion the spatial node density

and the relative size of the EDJ within the domain were also defined. To scale different sized

EDJs uniformly (e.g., when EDJs become smaller towards the cusp tip), two small triangles

placed in diagonally opposing corners were included in each level set conversion. To enhance

the visualization of the incremental lines in synchrotron reconstructions, three adjacent slices

were averaged.

**Simulations and analyses**

The main simulation parameters are listed in S1 Table. Simulation output for each step is an

image file. The pig trigonids and talonids were simulated separately (Fig 2B). To compensate

for the isolated entoconid cusp being larger (Fig 3) than when part of the talonid (Fig 2),

interfacial tension and number of iterations were decreased in the individual cusp simulations.

For simulations of multiple sections, all the sections of the analyzed step were merged into a

stack in Fiji. For the pig molar cusp (Fig 4), every second microCT slice (20 μm interval, 51

slices), and for the human molar (Fig 6C), every fifth slice (66 μm interval, 27 slices) was

simulated. We used the basal sink to approximate developmental progression in the vertical

simulations. Because intercuspal regions lack the sink, vertical simulations slightly exaggerate

enamel thickness in valleys relative to cusp tips. Geometric extrapolation of enamel thickness

was obtained in Fiji by fitting a fixed sized circle along the EDJ (Process/Morphology/Gray

Morphology/Dilate). To visualize enamel thickness in 3D, the EDJ and enamel surfaces were

exported from Fiji (surfaces exported as Wavefront.obj from 3D Viewer plugin) and imported

into Meshlab (http://meshlab.sourceforge.net/). In Meshlab, Hausdorff distance was used to

compare distances between two surfaces [34]. The distances were calculated after smoothing the

meshes with Laplacian smooth (3 steps). All 3D visualizations use orthographic projections.

We calculated box-counting dimensions from microCT slices of human and orangutan

molars. Binary contours were generated after thresholding in Fiji (Process/Binary/Outline) and

analyzed using Fiji FracLac 2.5 plugin [26]. Twelve different starting positions for the grids

were used, and the contours were rotated at 36 degree steps (resulting in total of 120 $D_B$ values

for each contour). Default sampling size was used for the scaling method. We analyzed the

contours with (S2 Table) and without the outer walls of the profiles, and the pattern of results

remained the same.

## Supporting information

**S1 Fig. Enamel area and surface perimeter in horizontal slices of a pig cusp.** Whereas both

the diffusion-limited simulation and geometric extrapolation of Fig. 4 approximate the amount

of real enamel, only the diffusion-limited simulation reproduces the length of the perimeter of

the real surface. The drop in enamel perimeter in the diffusion-limited simulation towards the

cusp tip relative to empirical data is due to the horizontal simulations not capturing the

relatively round EDJ tip. (PDF)

**S1 Table. EDJ conversion and simulation parameters.** Node density denotes the number of

nodes along the longest axes in the regular triangular mesh. 'Border' is the proportional size of

the margins surrounding the EDJ shape. 'Base' sets the y-axis position of the base in simulations

using a sink. All EDJ simulations are run with Neumann domain boundaries, except for the

simulations using basal sink for which we use mixed boundaries (Dirichlet/Neumann), with a

fixed time step 0.001. All parameters are dimensionless and their absolute values have no

meaning. (PDF)

**S2 Table. Box-counting dimensions calculated for human and orangutan molar teeth.** The

$D_B$ values are calculated for the binary outlines of the EDJs and enamel surfaces. The

background nutrient production values for the excess background nutrient, weak and strong

diffusion-limited effect simulations are 160, 75, and 30, respectively. (PDF)

**S1 Appendix. A detailed implementation of the diffusion-limited model.** (PDF)

## Acknowledgements

369  We thank H. Suhonen for help with microCT imaging, and J. Laakkonen for help with material.

370  We acknowledge the European Synchrotron Radiation Facility for provision of synchrotron

371  radiation facilities and thank P. Tafforeau for assistance in using beamline ID19. We thank M.

372  Fortelius and the members of the Center of Excellence in Experimental and Computational

373  Developmental Biology Research for discussions or advise.

## Author Contributions

**Conceptualization:** Teemu J. Häkkinen, S. Susanna Sova, Jukka Jernvall.

**Data curation:** Teemu J. Häkkinen, S. Susanna Sova, Jukka Jernvall.

**Funding acquisition:** Teemu J. Häkkinen, Jukka Jernvall.

**Investigation:** Teemu J. Häkkinen, S. Susanna Sova, Ian J. Corfe, Jukka Jernvall.

**Methodology:** Teemu J. Häkkinen, S. Susanna Sova, Antti Hannukainen, Jukka Jernvall.

**Project administration:** Jukka Jernvall.

**Resources:** S. Susanna Sova, Ian J. Corfe, Leo Tjäderhane, Jukka Jernvall.

**Software:** Teemu J. Häkkinen.

**Supervision:** Leo Tjäderhane, Antti Hannukainen, Jukka Jernvall.

**Validation:** Teemu J. Häkkinen, S. Susanna Sova, Jukka Jernvall.

**Visualization:** Teemu J. Häkkinen, S. Susanna Sova, Jukka Jernvall.

**Writing – original draft:** Teemu J. Häkkinen, S. Susanna Sova, Jukka Jernvall.

**Writing – review & editing:** Teemu J. Häkkinen, S. Susanna Sova, Ian J. Corfe, Leo

Tjäderhane, Antti Hannukainen, Jukka Jernvall.

## References

1.  Schwartz GT. Taxonomic and functional aspects of the patterning of enamel thickness

distribution in extant large-bodied hominoids. Am J Phys Anthropol. 2000; 111:221-244.

2.   Grine FE. Enamel Thickness of Deciduous and Permanent Molars in Modern *Homo*

*sapiens*. Am J Phys Anthropol. 2005; 126:14-31.

3.   Smith TM, Olejniczak AJ, Martin LB, Reid DJ. Variation in hominoid molar enamel

thickness. J Hum Evol. 2005; 48: 575-592.

4.   Dean MC. Tooth microstructure tracks the pace of human life-history evolution. *Proc R Soc*

*B*. 2006; 273:2799-2808.

5.   Lucas P, Constantino P, Wood B, Lawn B. Dental enamel as a dietary indicator in

mammals. BioEssays. 2008; 30:374-385.

6.   Olejniczak AJ, Smith TM, Skinner MM, Grine FE, Feeney RNM, Thackeray JF, et al.

Three-dimensional molar enamel distribution and thickness in Australopithecus and

Paranthropus. Biol Lett. 2008; 4:406-410.

7.   Skinner MM, Alemseged Z, Gaunitz C, Hublin J-J. Enamel thickness trends in Plio-

Pleistocene hominin mandibular molars. J Hum Evol. 2015; 85:35-45.

8   Pan L, Dumoncel J, de Beer F, Hoffman J, Thackeray JF, Duployer B, et al. Further

morphological evidence on South African earliest Homo lower postcanine dentition:

Enamel thickness and enamel dentine junction. J Hum Evol. 2016; 96:82-96.

9.   Kierdorf H, Breuer F, Witzel C, Kierdorf U. Pig enamel revisited – Incremental markings in

enamel of wild boars and domestic pigs. J Struc Biol. 2019; 205:48-59.

10.   Wright JT, Hart TC, Hart PS, Simmons D, Suggs C, Daley B, et al. Human and mouse

enamel phenotypes resulting from mutation or altered expression of AMEL, ENAM,

MMP20 and KLK4. Cells Tissues Organs. 2009; 189:224-229.

11.   Horvath JE, Ramachandran GL, Fedrigo O, Nielsen WJ, Babbitt CC, St. Clair EM, et al.

Genetic comparisons yield insight into the evolution of enamel thickness during human

evolution. J Hum Evol. 2014; 73:75-87.

12.   Daubert DM, Kelley JL, Udod YG, Habor C, Kleist CG, Furman IK, et al. Human enamel

thickness and ENAM polymorphism. *Int J Oral Sci*. 2016; 8:93-97.

13.   Skinner MM, Wood BA, Hublin J-J. Protostylid expression at the enamel-dentine junction

and enamel surface of mandibular molars of *Paranthropus robustus* and *Australopithecus africanus*. J Hum Evol. 2009; 56:76-85.

14. Skinner MM, Evans A, Smith T, Jernvall J, Tafforeau P, Kupczik K, et al, Contributions of enamel-dentine junction shape and enamel deposition to primate molar crown complexity. Am J Phys Anthropol. 2010; 142:157-163.

15. Guy F, Lazzari V, Gilissen E, Thiery G. To what extent is primate second molar enamel occlusal morphology shaped by the enamel-dentine junction? PLoS ONE. 2015 Sep 25. 10:e0138802. doi:10.1371/journal.pone.0138802.

16. Chen S, Merriman B, Osher S, Smereka PA. A simple level set method for solving Stefan problems. J Comp Physics. 1997; 135:8-29.

17. Chen H, Min C, Gibou F. A numerical scheme for the Stefan problem on adaptive Cartesian grids with supralinear convergence rate. J Com. Physics. 2009; 228:5803-5818.

18. Yoshida S, Oshima, H, Kobayashi S. Vascularization of the enamel organ in developing molar teeth of rats – Scanning electron microscope study of corrosion casts. Okajima Folia Anat jpn. 1989; 66:99-112.

19. Ida-Yonemochi H, Nakajima M, Saku T. Heparanase, heparan sulfate and perlecan distribution along with the vascular penetration during stellate reticulum retraction in the mouse enamel organ. *Arch Oral Biol.* 2010; 55:778-787.

20. Sova SS, Tjäderhane L, Heikkilä PA, Jernvall J. A microCT study of three-dimensional patterns of biomineralization in pig molars. *Front Physiol.* 2018 Feb 9. 9:71 doi: 10.3389/fphys.2018.00071.

21. Beynon AD, Dean MC, Reid DJ. On thick and thin enamel in Hominoids. Am J Phys Anthropol. 1991; 86:295-309.

22. Tafforeau P, Smith TM. Nondestructive imaging of hominoid dental microstructure using phase contrast X-ray synchrotron microtomography. J Hum Evol. 2008; 54:272-278.

23. Marchiori DF, Packota GV, Boughner JC. Third-molar mineralization as a function of available retromolar space. Acta Odont Scan. 2016; 74:509-517.

24. Renvoisé E, Kavanagh, KD, Lazzari V, Häkkinen TJ, Rice R, Pantalacci S, et al.

Mechanical constraint from growing jaw facilitates mammalian dental diversity. Proc Nat
Acad Sci USA. 2017; 114:9403-9408.

25. Halsey TC. Diffusion-limited aggregation: A model for pattern formation. Physics Today
2000; 53:36-41.

26. Karperien A. FracLac for ImageJ. Available from:
http://rsbweb.nih.gov/ij/plugins/fraclac/FLHelp/ Introduction.htm. 2012.

27. Boyde A. Enamel. In: Oksche A, Vollrath L (Eds.), Handbook of Microscopic Anatomy vol
V/6 Teeth. Springer; 1989. pp. 309-473.

28. Green DR, Oack G, Colman AS. Determinants of blood water $\delta^{18}O$ variation in a population
of experimental sheep: Implications for paleoclimate reconstruction. Chem Geol. 2018;
485:32-43.

29. Smith TM, Austin C, Green DR, Joannes-Boyau R, Bailey S, Dumitriu D, et al. Wintertime
stress, nursing, and lead exposure in Neanderthal children. Sci Adv. 2018; 4: 10:eaau9483.

30. Suwa G, Kono RT. A micro-CT based study of linear enamel thickness in the mesial cusp
section of human molars: reevaluation of methodology and assessment of within-tooth,
serial, and individual variation. *Anthropological Sci*. 2005; 113:273-289.

31. Vuik C. Some historical notes about the Stefan problem, Nieuw Archief voor Wiskunde.
1993; 11:157-167.

32. Häkkinen T. On Free Boundary Problems. MSc thesis, Aalto University. 2016. Available
from: https://aaltodoc.aalto.fi/handle/123456789/20347

33. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat methods. 2012; 9:676-682.

34. Cignoni P, Rocchini C, Scopigno R. Metro: measuring error on simplified surfaces Comput
Graph Forum. 1998; 17:167-174.

473