INTERPRETATIVE SUMMARY. Validation of luteolysis monitoring tool. Adriaens.

Monitoring of milk progesterone allows to detect luteolysis preceding a dairy cow's estrus. In this

research, two algorithms were compared using on-farm measured milk progesterone data with a

variable, cost-effective sampling rate. The first was the state of the art for progesterone monitoring,

a smoothing multiprocess Kalman filter compared with a fixed threshold. The second algorithm was

the recently developed progesterone-based monitoring algorithm using synergistic control. Both

algorithms detected approximately the same amount of luteolyses, but the second gave the alerts on

average 20 hours earlier. This proves that this algorithm can operate reliably in an on-farm setting.

9
VALIDATION OF LUTEOLYSIS MONITORING TOOL

Short communication: Validation of a novel milk progesterone-based tool to monitor

luteolysis in dairy cows using cost-effective, on-farm measured data

Ines Adriaens*[,1], Wouter Saeys*, Katleen Geerinckx†, Bart De Ketelaere*, Ben Aernouts*‡

* KU Leuven, Department of Biosystems, MeBioS, Kasteelpark Arenberg 30, 3001, Heverlee,

Belgium

† Province of Antwerp, Hooibeekhoeve, Hooibeeksedijk 1, 2440 Geel, Belgium

‡ KU Leuven, Department of Microbial and Molecular Systems, Cluster for Bioengineering

Technology, Campus Geel, 2440 Geel, Belgium

[1] Corresponding author: ines.adriaens@kuleuven.be, Kasteelpark Arenberg 30, box 2456, 3001,

Heverlee, Belgium, +32 16 32 88 73

The Progesterone (**P4**) monitoring algorithm using Synergistic control (**PMASC**) uses the

luteal dynamics to identify fertility events in dairy cows (Adriaens et al., 2017, 2018a). Hereto,

PMASC employs a combination of mathematical functions describing the increasing and decreasing

P4 concentrations during the development and regression of the corpus luteum and a statistical

control chart which allows to identify luteolysis. The mathematical model combines sigmoidal

functions from which the cycle characteristics can be calculated. Both the moment at which

luteolysis is detected and confirmed by PMASC, as well as the model features themselves can be

used to inform the farmer on the fertility status of the cows.

Until now, the PMASC methodology was designed and optimized using milk P4 data obtained

through experimental trials, collected at maximum frequency (e.g. every milking) and analyzed

post-hoc using the state-of-the-art laboratory ELISA technique. Moreover, this method was not yet

challenged on data sampled at reduced frequency to show its possible cost-effectiveness, nor was it

shown that PMASC can work on data measured with an on-line device proving its value as an on-

farm monitoring algorithm. The objective of this study was to show the capability of PMASC to

work in an on-farm setting, identifying luteolysis using smart-sampled data and analyzed via an on-

farm lateral flow immunoassay (**LFIA**). Moreover, we compared the PMASC algorithm with the

state of the art for P4 monitoring on farm, i.e. smoothing the raw P4 values, combined with a fixed

threshold for detection of luteolysis and alerts.

Datasets of on-farm measured milk P4 were collected on two farms in Flanders, Belgium. The

first dataset (dataset 1) was obtained from an experimental research farm between March 2017 and

January 2019. On this farm, the herd was split up in two groups, each consisting of on average 55

lactating dairy cows. Both groups were milked with an automated milking system (VMS, Delaval,

Tumba, Sweden). The second dataset (dataset 2) was collected on a commercial dairy farm between

November 2016 and December 2018. The herd consisted of on average 250 animals, milked by four

automated milking systems (VMS, Delaval, Tumba, Sweden). On both farms, animals were fed a

mixed ration of grass and corn silage, supplemented with concentrates through automated

concentrate feeders and in the milking robot.

For the P4 analysis, milk samples were taken automatically by the sampling unit integrated in

the Herd Navigator^{TM} (Delaval, Tumba, Sweden; Lattec, Hillerød, Denmark), and the milk P4

concentration was determined by the built-in LFIA milk analyzer. More details on the working

principle of this device can be found in Blom and Ridder, (2010) and Friggens et al., (2007). The

samples for P4 analysis were taken with a variable sampling frequency, of which the concept has

been described by Friggens and Chagunda, (2005). Basically, the time between samples is

determined by the days in milk, the likelihood of luteolysis, the P4 concentration of the previous

sample, the performed inseminations and their timing, the likelihood of pregnancy, and the

probability of prolonged postpartum anestrus or cysts. As the consumables are an important cost for

P4 monitoring on farm, this sampling scheme ensures that the necessary fertility information can be

gathered on crucial moments with a minimum number of samples.

Dataset 1 consisted of 10,791 P4 measurements originating from 144 lactations for 104 unique

cows (parity 1 to 6, 2.15±1.16, average milk production per day 36.47±12.66 kg, mean±standard

deviation). The second dataset consisted of 36,645 P4 measurements, originating from 460

lactations and 342 unique cows, with parities between 1 and 8 (2.3±1.26, average milk production

per day 34.24±10.10 kg).

The state of the art (milk) P4 monitoring in dairy cows builds on a multiprocess Kalman filter

(**MPKF**) to smooth the profiles and reduce the effect of potential outliers (Korsgaard and

Løvendahl, 2002; Friggens and Chagunda, 2005; Friggens et al., 2007). When the smoothed P4

value undercuts a 5 ng/mL threshold (**T**), after that it exceeded 5 ng/mL twice, luteolysis was

deemed confirmed. This algorithm, further referred to as MPKF+T, was implemented and used to

identify the reference for luteolysis which was then used to benchmark the performance of our

PMASC algorithm in terms of detecting the luteolysis events.

Progesterone profiles (i.e. the P4 time series of one lactation) with less than 20 measurements in

total (e.g. because they just started at the moment of data collection), or less than 5 measurements

below or above 5 ng/mL were discarded. Also, P4 data following a gap in the P4 data of at least 20

days were excluded. Accordingly, 719 and 5322 measurements originating from 47 and 317

lactations were deleted from each dataset respectively, leaving 10,072 and 31,213 measurements

from 133 and 409 lactations belonging to 98 and 316 unique cows. The characteristics of the

remaining profiles and related P4 data are summarized in Table 1.

[TABLE 1]

Adriaens et al., (2018a) describes the original PMASC to process time series of P4

measurements taken each milking, two to four times a day. To avoid false alerts caused by

coincidentally low P4 values (e.g. outliers), a first evidence of luteolysis (low P4) identified by the

statistical control chart needed to be confirmed by two additional successive measurements before a

luteolysis alert was raised by PMASC. However, in the datasets employed in this study, the P4

concentration was not available for every milking as the number of and moment of samplings

depended on the algorithm implemented in the analyzer. For example, when an upcoming estrus

was confirmed by this algorithm, no new milk samples were analyzed in the following 3 to 6 days.

In case of a steep drop in milk P4 in combination with very low absolute P4 concentrations (< 5

ng/mL), this often resulted in less than the three consecutive out-of-control measurements available

to confirm luteolysis with PMASC. Therefore, the main decision rule of PMASC needed to be

adjusted in function of these new datasets. In previous studies on the timing between luteolysis and

ovulation time (Roelofs et al., 2006; Adriaens et al., 2018b), it was found that ovulation followed

about 60 to 80 hours after the onset of luteolysis. To have the largest chance on successful

conception, insemination should take place 24 to 12 hours before ovulation (Roelofs, 2005). Based

on this information, we evaluated all the samples collected within 36 hours following an out-of-

control measurement to confirm or object luteolysis. More concretely, following rule was

implemented: if more than half of the measurements within this 36h window were also out of

control, luteolysis was assumed confirmed, an alert was raised and a new cycle started. In the other

case, the out-of-control measurement was considered an outlier. In practice, this would ensure that

luteolysis has taken place, while there would have been still sufficient time left to arrange the

insemination. Moreover, this rule gives some flexibility to deal with missing samples and outliers.

In future studies when the sampling frequency can be defined based on PMASC, other rules can be

implemented.

Next, PMASC was applied to the time series of P4 values of each lactation to evaluate its

performance when in this cost-effective setting controlled by the device's algorithm. The number of

and the timing of the alerts generated by PMASC were compared to those obtained using the

MPKF+T algorithm. Also, the number of measurements per detected cycle was calculated and the

cycle characteristics were summarized.

An overview of the results is given in Table 2. Dataset 1 consisted of 133 P4 profiles, with an

average sampling rate of 0.48±0.07 measurements per day in the period between the first and the

last measurement. This corresponds to one measurement each 2.07 days. For dataset 2, the 409

profiles had an average sampling rate of 0.50±0.07 measurements per day, corresponding to one

sample per 1.98 days. The MPKF+T algorithm gave 427 and 1416 luteolysis alerts (i.e. smoothed

time series undercutting the fixed threshold of 5 ng/mL), respectively for dataset 1 and 2, while

PMASC gave 420 and 1411 alerts. Alerts within 5 days of each other were taken as the same

luteolysis. Based on this timing, 25 (5.8%) and 54 (3.81%) of the alerts of MPKF+T could not be

matched with one of PMASC, and 18 (4.3%) and 49 (3.47%) of the alerts of PMASC could not be

matched with one given by MPKF+T. These unmatched alerts could be attributed to (1) a sampling

frequency too low to confirm the out-of-control samples, for example due to quite low luteal phase

P4 values, after which only one sample sufficed to generate an alert using MPKF+T. This was the

case for respectively 58% and 42% of the unmatched alerts for dataset 1 and 2; (2) high follicular

phase P4 values, resulting in enough evidence for PMASC to generate an alert, while MPKF+T did

not yet undercut the threshold value. In this case, sometimes the MPKF+T alert followed later (but

outside the time window for matching i.e. 5 days. This was the case for 42% and 58% of the

unmatched alerts in dataset 1 and 2, respectively. In the first case, the smart sampling algorithm

does not take a new confirmatory sample, while we set PMASC to evaluate at least two samples per

estrus, and PMASC thus failed to give an alert. When the sampling is based on decision rules

implemented in PMASC, outside the scope of this research, also these luteolyses can be detected by

PMASC. The intermediate values causing additional PMASC alerts in the second case (2), are

mainly due to the calibration technique of the online measurement device, as also discussed in

Adriaens et al., (2019).

[TABLE 2]

An example of a P4 profile and the corresponding alerts is shown in Figure 1. For the first cycle,

no alert was generated with PMASC. However, the sample for which the smoothed value is below

the 5 ng/mL threshold was also out-of-control in PMASC, but the next sample was only taken 6

days later. As such, luteolysis was not detected by PMASC because of the lack of confirmatory

samples out-of-control within 36 hours after the first out-of-control measurement. The final

PMASC models for which the detected luteolysis matches with the MPKF+T algorithm, while the

MPKF smoothed data are given in dark blue. The threshold of 5 ng/mL P4 is plotted in green. As

seen in the figure, the first sample detected as out of control by PMASC is often not yet below the 5

ng/mL threshold, and the model characteristics can be used to estimate luteolysis reliably and

independent of both the milking frequency and the absolute P4 values. The time lag between the

drop in P4 during luteolysis and the smoothed MPKF value undercutting the threshold is shown to

be dependent of sampling rate and the absolute P4 values (yellow box).

[FIGURE 1]

The first out-of-control sample detected by PMASC preceded the MPKF/T alarm on average

20.0±16.1 hours and 2.64 milkings for dataset 1 and 19.8±23.7 hours and 2.2 milkings for dataset 2.

The time-delay of the MPKF ensures robustness for outliers and measurement errors, and will

correct for unexpected variability. However, contrasting this MPKF smoothed value against a fixed

threshold results in alerts for which the timing strongly depends on the absolute P4 values measured

and the sampling rate. In cases where the sampling frequency is insufficiently high for timely

detection of luteolysis by the MPKF+T method, for example when the sampling or the analysis

fails, the alerts of the MPKF+T are less valuable because they arrive too late for timely

insemination. However, when these low P4 values are detected by PMASC, information calculated

from the model parameters (e.g. time of the inflection point) can be used to interpret these

detections and eventually, to adjust the decision process accordingly.

Using the model parameters, the P4 cycle characteristics were calculated. The average cycle

length calculated as the time between two successive luteolysis alerts was 27.33±10.0 days, with

56% of the cycles between 19 and 25 days. The average time between the two inflection points of

the increasing Hill function and the decreasing Gompertz function was 14.4±10.0 days, which is

slightly higher than the time reported in Adriaens et al., (2017), probably because in this previous

work only data of normal cycles and profiles were used. The average baseline and maximal P4

concentrations were 0.97±0.77 and 21.5±3.9 ng/mL respectively, and the decrease in P4 at

luteolysis was typically fast with a slope of 36.17±24.6 ng/mL per day. The latter means that a drop

in P4 from 25 to 2 ng/mL during luteolysis on average takes 14.56 hours, which is in agreement

with the values reported by Blavy et al., (2016) and Adriaens et al., (2018b).

In this study, we investigated the performance of PMASC on on-farm measured datasets using

less frequent sampling and the LFAI technique, and compared it with the current state of the art

being an MPKF+T algorithm. We showed that PMASC, on the condition of a small adjustment in

its decision requirements, was able to deal with less samples and the alternative measurement

technique, and accordingly, has the potential to work cost-effective in an on-farm setting. More concretely, this study showed that a limited number of samples was sufficient to fit the mathematical model used by PMASC, and did on average not influence the cycle characteristics extracted from it. The non-detected luteolyses were related to cases in which only one sample was taken to confirm estrus, which is rather rare and mainly occurs during the first estrus after calving, in which the maximum P4 concentration remains low. When the sampling rate would be optimized based on information extracted from PMASC (e.g. the time from the previous luteolysis, out-of-control samples etc.) these kinds of artifacts could be avoided. While it would be possible to introduce simple decision rules in the algorithm, the implementation and optimization of the P4 sampling rate based on PMASC was beyond the scope of this study.

ACKNOWLEDGEMENTS

This work was supported by the Institute for the Promotion of Innovation through Science and Technology in Flanders, Belgium (IWT) [IWT-LA project 110770]. Ines Adriaens is supported by the Fund for Scientific Research (FWO) Flanders, grant number 11ZG916N. We thank Lattec (Hillerod, Denmark) and the commercial dairy farmer who provided us with the second P4 dataset.

REFERENCES

Adriaens, I., T. Huybrechts, K. Geerinckx, D. Daems, J. Lammertyn, B. De Ketelaere, W. Saeys, and B. Aernouts. 2017. Mathematical characterization of the milk progesterone profile as a leg up to individualised monitoring of reproduction status in dairy cows. Theriogenology 103:44–51. doi:https://doi.org/10.1016/j.theriogenology.2017.07.040.

Adriaens, I., O. Martin, W. Saeys, B. De Ketelaere, N.C. Friggens, and B. Aernouts. 2019. Validation of a novel milk progesterone-based tool to monitor luteolysis in dairy cows. Timing of the alerts and robustness against missing values.

Adriaens, I., W. Saeys, T. Huybrechts, C. Lamberigts, K. Geerinckx, J. Leroy, B. De Ketelaere, B.

Aernouts, and K.U. Leuven. 2018a. A novel system for online fertility monitoring based on

milk progesterone. J. Dairy Sci. 101:1–14. doi:10.3168/jds.2017-13827.

Adriaens, I., W. Saeys, C. Lamberigts, M. Berth, K. Geerinckx, J. Leroy, B. De Ketelaere, and B.

Aernouts. 2018b. Short communication: Sensitivity of estrus alerts and their relation with the

timing of the LH surge. J. Dairy Sci. 102:1-5. doi:10.3168/jds.2018-15514

Blavy, P., M. Derks, O. Martin, J.K. Hoglund, and N.C. Friggens. 2016. Overview of progesterone

profiles　　in　　dairy　　cows.　　Theriogenology　　86:1061–1071.

doi:https://doi.org/10.1016/j.theriogenology.2016.03.037.

Blom, J.Y., and C. Ridder. 2010. Reproductive management and performance can be improved by

use of Delaval Herd Navigator. Page in Proceedings of First North American Conference on

Precision Dairy Management.

Friggens, N.C., and M.G.G. Chagunda. 2005. Prediction of the reproductive status of cattle on the

basis　of　milk　progesterone　measures:　model　description.　Theriogenology　64:155–90.

doi:10.1016/j.theriogenology.2004.11.014.

Friggens, N.C., K.L. Ingvartsen, I.R. Korsgaard, T. Larsen, P. Løvendahl, C. Ridder, and N.I.

Nielsen. 2007. System and a method for observing and predicting a physiological state of an

animal. US Pat. No. US7302349 B2.

Korsgaard, I.R., and P. Løvendahl. 2002. An introduction to multiprocess class II mixture models.

Pages 185–188 in Proceedings of the 7th World Congress on genetics applied to livestock

production. Département de Génétique Animale INRA, Castanet-Tolosan.

Roelofs, J.B. 2005. When to inseminate the cow? Insemination, ovulation and fertilization in dairy

cattle. PhD thesis, Wageningen Institute of Animal Sciences, Wageningen, the Netherlands,.

Roelofs, J.B., F.J.C.M. Van Eerdenburg, W. Hazeleger, N.M. Soede, and B. Kemp. 2006. Short

communication: Relationship between progesterone concentrations in milk and blood and time

of   ovulation   in   dairy   cattle.   Anim.   Reprod.   Sci.   91:337–343.

doi:doi:10.1016/j.anireprosci.2005.04.015.

Figure 1. A profile of raw milk progesterone (P4) measurements for one lactation are indicated in

black. The light blue crosses are the first out-of-control measurements detected by PMASC, and the

light blue lines give final models fitted by PMASC. The first luteolysis was not detected because of

an insufficiently high sampling rate (red box), although the measurements pushing the MPKF

(indicated in blue) below the threshold (indicated in green) were also out of control for PMASC

(indicated with purple cross). The (inconsistent) time-lag between the MPKF dropping below the

threshold and actual low measurements of P4 can be seen in the yellow box.

Table 1. Summary of the datasets of farm 1 and farm 2

Table 2. Overview of the detected luteolysis by each algorithm and the corresponding cycle

characteristics

241   Adriaens, Table 1

Dataset	No. of cows	No. of profiles	MY[1] (kg)	MI[2] (h)	P4[3]	
					Concentration (ng/mL)	No. of meas. per profile
Dataset 1	98	133	9.98 – 65.5 (36.47 ± 12.66)	5.6 - 29.7 (7.9 ± 2.62)	0.03-27.98 (14.71±10.23)	23 - 168 (75.7 ± 29.97)
Dataset 2	321	421	10.1-66.4 (32.0±11.7)	5.0-29.05 (9.36±3.19)	0.01-27.99 (15.36±9.69)	21-232 (80.2±35.39)

242   [1] Daily milk yield in kg; [2] Average milking interval in hours; [3]P4 milk progesterone;
243

244   Adriaens, Table 2

	Dataset 1	Dataset 2	Total
Number of profiles	133	409	**542**
Number of samples/day	0.48	0.5	
Number of days/sample	2.1	1.98	
Number of alerts MPKF+T	427	1416	**1843**
Number of alerts PMASC	420	1411	**1831**
Number of unmatched alerts PMASC	18 (4.3%)	49 (4.3%)	**72**
Number of unmatched alerts MPKF+T	25 (5.8%)	54 (3.83%)	**74**
Total number of unmatched alerts	43	103	**146**
Infrequent sampling / low luteal phase	25	43	**68**
Intermediate and noisy P4	18	60	**78**
Average cycle length (days)	27.33±10.02	26.37±11.27	**26.58±9.62**
% normal cycle length 19-25 days	56.33%	55.21%	**56.47%**
Average baseline concentration	1.09±0.91	0.93±0.73	**0.97±0.77**
Average luteal concentration	21.89±4.12	21.42±3.9	**21.53±3.96**
Decreasing slope during luteolysis[1]	34.26±23.33	37.41±24.94	**36.7±24.61**

245   [1] Calculated using the decreasing Gompertz model implemented in PMASC