INTERPRETATIVE SUMMARY. Validation of luteolysis monitoring tool Adriaens.

Recently, a novel progesterone-based monitoring algorithm using synergistic control, PMASC, was

developed. This algorithm employs the known luteal dynamics of the progesterone profile to

estimate the moment of luteolysis with minimal dependence on the absolute measurement values.

As its value and implementability strongly dependends on the number of samples it requires to

obtain the desired accuracy, its performance on cost-effective, 'smart' sampled and on-farm

measured data is demonstrated.

VALIDATION OF LUTEOLYSIS MONITORING TOOL

SHORT COMMUNICATION: Validation of a novel milk progesterone based tool to monitor

luteolysis in dairy cows. Performance on cost-effective, on-farm measured data

Ines Adriaens*[,1], Wouter Saeys*, Katleen Geerinckx†, Bart De Ketelaere*, Ben Aernouts*‡

* KU Leuven, Department of Biosystems, MeBioS, Kasteelpark Arenberg 30, 3001, Heverlee,

Belgium

† Province of Antwerp, Hooibeekhoeve, Hooibeeksedijk 1, 2440 Geel, Belgium

‡ KU Leuven, Department of Microbial and Molecular Systems, Cluster for Bioengineering

Technology, Campus Geel, 2440 Geel, Belgium

[1] Corresponding author: ines.adriaens@kuleuven.be, Kasteelpark Arenberg 30, box 2456, 3001,

Heverlee, Belgium, +32 16 32 88 73

[ABSTRACT, max 2500 keystrokes, no statistics, no references, ..., only for submission, not for the

manuscript in SC]

*[**Key words**. milk progesterone, fertility, dairy cow, monitoring tool; only for submission, not for the*

manuscript in SC]

The Progesterone (**P4**) monitoring algorithm using synergistic control (**PMASC**) enables to identify fertility events stooled on the underlying physiological basis of the related progesterone dynamics (Adriaens et al., 2017, 2018b). Therefore, PMASC employs a combination of mathematical functions describing the increasing and decreasing P4 concentrations during the development and regression of the corpus luteum (**CL**) and a statistical control chart which allows to identify luteolysis. The mathematical model is a combination of sigmoidal functions, from which the cycle characteristics can be calculated. Both the moment that luteolysis is detected and confirmed by PMASC, and the model features might possibly be used to inform the farmer on the fertility status of the cows.

Until now, the PMASC system was designed and optimized on data obtained through experimental trials and analyzed *post-hoc* via the state-of-the-art laboratory ELISA technique. As a result, it was not yet tested on data sampled according to a reduced frequency to show its possible cost-effectiveness, nor was it shown that PMASC can work on data measured with an on-line device proving its value as an on-farm monitoring algorithm. The objective of this study was to show the capability of PMASC to work in an on-farm setting, identifying luteolysis using smart-sampled data and analyzed via an on-farm lateral flow immunoassay (**LFIA**).

To this end, datasets of on-farm measured milk P4 were collected on two farms in Flanders, Belgium. The first dataset (dataset 1) was obtained from an experimental research farm between December 2016 and March 2018. On this farm, the herd was split up in two groups, each consisting of on average 55 lactating dairy cows. Both groups were milked with an automated milking system (VMS, Delaval, Tumba, Sweden). The second dataset (dataset 2) was collected on a commercial dairy farm between November 2016 and December 2018. The herd consisted of on average 250 animals, milked by four automated milking systems of Delaval (VMS, Delaval, Tumba, Sweden). On both farms, animals were fed a mixed ration of grass and corn silage, supplemented with concentrates through automated concentrate feeders and in the milking robot.

For the P4 analysis, milk samples were taken automatically by the sampling unit integrated in

the 'Herd Navigator[TM]' (Delaval, Tumba, Sweden; Lattec, Hillerød, Denmark), and the milk P4

concentration was determined via an on-farm LFIA. More details on the working principle of this

device can be found in Blom and Ridder, (2010) and in the patent of Friggens et al., (2007). The

samples for P4 analysis were taken with a variable sampling frequency. While the algorithm behind

this sampling scheme is confidential, its basis can be found in Friggens and Chagunda, (2005).

Basically, the time between samples is dependent on the days in milk, the likelihood of luteolysis,

the P4 concentration of the previous sample, the performed inseminations and their timing, the

likelihood of pregnancy, prolonged postpartum anestrus or cysts and so on. As the consumables are

one of the most important costs for P4 monitoring on farm, this 'intelligent' sampling frequency

ensures that as much information can be gathered on crucial moments with as little samples as

possible.

Dataset 1 consisted of 10,958 P4 measurements originating from 151 lactations and 102 cows.

However, the collection period enclosed two months in which the sampling frequency was

increased for several cows to one sample per milking in the context of a research trial, as described

in Adriaens et al., (2018a). A total of 3496 samples (31.9%) was taken in the context of this

research trial. On the second farm, P4 samples were collected according to the variable 'smart'

sampling scheme during the whole period. This dataset consisted of 36,535 P4 measurements,

originating from 457 lactations and 342 unique cows, with parities between 1 and 8 (2.3±1.6,

mean±standard deviation).

The raw P4 measurements were smoothed using a multiprocess Kalman filter (**MPKF**) as

described in Friggens and Chagunda, (2005) and Friggens et al., (2007, 2008), and estrus alerts

were generated when this smoothed P4 value undercut a 5 ng/mL threshold (**T**). Due to the absence

of a real gold standard for estrus and ovulation, these estrus alerts (MPKF+T) were taken as the

reference for luteolysis against which the performance of PMASC was tested.

Progesterone profiles (i.e. the P4 time series of one lactation) with less than 15 measurements in

total, or less than 5 measurements below or above 5 ng/mL were discarded. Also, P4 data following

a gap in the P4 data of at least 20 days were excluded. Accordingly, 870 and 2780 measurements

originating from 52 and 214 lactations were deleted for each dataset respectively, leaving 10,088

and 33,755 measurements from 99 and 421 lactations belonging to 52 and 321 unique cows. The

characteristics of the remaining profiles and related P4 data are summarized in Table 1. Here, the P4

measurements taken during the trial period for dataset 1 were not included in the calculation of the

number of measurements per profile because this was not the commercial cost-effective setting.

[TABLE 1]

In Adriaens et al., (2018a), PMASC was described for its use on time series of P4

measurements taken each milking. More specifically, a rule was introduced that when a first

evidence of luteolysis was obtained via the statistical control chart, two additional measurements

should be taken to ensure luteolysis has taken place and to avoid false alerts caused by

coincidentally low P4 values (e.g. outliers). However, in the datasets presently employed, the

number of samples taken was not dependent on PMASC's detection, but on the smoothed value

produced by the MPKF undercutting the threshold of 5 ng/mL. The latter is in its turn dependent on

how fast P4 decreases during luteolysis, which may vary from 12 to 72 hours (Blavy et al., 2016),

the absolute P4 concentrations measured and the milking interval. Accordingly, when luteolysis

went fast and the final raw P4 concentrations were considerably below 5 ng/mL, repeatedly less

than the three measurements needed to confirm luteolysis with PMASC were available. As also

discussed in Adriaens et al., (2018a), this rule is certainly adjustable, and for example the size of the

residual from the fitted mathematical model might be used in practice to decide how many samples

should confirm luteolysis. However, for the current validation, we were dependent on the samples

taken by the online device. Therefore, an adaption of PMASC was legitimized. In previous studies

on the timing between luteolysis and ovulation time (Roelofs et al., 2006; Adriaens et al., 2018c), it

was found that ovulation followed about 60 to 80 hours after the onset of luteolysis. To have the

largest chance on successful conception, insemination should take place 24 to 12 hours before

ovulation. Based on this information, it was assumed reasonable to take all the samples taken within

48 hours from the first out-of-control measurement into account to confirm or object luteolysis. In

practice, this would allow to ensure that luteolysis has taken place, while there is still sufficient time

left to arrange the inseminator. More concretely, the following rule was implemented: if a

measurement was out of control, all measurements following within a period of 48 hours were

evaluated using the current, non-updated model parameters of the increasing Hill function

(Adriaens et al., 2017). If in that period another residual was also below the current LCL, we

assumed luteolysis had taken place and an alert was raised. Accordingly, this rule gives some

flexibility to deal with missing samples and outliers. In practice, the sampling frequency would be

dependent on the PMASC itself, and more intelligent rules can be implemented.

Next, PMASC was run on the time series of P4 values of each lactation to evaluate its

performance when a reduced sampling scheme, shown cost-effective, was applied. The number of

and the timing of the alerts generated by PMASC were compared to those obtained from the

currently implemented algorithm (MPKF+T). Also, the number of measurements per detected cycle

was calculated and the profile characteristics (length postpartum anestrus, interval calving to

pregnancy and cycle properties) were summarized.

In total, the datasets consisted of 520 P4 profiles, with an average sampling rate of 0.51±0.08

measurements per day in the period between the first and the last measurement. This corresponds to

one measurement each 1.9 days. The on-farm devices gave 322 and 1664 estrus alerts respectively

(i.e. smoothed time series undercutting the fixed threshold of 5 ng/mL). From these, 309 and 1559

were also detected by PMASC, while 13 (4.0%) and 105 (6.3%) were not. These non-detected

estruses were all related to infrequent sampling, for example because the smoothed P4

concentration only slightly exceeded the 5 ng/mL threshold, after which only one sample was

sufficient to bring the smoothed value (MPKF) below 5 ng/mL and generate an alert. After this, the

smart sampling algorithm does not take a new confirmatory sample, while we set PMASC to

evaluate at least two samples per estrus, and PMASC thus failed to give an alert. This mainly occurs

during the first estrus after calving, also shown by the fact that 65% of the cases not detected by

PMASC in dataset 2 concerned the first alert of that profile, with an average alert time of 47±21

days in lactation. An example is shown in Figure 1. Here, the first luteolysis was not detected by

PMASC because of the lack of confirmatory samples out-of-control within 48 hours. The sample

for which the smoothed value is below the 5 ng/mL threshold was also out-of-control in PMASC,

but the next sample was only taken 6 days later, providing not enough evidence for PMASC to

indicate luteolysis. The model of the detected cycles is plotted in light blue, while the MPKF

smoothed data is given in dark blue. The threshold of 5 ng/mL P4 is plotted in green. As seen in the

figure, the first samples detected as out of control by PMASC is often not yet below the 5 ng/mL

threshold, and the model characteristics can be used to estimate luteolysis reliably and independent

of the milking frequency and absolute P4 values. The time lag between the drop in P4 during

luteolysis and the smoothed MPKF value undercutting the threshold is shown to be dependent of

sampling rate and the absolute P4 values (yellow box).

[FIGURE 1]

The first out-of-control sample detected by PMASC preceded the MPKF/T alarm on average

20.0±16.1 hours and 2.6±2.1 milkings for dataset 1 and 20.2±15.3 hours and 2.5±1.9 milkings for

dataset 2. The time-delay of the MPKF ensures robustness for outliers and measurement errors, and

will correct for unexpected variability. However, by combining this MPKF smoothed value with a

fixed threshold, this method strongly depends on the absolute P4 values measured and the sampling

rate. In cases where the sampling frequency is insufficiently high for timely detection of luteolysis

by the MPKF+T method, for example when the sampling algorithm fails, the alerts of the MPKF+T

are less valuable. However, when these low P4 measurements are detected by PMASC, information

calculated from the model parameters (e.g. time of the inflection point) can be used to interpret

these detections and eventually, adjust the decision process accordingly.

Using the model parameters, the P4 cycle characteristics were calculated. The average cycle

length calculated as the time between two successive luteolysis alerts was 27.3±10.4 days, with

21% of the cycles between 22 and 25 days, and 11% between 19 and 22 days. The average time

between the two inflection points of the increasing Hill function and the decreasing Gompertz

function was 14.9±10.5 days, which is slightly higher than the time reported in Adriaens et al.,

(2017), probably because here only data of normal cycles and profiles were used. The average

baseline and maximal P4 concentrations were 0.98±0.9 and 21.6±4 ng/mL respectively, and the

decrease in P4 at luteolysis was typically fast with a slope of 37.1±23.9 ng/mL per day. The latter

means that a drop in P4 from 25 to 2 ng/mL during luteolysis on average lasts 14.6 hours, which

agrees with the results reported by Blavy et al., (2016) and Adriaens et al., (2018b).

In this study, we investigated the performance of PMASC on on-farm measured datasets using

less frequent sampling and the LFAI technique, and compared it with the current state of the art

being an MPKF+T algorithm. We showed that PMASC, on the condition of a small adjustment in

its decision requirements, was able to deal with less samples and the alternative measurement

technique, and accordingly, has indeed the potential to work cost-effective in an on-farm setting.

More concretely, this study showed that a limited number of samples sufficed to fit the

mathematical model used by PMASC, and did on average not influence the cycle characteristics

extracted from it. The non-detected luteolyses were related to cases in which only one sample was

taken to confirm estrus, which is rather rare and mainly occurs during the first estrus after calving,

in which the maximum P4 concentration remains low. In practice, the sampling rate would be

optimized based on information extracted from PMASC (e.g. the time from the previous luteolysis,

out-of-control samples etc.), and these kinds of artifacts can be avoided. Although it is possible by

introducing simple decision rules in the algorithm, the implementation and optimization of the P4

sampling rate based on PMASC was considered out of the scope of this research.

ACKNOWLEDGEMENTS

This work was supported by the Institute for the Promotion of Innovation through Science

and Technology in Flanders, Belgium (IWT) [IWT-LA project 110770]. Ines Adriaens is supported

by the Fund for Scientific Research (FWO) Flanders, grant number 11ZG916N. We thank Lattec

(Hillerod, Denmark) and the commercial dairy farmer who provided us with the second P4 dataset.

REFERENCES

Adriaens, I., T. Huybrechts, K. Geerinckx, D. Daems, J. Lammertyn, B. De Ketelaere, W. Saeys,

and B. Aernouts. 2017. Mathematical characterization of the milk progesterone profile as a leg

up to individualised monitoring of reproduction status in dairy cows. Theriogenology 103:44–

51. doi:https://doi.org/10.1016/j.theriogenology.2017.07.040.

Adriaens, I., W. Saeys, T. Huybrechts, C. Lamberigts, L. François, K. Geerinckx, J. Leroy, B. De

Ketelaere, and B. Aernouts. 2018a. A novel system for on-farm fertility monitoring based on

milk progesterone.

Adriaens, I., W. Saeys, T. Huybrechts, C. Lamberigts, K. Geerinckx, J. Leroy, B. De Ketelaere, B.

Aernouts, and K.U. Leuven. 2018b. A novel system for online fertility monitoring based on

milk progesterone. J. Dairy Sci. 1–35. doi:10.3168/jds.2017-13827.

Adriaens, I., W. Saeys, C. Lamberigts, M. Berth, K. Geerinckx, J. Leroy, B. De Ketelaere, and B.

Aernouts. 2018c. Short communication: Sensitivity of estrus alerts and their relation with the

timing of the LH surge.

Blavy, P., M. Derks, O. Martin, J.K. Hoglund, and N.C. Friggens. 2016. Overview of progesterone

profiles   in   dairy   cows.   Theriogenology   86:1061–1071.

199 doi:https://doi.org/10.1016/j.theriogenology.2016.03.037.

200 Blom, J.Y., and C. Ridder. 2010. Reproductive management and performance can be improved by

201  use of Delaval Herd Navigator. Page in Proceedings of First North American Conference on

202  Precision Dairy Management.

203 Friggens, N.C., M. Bjerring, C. Ridder, S. Højsgaard, and T. Larsen. 2008. Improved detection of

204  reproductive status in dairy cows using milk progesterone measurements. Reprod. Domest.

205  Anim. 43:113–121.

206 Friggens, N.C., and M.G.G. Chagunda. 2005. Prediction of the reproductive status of cattle on the

207  basis of milk progesterone measures: model description. Theriogenology 64:155–90.

208  doi:10.1016/j.theriogenology.2004.11.014.

209 Friggens, N.C., K.L. Ingvartsen, I.R. Korsgaard, T. Larsen, P. Løvendahl, C. Ridder, and N.I.

210  Nielsen. 2007. System and a method for observing and predicting a physiological state of an

211  animal. US Pat. No. US7302349 B2.

212 Roelofs, J.B., F.J.C.M. Van Eerdenburg, W. Hazeleger, N.M. Soede, and B. Kemp. 2006. Short

213  communication: Relationship between progesterone concentrations in milk and blood and time

214  of ovulation in dairy cattle. Anim. Reprod. Sci. 91:337–343.

215  doi:doi:10.1016/j.anireprosci.2005.04.015.

Figure 1. An example profiles of real progesterone (P4) data indicated in black. The light blue

crosses are the first out-of-control measurements detected by PMASC, and the light blue lines give

final models fitted by PMASC. The first cycle was not detected because an insufficiently high

sampling rate, although the measurements pushing the MPKF (indicated in blue) below the

threshold (indicated in green) were also out of control for PMASC (indicated with purple cross).

The (inconsistent) time-lag between the MPKF dropping below the threshold and actual low

measurements of P4 can be seen in the yellow box.

Table 1. Summary of the datasets of farmer 1 and farmer 2

232   Adriaens, Table 1

Dataset	No. of cows	No. of profiles	MY[1] (kg)	MI[2] (h)	P4[3]		Estrus alerts
					Concentration (ng/mL)	No. of meas. per profile	
Dataset 1	52	99	20.5-58.2 (39.0±9.2)	5.6-11.8 (8.4±1.3)	0.1-28.0 (14.5±10.4)	24 -325 (137.1±56.8)[4]	322
Dataset 2	321	421	10.1-66.4 (32.0±11.7)	5.0-29.05 (9.36±3.19)	0.01-27.99 (15.36±9.69)	21-232 (80.2±35.39)	1656

233   [1] Daily milk yield in kg; [2] Average milking interval in hours; [3]P4 milk progesterone; [4] excluding the measurements taken during the
234   research trial in spring 2017 as described in Adriaens et al., (2018a)