Interspecies metabolite transfer and aggregate formation in a co-culture of

***Dehalococcoides* and *Sulfurospirillum* dehalogenating tetrachloroethene to**

ethene

Stefan Kruse[1,§], Dominique Türkowsky[2], Jan Birkigt[3], Bruna Matturro[4], Steffi Franke[3,&], Nico

Jehmlich[2], Martin von Bergen[2,6], Martin Westermann[5], Simona Rossetti[4], Ivonne Nijenhuis[3,8],

Lorenz Adrian[3,7], Gabriele Diekert[1], Tobias Goris[1,$,#]

[1]Department of Applied and Ecological Microbiology, Institute of Microbiology, Friedrich Schiller

University, Jena, Germany

[2]Department Molecular Systems Biology, Helmholtz Centre for Environmental Research – UFZ,

Leipzig, Germany

[3]Department of Isotope Biogeochemistry, Helmholtz Centre for Environmental Research – UFZ,

Leipzig, Germany

[4]Water Research Institute, IRSA-CNR, Monterotondo, Rome, Italy

[5]Center for Electron Microscopy of the University Hospital Jena, Jena, Germany

[6]Institute of Biochemistry, Faculty of Life Sciences, University of Leipzig, Leipzig, Germany

[7]Chair of Geobiotechnology, Technische Universität Berlin, Germany

[$]Current affiliation: German Institute of Human Nutrition, Department Molecular Toxicology, Research

Group Intestinal Microbiology, Potsdam-Rehbrücke

[§]Current affiliation: Novartis Pharma GmbH, Trial Monitoring Organization

[&]Current affiliation: Eurofins Institute Dr. Appelt Leipzig

[#]Corresponding author: Tobias Goris, tobias.goris@uni-jena.de

Abstract

Microbial communities involving dehalogenating bacteria assist in bioremediation of areas contaminated with halocarbons. To understand molecular interactions between dehalogenating bacteria, we co-cultured *Sulfurospirillum multivorans*, dechlorinating tetrachloroethene (PCE) to *cis*-1,2-dichloroethene (*c*DCE), and *Dehalococcoides mccartyi* strains BTF08 or 195, dehalogenating PCE to ethene. The co-cultures were cultivated with lactate as electron donor. In this co-culture, the bacterial cells formed aggregates and *D. mccartyi* established an unusual, barrel-like morphology. An extracellular matrix surrounding bacterial cells in the aggregates enhanced cell-to-cell contact. PCE was dehalogenated to ethene at least three times faster in the co-culture. The dehalogenation was carried out via PceA of *S. multivorans*, and PteA (a recently described PCE dehalogenase) and VcrA of *D. mccartyi* BTF08, as supported by protein abundance. The co-culture was not dependent on exogenous hydrogen and acetate, suggesting a syntrophic relationship in which the obligate hydrogen consumer *D. mccartyi* consumes hydrogen and acetate produced by *S. multivorans*. The cobamide cofactor of the reductive dehalogenase - mandatory for *D. mccartyi* - was also produced by *S. multivorans*. *D. mccartyi* strain 195 dechlorinated *c*DCE in the presence of norpseudo-B$_{12}$ produced by *S. multivorans*, but *D. mccartyi* strain BTF08 depended on an exogenous lower cobamide ligand. This observation is important for bioremediation, since cofactor supply in the environment might be a limiting factor for PCE dehalogenation to ethene, described for *D. mccartyi* exclusively. The findings from this co-culture give new insights into aggregate formation and the physiology of *D. mccartyi* within a bacterial community.

Introduction

Microbial communities interact in numerous ways, involving the transfer and consumption of metabolic products. Molecular hydrogen, for example, is an important electron carrier in syntrophic communities, in which hydrogen is produced by e.g. fermentation and taken up by hydrogen-consuming prokaryotes. This hydrogen consumption leads to a lower hydrogen partial pressure, which allows otherwise thermodynamically unfavorable reactions to proceed. Therefore, the involved bacteria are physiologically dependent on each other [1, 2, 3]. The probably carcinogenic groundwater pollutant tetrachloroethene (PCE), a remnant of e.g. bleaching, dry-cleaning and metal degreasing, is

completely, but often slowly, dechlorinated to ethene in communities involving hydrogen transfer from

fermenting bacteria to the obligately hydrogen-consuming *D. mccartyi* [4, 5, 6, 7]. PCE and other

organohalides such as hexachlorobenzene or polychlorinated biphenyls are used as terminal electron

acceptors by *D. mccartyi* and other bacteria in an anaerobic respiratory chain coupling the

dehalogenation to energy conservation via electron transport phosphorylation [6]. This process is

termed organohalide respiration (OHR) and involves the corrinoid-containing reductive dehalogenases

(RDases) as terminal reductases [8, 9, 10]. However, *D. mccartyi* is characterized by low growth rates

and dehalogenates PCE slower [11] compared to other dehalogenating species such as

Sulfurospirillum multivorans. *D. mccartyi* is strictly dependent on several specific vitamins, most

importantly cobamides [12]. Besides being restricted to hydrogen as electron donor, *D. mccartyi* uses

only acetate plus bicarbonate as carbon source and organohalides as electron acceptors. In addition,

these bacteria are not able to *de novo* synthesize corrinoids, the obligate cofactor of RDases [11, 13].

While proteins for complete corrinoid biosynthesis are usually not encoded in the genomes of *D.*

mccartyi [14], different studies revealed its ability to salvage and remodel corrinoids, enabling *D.*

mccartyi to restore dechlorination [15, 16, 17]. The functionality of the corrinoid and thus the RDase is

directly dependent on the type of the lower base in *D. mccartyi*. Only three types of corrinoids have so

far been described to be functional in *D. mccartyi* strain 195: 5,6-dimethylbenzimidazolyl-cobamide

([DMB]-Cba), 5-methylbenzimidazolyl-cobamide and 5-methoxybenzimidazolyl-cobamide ([5-

OMeBza]Cba). Nonfunctional corrinoids e.g. 5-hydroxybenzimidazolyl-cobamide or 7-adeninyl-

cobamide ([Ade]Cba) can be converted to functional ones by replacement of the lower ligand in *D.*

mccartyi when 5,6-dimethylbenzimidazole (DMB) is provided [16, 17].

Several studies on dechlorinating communities containing *D. mccartyi* in association with fermenting,

acetogenic and/or methanogenic bacteria revealed higher dechlorination and growth rates than those

of pure cultures [16, 18, 19, 20, 21, 22, 23, 24, 25, 26]. It is hypothesized that cross-feeding and a

constant supply of growth factors such as corrinoids and biotin enhance growth [27]. Within these

communities, non-dechlorinating fermenting bacteria provide hydrogen, acetate and CO_2 from e.g.

lactate or butyrate fermentation. The fermenting bacteria are dependent on hydrogen consumers

which keep the hydrogen partial pressure low [18, 21, 22, 28]]. For example, co-culture experiments

revealed an interspecies hydrogen transfer between *Desulfovibrio desulfuricans* fermenting lactate

and *D. mccartyi* dechlorinating TCE [15]. Acetogens like *Sporomusa ovata* and sulfidogens

(*Desulfovibrio*), produce different types of corrinoids which can be used by *D. mccartyi* [29, 30]. An

interspecies cobamide transfer was also shown between *Methanosarcina barkeri* strain Fusaro and *D.*

mccartyi strain BAV1, GT and FL2, when DMB was present [25, 26]. In the only co-culture of *D.*

mccartyi (strains BAV1 and FL2) with a PCE-dechlorinating bacterium, *Geobacter lovleyi*, a corrinoid

transfer was also observed [31]. In this co-culture, however, hydrogen had to be supplemented. All

these studies showed more robust growth of *D. mccartyi* in co-cultures, resulting in higher

dechlorination rates and cell yields than in pure cultures and sometimes, depending on the cobamide

metabolism of the co-culture, a different degree of the dechlorination extent [32]. However, a PCE-

dechlorinating bacterium providing hydrogen, acetate and corrinoids to *D. mccartyi*, to our knowledge,

has not been described in the literature. Thus, further information on *D. mccartyi* co-cultures is needed

to gain insights into complex microbial communities involving *D. mccartyi*. Relatively simple co-

cultures provide a model community in which metabolic transfers are easier to follow. Recently, the

PCE to *c*DCE-respiring *Sulfurospirillum multivorans*, capable of *de novo* corrinoid production [33, 34,

35], was shown to produce hydrogen and acetate under fermentative growth conditions, e.g. with

pyruvate or lactate and without electron acceptor present [36]. In the same study, a co-culture of *S.*

multivorans with a methanogen dependent on hydrogen as electron donor was established. Therefore,

S. multivorans is a promising partner for a co-cultivation with *D. mccartyi*. In addition, it is of interest,

whether *D. mccartyi* is able to take up and utilize norpseudo-B_{12} ([Ade]NCba), so far known to be

produced exclusively by *S. multivorans* and *S. halorespirans* [35, 37].In this study, we investigated the

physiological interaction between *D. mccartyi* strain BTF08 and 195 in co-culture with *S. multivorans*

dechlorinating PCE to ethene by using dechlorination analytics, microscopy and proteomics.

Materials and Methods

Cultivation of pure cultures

D. mccartyi cultures BTF08 and 195 (maintained at the UFZ Leipzig) were cultivated in 200 mL serum

bottles containing 100 mL bicarbonate-buffered mineral salt medium with 5 mM acetate and 148 nM

vitamin B_{12} (cyanocobalamin, ca. 200 µg/L), reduced with $Na_2S \cdot 7 H_2O$ (final concentration, 0.25 g/L)

[38]. Anoxic atmosphere was established by 30 cycles of gassing and degassing with nitrogen, and

CO_2 was added to a final atmosphere of $N_2:CO_2$ (75:25 v/v 150 kPa). After autoclaving the medium

and adjusting the pressure to approximately 100 kPa with nitrogen and CO_2, we applied, in addition,

hydrogen [50 kPa] to avoid possible electron donor limitation. PCE (>99% purity, Sigma Aldrich,

Steinheim, Germany) and *c*DCE (97% purity, Sigma Aldrich, Steinheim, Germany) served as electron

acceptors and were added with a microliter syringe (Hamilton, Bonaduz, Switzerland) to a final

concentration of 0.35 mM (aqueous phase concentration). Re-feeding of the cultures was done with

the same amount of PCE or *c*DCE after complete conversion to ethene. After maximally three re-

feeding steps, cultures were transferred [10% (v/v)] into fresh medium. To evaluate the effect of

different types and concentrations of B_{12} on dechlorination activities, *D. mccartyi* cultures received 54

nM norpseudo-B_{12} ([Ade]Cba) or 5-OMeBza-B_{12} ([5-OMeBza]Cba), each. Norpseudo-B_{12} was

extracted as described previously [39] from 6 L of *S. multivorans* grown anaerobically with 40 mM

pyruvate and 10 mM PCE as described elsewhere [40]. In brief, crude cell extract was acidified to a

pH of ≤ 5 with acetic acid before adding 0.1 M potassium cyanide and boiling the sample for 15 min.

Afterwards, the sample was centrifuged and the supernatant kept on ice. The cell pellet was

reuspendend in 10 mL UPW and the same extraction procedure was applied twice. The combined

supernatant was mixed with Amberlite XAD4 (0.25 g XAD4 per ml extract; Sigma-Aldrich) and

incubated on a shaker overnight. The supernatant was removed and the cobamide-loaded XAD4

material was washed with 10 volumes of UPW. Elution of cobamide was done for 1 h on a shaker with

1 volume of methanol which was repeated twice. A vacuum concentrator was used to completely dry

the eluates which were then resuspended in 2 mL UPW. The sample was applied on a column

containing 3 g aluminium oxide, recovered with 50 mL UPW and dried. Resuspension of purified

Norpseudo-B12 was done with UPW. 5-OMeBza-B_{12} was obtained in the same way from 6L of

Desulfitobacterium hafniense DCB2 grown anaerobically with 40 mM pyruvate and 10 mM ClOHPA (3-

chloro-4-hydroxy-phenylacetate) on a medium amended with 25 µM 5-OMeBza [41]. *S. multivorans*

(DSMZ 12446) was cultured in the same mineral salt medium as *D. mccartyi* [38] with 40 mM lactate

and 10mM nominal PCE in hexadecane (2 mL PCE/hexadecane per 100 mL medium) but without the

addition of acetate and hydrogen [38]. All cultivation experiments were performed statically at 28°C in

the dark and in biological triplicates.

Co-cultivation of *S. multivorans* and *D. mccartyi*

Co-cultures of *S. multivorans* and *D. mccartyi* strain BTF08 or 195 were maintained in the same mineral salt medium as the pure cultures, without acetate and hydrogen. Initially, the inoculum of the pure cultures into the co-culture was 10 % of each pure culture. The medium contained 25 mM lactate (D/L, > 85% purity, Sigma-Aldrich, Steinheim, Germany) as the electron donor for *S. multivorans* and 0.35 mM PCE (aqueous-phase concentration) as the electron acceptor. Lactate (25 mM) and PCE (0.35 mM) was re-fed after depletion of substrates was measured. The B_{12}-dependance of the co-cultures was tested with 148 nM vitamin B_{12} (>98% purity, Sigma-Aldrich, Steinheim, Germany) serving as the positive control and without vitamin B_{12} amendments. Cultures without vitamin B_{12} received 1 µM DMB (>99% purity, Sigma-Aldrich, Steinheim, Germany), where indicated. The DMB dose used had only minimal negative effect on *S. multivorans* growth [39]. For isotope fractionation experiments, *S. multivorans* and *Sm*/BTF08 co-cultures were cultivated in 50 mL serum bottles with 25 mL medium. Replicate bottles (triplicates for each time point) were inoculated at the same time and the dehalogenation process was stopped at different time points by addition of 3 mL 2 M Na_2SO_4 (pH 1.0).

Quantitative (q)PCR analysis of cell growth

Quantitative PCR (qPCR) was applied to count *Sulfurospirillum* and *Dehalococcoides* 16S rRNA gene copies. DNA was extracted from 1 mL co-culture taken from different time points during the cultivation experiment using the NucleoSpin Tissue DNA extraction kit according to the manufacturer's instructions (Macherey-Nagel, Düren, Germany). Cell numbers were quantified by direct epifluorescence microscopy of Sybr-green stained cells immobilized on agarose-coated slides to generate standard curves for *q*PCR as described previously [42]. For the corresponding standard curves, *D. mccartyi* and *S. multivorans* cell suspensions with known cell numbers ($1x10^4$ to $1x10^8$ and $1x10^5$ to $1x10^9$ respectively) were subjected to qPCR. The qPCR reaction mixture contained 1 µl of gDNA or DNA standard, 6.25 µl 1x KAPA SYBR Fast master mix (Sigma Aldrich, Steinheim, Germany) and 0.208 µM forward and reverse primer. Primers used were Dhc_sp_16S_fw (5'-GTATCGACCCTCTCTGTGCCG-3') and Dhc_sp_16S_rev (5'-GCAAGTTCCTGACTTAACAGGTCGT-3') for *D. mccartyi* and Smul_16S_fw (5'-AGGCTAGTTTACTAGAACTTAGAG-3') and Smul_16S_rev (5'-CAGTCTGATTAGAGTGCTCAG-3') for *S. multivorans*. The conditions of the PCR program were

as followed: 95°C for 2 min (initial denaturation) followed by 40 cycles of 55°C (*S. multivorans* primer)

or 60°C (*D. mccartyi* primer) for 20s (annealing), 72°C for 30s (elongation) and 95°C for 10 s

(denaturation). Each *q*PCR program included a melting curve for verification of specific target DNA

amplification and primer efficiency was tested for used primers (R^2 *Sm*: 99.7%; R^2 *Dhc*: 98.8%, see

Supplementary Figure S1). The cell numbers of samples were calculated by comparing the sample C_T

values with the C_T values of the standard curve obtained from microscopical cell counting

(Supplementary Figure S1). All samples were conducted in three biological replicates with two

corresponding technical replicates and three technical replicates for the calibration curve.

Analytical methods

Ethene and chlorinated ethenes were quantified gas-chromatographically with a flame ionization

detector (Clarus 500, Perkin Elmer, Rodgau, Germany) and a CP-PoraBOND Q FUSED SILICA 25 m

x 0.32 mm column (Agilent Technologies, Böblingen, Germany). Samples from the cultures (1 mL of

the gas phase and 1 mL of the liquid phase) were transferred to GC vials with gast tight syringes, the

sampling proceeded via a headspace sampler (HS 40, Perkin Elmer; needle temperature 100°C, vial

pressurized with 0.1 kPa nitrogen for 2 mins). The GC vial with the liquid phase was heated in the HS

40 to 80°C for 5 min. Samples were transferred through the transfer line (100°C) into the GC. Ethene

and chlorinated ethenes were separated with 4 min at 60°C rising to 280°C in 10°C min^{-1} steps. The

injector temperature was fixed at 250°C and the detector temperature at 300°C. Standard curves of

ethene and each chlorinated ethene were recorded for peak area quantification and retention times

were compared to known standards. Nonane served as an internal standard during measurements.

The retention times were: PCE (detection limit [dl] 2.1 µM), 15.1 min; TCE (dl 2.5 µM) 13.2 min; *c*DCE

(dl 3.1 µM), 11.3 min; VC (dl 3 µM), 6.0 min; ethene (dl 2.5 µM), 1.3 min. Due to the low solubility of

PCE, it was added to the medium followed by stirring for at least 8 hours to allow it to completely

dissolve. This lead to a reliable quantification of only the initial PCE concentration, but not of the re-fed

PCE, especially since a portion of the re-fed PCE is rapidly degraded by the now higher abundant *S.*

multivorans cells. The concentrations of chlorinated ethenes (in µM or mM) were calculated by

summing up the concentrations in the gas phase and liquid phase. Dechlorination rates were

calculated according to the average amount of stoichometrically converted PCE to the corresponding

end products VC (in the case of *D. mccartyi* strain 195, where the formation of co-metabolically

produced ethene was negligible) or ethene (for *D. mccartyi* strain BTF08) divided by time in days. For

example the conversion of 400 µM PCE into 400 µM ethene over 10 days resulted in a dechlorination

rate of 40 µM PCE/day. Hydrogen (dl 0.8 µM) was measured gas-chromatographically using a thermal

conductivity detector (AutoSystem, Perkin Elmer, Rodgau, Germany) equipped with a Carboxen-1010

PLOT capillary column 30 m x 0.53 mm (Supelco, München, Germany) column. Separation of

analytes was done at a temperature of 35°C (detector at 150°C). For this, the culture headspace (1

mL) was sampled with gas-tight syringes (Hamilton, Bonaduz, Switzerland) and injected directly into

the injection valve of the GC. Concentrations were calculated using calibration curves ranging from 4 –

50 µmol hydrogen. Organic acids (lactate, acetate, pyruvate and succinate) were analyzed using a

reversed-phased HPLC (Merck-Hitachi, Darmstadt, Germany) equipped with a diode array detector (L-

7450) and separated on an AMINEX HPX-87H column (7.8 x 300 mm; BioRad, Munich, Germany)

with a cation H guard pre-column. Samples (1 mL) from the liquid phase of the cultures were filtered

(MiniSart RC4, 0.2 µm, Sartorius, Göttingen, Germany) and acidified with 2.5 µL H_2SO_4. Of this

sample, 20 µL was injected into the HPLC. Monitoring of organic acids was based on their absorption

at 210 nm. The column was operated at a temperature of 50°C and 5 mM (v/v) H_2SO_4 served as

mobile phase with a flow rate of 0.7 mL min^{-1}. The retention times were: lactate (dl 0.2 mM), 10.61

min; pyruvate (dl 0.01 mM) 7.92 min; acetate (dl 0.4 mM), 12.88 min; succinate (dl 0.15 mM), 9.81

min. Peak areas and retention times were related to standards ranging from 1 to 10 mM or 0.5 to 2

mM (for pyruvate).

Compound-specific stable isotope analysis

Determination of the carbon isotope composition of the chlorinated ethenes in pure culture of *S.*

multivorans and in co-culture of *S. multivorans* and *D. mccartyi* strain BTF08 was done using gas

chromatography combustion isotope ratio mass spectrometry (GC-C-IRMS; Thermo GC Trace 1320

combined with Thermo-Finnigan MAT 253 IRMS, Bremen, Germany) [43]. All samples were analyzed

in technical triplicates. 2 mL liquid phase were taken from the respective sample and transferred to a

He-flushed 10 mL crimped vial. Of these, 1 mL was taken from the headspace via an autosampler

(Thermo TriPlus RSH Autosampler, Bremen, Germany) and injected in a gas chromatograph with a

split ratio of 1:5. Chlorinated ethenes were separated on a DB-MTBE column (60 m x 0.32 mm x 1.8

µm, J&W Scientific, Waldbronn, Germany) and the following temperature program: 40°C for 5 min,

increase to 250°C by 20°C min^{-1} and hold for 5 min using helium as carrier gas at a flow rate of 2.0 mL

min^{-1} (Injector at 250°C).

The carbon isotope composition is given in the δ-notation (‰) relative to the Vienna Pee Dee

Belemnite standard [44]. Carbon isotope fractionation was calculated using the Rayleigh equation (eq

1) where R_0 and R_t represent the isotope values and C_0 and C_t the concentrations at time 0 and t [45,

46].

232
$$ln \frac{Rt}{R0} = \varepsilon C * ln \frac{Ct}{C0} \qquad (1)$$

The carbon isotope enrichment factor (ε_C) relates changes in the concentration of the isotopes to

changes in their isotope composition. A two-tailed T-Test was used to calculate the 95 % confidence

interval based on the slope. Standard deviations were obtained from at least triplicate measurements

(< 0.5 ‰).

B_{12} extraction and MS analysis

The B_{12} content of three independently cultivated 100 mL cultures was analyzed. For this, the culture

volume was reduced to 20 mL using a vacuum concentrator and 0.1 M potassium cyanide was added.

After boiling the samples for 20 min, cell debris was removed by centrifugation (10 min, 6700 x g,

8°C). The supernatant was applied onto a C-18 column (CHROMABOND C-18 ec, Macherey-Nagel,

Düren, Germany) equilibrated with 5 mL 100% (v/v) methanol and 5 mL ultrapure water (UPW).

Washing of the column was done twice with 5 mL UPW and B_{12}-types were eluted with 5 mL 100%

(v/v) methanol. The eluate was completely dried in a vacuum dryer and dissolved in UPW prior to

analysis. The extract was injected to ultrahigh performance nano-flow liquid chromatography (UHPLC)

(Ultimate 3000, Thermo Fisher, Waltham, USA) coupled to mass spectrometer (LC-MS, Orbitrap

Fusion, Thermo Fisher, Waltham, USA) via heated electrospray ionization (HESI-II, Thermo Fisher,

Waltham, USA). The UHPLC was equipped with a Hypersil Gold C18 column (150 x 2.1 mm, 3 µm

film thickness; Thermo Fisher, Waltham, USA) and a C18 guard column (10 x 2.1 mm, Waters,

Milford, USA). Chromatographic separation using a gradient method with 0.1% formic acid (A) and

methanol (B) as mobile phase was applied as following: 5% B for 1 min, 60 min gradient to 90% B, 4

min at 90% B, 1 min gradient to 5% B and 4 min at 5% B with a constant flow of 0.2 mL min^{-1} and

25°C column oven temperature. Injection volume was 5 µL. Ionization was set to positive ion mode at

3.5 kV, 35 arbitrary units (Arb) sheath gas, 10 Arb auxillary gas, 325°C ion transfer tube temperature

and 275°C vaporizer temperature. Orbitrap resolution for precursor scan (MS1) was set to 120.000

with a scan range of 300-1600 *m/z*. Data evaluation was done on the original mass spectra comparing

with predicted masses of corrinoids (Supplementary Tables S1 and S2, for cobalamin standards see

Supplementary Figure S2).

Scanning Electron Microscopy and fluorescence in situ hybridization

Field emission-scanning electron microscopy (FE-SEM) was performed with co-cultures of *S.*

multivorans and *D. mccartyi* strains BTF08 and 195 and pure *S. multivorans* and *D. mccartyi* strain

BTF08 cultures. All of the cultures were treated the same way for FE-SEM as follows. Cells of 5 mL

culture were incubated for 15 min with 2.5% glutaraldehyde and pre-fixed for 2 h on poly-L-lysine

coated coverslides (12 mm, Fisher Scientific, Schwerte, Germany). Cover slides were washed three

times with 0.1 M sodium cacodylate (pH 7.2) (>98% purity, Sigma Aldrich, Steinheim, Germany) and

post-fixed for 1 h with 1% osmium tetroxyde in the same cacodylate buffer. After fixation, samples

were dehydrated using different ethanol concentrations (35%/50%/70%/80%/95%/100% v/v) and

incubated for 10 min at each step. Critical point drying was done in a Leica EM CPD200 Automated

Critical Point Dryer (Leica, Wetzlar, Germany), followed by coating with 6 nm platinum in a BAL-TEC

MED 020 Sputter Coating System (BAL-TEC, Balzers, Liechtenstein). Imaging of the samples was

done with a Zeiss-LEO 1530 Gemini field emission scanning electron microscope (Carl Zeiss,

Oberkochen, Germany) at different magnifications and at 10 kV acceleration voltage. Fluorescence in

situ hybridization (FISH) was performed as described previously [47, 48]. In brief, samples were fixed

with formaldehyde (2% v/v final concentration) for 2 h at 4°C and filtered on polycarbonate membrane

filters (47 mm diameter, 0.2 µm pore size, Nucleopore). Filters were stored at -20°C until further

processing. FISH detection of *D. mccartyi* strain BTF08 and 195 was done with Cy3-labeled

DHC1259t and DHC1259c probes and *S. multivorans* detection with FITC-labeled probes SULF

F220ab [49]. Imaging of un-aggregated cells was done with an epifluorescence microscope (Olympus,

BX51) combined with an Olympus XM10 camera. Images were analyzed via Cell-F software.

Aggregates were visualized using a confocal laser scanning microscopy (CSLM, Olympus FV1000).

Protein extraction and proteome analysis

Samples for proteome analyses were taken approximately two weeks after one re-feeding with PCE or

*c*DCE (pure *D. mccartyi* strain BTF08 cultures) or 12 hours after the third re-feeding with PCE (co-

cultures *Sm*/BTF08, Supplementary Figure S3). When harvested, PCE, TCE and *c*DCE were the

dominant chlorinated ethenes in the pure cultures, while *c*DCE and VC were the dominant chlorinated

ethenes in the co-cultures (Supplementary Figure S3). Samples were processed as described

previously [50]. Briefly, protein extraction was performed in lysis buffer (20 mM HEPES, 1 mM sodium

vanadate, 1 mM β-glycerol phosphate, 2.5 mM sodium pyrophosphate, 8 M urea [51]) by three

freeze/thaw-cycles and ultrasonic bath treatments. Ten µg protein dissolved in 100 µL ultra pure water

was precipitated with the five-fold amount (v/v) of ice-cold (-20°C) acetone. Protein pellets were

dissolved in 50 µL SDS sample buffer (2% SDS, 2 mM beta-mercaptoethanol, 4% glycerol, 40 mM

Tris-HCl pH 6.8, 0.01% bromophenolblue), heated to 90 °C for 5 min and separated on a SDS gel

(12.5%). The gel was run until the samples entered the separating gel. Afterwards, a 3-5 mm protein

band from each sample was cut out, destained, dehydrated, reduced with 10 mM dithiothreitol,

alkylated with 100 mM iodoacetamide and proteolytically cleaved over night at 37°C using trypsin

(Promega, Madison, WI, USA). Peptides were extracted, desalted using C18 ZipTip columns (Merck

Millipore, Darmstadt, Germany) and resuspended in 0.1% (v/v) formic acid before LC–MS analysis.

Proteolytic lysates were separated using an Ultimate 3000 RSLCnano liquid chromatographic

instrument (Thermo Scientific, Germany). Mass spectrometry was performed on an Orbitrap Fusion

mass spectrometer (Thermo Scientific, San Jose, CA, USA) coupled to a TriVersa NanoMate (Advion,

Ltd., Harlow, UK). Samples of 5 µL were loaded onto a trapping column (Acclaim PepMap100 C18,

75 µm × 2 cm, Thermo Scientific) using 96% eluent A (0.1% formic acid) and 4% eluent B (0.08%

formic acid, 80% acetonitrile) at a flow rate of 5 µL min^{-1} and separated via a 25 cm analytical column

(Acclaim PepMap100 C18, 75 µm × 25 cm, Thermo Scientific) at 35°C using a constant flow rate of

300 nL/min. Peptide separation was achieved by applying a linear gradient of eluent B from 4% to

50% within 100 min. Full MS scans were measured in the Orbitrap mass analyzer within the mass

range of 400-1,600 *m/z* at 120,000 resolution using an automatic gain control (AGC) target of $4x10^5$

and maximum fill time of 60 ms. The MS instrument measured in data-dependent acquisition (DDA)

mode using the highest intense ion (top speed, 3 sec cycle time). Positive ion charge states between 2

and 7 were selected for MS/MS. Precursor masses for MS/MS were selected based on the highest

intensity and excluded from further MS/MS for 30 s to prevent redundancy in MS/MS acquisition. After

higher energy collisional (HCD) fragmentation at normalized collision induced energy of 30%, fragment

masses were scanned in the Orbitrap mass analyzer at a resolution of 15,000 with $5x10^4$ AGC target

and a maximum injection time of 150 ms.

LC-MS/MS data were analyzed using Proteome Discoverer (v2.1, Thermo Scientific). MS/MS spectra

were searched against a combination of a *S. multivorans* (3,233 non-redundant protein-coding

sequences, downloaded January 2017 from NCBI GenBank, accession number CP007201.1) and a

D. mccartyi strain BTF08 database (1,535 non-redundant protein-coding sequences, downloaded

December 2016 from NCBI, accession number CP004080.1). A "common repository of adventitious

proteins database" (cRAP) was integrated to ensure correct protein identifications. The SEQUEST HT

algorithm was used with the following settings: trypsin as cleavage enzyme, oxidation on methionine

as dynamic and carbamidomethylation on cysteine as static modification, up to two missed cleavages,

precursor mass tolerance set to 10 ppm and fragment mass tolerance to 0.02 Da, respectively. Only

peptides with a false discovery rate (FDR) 1%, XCorr ≥2, q-value and the posterior error probability

(PEP) ≤0.01 were considered as identified (Supplementary Excel File S1). Quantification of proteins

was performed using the average of the top three peptide MS1 areas. After log10 transformation, the

protein values were normalized to the median of all proteins of a sample, to the median of all proteins

of the respective organism in that sample and the median of all samples and scaled so that the global

minimum is zero. Two outliers of the co-culture proteome replicates with only 11 (C3) and 92 (P1, see

Supplementary Excel File S1) protein identifications, compared to at least 317 protein identifications

from the other *D. mccartyi* pure cultures, were excluded from the analysis. Proteins with only one out

of three possible quantitative values per sample were considered as identified only. P-values were

calculated using a two-tailed, homoscedastic student's t-test and multiple corrected with the

Benjamini−Hochberg method. Figures were created using an in-house written R-script with the

packages gplots, ggplot2, ggbiplot, dplyr, miscTools and vegan. The non-parametrical multiple

dimensional scaling (nMDS)-analysis was performed with the anosim-function of the vegan package in

R (v3.4.1). *Pairwise Indicator Species Analysis* was used to identify proteins that were significantly

associated with the different cultivation conditions [52, 53]. Indval scores and significances were

calculated according to Malik et al. [54].

Results

1. Growth and dechlorination in *Sulfurospirillum* (*Sm*)/*Dehalococcoides* (*Dhc*) co-cultures

The growth characteristics and PCE to *c*DCE dechlorination pattern of *S. multivorans* cultivated in the co-culture medium (Supplementary Figure S4) was comparable to those in the routinely used medium [55]. With lactate as electron donor and PCE as electron acceptor, a co-culture of *S. multivorans* and *D. mccartyi* BTF08 maintained over 10 transfers under the same conditions dechlorinated PCE to stoichiometric amounts of ethene within eight days after inoculation (Figure 1A). The dechlorination rate (PCE to ethene) of the *Sm*/BTF08 co-culture was 4.5-fold higher compared to the *D. mccartyi* strain BTF08 pure culture (Table 1). In the latter, PCE was completely reduced to ethene within 35 days at a rate of 9 ± 0.3 µM/day (Supplementary Figure S5A). The culture was re-fed with PCE at day 8 and day 12 after inoculation. After re-feeding, the dechlorination increased (Figure 1A) and complete dechlorination of PCE occurred within three days. PCE was dechlorinated to *c*DCE within two days. To check whether *S. multivorans*, known for its high dehalogenation rate [55], was responsible for this fast dechlorination, the stable carbon isotope fractionation patterns (which largely differ between *D. mccartyi* and *S. multivorans*) of pure and co-cultures were compared. The isotope fractionation pattern of PCE in the co-culture did not change significantly, and ranged from -30.2 ± 0.09 to -29.5 ± 0.12 ‰ at 70% of transformed PCE (Supplementary Figure S6A). A low but significant fractionation was measured in the pure culture of *S. multivorans* (from -29.2 ± 0.05 to -27.6 ± 0.11 ‰ at 56% of transformed PCE, Supplementary Figure S6A). The isotope enrichment factors for the *S. multivorans* pure and the co-culture were in the same range while differing largely from that of a *D. mccartyi* strain BTF08 pure culture (*Sm*/BTF08 co-culture: ε_C = -0.4 ± 0.3 ‰; *S. multivorans* pure culture: ε_C = -2.0 ± 0.4 ‰, Supplementary Figure S6B; *D. mccartyi* strain BTF08: ε_C = -5.0 to -9.0 ‰ [56]). This high similarity in the PCE isotope fractionation patterns of *S. multivorans* pure and co-cultures is likely due to PCE to cDCE dechlorination performed nearly exclusively by *S. multivorans*. The fast dechlorination of PCE to *c*DCE in less than two days was also reflected in the increase of the *S. multivorans* cell number (at least one cell doubling per day), until PCE was completely dechlorinated to *c*DCE (Figure 1C). After this respiratory growth, a weaker growth (less than one cell doubling in 20 days) was observed, probably correlating to lactate fermentation of *S. multivorans* (Figures 1E, 1F). *D. mccartyi* strain BTF08 needed seven days for one cell doubling in the co-culture. For both organisms, a

correlation between dechlorination and growth was observed. The ratio between *S. multivorans* and *D.*

mccartyi strain BTF08 cells changed from initially 2.6 to 5.9 after 2 days and 4.6 after 15 days and two

re-feeding steps. After 12 days, when the second dose of PCE was completely dechlorinated to

ethene, lactate was completely consumed, therefore lactate and PCE were re-fed. Acetate production

occurred continuously during the whole dechlorination process (Figure 1E), other organic acids such

as succinate were not detected. No hydrogen was detected in the gas phase at any point. Because

the concentration of hydrogen and succinate which were shown to be produced during pyruvate

fermentation of *S. multivorans* pure cultures [36] might be below the detection limit in our setup and

pure *S. multivorans* cultures could not grow via lactate fermentation, a stoichiometry could not be

calculated.

Similar growth characteristics and dechlorination behavior were observed in the *Sm*/195 co-culture

cultivated under the same conditions, except that vinyl chloride (VC) was the major dechlorination

product. The first dose of PCE was dechlorinated within 7 days stoichiometrically to VC (Figure 1B).

The increase of cell number was slightly lower for *S. multivorans* and slightly higher for *D. mccartyi*

strain 195 compared to the *Sm*/BTF08 co-culture (Figure 1D). After the second dose of PCE was

dechlorinated to VC, a low amount of ethene was produced starting on day 10, reaching 104 µM

ethene after day 15. The *Sm*/195 co-culture reduced PCE to VC more than 3-fold faster than the *D.*

mccartyi strain 195 pure culture, which completed dechlorination to VC within 18 days (Table 1,

Supplementary Figure S5B).

2. Corrinoid transfer in co-cultures and the effect of the lower corrinoid ligand

D. mccartyi strains rely on externally provided corrinoids for dehalogenation and growth. This was also

tested for *D. mccartyi* strains BTF08 and 195, which only showed negligible *c*DCE dechlorination after

100 days without addition of corrinoid (Figure 2A and D). Therefore, it was of interest whether *S.*

multivorans is able to provide functional corrinoids for *D. mccartyi* strains BTF08 and 195. In

Sm/BTF08 co-cultures without the amendment of vitamin B_{12}, stoichiometric dechlorination of PCE to

*c*DCE was obtained (~30 µmol/bottle/day) (Figure 2B). No further dechlorination of *c*DCE to VC or

ethene was detected, indicating that *S. multivorans* alone was responsible for the dechlorination.

When *c*DCE dechlorination stalled in the *Sm*/BTF08 co-culture, 1 µM 5,6-dimethylbenzimidazole

(DMB) was added, which resulted in subsequent *c*DCE dechlorination (Figure 2C). Ethene production

rates from *c*DCE (36 ± 2 µM/day) were similar to the co-culture amended with vitamin B_{12} (41 ± 2

µM/day). The dechlorination rate increased after re-feeding with PCE to 8 µmol/bottle/day. In the

Sm/BTF08 co-culture with DMB but without vitamin B_{12} amendment, three different types of corrinoids

were found by mass spectrometric analysis: [Ade]NCba, [DMB]NCba and [DMB]Cba (Supplementary

Figure S7).

Interestingly and in contrast to the *Sm*/BTF08 co-culture, the *Sm*/195 co-culture without vitamin B_{12}

amendment (-B_{12}) dechlorinated PCE to VC, although at low rates (Figure 2E). After 35 days, PCE

was re-fed, and the *c*DCE to VC dechlorination rate increased 4-fold (3.6 µmol/bottle/day). The only

corrinoid detected in the *Sm*/195 co-culture -B_{12} was [Ade]NCba (Supplementary Figure S8). No

significant increase in *c*DCE dechlorination was observed when 1 µM DMB was added to the *Sm*/195

co-culture (Figure 2F). Similar to the *Sm*/BTF08 co-culture, [Ade]NCba, [DMB]NCba and [DMB]Cba

were found by mass spectrometric analysis when DMB was added to the *Sm*/195 co-culture

(Supplementary Figure S9). To confirm that *D. mccartyi* strain 195 can use [Ade]NCba for

dechlorination of *c*DCE to VC, we amended a pure culture with [Ade]NCba isolated from *S.*

multivorans (Supplementary Figure S10A). Dechlorination of 50 µmol *c*DCE to VC was completed

within 22 days (Supplementary Figure S10A). *D. mccartyi* strain 195 pure cultures amended with [5-

OMeBza]Cba showed slightly faster dechlorination and the *c*DCE was converted within 16 days into

VC (Supplementary Figure S10B). Only the amended corrinoid types were detected in the cultures via

MS, impurities and rearrangement of corrinoids could be therefore excluded (Supplementary Figure

S11 and S12).

3. Electron microscopy and FISH analysis of formed cell aggregates

After about 25 transfers on lactate and PCE, all co-cultures formed spherical aggregates up to 2 mm in

diameter (Figure 3A). Field emission-scanning electron microscopy (FE-SEM) was applied to uncover

the cell morphology and cell distribution in these aggregates. After preparation for FE-SEM, the sizes

of the aggregates were lower (30 to 200 µm, Figure 3B and C). Electron micrographs of both co-

cultures revealed a compact network of *S. multivorans* and *D. mccartyi* cells coiled around net-forming

filament-like structures. The cells were embedded in an extracellular matrix (ECM) which might aid

cell-to-cell contact (Figure 3D, Supplementary Figure S13). *S. multivorans* could be distinguished from

D. mccartyi by FISH with specific oligonucleotide probes targeting 16S rRNA and 3-dimensional

imaging showed a spatial organization and an almost equal distribution of both species within the

aggregates (Figure 3E, Supplementary Figure S14). The high resolution of the confocal laser scanning

microscopy enabled visualization of single cells and revealed the same morphologies as in the

electron micrographs. In the FISH pictures, the sizes of the aggregates ranged from 50 to 100 µm in

diameter.

S. multivorans revealed a typical helical rod-shaped cell structure and a size of 2 to 5 µm by 0.5 µm at

magnifications of around 10.000x and 20.000x in FE-SEM as described previously [55] (Figure 4,

Supplementary Figure S13). However, the typical polar flagellum was only observed in a few cells and

several flagella seemed to be detached from cells, possibly a part of the ECM (Figure 4,

Supplementary Figure S13). *D. mccartyi* showed an atypical cell morphology in the co-cultures.

Microscopic analysis of the pure culture revealed a disc-shaped irregular coccus of 0.5 µm diameter

(Figure 4a, Supplementary Figure S15), as previously described [11], whereas the *D. mccartyi* strains

in the co-culture showed 0.5 µm large barrel-like cells with a flattened cell pole at one side and a ring-

shaped septum (Figure 4, Supplementary Figure S13). Visual analysis of cells separated from

aggregates in the co-culture identified unequivocally both organisms and confirmed the presence and

unusual morphology of *D. mccartyi* (Supplementary Figure S15 and S16).

4. Proteomics of pure and co-cultures

We applied a label-free shotgun proteomics approach to identify changes in protein abundances

during PCE dechlorination in pure and co-cultures (with and without vitamin B_{12}) with *D. mccartyi* strain

BTF08 and *S. multivorans*. The clustering approach nMDS based on quantified proteins revealed a

significant separation between pure and co-cultures (*Dhc* BTF08 proteins p=0.001, *Sm* proteins

p=0.004, Supplementary Figure S17). A multi-level pattern analysis was applied to determine indicator

proteins for a given condition (i.e. co-culture vs. pure culture) to better understand the functional

changes occurring during the dehalogenation in the co-culture (Supplementary Excel File S2).

Of the 20 reductive dehalogenases encoded in the genome of *D. mccartyi* strain BTF08 [57], only two

RDase proteins were identified, the gene products of btf_1393 (a hitherto not annotated RdhA) and

btf_1407 (the vinyl chloride reductase VcrA). The two reductive dehalogenases PceA (PCE reductive

dehalogenase) and TceA (TCE reductive dehalogenase), putatively involved in dechlorination of PCE

to *c*DCE, were not identified under any condition. While VcrA was one of the most abundant proteins

under all tested conditions (Figure 5, Supplementary Figure S18, Supplementary Excel File S1), the

gene product of btf_1393 was more abundant in the two co-cultures (Figure 5, Supplementary Excel

File S2). A BLASTp search against the NCBI nr database revealed that the btf_1393 amino acid

sequence was almost identical (99% or 497/498 amino acid sequence identity over the whole length)

to an RdhA from *D. mccartyi* 11a5, encoded by 11a5_1355 and characterized as a novel PCE

reductive dehalogenase PteA [56,58]. In *S. multivorans*, the proteins encoded in the organohalide

respiratory gene were present among all conditions as described previously [59, 60] (Supplementary

Excel File S1).

Only a few proteins encoded in the genomes of *S. multivorans* or *D. mccartyi* are annotated to play a

potential role in the formation of aggregates, e.g. via an ECM. The proteins which are part of a putative

type II pili system in *D. mccartyi* (encoded by btf_1229 to btf_1240) were present in low to medium

amounts among all conditions (about median [Med] to lower than Med minus one standard deviation

[SD]) but not significantly more abundant in the co-cultures. Many proteins involved in flagellar motility

of *S. multivorans* were lower abundant in the co-culture, with most of the Flg and Fli proteins not

detected (Supplementary Excel File S1). Protein indicator analysis revealed further that several outer

membrane porins of *S. multivorans* (encoded by SMUL_0494, SMUL0693, SMUL_0926 and

SMUL_2351) were relevant proteins for the co-cultures. In addition, several molybdopterin

oxidoreductases (e.g. a nitrate reductase) of *S. multivorans* were highly abundant only in the co-

culture (Supplementary Table S3). Proteins related to cell division (tubulin/GTpase encoded by

btf_0551, FtsZ, btf_0595, FtsH, btf_357, a cell division trigger factor, btf_0631) were more abundant or

exclusively quantified in pure cultures (Figure 5, Supplementary Table S4, Supplementary Excel File

S2). A hypothetical protein encoded close to the tubulin (btf_0548) was also highly abundant

(>Med+SD or >Med) only in cells of the pure culture, while it could not be quantified in any of the co-

cultures.

The amidohydrolase CbiZ (btf_610), responsible for cleavage of the nucleotide loop of corrinoids, was

quantified only in both co-culture conditions in average abundance (Figure 5). The L-threonine 3-O-

phosphate decarboxylase CobD, which synthesizes the linker of the lower base and the corrinoid ring

of cobamides, could not be quantified in any culture.

Discussion

In this study, we investigated the dechlorination of PCE to ethene (or vinyl chloride) in co-cultures of *S.*

multivorans with *D. mccartyi* BTF08 and 195. Dechlorination profiles, metabolite analysis and growth

studies point to a biphasic physiology of the co-culture. In the first stage, when PCE or TCE serve as

electron acceptors, *S. multivorans* grows by organohalide respiration with lactate as electron donor. In

this phase, *D. mccartyi* does not contribute significantly to PCE dechlorination as indicated by the

stable isotope fractionation pattern and the corresponding enrichment factors. This can be explained

by the insufficient supply of *D. mccartyi* with hydrogen as electron donor due to the lack of (or very

low) hydrogen production by *S. multivorans* during respiratory growth. In the second phase of the co-

culture*,* when PCE has been completely converted to *c*DCE after less than two days, *S. multivorans*

grows via fermentation of lactate mainly to acetate, CO_2 and hydrogen [36]. The fate of a part of the

organic carbon and of the electrons generated by lactate oxidation could not be elucidated in this

study. This part might have been used for biomass production and the generation of small amounts of

organic acids, such as succinate, below the detection limit of the HPLC used here. It has been shown

earlier that, for thermodynamic reasons, lactate can only be utilized fermentatively by *S. multivorans* in

co-cultures, where hydrogen concentration is kept low by a syntrophic hydrogen consumer, while *S.*

multivorans does not grow fermentatively on lactate in pure cultures [36]. In the co-culture described

here, *D. mccartyi* utilizes H_2 produced by *S. multivorans* as electron donor for reductive

dehalogenation of *c*DCE or VC (Figure 6). As expected, PCE was completely dechlorinated to ethene

in the *Sm*/BTF08 co-culture and mainly to VC and minor amounts of ethene in *Sm*/195. A similar slow

and incomplete VC dechlorination was found for pure cultures of the latter strain [61] and co-cultures

of *D. mccartyi* strain 195 with either *Desulfovibrio vulgaris* Hildenborough or *Syntrophomonas wolfei*

[21, 22]. *c*DCE was presumably converted to ethene by VcrA of BTF08. This is supported by the

proteomic analysis described here, in which VcrA of *D. mccartyi* strain BTF08 was highly abundant

and the predominant RDase in all cultures. A highly similar VcrA ortholog of *D. mccartyi* strain VS

(99% amino acid sequence identity) has already been biochemically characterized and described to

dechlorinate *c*DCE and VC [62]. Interestingly, neither TceA, nor PceA, which had been suggested to

be responsible for dechlorination of PCE to VC [57], were detected in any of the samples. An identical

ortholog of the only other detected RDase (encoded by btf_1393) was recently characterized as PteA,

a novel PCE reductive dehalogenase dechlorinating PCE to TCE in *D. mccartyi* strain 11a5 [58]. *D.*

mccartyi strain 195 was not subject to proteomic investigation in our study. However, previous studies

suggested that TceA of this organism is involved in *c*DCE to VC dechlorination and that VC is further

dechlorinated slowly by the same enzyme [61, 63].

Norpseudo-B$_{12}$, produced by *S. multivorans*, is used differently by the two *D. mccartyi* strains

Without amendment of vitamin B$_{12}$ the *Sm*/BTF08 co-culture was not capable of *c*DCE dechlorination

to ethene, indicating that *D. mccartyi* strain BTF08 cannot use the norpseudo-B$_{12}$ ([Ade]NCba)

synthesized by *S. multivorans* for *c*DCE dechlorination. Complete dechlorination to ethene by *D.*

mccartyi strain BTF08 was restored by the addition of the lower ligand DMB, indicating that the non-

functional adenine ligand of [Ade]NCba could be replaced by DMB. This assumption is supported by a

previous study, in which *D. mccartyi* was shown to incorporate different lower ligands into cobamides

[17]. Therefore, it was not surprising that DMB-containing cobalamin were detected in the *Sm*/*Dhc* co-

cultures. Most probably, after addition of DMB the detected [DMB]NCba was synthesized by *S.*

multivorans besides [Ade]NCba, since the organism is able to incorporate different benzimidazoles

and generate the corresponding cobamides [35, 64]. [DMB]Cba could then be produced during

salvaging and remodeling of available [Ade]NCba and/or [DMB]NCba by *D. mccartyi*. It can, however,

not be excluded from our data that [DMB]NCba can be used as well by *D. mccartyi* strain BTF08 for

cDCE or VC dechlorination. The nucleotide loop cleavage is most likely mediated by the

adenosylcobinamide hydrolase CbiZ, as shown for CbiZ of *Rhodobacter sphaeroides* [65, 66]. Of the

two CbiZ encoded in the genome of *D. mccartyi* strain BTF08, one was detected in the proteome and

showed a higher abundance in the co-culture compared to the pure culture. These results present

evidence of the involvement of CbiZ in exchanging the complete nucleotide loop in *D. mccartyi*.

Different from the *c*DCE accumulation in *Sm*/BTF08 co-cultures grown without DMB, the *Sm*/195 co-

culture dechlorinated PCE to VC in media without vitamin B$_{12}$ or DMB. This indicates *c*DCE to VC

conversion by strain 195 with [Ade]NCba produced by *S. multivorans*. However, previous studies

showed [Ade]Cba (or other cobalamins with adenine as lower ligand) to be non-functional in *D.*

mccartyi strain 195 with TCE as electron acceptor [17]. Trace amounts of DMB or cobalamin could be

an explanation, but this explanation is unlikely given the same treatment but different dechlorination

pattern of the co-cultures. Since RDase gene expression is often dependent on the electron acceptor

present [14], *c*DCE in the *Sm*/195 co-culture might have induced a yet unknown RDase dechlorinating

*c*DCE to VC with [Ade]NCba as cofactor. Alternatively, TceA containing [Ade]NCba could catalyze

cDCE to VC dechlorination, but not TCE dechlorination to *c*DCE. The mechanism of cobalamin

transfer from *S. multivorans* to *D. mccartyi* is not yet known. Most likely, cobalamin is released upon

cell lysis or degradation of periplasmic PceA and possibly excreted through porins of *S. multivorans*,

some of which were higher abundant in co-cultures.

Cell aggregates formed by *D. mccartyi* and *S. multivorans* could enhance interspecies

hydrogen transfer

The cell aggregates observed in all studied co-cultures but not in pure cultures are common for

obligate syntrophic interactions often found in e.g. acetogenic and methanogenic communities [67, 68,

69]. The decrease of intermicrobial distances and establishment of cell-to-cell contacts, which were

observed in co-cultures via FE-SEM, should lead to increased metabolite (e.g. hydrogen) fluxes

between species according to Fick's law [70], ultimately enhancing growth and dechlorination rates.

Experimentally, this was shown, for example, during syntrophic propionate conversion of

Pelotomaculum thermopropionicum SI and *Methanothermobacter thermoautotrophicus* ΔH where

interspecies hydrogen transfer was calculated to be optimal in aggregates [71]. EPS-like substances

and flagella most likely contribute to a stabilization of aggregates by adhesion and attachment of the

cells [67, 72]. In addition, cellular extrusions such as pili and flagella might contribute to direct

interspecies electron transfer (DIET). While *S. multivorans* produces flagella in pure cultures [55],

several structural flagellar proteins were down-regulated in the co-culture, speaking against a use of

flagellar for DIET in the co-culture of *D. mccartyi* and *S. multivorans*. Other proteins putatively involved

in DIET are not encoded in the genomes of *D. mccartyi* or *S. multivorans*, arguing for an exclusive

interspecies transfer of electrons via hydrogen. Aggregated *D. mccartyi* cells embedded in an ECM

were observed in a bioreactor community dechlorinating trichloroethene to ethene [73]. However in

this study, *D. mccartyi* formed typical disc-shaped cells, in contrast to the unusual, never before

observed barrel-like morphology of *D. mccartyi* in the co-cultures investigated in our study. This barrel-

like morphology might be caused by the down-regulation of proteins involved in cell division in the co-

culture. One of these proteins, FtsZ, localized at the cell division site, was shown to play a key role in

cytokinesis in *E. coli*. It was shown to be responsible for septal invagination of the cell wall and

cytoplasmic membrane by forming a ring-shaped septum followed by cell division [74, 75, 76]. Its

down-regulation might hamper a complete membrane constriction resulting in slower cell division that

might cause the observed barrel-like cell morphologies. However, the reasons for this down-regulation

and the different morphology remain obscure. Possibly the less frequent cell divisions and/or the

barrel-like morphology give *D. mccartyi* an advantage in cell-to-cell-contact. In addition, this shape

could represent a "sessile" state in which *D. mccartyi* adjusts its metabolism to the given condition in

the aggregates, including a downregulation of cell division. Since electron micrographs of co-cultures

with *D. mccartyi* are scarce, it is not possible to state whether the unusual cell morphology is specific

for the partnership with *S. multivorans* or whether it is found frequently for co-cultures containing *D.*

mccartyi. With *S. wolfei* as syntrophic partner, disc-like cell structures typical for *D. mccartyi* strain 195

were observed [21].

Conclusion

This study provides first insights into the interactions of *S. multivorans* in association with *D. mccartyi*

and the formation of aggregates in which *D. mccartyi* showed an unusual barrel-like morphology.

Dechlorinating microbial communities sometimes reveal the presence of *Sulfurospirillum* spp., but the

functional role of these Campylobacterota (formerly Epsilonproteobacteria) is unexplored [77]. We

observed that interspecies hydrogen and cobamide transfer in the co-culture resulted in fast and

complete dechlorination of PCE to ethene. *S. multivorans* could provide all growth factors required by

D. mccartyi, including hydrogen and *c*DCE as energy source, acetate as carbon source and

cobamides as RDase cofactors (Figure 6). It is the first study in which *D. mccartyi* was provided with

all nutrients required for growth by its syntrophic partner. Additionally, PCE to *c*DCE dechlorination

was sped up by the fast dechlorination rate of *S. multivorans*. This is of high interest for bioremediation

attempts using *Dehalococcoides*-containing mixed cultures, since electron donor and cobalamin

limitations may impede *Dehalococcoides* dechlorination activities.

Acknowledgement

This work was supported by the Jena School for Microbial Communication (JSMC) and the DFG

Research Unit FOR 1530 (sub-projects 1, 2, 5 and 6 were involved). The authors would like to thank

Benjamin Scheer and Kathleen Eismann (Helmholtz Centre for Environmental Research, Leipzig) for

skillfull technical assistance in the lab, Susanne Linde (University Hospital Jena, Center for Electron

Microscopy) for the field-emission scanning electron microscopic analysis and Dr. Elena Romano

(University of Rome Tor Vergata, Department of Biology) for invaluable assistance in the confocal

laser scanning microscope image acquisition and processing. Dr. Torsten Schubert (Friedrich Schiller

University Jena) is gratefully acknowledged for insightful discussions on corrinoid metabolism.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1.   Morris, B.E., Henneberger, R., Huber, H., Moissl-Eichinger, C. 2013. Microbial syntrophy:
Interaction for the common good. *FEMS Microbiol Rev*, **37**(3), 384-406.
2.   Schink, B., Stams, A. 2013. Syntrophism Among Prokaryotes. in: *The Prokaryotes :*
Prokaryotic Communities and Ecophysiology, (Eds.) E. Rosenberg, E. DeLong , S. Lory, E.
Stackebrandt , F. Thompson, Springer. Berlin Heidelberg, pp. 471-493.
3.   Stams, A.J., Plugge, C.M. 2009. Electron transfer in syntrophic communities of anaerobic
bacteria and archaea. *Nat Rev Microbiol*, **7**(8), 568-77.
4.   Cichocka, D., Nikolausz, M., Haest, P.J., Nijenhuis, I. 2010. Tetrachloroethene conversion to
ethene by a *Dehalococcoides*-containing enrichment culture from Bitterfeld. *FEMS Microbiol*
Ecol, **72**(2), 297-310.
5.   Duhamel, M., Edwards, E.A. 2006. Microbial composition of chlorinated ethene-degrading
cultures dominated by *Dehalococcoides*. *FEMS Microbiology Ecology*, **58**(3), 538-549.
6.   Jugder, B.E., Ertan, H., Bohl, S., Lee, M., Marquis, C.P., Manefield, M. 2016. Organohalide
Respiring Bacteria and Reductive Dehalogenases: Key Tools in Organohalide Bioremediation.
Front Microbiol, **7**, 249.
7.   Wei, K., Grostern, A., Chan, W.W.M., Richardson, R.E., Edwards, E.A. 2016. Electron
Acceptor Interactions Between Organohalide-Respiring Bacteria: Cross-Feeding, Competition,
and Inhibition. in: *Organohalide-Respiring Bacteria*, (Eds.) L. Adrian, F.E. Löffler, Springer
Berlin Heidelberg. Berlin, Heidelberg, pp. 283-308.
8.   Leys, D., Adrian, L., Smidt, H. 2013. Organohalide respiration: microbes breathing chlorinated
molecules. *Philos Trans R Soc Lond B Biol Sci*, **368**(1616), 20120316.
9.   Schubert, T., Adrian, L., Sawers, R.G., Diekert, G. 2018. Organohalide respiratory chains:
composition, topology and key enzymes. *FEMS Microbiology Ecology*, **94**(4), fiy035-fiy035.
10. Schubert, T., Diekert, G. 2016. Comparative Biochemistry of Organohalide Respiration. in:
Organohalide-Respiring Bacteria, (Eds.) L. Adrian, F.E. Löffler, Springer Berlin Heidelberg.
Berlin, Heidelberg, pp. 397-427.
11. Löffler, F.E., Yan, J., Ritalahti, K.M., Adrian, L., Edwards, E.A., Konstantinidis, K.T., et al.
2013. *Dehalococcoides mccartyi* gen. nov., sp. nov., obligately organohalide-respiring
anaerobic bacteria relevant to halogen cycling and bioremediation, belong to a novel bacterial
class, *Dehalococcoidia* classis nov., order *Dehalococcoidales* ord. nov. and family
Dehalococcoidaceae fam. nov., within the phylum Chloroflexi. *Int J Syst Evol Microbiol*, **63**(Pt
2), 625-35.
12. Zinder, S. 2016. The Genus *Dehalococcoides*. in: *Organohalide-respiring Bacteria*, (Eds.) L.
Adrian, F. Löffler, Springer. Berlin Heidelberg.
13. Schipp, C.J., Marco-Urrea, E., Kublik, A., Seifert, J., Adrian, L. 2013. Organic cofactors in the
metabolism of *Dehalococcoides mccartyi* strains. *Philos Trans R Soc Lond B Biol Sci*,
368(1616), 20120321.
14. Türkowsky, D., Jehmlich, N., Diekert, G., Adrian, L., von Bergen, M., Goris, T. 2018. An
integrative overview of genomic, transcriptomic and proteomic analyses in organohalide
respiration research. *FEMS Microbiol Ecol*, **94**(3).
15. He, J., Holmes, V.F., Lee, P.K., Alvarez-Cohen, L. 2007. Influence of vitamin B_{12} and
cocultures on the growth of *Dehalococcoides* isolates in defined medium. *Appl Environ*
Microbiol, **73**(9), 2847-53.

16. Men, Y., Seth, E.C., Yi, S., Allen, R.H., Taga, M.E., Alvarez-Cohen, L. 2014. Sustainable
Growth of *Dehalococcoides mccartyi* 195 by Corrinoid Salvaging and Remodeling in Defined
Lactate-Fermenting Consortia. *Applied and Environmental Microbiology*, **80**(7), 2133-2141.
17. Yi, S., Seth, E.C., Men, Y.-J., Stabler, S.P., Allen, R.H., Alvarez-Cohen, L., et al. 2012.
Versatility in Corrinoid Salvaging and Remodeling Pathways Supports Corrinoid-Dependent
Metabolism in *Dehalococcoides mccartyi*. *Applied and Environmental Microbiology*, **78**(21),
7745-7752.
18. Cheng, D., Chow, W.L., He, J. 2010. A *Dehalococcoides*-containing co-culture that
dechlorinates tetrachloroethene to *trans*-1,2-dichloroethene. *ISME J*, **4**(1), 88-97.
19. He, J., Ritalahti, K.M., Aiello, M.R., Löffler, F.E. 2003a. Complete detoxification of vinyl
chloride by an anaerobic enrichment culture and identification of the reductively dechlorinating
population as a *Dehalococcoides* species. *Appl Environ Microbiol*, **69**(2), 996-1003.
20. He, J., Ritalahti, K.M., Yang, K.L., Koenigsberg, S.S., Löffler, F.E. 2003b. Detoxification of
vinyl chloride to ethene coupled to growth of an anaerobic bacterium. *Nature*, **424**(6944), 62-5.
21. Mao, X., Stenuit, B., Polasko, A., Alvarez-Cohen, L. 2015. Efficient Metabolic Exchange and
Electron Transfer within a Syntrophic Trichloroethene-Degrading Coculture of
Dehalococcoides mccartyi 195 and *Syntrophomonas wolfei*. *Appl Environ Microbiol*, **81**(6),
2015-2024.
22. Men, Y., Feil, H., Verberkmoes, N.C., Shah, M.B., Johnson, D.R., Lee, P.K., et al. 2012.
Sustainable syntrophic growth of *Dehalococcoides ethenogenes* strain 195 with *Desulfovibrio*
vulgaris Hildenborough and *Methanobacterium congolense*: global transcriptomic and
proteomic analyses. *ISME j*, **6**(2), 410-21.
23. Murray, A., Maillard, J., Rolle, M., Broholm, M., Holliger, C. 2020. Impact of iron- and/or
sulfate-reduction on a cis-1,2-dichloroethene and vinyl chloride respiring bacterial consortium:
experiments and model-based interpretation. *Environ Sci Process Impacts*, **22**(3), 740-750.
24. Wang, S., Chen, C., Zhao, S., He, J. 2019, Microbial synergistic interactions for reductive
dechlorination of polychlorinated biphenyls. Sci Total Environ. **666**, 368-376
25. Yan, J., Im, J., Yang, Y., Löffler, F.E. 2013. Guided cobalamin biosynthesis supports
Dehalococcoides mccartyi reductive dechlorination activity. *Philos Trans R Soc Lond B Biol*
Sci, **368**(1616), 20120320.
26. Yan, J., Ritalahti, K.M., Wagner, D.D., Löffler, F.E. 2012. Unexpected Specificity of
Interspecies Cobamide Transfer from *Geobacter* spp. to Organohalide-Respiring
Dehalococcoides mccartyi Strains. *Applied and Environmental Microbiology*, **78**(18), 6630-
6636.
27. Richardson, R. 2016. Organohalide-Respiring Bacteria as Members of Microbial Communities:
Catabolic Food Webs and Biochemical Interactions. in: *Organohalide-respiring Bacteria*,
(Eds.) L. Adrian, F. Löffler, Springer. Berlin Heidelberg.
28. Richardson, R.E., Bhupathiraju, V.K., Song, D.L., Goulet, T.A., Alvarez-Cohen, L. 2002.
Phylogenetic characterization of microbial communities that reductively dechlorinate TCE
based upon a combination of molecular techniques. *Environ Sci Technol*, **36**(12), 2652-62.
29. Duhamel, M., Edwards, E.A. 2007. Growth and yields of dechlorinators, acetogens, and
methanogens during reductive dechlorination of chlorinated ethenes and dihaloelimination of 1
,2-dichloroethane. *Environ Sci Technol*, **41**(7), 2303-10.
30. Guimaraes, D., Weber, A., Klaiber, I., Vogler, B., Renz, P. 1994. Guanylcobamide and
hypoxanthylcobamide: corrinoids formed by *Desulfovibrio vulgaris*. *Archives of Microbiology*,
162(4), 272-276.
31. Lu, Q., Zou, X., Liu, J., Liang, Z., Shim, H., Qiu, R., et al. 2020. Inhibitory effects of metal ions
on reductive dechlorination of polychlorinated biphenyls and perchloroethene in distinct
organohalide-respiring bacteria. *Environ Int*.**135**, 105373.
32. Yan, J., Şimşir, B., Farmer, A.T., Bi, M., Yang, Y., Campagna, S.R., et al. 2016. The corrinoid
cofactor of reductive dehalogenases affects dechlorination rates and extents in organohalide-
respiring *Dehalococcoides mccartyi*. *ISME J.* **10**:1092-101.
33. Kräutler, B., Fieber, W., Ostermann, S., Fasching, M., Ongania, K.-H., Gruber, K., et al. 2003.
The Cofactor of Tetrachloroethene Reductive Dehalogenase of *Dehalospirillum multivorans* Is
Norpseudo-B$_{12}$, a New Type of a Natural Corrinoid. *Helvetica Chimica Acta*, **86**(11), 3698-
3716.
34. Neumann, A., Wohlfarth, G., Diekert, G. 1996. Purification and Characterization of
Tetrachloroethene Reductive Dehalogenase from *Dehalospirillum multivorans*. *Journal of*
Biological Chemistry, **271**(28), 16515-16519.

35. Schubert, T. 2017. The organohalide-respiring bacterium *Sulfurospirillum multivorans*: a natural source for unusual cobamides. *World Journal of Microbiology and Biotechnology*, **33**(5), 93.

36. Kruse, S., Goris, T., Westermann, M., Adrian, L., Diekert, G. 2018. Hydrogen production by *Sulfurospirillum* spp. enables syntrophic interactions of Epsilonproteobacteria. *Nat Comm*. **9**, 4872.

37. Goris, T., Schenz, B., Zimmermann, J., Lemos, M., Hackermüller, J., Schubert, T., et al. 2017. The complete genome of the tetrachloroethene-respiring Epsilonproteobacterium *Sulfurospirillum halorespirans. Journal of Biotechnology*, **255**, 33-36.

38. Maymó-Gatell, X., Anguish, T., Zinder, S.H. 1999. Reductive dechlorination of chlorinated ethenes and 1, 2-dichloroethane by "Dehalococcoides ethenogenes" 195. *Appl Environ Microbiol*, **65**(7), 3108-13.

39. Keller, S., Ruetz, M., Kunze, C., Kräutler, B., Diekert, G., Schubert, T. 2014. Exogenous 5,6-dimethylbenzimidazole caused production of a non-functional tetrachloroethene reductive dehalogenase in *Sulfurospirillum multivorans*. *Environmental Microbiology*, **16**(11), 3361-3369.

40. Kruse, S., Goris, T., Wolf, M., Wei, X., Diekert, G. 2017. The NiFe Hydrogenases of the Tetrachloroethene-Respiring Epsilonproteobacterium *Sulfurospirillum multivorans*: Biochemical Studies and Transcription Analysis. *Front Microbiol*, **8**, 444.

41. Schubert, T., von Reuß, S.H., Kunze, C., Paetz, C., Kruse, S., Brand-Schön, P., et al. 2018. Guided cobamide biosynthesis for heterologous production of reductive dehalogenases. *Microbiol Biotechnol*, 12(2), 346-359.

42. Marco-Urrea, E., Nijenhuis, I., Adrian, L. 2011 Transformation and carbon isotope fractionation of tetra- and trichloroethene to trans-dichloroethene by *Dehalococcoides* sp strain CBDB1, Environ. Sci. Technol. 45, 1555-1562

43. Schmidt, M., Lege, S., Nijenhuis, I. 2014. Comparison of 1,2-dichloroethane, dichloroethene and vinyl chloride carbon stable isotope fractionation during dechlorination by two *Dehalococcoides strains. Water Res*, **52**, 146-54.

44. Coplen, T.B., Brand, W.A., Gehre, M., Gröning, M., Meijer, H.A., Toman, B., et al., 2006. After two decades a second anchor for the VPDB delta13C scale. *Rapid Commun Mass Spectrom*, **20**(21), 3165-6.

45. Elsner, M. 2010. Stable isotope fractionation to investigate natural transformation mechanisms of organic contaminants: principles, prospects and limitations. *J Environ Monit*, **12**(11), 2005-31.

46. Elsner, M., Chartrand, M., Vanstone, N., Couloume, G.L., Lollar, B.S. 2008. Identifying abiotic chlorinated ethene degradation: characteristic isotope patterns in reaction products with nanoscale zero-valent iron. *Environ Sci Technol*, **42**(16), 5963-70.

47. Rossetti, S., Aulenta, F., Majone, M., Crocetti, G., Tandoi, V. 2008. Structure analysis and performance of a microbial community from a contaminated aquifer involved in the complete reductive dechlorination of 1,1,2,2-tetrachloroethane to ethene. *Biotechnol Bioeng*, **100**(2), 240-9.

48. Fazi, S., Aulenta, F., Majone, M., Rossetti, S. 2008. Improved quantification of Dehalococcoides species by fluorescence in situ hybridization and catalyzed reporter deposition. *Syst Appl Microbiol*, **31**(1), 62-7.

49. Matturro, B., Tandoi, V., Rossetti, S. 2013. Different activity levels of Dehalococcoides mccartyi revealed by FISH and CARD-FISH under non-steady and pseudo-steady state conditions. *N Biotechnol*, **30**(6), 756-62.

50. Jehmlich, N., Schmidt, F., von Bergen, M., Richnow, H.-H., Vogt, C. 2008. Protein-based stable isotope probing (Protein-SIP) reveals active species within anoxic mixed cultures. *The Isme Journal*, **2**, 1122.

51. Türkowsky (b), D., Esken, J., Goris, T., Schubert, T., Diekert, G., Jehmlich, N., et al. 2018. A Retentive Memory of Tetrachloroethene Respiration in *Sulfurospirillum halorespirans* - Involved Proteins and a Possible Link to Acetylation of a Two-Component Regulatory System. *J Proteom*, **181,** 36-46.

52. Kanehisa, M., Sato, Y., Morishima, K. 2016. BlastKOALA and GhostKOALA: KEGG Tools for Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol*, **428**(4), 726-731.

53. Marc, D., Pierre, L. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs*, **67**(3), 345-366.

54. Malik, A., Puissant, J., Buckeridge, K., Goodall, T., Jehmlich, N., Chowdhury, S.,et al. 2017.
Land use driven change in soil pH affects microbial carbon cycling processes. *Nat Comm*.
9(1), 3591.
55. Scholz-Muramatsu, H., Neumann, A., Meßmer, M., Moore, E., Diekert, G. 1995. Isolation and
characterization of *Dehalospirillum multivorans* gen. nov., sp. nov., a tetrachloroethene-
utilizing, strictly anaerobic bacterium. *Archives of Microbiology*, **163**(1), 48-56.
56. Franke, S., Seidel, K., Adrian, L., Nijenhuis, I. 2020. Dual Element (C/Cl) Isotope Analysis
Indicates Distinct Mechanisms of Reductive Dehalogenation of Chlorinated Ethenes and
Dichloroethane in Dehalococcoides mccartyi Strain BTF08 With Defined Reductive
Dehalogenase Inventories. Front Microbiol. 11:1507.
57. Pöritz M, Goris T, Wubet T, Tarkka MT, Buscot F, Nijenhuis I, et al. 2013. Genome sequences
of two dehalogenation specialists – *Dehalococcoides mccartyi* strains BTF08 and DCMB5
enriched from the highly polluted Bitterfeld region. *FEMS Microbiol Lett*.;343(2):101-104.
58. Zhao, S., Ding, C., He, J. 2017. Genomic characterization of *Dehalococcoides mccartyi* strain
11a5 reveals a circular extrachromosomal genetic element and a new tetrachloroethene
reductive dehalogenase gene. *FEMS Microbiology Ecology*, **93**(4), fiw235.
59. Goris, T., Schiffmann, C.L., Gadkari, J., Schubert, T., Seifert, J., Jehmlich, N., et al. 2015.
Proteomics of the organohalide-respiring Epsilonproteobacterium *Sulfurospirillum multivorans*
adapted to tetrachloroethene and other energy substrates. *Scientific Reports*, **5**, 13794.
60. Gadkari, J., Goris, T., Schiffmann, C.L., Rubick, R., Adrian, L., Schubert, T., et al. 2018.
Reductive tetrachloroethene dehalogenation in the presence of oxygen by *Sulfurospirillum*
multivorans: physiological studies and proteome analysis. *FEMS Microbiology Ecology*, **94**(1),
fix176-fix176.
61. Maymó-Gatell, X., Nijenhuis, I., Zinder, S.H. 2001. Reductive Dechlorination of cis-1,2-
Dichloroethene and Vinyl Chloride by "*Dehalococcoides ethenogenes*". *Environmental*
Science & Technology, **35**(3), 516-521.
62. Parthasarathy, A., Stich, T.A., Lohner, S.T., Lesnefsky, A., Britt, R.D., Spormann, A.M. 2015.
Biochemical and EPR-Spectroscopic Investigation into Heterologously Expressed Vinyl
Chloride Reductive Dehalogenase (VcrA) from *Dehalococcoides mccartyi* Strain VS. *Journal*
of the American Chemical Society, **137**(10), 3525-3532.
63. Magnuson, J.K., Romine, M.F., Burris, D.R., Kingsley, M.T. 2000. Trichloroethene reductive
dehalogenase from *Dehalococcoides ethenogenes*: sequence of *tceA* and substrate range
characterization. *Appl Environ Microbiol*, **66**(12), 5141-7.
64. Keller, S., Kunze, C., Bommer, M., Paetz, C., Menezes, R.C., Svatoš, A., et al. 2018.
Selective utilization of benzimidazolyl-norcobamides as cofactors by the tetrachloroethene
reductive dehalogenase of *Sulfurospirillum multivorans*. *Journal of Bacteriology*.
65. Gray, M.J., Escalante-Semerena, J.C. 2009a. The cobinamide amidohydrolase (cobyric acid-
forming) CbiZ enzyme: A critical activity of the cobamide remodeling system of *Rhodobacter*
sphaeroides. *Molecular microbiology*, **74**(5), 1198-1210.
66. Gray, M.J., Escalante-Semerena, J.C. 2009b. In Vivo Analysis of Cobinamide Salvaging in
Rhodobacter sphaeroides Strain 2.4.1. *Journal of Bacteriology*, **191**(12), 3842-3851.
67. Hulshoff Pol, L.W., de Castro Lopes, S.I., Lettinga, G., Lens, P.N.L. 2004. Anaerobic sludge
granulation. *Water Research*, **38**(6), 1376-1389.
68. Ishii, S., Kosaka, T., Hori, K., Hotta, Y., Watanabe, K. 2005. Coaggregation facilitates
interspecies hydrogen transfer between *Pelotomaculum thermopropionicum* and
Methanothermobacter thermautotrophicus. *Appl Environ Microbiol*, **71**(12), 7838-45.
69. Stams, A.J.M., Worm, P., Sousa, D.Z., Alves, M.M., Plugge, C.M. 2012. Syntrophic
Degradation of Fatty Acids by Methanogenic Communities. in: *Microbial Technologies in*
Advanced Biofuels Production, (Ed.) P.C. Hallenbeck, Springer US. Boston, MA, pp. 127-142.
70. Schink, B., Thauer, R.K. 1988. Energetics of syntrophic methane formation and the influence
of aggregation. in: *Granular anaerobic sludge: microbiology and technology*, (Eds.) L. G, Z.
AJB, G. JTC, H. LW. Puduc, Wageningen, pp. 5-17.
71. Ishii, S.i., Kosaka, T., Hotta, Y., Watanabe, K. 2006. Simulating the Contribution of
Coaggregation to Interspecies Hydrogen Fluxes in Syntrophic Methanogenic Consortia.
Applied and Environmental Microbiology, **72**(7), 5093-5096.
72. Grotenhuis, J.T.C., Smit, M., van Lammeren, A.A.M., Stams, A.J.M., Zehnder, A.J.B. 1991.
Localization and quantification of extracellular polymers in methanogenic granular sludge.
Applied Microbiology and Biotechnology, **36**(1), 115-119.

73. Delgado, A.G., Fajardo-Williams, D., Bondank, E., Esquivel-Elizondo, S., Krajmalnik-Brown, R. 2017. Coupling Bioflocculation of *Dehalococcoides mccartyi* to High-Rate Reductive Dehalogenation of Chlorinated Ethenes. *Environ Sci Technol*, **51**(19), 11297-11307.

74. Bi, E., Lutkenhaus, J. 1991. FtsZ ring structure associated with division in *Escherichia coli*. *Nature*, **354**, 161.

75. de Boer, P., Crossley, R., Rothfield, L. 1992. The essential bacterial cell-division protein FtsZ is a GTPase. *Nature*, **359**, 254.

76. Lutkenhaus, J. 1993. FtsZ ring in bacterial cytokinesis. *Molecular Microbiology*, **9**(3), 403-409.

77. Goris, T., Diekert, G. 2016. The Genus *Sulfurospirillum*. in: *Organohalide-Respiring Bacteria*, (Eds.) L. Adrian, F.E. Löffler, Springer-Verlag. Berlin, Heidelberg, pp. 209-234.

Figures

Figure 1: Dechlorination of chlorinated ethenes, growth and metabolite analysis of *S. multivorans*/*D. mccartyi* co-cultures with vitamin B$_{12}$ amendment. (A) PCE dechlorination of *Sm*/BTF08 and (B) *Sm*/195. (C) Growth curve of *Sm*/BTF08 and (D) *Sm*/195. (E) Lactate consumption and acetate production of *Sm*/BTF08 and (F) *Sm*/195. Arrows indicate the time points of re-feeding the culture with PCE (A-D) or lactate (E and F). Broken red lines with open symbols represent theoretical, not analytical, values of the PCE concentration as added to the culture. Please note the secondary y-axes for *D. mccartyi* cell numbers in C and D. Negative controls were run with autoclaved cells (abiotic controls, C). Standard deviation of three independent biological replicates (N=3) are represented by error bars (not visible when smaller than the used symbol). ∑ - mass balance; sum of PCE, TCE, *cis*-DCE, VC and ethene.

Figure 2: Dechlorination of *S. multivorans*/*D. mccartyi* co-cultures without addition of vitamin B$_{12}$. (A) Strain BTF08 with *c*DCE as electron acceptor (negative control). (B) *Sm*/BTF08 with PCE as the electron acceptor. (C) *Sm*/BTF08 with PCE as electron acceptor and amendment of 1 µM DMB. (D) Strain 195 with *c*DCE as electron acceptor (negative control). (E) *Sm*/195 with PCE as the electron acceptor. (F) *Sm*/195 with PCE as electron acceptor and amendment of 1 µM DMB. Please note the different time scales. All growth experiments were conducted in biological triplicates (N=3). Arrows indicate re-feeding of PCE. ∑ - mass balance; sum of PCE, TCE, *c*DCE, VC and ethene.

Figure 3: Microscopic analysis of cell aggregates in co-cultures of *S. multivorans* and *D. mccartyi* strains 195 and BTF08. (A) Serum bottle (200 mL) of a *Sm*/BTF08 co-culture showing cell aggregates. (B) Light microscopic image of a *Sm*/BTF08 aggregate. (C, D) Scanning electron micrographs of an aggregate of *Sm*/195. (E) Confocal laser scanning image of FISH stained aggregates of *Sm*/BTF08. Red - *S. multivorans*, green - *D. mccartyi*.

Figure 4: Different cell morphologies of *D. mccartyi* strain BTF08 cells in pure culture (A) and the hitherto unknown barrel-like morphology in co-culture with *S. multivorans* (B, C). White arrows indicate flagella and yellow arrows indicate ring-shaped septum.

Figure 5: Protein abundances of reductive dehalogenases and a set of significant indicator proteins of *Dhc* BTF08 in pure and co-cultures. Average protein abundance values under different cultivation conditions are shown. Abundances represent log10 fold changes of the median, scaled to zero. The median of all proteins is at 2.3 (see Supplementary Figure S18, which displays each replicate in a scatter plot graph). Four replicates (N=4) were used for proteomic analyses; one of the four replicates were discarded from each of the *D. mccartyi* pure cultures (see Methods). Error bars represent standard deviation, which is covered completely by the symbol if <0.1. P – *D. mccartyi* strain

878 BTF08 pure culture, C – *Sm*/BTF08 co-culture, CbiZ - adenosylcobinamide amidohydrolase (btf_610),
879 FtsZ - cell division protein (btf_0595), FtsZ/Tubulin GTPase (btf_0551), VcrA - vinyl chloride reductive
880 dehalogenase (btf_1407), btf_1393 - reductive dehalogenase homolog to 11a5_1355 of *D. mccartyi*
881 11a5.
882

Figure 6: Interspecies metabolite transfer of *Sulfurospirillum multivorans* and *Dehalococcoides*
***mccartyi*.** PCE is dechlorinated to *c*DCE by *S. multivorans* with electrons from lactate oxidation. After
depletion of PCE, *S. multivorans* switches to fermentative metabolism, thereby generating hydrogen,
which is consumed by *D. mccartyi* as electron donor. The electron acceptor for *D. mccartyi* is *c*DCE
and is further dechlorinated to ethene or VC by *D. mccartyi*. *S. multivorans* synthesizes norpseudo B_{12}
([Ade]NCba, ●) and [DMB]NCba (●) when DMB (maximally 1 μM) is amended. Both can be salvaged
and remodeled by *D. mccartyi* into [DMB]Cba (● with a y-shaped linker). The interspecies cobamide
transfer is indicated by yellow arrows. A discrimination between norcobamide or cobamide
incorporation into the *D. mccartyi* RDases is not depicted here, since the data do not allow a
conclusion on the cobalamin nucleotide loop type.

Table 1: Dechlorination rates of different co-culture set-ups *of S. multivorans* and *D. mccartyi* BTF08 or 195 from PCE to the products VC (*D. mccartyi* strain 195) or ethene (*D. mccartyi* strain BTF08).

Organism(s)	Electron donor	Dechlorination rate (μM day^{-1})
With vitamin B$_{12}$		
D. mccartyi BTF08	H$_2$	9 ± 0.3
D. mccartyi 195	H$_2$	15 ± 0.4
Sm/BTF08	Lactate	41 ± 2
Sm/195	Lactate	48 ± 1
Without vitamin B$_{12}$		
Sm/BTF08	Lactate	-
Sm/195	Lactate	9 ± 0.04
Without vitamin B$_{12}$ / with DMB		
Sm/BTF08	Lactate	36 ± 2
Sm/195	Lactate	8 ± 0.5

-: No dechlorination to VC or ethene observed.