

The cytoprotective role of antioxidants in mammalian cells under rapidly varying temperature, pressure and UV conditions during stratospheric balloon campaign

The cytoprotective role of antioxidants in mammalian cells during stratospheric balloon campaign

Dawid Przystupski^{1▲►■}, Agata Górski^{2►■}, Paulina Rozborska^{3►■}, Weronika Bartosik^{4►■}, Olga Michel^{5□}, Joanna Rossowska^{6■}, Anna Szewczyk^{7■}, Małgorzata Drąg-Zalesińska^{8■}, Paulina Kasperkiewicz^{9■}, Jędrzej Górski^{10▼}, Julita Kulbacka^{11●}

¹ Faculty of Medicine, Wrocław Medical University, J. Mikulicza-Radeckiego 5, 50-345 Wrocław, Poland

² Department of Biological Sciences, Institute of Experimental Biology, University of Wrocław, Kanonia 6/8, 50-328 Wrocław, Poland

³ Faculty of Chemistry, Wrocław University of Science and Technology, Wrocław, Poland, Norwida 4/6, 50-373 Wrocław

⁴ Faculty of Biotechnology, University of Wrocław, Joliot-Curie 14a, 50-385 Wrocław, Poland

⁵ Department of Medical Biochemistry, Wrocław Medical University, Chalubinskiego 10, 50-368, Wrocław, Poland

⁶ Institute of Immunology and Experimental Therapy Polish Academy of Sciences, Rudolfa Weigla 12, 53-114 Wrocław, Poland

⁷ Department of Animal Developmental Biology, Institute of Experimental Biology, University of Wrocław, Sienkiewicza 21, 50-357 Wrocław, Poland

⁸ Department of Histology and Embryology, Wrocław Medical University, Chalubinskiego 6a, 50-368 Wrocław

⁹ Department of Bioorganic Chemistry, Faculty of Chemistry, Wrocław University of Science and Technology, Wyb. Wyspińskiego 27, 50-370, Wrocław, Poland

¹⁰ Faculty of Mechanical Engineering, Wrocław University of Science and Technology, Wybrzeże Wyspińskiego 27, 50-370 Wrocław, Poland

¹¹ Department of Molecular and Cellular Biology, Wrocław Medical University, Borowska 211A, 50-556, Wrocław, Poland

▲ - project administration; ► - writing –original draft preparation; ●writing – review & editing, supervision; ■ investigation, □ formal analysis, ▼software

Correspondence should be addressed to Julita Kulbacka, PhD, Eng.
email: julita.kulbacka@umed.wroc.pl

1. Abstract

Currently ongoing age of the dynamic development of the space industry brings the mankind closer to the routine manned space flights and space tourism. That progress leads to a demand for intensive astrobiological research aimed at improving strategies of the pharmacological protection of the human cells against extreme conditions. Although routine research in space remain out of our reach, it is worth noticing that unique severe environment of the Earth's stratosphere have been found to mimic subcosmic conditions, giving rise to the opportunity for use of stratospheric surface as a research model for the astrobiological studies. Our study included launching balloon into the stratosphere containing the human normal and cancer cells treated with various compounds to examine whether these medicines are capable to protect the cells against the stress caused by rapidly varying temperature, pressure and radiation, especially UV. Due to oxidative stress caused by irradiation and temperature shock, we used natural compounds which display antioxidant properties, namely catechin isolated from green tea, honokiol derived from magnolia, curcumin from turmeric and cinnamon extract. "After-flight" laboratory tests displayed the most active antioxidants as potential agents which can minimize harmful impact of extreme conditions to the human cells.

2. Introduction

The world's first hydrogen balloon was created over 200 years ago in Paris, which started the era of scientific ballooning(1). Dynamic development of this technique has enabled the advanced research in atmospheric science, aerobiology, and meteorology (2), remaining promising tool in astrobiology and space biology. In stratosphere the temperature generally drops down -40°C, atmospheric pressure is at 1 kPa, the relative humidity of air is lower than 1%, solar UV irradiance is about 100 W/m² and cosmic radiation is at the level of 0.1 mGy/d (3–5). Stratospheric flights could potentially provide us with a valuable information about the stress response in living organisms after exposure to different severe environmental factors rapidly varying at the same time, which is hard to be mimicked in the laboratory. Moreover, we can examine if some medicines are able to support the viability of living organisms and cells in such an extreme environment. The exposure of varied biological samples to the stratospheric conditions opens the possibility to observe many changes in the cells' functioning e.g. decreased viability, dysfunction of cellular organelles and their localization, cell cycle arrest, changes of gene expression and DNA damage (6,7). Additionally, irradiation in the stratosphere affects the cells either by direct or indirect pathways (water radiolysis, therefore exacerbating the oxidative stress (8)), causing e.g. mitochondrial function disturbance (9), DNA damage, proteins and lipids peroxidation correlated with disruption of the cell membrane (10), that altogether may lead to the cell death (8,10) (Fig 1). Furthermore, stratospheric flights provide unique cyclic changes of linked environmental factors, including radiation, overload, pressure, temperature, wind and vibrations, which are impossible to be simulated altogether in laboratory. The wide use of balloons provides numerous advantages including lower overall project costs, recoverable and massive payloads (up to 3600 kg) and more rapid and flexible flight(3).

Fig 1. Effect of the radiation on the DNA.

High-energy particles could affect DNA damage either by direct pathway or indirect pathway. Free radicals and by-products of the process of water radiolysis contribute to an increase of the oxidative stress in the cell affecting the DNA. Survival of the cell depends on the extent of the DNA damage (8–10). This figure was prepared using Servier Medical Art, available from www.servier.com/Powerpoint-image-bank.

It is known that natural substances derived from plants demonstrated positive impact on living cells including their protective role against diverse factors generating oxidative stress. Green tea polyphenols, such as epigallocatechin gallate, epicatechin gallate, epigallocatechin, epicatechin and catechin (11), are one of the strongest antioxidants and the long-term consumption of green tea extracts increases the activity of oxidative stress enzymes (12) (e.g. superoxide dismutase), inhibits lipo- and cyclooxygenase, xanthine oxidase, activator protein 1 and NF- κ B transcription factors activity (13) and protects cells against radiation(14). Previous meta-analyses suggest a protective role of green tea against numerous cancer types (15–20). Similar properties are displayed by honokiol. This magnolia-derived compound inhibits UVB-induced immunosuppression and induces apoptosis in malignant cells (21). Chilampalli et al. (2010) showed that pretreatment with honokiol is efficient in preventing skin carcinogenesis induced by UV (22). Furthermore, honokiol inhibits photocarcinogenesis by targeting UVB-induced inflammatory mediators and cell cycle regulator (23,24). Turmeric compounds are another free radicals' scavengers. A yellow spice curcumin displays anti-inflammatory (25), antibacterial (26), antiviral (27), anti-cancer (28), proapoptotic (29), neuroprotective (30,31), hepatoprotective (32) activity, promotes autophagy (33), affects the cytochrome P-450 enzyme system (34) and phase II enzymes (35). Curcumin is able to deactivate different forms of free radicals, such as reactive oxygen and nitrogen species (36), modulate the activity of GSH, catalase, and SOD enzymes which reduce oxidative stress (37,38). It can inhibit ROS-generating enzymes such as lipoxygenase/cyclooxygenase and xanthine hydrogenase/oxidase (39–41). Additionally, curcumin shows photosensitizing properties - it can absorb radiation of the appropriate wavelength damaging the tissue by free radicals and facilitating cell death (42). Different forms of free radicals can be scavenged by turmeric compounds. Similar properties are displayed by organic compounds found in cinnamon extract. Beside the antioxidant activity, cinnamon presents radioprotective potential. Azab et al. (2011) have revealed that extract of cinnamon improves the disturbance of the antioxidant system. Furthermore, this substance triggers protective action against protein and lipid oxidation via alteration of membrane structure (43).

Thus, the main aim of this study was to evaluate the effect of stratospheric environment on two living cell lines: human ovarian cancer cells (SKOV-3) and Chinese hamster ovary cells (CHO-K1) in the presence of various natural compounds and investigate the protective role of these drugs. Additionally, the effect of radiation on the examined cells was observed. First, the adherent cells were seeded at equal density (500 000 cells/well) on 6-well plates. After 24-hour incubation with various antioxidant the cells were detached, suspended in freezing medium Bambanker™ (Nippon Genetics, Cat. no. BB01) ($1,5 \times 10^6$ cells/300 μ L) and placed in

microtubes 30 minutes before the balloon flight. Then the samples were transported on ice to the starting point and placed in a radiation transmitting gondola, located on the upper side of environmental measurement unit with accelerometer and temperature, pressure and UV sensors. One half of the samples was covered with aluminum foil to protect the cells against irradiation – mostly UV (the samples described in the study as “protected against radiation”), another half was sent into the stratosphere without the protective layer (described as “not protected against radiation”). As a result, we were able to evaluate the effect of radiation on examined cells in the presence of various antioxidants. As a control we used the appropriate number of cells incubated at 37°C in a humidified incubator with 5% CO₂. Directly after landing, the biological samples were transported on ice to the specialized laboratory, where after-light tests were performed. The cells were seeded on 96-well plates (10 000 cells/well) and incubated in an appropriate drug solution for 24 hours to perform membrane permeabilization assay and intracellular reactive oxygen species generation assay, and for 24, 48 and 72 hours to evaluate mitochondrial activity in MTT assay. Modified versions of these assays (adding reagents into the cells’ suspension, appropriate incubation, centrifugation and seeding on 96-well plates) were used to analyze the suspended (not adherent to multiwell plates) cells directly after the balloon landing (0 h). Additionally, the cells were plated on 6-well plates and incubated in the appropriate drug solution for 24 hours to perform cell cycle assay, and for 7 days to carry out clonogenic assay. To analyze the expression of manganese-dependent superoxide dismutase (SOD2) the cells were seeded on 10-well diagnostic microscopic slides and fixed after 24 hours. Immunocytochemical staining was carried out in the two following days. Furthermore, neutral comet assay was performed to analyze DNA damages associated with the balloon flight. The scheme of the completed experiment is provided below (Fig 2).

Fig 2. The schematic representation of the procedure of balloon flight and the preparation of biological samples.

This figure was prepared using Servier Medical Art, available from www.servier.com/Powerpoint-image-bank.

3. Material and Methods

3.1. Cell culture

Human ovarian cancer cells (SKOV-3) and Chinese hamster ovary cells (CHO-K1) were obtained from the American Type Culture Collection (ATCC, London, UK). Cells were cultured as a monolayer in Dulbecco’s modified Eagle medium (DMEM, Sigma-Aldrich, USA; for SKOV-3 cells) and Ham’s F-10 Nutrient Mix (F-10, Sigma-Aldrich, USA; for CHO-K1 cells) containing 2 mM L-glutamine, 10% fetal bovine serum (FBS, Sigma-Aldrich) and 20 units penicillin and 20 µg streptomycin/mL (Sigma-Aldrich) at 37°C in a humidified incubator with 5% CO₂. For the experiments, the cells were washed with Dulbecco’s Phosphate Buffered Saline (PBS, Bioshop, UK) and detached from the flask’s surface using 0.25% trypsin with 0.02% EDTA (Sigma-Aldrich, Poland).

3.2. Chemical substances

For the experiments, we used four compounds with antioxidative potential: (±)-catechin hydrate (Sigma, Cat. no. C1788), curcumin (Sigma-Aldrich, Cat. no. C1386), honokiol (Sigma-Aldrich, Cat. no. H4914) and cinnamon oil (Avicenna-Oil, Cat. no. 1011/5). Each drug solution was freshly prepared before the experiment. Catechin was firstly dissolved in 96% ethanol to give a stock solution of 5 mM concentration and afterwards diluted to 10 µM concentration with culture medium. Honokiol and curcumin were dissolved in dimethyl sulfoxide (DMSO, Sigma-Aldrich, Cat. no. D8418) to the 5 mM concentration and subsequently diluted to 10 µM (curcumin) and 2 µM (honokiol) concentrations with growth medium. Cinnamon was prepared by dilution in the PBS buffer to the 1000 µg/L concentration and then diluted again to 2 µg/L concentration with culture medium.

3.3. MTT assay

The viability of cells was determined using the standard MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay (MTT, Sigma-Aldrich, PL). The cells were incubated on 96-well plate (Perkin Merkel) in the concentration 10 000 cells/well for 3 hours (37°C) in 0,5 mg/ml MTT solution in PBS buffer (100 µl/well). The absorbance was measured in 0h (directly after the balloon flight), 24h, 48h and 72h incubation after balloon landing at 570 nm (Multiplate reader EnSpire, Perkin Elmer). Acidified isopropanol (100 µL/well, 0.04 M HCl in absolute isopropanol per well) was used to dissolve the formed formazan crystals.

3.4. Cell death assay

After landing, the cells were plated on 6-well plates (200 000 cells/well) and incubated for 24 hours in the appropriate drug solution. Following incubation, the culture medium was removed cells were washed with PBS buffer and detached from the surface using trypsin-EDTA solution. Subsequently, they were stained with Annexin V-APC Apoptosis Kit with PI (BioLegend, Cat. no. 640932) and analyzed with FACS Calibur flow cytometer (Becton Dickinson) in order to indicate the percentage ratio of early and late apoptotic and necrotic cells.

3.5. Membrane permeabilization assay

Fluorescent dye YO-PRO-1 (Invitrogen, Cat. no. Y3603) was used to evaluate the plasma membrane permeabilization resulting from the low-temperature and radiation cell damage during the balloon flight. The cells were seeded on black 96-well plates (Perkin Elmer) and stained in 1 µM YO-PRO-1 diluted in growth medium for 10 minutes. After 10 min. of incubation and washing with PBS, the intracellular fluorescence of YO-PRO-1 was measured with excitation wavelength of 491 nm and the emission wavelength of 509 nm. Membrane permeabilization assay was evaluated directly after balloon flight (0h) and 24h after landing (24h).

3.6. Intracellular ROS generation assay

The level of the reactive oxygen species (ROS) in cells was determined with the DCF (2,7-dichlorofluorescein) assay (Life Technologies, Poland). For experiments, the stock solution of

carboxy-H₂DCFDA (50 µg/mL in sterile DMSO; Sigma, Poland) was established at the RT in the dark and then diluted in a cell culture medium without FBS. The cells were seeded on black 96-well plates. After washing with PBS the reagent was added to the cell culture to a final concentration of 5 µM and cells were incubated at 37°C in darkness for 30 min. After the incubation, the fluorescence of DCF in wells was measured every 30 minutes for 90 minutes total (with excitation wavelength of 495 nm and the emission wavelength of 530 nm). Intracellular ROS generation assay was evaluated directly after balloon flight (0h) and 24 hours after landing (24h).

3.7. Immunocytochemical staining

Immunocytochemical staining was performed using the ABC method in order to investigate the effect of stratospheric conditions and antioxidative drugs on the expression of manganese-dependent superoxide dismutase (SOD2) in SKOV-3 and CHO-K1 cells. After landing, the cells were seeded on 10-well diagnostic microscopic slides (Thermo Fisher Scientific) and incubated for 24 hours in the appropriate drug solution. After incubation, cells were fixed and dehydrated using 4% paraformaldehyde (PFA, Sigma-Aldrich,) for 10 minutes. The, the cells were stained using the EXPOSE Mouse and Rabbit Specific HRP/DAB Detection IHC kit (Abcam, United States; Cat. no. ab80436). The enzyme expression was visualized with the mouse monoclonal antibody anti-SOD2 (Santa Cruz, USA, Cat. no. sc-362300) diluted 1:200 with the PBS buffer. After the overnight incubation with the primary antibody, cells were incubated with the secondary antibody conjugated with horseradish peroxidase (HRP). Next, samples were incubated with a diaminobenzidine-H₂O₂ mixture in order to visualize the HRP label. Between particular steps the samples were rinsed using 1% Triton X-100 in PBS. The cells were stained with hematoxylin for 3 minutes to visualize nuclei. The immunocytochemical reaction was evaluated with double-blinded method using upright microscope (Olympus BX51, Japan). Then the percentage of stained cells was estimated and the intensity of immunocytochemical reaction evaluated according the scale: (-) negative, (+) weak, (++) moderate and (+++) strong.

3.8. Clonogenic assay

Clonogenic assay is a technique allowing for the assessment of cell survival and proliferation following the exposition to the tested compounds. After balloon landing, the cells were plated in appropriate dilutions (150 cells/well) into 6-well-plates at appropriate drug solution. Multi-well-plates were placed in an incubator and left there for 7 days until large colonies (> 1 mm) were formed (50 cells or more). After incubation, growth medium was removed, and the cells were washed with PBS. Fixation and staining of clones was done with a mixture of 0.5% crystal violet in 4% paraformaldehyde (PFA, Sigma-Aldrich, USA) for 10 minutes. Then, the plates were rinsed with water and left to dry at room temperature. Counting of clones was performed the following day.

3.9. Neutral comet assay

For detection of DNA damages associated with the exposure to extreme environment during the balloon flight, neutral comet assay method described by Collins (44) was used. Directly after the balloon flight, CHO-K1 and SKOV-3 cells were suspended in freezing medium Bambanker™ (Nippon Genetics, Cat. no. BB01) (1×10^5 cells /50 μ L) and frozen for further analysis. After defrosting and centrifugation with PBS, the cells at the concentration 1×10^5 /ml were mixed with low temperature melting agarose (Sigma) at ratio 1:10 (v/v) and spread on a slide. Slides were submerged in precooled lysis solution (2.5 M NaCl, 100 mM EDTA, pH 10, 10 mM Tris base and 1 % Triton X-100) at 4°C for 60 min. After lysis and rinsing, slides were equilibrated in TBE solution (40 mM Tris/boric acid, 2 mM EDTA, pH 8.3), electrophoresed at 1.2 V/cm for 15 min and then silver staining was performed (45). For scoring the comet patterns, 200-300 nuclei from each slide were assessed. Additionally, the samples were stained using fluorescent dye Green Dead Cell Stain (Thermo Fisher Scientific, Cat. no. S34860). The images were acquired on a fluorescence microscope (Olympus Nikon). For each measurement, at least 50 cells per sample were counted and the CometScore 2.0 software was used to analyze the comets. The tail DNA percentage was taken as a quantified index of DNA damage.

3.10. Confocal microscopy and cell morphology analysis

Confocal laser scanning microscopy (CLMS) was used for the visualization of cell membrane damage and morphology. After the balloon flight, the cells were grown on coverslips for 24 hours. Subsequently, the cells were washed 3 times with PBS and quickly submerged in staining solution CellMask™ Deep Red Plasma Membrane Stains (at a concentration of 0.5 μ g/ml dissolved in growth medium; Molecular Probes, Cat. no. C10046, Ex./Em. 649/666 nm) for 10 minutes at 37°C. Next, the staining solution was removed and the coverslip were rinsed with PBS three times. After washing out with PBS, nuclear DNA was stained with DAPI (4,6-diamidino-2-phenylindole; 0.2 μ g/ml, Ex./Em. 358/461 nm). In the end the cells were mounted in fluorescence mounting medium (DAKO). For imaging, Olympus FluoView FV1000 confocal laser scanning microscope (Olympus) was used. The images were recorded by employing a Plan-Apochromat 60 \times oil-immersion objective.

3.11. Stratospheric equipment

The balloon was filled with helium and its position was tracked by APRS-supported services: www.habhub.com and www.aprs.fi. The telemetry was emitted through RTTY 70 cm/APRS 2 m signal, supported with GPS/GSM tracker. On-board computer (ATMEGA328P) was equipped with sensors located at the top wall (for UV radiation (ML8511), pressure (BMP085), and temperature (DHT22) measurements), additional UV sensors at the side wall of gondola and accelerometer (MPU6050) inside the box. To compare stratospheric and Earth UV irradiation, the UV ML8511 sensor was used which is designed especially for surface measurement of UV spectra (UVA (280-400 nm) and UVB (260-280 nm)). As it is mentioned in the specification, this sensor has a nominal operation range of -20°C to 70°C suggesting the monitoring of UV would only be valid up to roughly the minute 30 of the flight. This means that some of the random variation seen in Fig 1C may be an artefact of the electronics which is not operating adequately at

these low temperatures. It often happens that the electronic produces too much noise when it goes below certain temperature thresholds and furthermore here is exposed to moisture freezing. However, grounded analyses displayed that the sensor may be used in temperatures below -20 which cause insignificant disturbances of measurements.

3.12. Statistics

Statistical significance was determined by unpaired T-Student tests for cytotoxicity tests and two-way analysis of variance (ANOVA). Differences between treated samples and control cells with p values ≤ 0.05 were assumed to be statistically significant. The results were analyzed with the Microsoft Office Excel 2017 and GraphPad Prism 7.0 software.

4. Results

4.1. Balloon flight and stratospheric conditions analysis

The biological samples were launched to the stratosphere on 30th of April 2018, from Wrocław, Poland (51°06'23.6" N 17°03'32" E) at 11:30 AM. The balloon reached the stratosphere at maximal altitude of 30 298 m. The mission lasted about 2 hours: 90 min of ascent and 25 min of descent and ended at 1:25 PM. The biological samples were collected directly after landing in Sulisław, Poland (52°23'49.9" N 18°45'52.8" E) and transported to the laboratory. During the first stage of ascending phase recorded ambient temperature dropped to -22°C. Subsequently, when the balloon reached the ozonosphere, the ambient temperature increased to -2°C. At the highest altitude, the temperature reached the lowest level of -35°C (Fig 3A) and the lowest pressure (1252 Pa) was measured (Fig 3B). Measurements provided by two UV sensors show rapid variations of UV radiation, indicating continuous rotations of the gondola (Fig 3C). Voltage level of 1170 mV correlates with the highest score (11) in the UV Index-exposition scale, which shows extreme exposure to the UV radiation causing immediate damage of unprotected human skin and eyes (46). On the right side of the chart there is a clearly visible period when the parachute was opened, and the gondola was stabilized. In the upper parts of the atmosphere, the UV dose was greater more than two times than the dose correlating with the maximum dose in the UV-Index scale (reaching nearly 2463 mV).

Fig 3. The environmental analysis.

(A) The temperature fluctuations [°C] during the balloon flight outside the payload; the sensor was located at the top wall of gondola. (B) The pressure fluctuations [Pa] during the balloon flight, the sensor was located at the top wall of gondola. (C) UV radiation fluctuations [mV] during the balloon flight. The 1st UV sensor was located at the top wall of gondola, the 2nd UV sensor measured UV radiation at the side wall.

4.2. Cells viability

Fig 4. Cells viability verified by MTT assay after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells

protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Following the flight, we observed a decrease of the mitochondrial activity of CHO-K1 and SKOV-3 cells not protected against radiation and not incubated with antioxidants (Fig 4A, 4C). The presence of antioxidants, except for curcumin, caused increase in the CHO-K1 cells' viability. The most protective effect of antioxidants was observed in CHO-K1 cells incubated with catechin and honokiol. Studies have shown a slight increase of mitochondrial activity in normal cells after treating with cinnamon. Curcumin was the most lethal for these cells (72h - 25%) (Fig 4A). There was also observed a significant increase in the mitochondrial activity in CHO-K1 cells protected against radiation, especially after catechin and honokiol treatment (Fig 4B).

The results have shown that SKOV-3 cells not protected against radiation display similar mitochondrial activity after balloon flight - about 70% in comparison to CHO-K1 cells (Fig 4C). Furthermore, it was observed that curcumin and cinnamon were the most lethal for SKOV-3 cells, as their mitochondrial curcumin and cinnamon activity was decreasing repetitively approximately to the 18% and 30%. Other substances used in this experiment exert the less lethal effect on SKOV-3 cell line. The most protective antioxidant for that cells was catechin. However, there was no increase of mitochondrial activity in SKOV-3 cells incubated with honokiol. It was observed that curcumin significantly diminished mitochondrial metabolism in both cell lines.

The mitochondrial activity of cancer cells incubated without antioxidants and protected against radiation remained unchanged (approximately 100%) in all 72 hours (Fig 4D). Directly after the flight we observed the highest mitochondrial activity when the antioxidants were applied. A significant decrease was observed after usage of curcumin – 88% after 24 hours, 79% after 48 hours and 71% after 72 hours. However, a smaller but still significant drop was observed after cells' incubation with cinnamon. Catechin and honokiol application generated results similar to the control cells and cells without antioxidants.

4.3. Intracellular reactive oxygen species (ROS) generation

Directly after the stratospheric flight (0h), a significant decrease in intracellular ROS generation in CHO-K1 cells not protected against radiation in case of pre-treatment with catechin and honokiol was observed (Fig 5A). However, after 24 hours, an increased level of ROS was detected. In case of CHO-K1 not protected against radiation and pretreated with catechin, the lowest and relatively constant level of ROS was detected. Those results confirmed that catechin displays the most valuable protective role from all of the tested compounds.

Furthermore, directly after the balloon flight and 24h incubation there was a reduced level of oxidative stress after pretreatment with various antioxidants in CHO-K1 cells protected against

radiation measured (Fig 5B). However, it was still similar to the cells incubated in antioxidant-free medium.

Moreover, our studies revealed that in SKOV-3 cells not protected against radiation, the level of oxidative stress after pretreatment with curcumin and catechin is almost the same as in the cells incubated without antioxidant (Fig 5C). Treatment with honokiol and cinnamon resulted in the increase of ROS generation.

In the case of SKOV-3 cells protected against radiation, a higher level of ROS in cells incubated with curcumin and cinnamon was observed (Fig 5D). The results indicated that those compounds could have display antitumor activity by generating reactive oxygen species in cancer cells, which was reflected in our research. After 24 hours, the level of oxidative stress for all the tested compounds and cells without antioxidant was at 80%.

In summary, our results have shown that intracellular ROS generation is related to the exposure to stratospheric conditions, especially radiation, and treatment with the examined antioxidants which act differently in various cells.

Fig 5. Intracellular reactive oxygen species generation verified by DCF assay after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

4.4. Membrane permeabilization

Reduced membrane permeabilization is crucial due to the greater cells protection against harmful radiation and temperature shock. Directly after the stratospheric balloon flight (0h), high membrane permeabilization in CHO-K1 cells not protected against radiation was observed (Fig 6A). After 24 hours the membrane permeabilization for all samples decreased significantly. However, it was observed that the cells after preincubation with catechin have less permeable cell membranes when compared to the other samples. In the case of CHO-K1 cells not protected against radiation, a constant level of cell membrane permeabilization regardless to the type of antioxidant was observed (Fig 6B).

Our studies indicated a decreased membrane permeabilization in SKOV-3 cells not protected against radiation and incubated with antioxidants (Fig 6C). The same dependency was observed in case of SKOV-3 cells protected against UV radiation (Fig 6D). Of all the tested compounds, once again catechin showed the best protective effect due to the reduced permeability of the plasma membrane.

Fig 6 Cell membrane permeabilization determined by YO-PRO-1 fluorescence after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

4.5. Cell death

Cell death assay revealed different values of apoptotic and necrotic cells after the balloon flight. We noticed the increased percentage ratio of necrotic cells in case of CHO-K1 cells exposed to radiation in comparison to CHO-K1 cells protected against radiation (Fig 7A-B, Tab. 1, Fig 8). Incubation with catechin, honokiol or curcumin resulted in a reduced level of total apoptotic cells, whereas the cinnamon revealed the strongest proapoptotic agent.

The highest percentage ratio of late apoptotic cells was observed after curcumin treatment of CHO-K1 cells protected against radiation (Fig 7B). Interestingly, incubation with honokiol and cinnamon resulted in the increased level of early apoptotic cells. The lowest level of total apoptotic cells was observed after the catechin treatment.

SKOV-3 cells not protected against radiation showed the increased percentage ratio of apoptotic cells (Fig 7C) in comparison to the cells protected against radiation. The highest level of dead cells was observed after treatment with curcumin and cinnamon, whereas honokiol exhibited antiapoptotic properties.

In case of SKOV-3 cells protected against radiation, similar percentage ratio of apoptotic cells, for the cells both incubated with and without antioxidants was observed. (Fig 7D). The highest level of necrotic cells was revealed for the cells treated with honokiol and curcumin. However, the reduced percentage ratio of apoptotic cells was detected only for the cells incubated with cinnamon.

Fig 7. The percentage ratio of undamaged, necrotic, early and late apoptotic cells evaluated by Annexin V-APC/PI staining 24-hour after balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation.

Tab. 1. Cell death evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, after 24-hour incubation after balloon flight. Results expressed as percentage of cells.

CHO-K1 cells	undamaged	necrosis	total apoptosis	SKOV-3 cells	undamaged	necrosis	total apoptosis
<i>not protected against radiation</i>				<i>not protected against radiation</i>			
control	84	6	10	control	85	5	10
w/o antioxidant	67	12	22	w/o antioxidant	75	8	17
catechin	73	10	17	catechin	71	9	20
honokioli	71	9	20	honokioli	77	9	15
curcumin	72	10	18	curcumin	64	12	24
cinnamon	61	14	25	cinnamon	65	11	24
<i>protected against radiation</i>				<i>protected against radiation</i>			
control	84	6	10	control	85	5	10
w/o antioxidant	83	7	10	w/o antioxidant	89	3	8
catechin	84	8	8	catechin	85	4	11
honokioli	80	5	15	honokioli	86	5	9
curcumin	69	7	24	curcumin	87	5	8
cinnamon	77	9	14	cinnamon	90	4	6

Fig 8. The comparison of early and late apoptotic CHO-K1 and SKOV-3 cells protected by aluminum foil and not protected against radiation, treated with various antioxidants, after 24-hour incubation after the balloon flight.

Results are represented as dot plots.

4.6. SOD2 expression

Immunocytochemical staining method enabled detection of the significant differences in SOD2 expression after the balloon flight in both examined cell lines (Fig 9, Tab. 2). After the stratospheric flight, we observed a weaker expression of the manganese-dependent superoxide dismutase in SKOV-3 cells in comparison to CHO-K1 cells in the presence of the same antioxidant compounds. CHO-K1 cells treated with catechin and cinnamon extract were the most immunoreactive. After the balloon flight only CHO-K1 cells exhibited the increased SOD2 reactivity in comparison to control cells whereas the opposite changes were observed in SKOV-3 cells. The obtained results indicate that the examined antioxidants induced SOD2 activity in normal cells more effectively than in cancer cells.

Fig 9. The representative microphotographs of the immunoreactivity of the manganese-dependent superoxide dismutase (SOD2) in CHO-K1 and SKOV-3 cells protected and not protected against radiation.

Tab. 2. The immunocytochemical reaction with SOD2 antibody in CHO-K1 and SKOV-3 cell lines after the balloon flight in the presence of various compounds, taking into account the exposure and the protection against radiation (% - percentage of stained cells, RI – Reaction Intensity).

	control		w/o antioxidant		catechin		honokiol		curcumin		cinnamon	
	%	RI	%	RI	%	RI	%	RI	%	RI	%	RI
CHO-K1 protected against radiation	90%	+/++	100%	++	95%	++/ +++	95%	+++	100%	+++	100%	++/ +++
CHO-K1 not protected against radiation	80%	++	85%	+/++	90%	+++	100%	++/++ +	100%	++/++ +	100%	+++
SKOV-3 protected against radiation	95%	++/++ +	50%	+	70%	+/++	80%	+/++	90%	++	70%	+/++
SKOV-3 not protected against radiation	95%	++/++ +	70%	+/++	100%	+/++	95%	++	100%	++	100%	++

4.7. Clonogenic assay

The clonogenic assay confirmed the protective role of selected antioxidants which enhanced cell survival after exposure to the stratospheric environment (Fig 10, 11). Differences in number of colonies between the protected and unprotected cells indicate the cytotoxic properties of radiation. Furthermore, this assay confirmed that CHO-K1 were more sensitive to high energy particles in the stratosphere. We observed more and bigger colonies in CHO-K1 cells, especially among the protected cells. The least efficient in cell protection was curcumin whereas the highest number of colonies was observed after cinnamon and catechin pretreatment.

Comparison of MTT, cell death and clonogenic assay for SKOV-3 cells not protected against radiation and treated with cinnamon revealed a decrease of cell viability after the balloon flight, however, proliferation ability was not affected. It suggests that cinnamon first, may cause increased cell death after the flight and then promote cells' multiplying. We did not observe this effect for curcumin treatment.

Fig 10. Number of colonies after 7-day incubation.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Fig 11. Colony formation in clonogenic assay after 7 day incubation.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation, 1 – control cells, 2 – w/o antioxidant, 3 – catechin, 4 – honokiol, 5 – curcumin, 6 – cinnamon.

4.8. DNA damage in the stratosphere

In neutral comet assay we were able to distinguish four types of comets. **Figure 12** shows the detailed categorization method used for the evaluation of our samples that allows us to divide the observed nuclei into three groups: undamaged, apoptotic and intermediately damaged (**Tab. 3**). Comparison of CHO-K1 and SKOV-3 cells not protected against radiation revealed protective role of antioxidants in DNA damage among CHO-K1 cells, however, curcumin and cinnamon promoted DNA destruction in SKOV-3 cells. In the cells protected against radiation antioxidants acted differently. In the normal cells we observed more undamaged cells after preincubation with catechin and honokiol, whereas in the cancer cells cinnamon additionally reduced the percentage ratio of cells with DNA damages. These results suggest that the protective role of antioxidants in DNA damages depends on the presence of radiation resulting in various activity of these substances in cells.

Tab. 3. DNA damage evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, directly after balloon flight. Results expressed as percentage of cells.

CHO-K1 cells	undamaged	apoptotic damaged	intermediately damaged	SKOV-3 cells	undamaged	apoptotic damaged	intermediately damaged
<i>not protected against radiation</i>				<i>not protected against radiation</i>			
w/o antioxidant	79	9	12	w/o antioxidant	71	1	28
catechin	84	3	13	catechin	82	1	18
honokiol	84	1	15	honokiol	83	1	16
curcumin	91	1	9	curcumin	64	0	36
cinnamon	83	7	10	cinnamon	61	1	38
<i>protected against radiation</i>				<i>protected against radiation</i>			
w/o antioxidant	90	2	8	w/o antioxidant	54	1	45
catechin	91	0	9	catechin	74	0	25
honokiol	93	2	5	honokiol	83	1	16
curcumin	83	0	17	curcumin	52	0	48
cinnamon	89	2	8	cinnamon	70	1	29

Fig 12. Different types of DNA damage visualized by comet assay.

(A) comet type 0 and 2; (B) comet type 0, 2 and 3; (C) comet type 0 and 1; (D) comet type 0, 1 and 3. Type 0 – size of the head the normal nucleus size, tail absent; type 1 – size of the head the normal nucleus size, tail size less than normal nucleus size, type 2 – size of the head less than half size of normal nucleus, tail size more than 2 times the normal nucleus size, type 3 – size of the head the normal nucleus size, tail size about 2 times the normal nucleus size.

Furthermore, using CometScore software we were able to perform the analysis of the pictures of comets and determine the percentage ratio of the damaged DNA accompanying the flight (Fig 13). Our studies showed that cancer cells were more vulnerable to UV radiation exposure causing an increase of the presence of DNA in tail. The most active compound preventing DNA damage was cinnamon extract, both for normal and cancer cells. In the case of protected cells, honokiol and curcumin enabled the most efficient preservation.

Fig 13 DNA damage evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, directly after balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation (* $p \leq 0.05$). Results expressed as percentage of damaged DNA presented in tail of comet.

4.9. Confocal microscopy and cells' morphology analysis

Fluorescence staining revealed slightly differences in the cell morphology after the balloon flight in both examined cell lines (Fig 14). After the stratospheric flight, we observed more slender spikes in both cell lines. CHO-K1 and SKOV-3 cells treated with honokiol, curcumin and cinnamon extract were morphologically similar to the control cells. The cells exposed to radiation were enlarged, especially CHO-K1 cells incubated without antioxidant and protected against radiation. Some nuclei were fragmented, in particular in the cells not protected against radiation. However, the incubation with antioxidant resulted in an increased percentage ratio of cells with condensed nuclei with many nucleoli.

Fig 14 The representative photographs of the morphology of CHO-K1 and SKOV-3 cells protected and not protected against radiation stained with CellMask™ Deep Red (cell membranes) and DAPI (nuclei).

5. Discussion

During the balloon flight the cell samples were exposed to several fluctuating stressful factors, namely radiation, temperature, pressure and overload. However, as long as the exposure to these factors does not reach extreme levels, most of the cells are still able to survive due to their natural capability of activating a variety of specific defensive pathways. The stress response could be dramatically different among various types of cells, notably normal and cancer cells. Therefore, the application of meticulously selected compounds could either alleviate or exacerbate the consequences of the exposure to rapidly varying temperature, pressure and UV conditions during stratospheric balloon campaign – depending on their various properties and type of cells – hence influencing not only the survival but also recovery of cells after the experiment. Natural medicines, such as catechin, curcumin, cinnamon extract and honokiol have been recently gaining much attention as therapeutic compounds for cancer prevention and treatment. Our study represents the first attempt at discovering compounds exhibiting protective properties in space by launching the human normal and cancer cells into the stratosphere on a meteorological balloon.

There is a huge ambiguity among the available data concerning quantification of the absorbed dose of cosmic rays depending on the length of the flight and height reached during the meteorological balloon mission. The intensity of the radiation varies depending on parameters such as location and time unit, latitude and longitude, ozone depth and solar activity (47). The average dose of ionizing radiation reaching the atmosphere is estimated at around 2-20 μSv (48). In the stratosphere, 30 000 meters above the Earth's surface, the UV radiation is much higher than on the sea level (6). The lower the height, the thicker the protective layer of the atmosphere and the higher the radiation exposure (49). During the flight the cells were deprived of the protective effect of the ozonosphere and troposphere (which naturally protect Earth against radiation), resulting in decreased cell viability after the flight.

During the experiment half of the samples was covered by aluminum foil. Therefore, we were able to evaluate the effect of radiation (mainly UV and high energy particles such as α , β) on examined cells in the presence of various compounds. We observed an increased mitochondrial activity and decreased level of apoptotic cells after pretreatment with various antioxidants, especially among CHO-K1 cells. Catechin was the most efficient in radioprotection of normal cells, whereas honokiol appeared to be the most protective for cancer cells. These results were reflected by intracellular ROS measurements and comet assay. We did not observe significant differences in DCF fluorescent signal between SKOV-3 cells protected and not protected against radiation. However, in MTT and cell death assay and immunocytochemical staining the results were altered. It could indicate that radiation-related death in SKOV-3 cells was not strictly related to the oxidative stress and caused direct cell damage.

According to the time of cell death after the irradiation, two types of the apoptosis can be distinguished: fast apoptosis occurring straight after G2 arrest and late apoptosis, when cell death is preceded by one or more cell divisions. Widely understood radiosensitivity has been demonstrated to vary among different cell types or even populations (10). For instance, radiosensitive cells such as thymocytes (50) or lymphocytes (50) tend to undergo fast apoptosis, whereas Yanagihara et al. (1995) described human gastric epithelial tumor cells which exhibited delayed radiation-induced apoptosis scarcely 12h after irradiation, reaching maximum from 72 to 96h (51). We observed a higher percentage ratio of apoptotic cells among normal cells (CHO-K1) as compared to cancer cells (SKOV-3). Our studies suggest that normal cells are more radiosensitive to fast apoptosis than tumorous cells, which resulted in earlier and more intensive apoptosis among CHO-K1 cells. The significant differences were also shown in response to different antioxidants of each cell type, which should be highlighted as particularly interesting in case of curcumin, which acted as light-sensitizer and induced cell death more effectively in SKOV-3. Similar results were observed after treatment with catechin, which promoted apoptosis more frequently in SKOV-3 than CHO-K1. Furthermore, we noticed less and smaller SKOV-3 colonies in comparison to CHO-K1 in clonogenic assay that indicated a delayed radiation-induced death of cancer cells. Additionally, the performed tests highlighted the considerable impact of honokiol and catechin on recovery of the cells after the flight, proving cytoprotective role of antioxidant after radiation exposure.

Radiation-related DNA strand breaks, confirmed in comet assay, have been demonstrated to initiate the expression of p53 (52), which promotes cytochrome C release from the mitochondria (53) and activation of caspase cascade leading to apoptosis (54). SOD2 is the essential mitochondrial antioxidant enzyme which plays a crucial role in protection against radiation in cells (55,56) by scavenging the free radical superoxide (57). Overexpression of SOD2 stabilizes mitochondrial membrane and protect complexes I and III of the respiratory chain from radiation-induced damages (58), decreases the release of cytochrome C, resulting in the apoptosis blockage (59). Due to the intranuclear localization of *SOD2* gene, SOD2 is synthesized in the cytoplasm and subsequently imported into the mitochondria via the mitochondrial protein influx (MPI), where a manganese ion is combined with SOD2 to make active enzyme (60). The research conducted on mammalian cells has demonstrated a decrease of the mitochondrial protein import through MPI caused by radiation which leads to the accumulation of precursor proteins outside mitochondria degraded by proteasomes (61) and the reduced number of mitochondrial proteins. Furthermore, Azzam et al. (2012) revealed that MPI proteins may be damaged by irradiation, resulting in deficient protein import (62). Accordingly, mitochondria play a critical role in radioprotection by affecting SOD2 activity through Cdk1-p53-mediated SOD2 regulation. Transcriptional factors promoted by reduction of ROS production induce transcription of SOD2-regulated genes causing adaptive response to the oxidative stress. Thus, medicines that increase SOD2 activity enhance mitochondrial membrane stability and block the activation of apoptosis. Natural derivatives used in our study have been displayed as radioprotective substances (22,63,64) promoting SOD2 activity (65–68). Our studies revealed a significantly increased amount of SOD2 in normal cells treated with the analyzed antioxidants whereas we observed less viable cancer cells and increased ROS generation among cancer cells treated with these compounds. Curcumin occurred to be the most efficient radioprotective substance in normal cells while it displayed radio- and photosensitive activity in SKOV-3 cells.

According to the data recorded by on-board sensors, the temperature varied between -36°C and 20°C. Lower cultivation temperature is contributed to a decreased cellular metabolism (69), projecting on prolonged preservation of cell viability. However, moderately cold temperature conditions (-60 to -15°C) remain the main limiting factor affecting the cell viability (70). During the flight, the biological samples were exposed to this stressful temperature zone for 2 hours and experienced repeated cycle of chilling and warming while the balloon was ascending and descending (Fig 3): first when temperature dropped down to -21°C, second when it raised to -2°C and third when it dropped down to -36°C before raising to 20°C in the landing area. However, the air density in the stratosphere is about 1000 times lower than on sea level and because of that the temperature exchange between the biological samples and air was limited, resulting in the relative temperature stability of cells. Along with extensive structural damage of the cell, freezing and thawing cycles are associated with oxidative stress and consequently mitochondrial disturbance, cell membrane permeabilization or DNA damage (71), finally leading to death either on apoptotic or necrotic pathway (72,73), which was reflected in our research. The biological samples after the balloon flight were characterized with increase of ROS generation and percentage ratio of dead cells; however, treatment with selected antioxidants limited these

processes. We speculate that antioxidants could have reorganized the chemical structure of cell membrane resulting in its increased stability in lower temperature and decreased permeabilization, which was measured in YO-PRO-1 assay directly after the balloon flight and highlighted in CLSM analysis. In this process catechin was the most efficient in CHO-K1 cells. However, we noticed that membrane permeability increased after incubation with curcumin in CHO-K1 cells not protected against radiation after the flight while it was strongly reduced in SKOV-3 cells exposed to the stratospheric conditions.

A number of studies suggest that cryopreservation and low temperature affect the cell sensitivity to radiation (74). According to this, the data concerning utilization of frozen Chinese-hamster-fibroblasts revealed the decrease in X-ray-induced damage of cells stored at (-196)°C in comparison to cells irradiated at room temperature (75). Information from cryocrystallography demonstrates that very low temperatures (-173°C) stop the diffusion of free radicals caused by irradiation which leads to less harmful cell damages induced by X rays (76). Due to this fact, the increased survival of cells affected by freezing may be associated with reduced oxidative stress and damage at low temperature (74). Altogether, the data suggest that freezing can protect cells from radiation-induced damage and apoptosis, affecting cell viability after the flight.

There is a huge ambiguity between different cell types in their response to the temperature-related stress. Our experiment confirms that mechanisms of cellular responses to oxidative stress are strongly associated with the type of a cell. In general, treatment with antioxidants resulted in increased viability in normal cells whereas oxidative stress was intensified in cancer cells. However, the SOD2 expression was higher in normal cells. It shows that molecular antioxidative mechanism of different antioxidants varies in diverse types of cells: the genomic mechanism associated with the increased expression of antioxidative enzymes is predominant in normal cells whereas in cancer cells the antioxidants and their metabolites work as scavengers of free radicals and do not induce intensively the expression of protective proteins.

It is worth noting that the atmospheric pressure during the flight decreased to 1hPa. However, it is not possible to draw indisputable conclusions about the influence of the pressure on the samples due to a fact that in our research the cells were surrounded by the layer of the liquid medium, which remained frozen after the temperature drop. Despite its shortcomings, this method still provides exposure for the decreased pressure to a relatively large extent, which probably influenced the cells homeostasis and hence, the presented data. Notably, among available literature low-pressure cultivating have been proven to influence homeostasis, morphology and proliferation of mammalian cells (77), which become more rounded and less spindle-shaped and show numerous blebs around their margins. Mitochondria and nucleus take more rounded shape due to edema and cell membrane creating bubbles causing cell death (78).

Remarkably, stratospheric balloon flights are accompanied with overloads. Accelerometer in the gondola shows that ascending and descending velocity is not permanent and because of that the biological samples were exposed to overloads. Numerous studies have investigated the effect of overloads on cells. For example, when animal cells are cultured under 10g (10g means 10 times higher force than Earth gravity), proliferation is increased by 20-30%, glucose consumption is

reduced, cell migration is inhibited by high overloads. At gravitational stress, cell may move to other metabolic pathways (79). On the other hand, overloads work as a natural antioxidant and reduce lipid membrane peroxidation (LPO). Induction of the passive mechanisms of biomembrane protection associated with changes in the phase status of the membrane is the most plausible explanation for the phenomenon being observed (80). In our experiment we noted changes in membrane permeability, which could have been affected by overloads. Zhan et al. (1999) revealed a protective effect of green tea polyphenols on 10g stress in rats (81) and our studies highlighted the value of natural antioxidants as protective agents in the gravitational stress. Furthermore, altered gravity deeply influences numerous biological processes in living organisms. Changes in gravitational values affect cell survival, development, and spatial organization. In addition, the indirect effects of altered gravity, such as those associated with hydrostatic pressure and fluid shear, strongly affect both *in vitro* and *in vivo* systems (82). Hence, in order to understand these phenomena, further research is necessary.

5.1. Conclusions

In our study, we analyzed changes in the functions of normal and cancer cells that occurred due to exposure to high radiation, overloads as well as low temperature and pressure during stratospheric flight. Our work has led us to conclude that the application of the carefully selected medicines enables us to manipulate cellular stress response depending on the type of cells. Altogether, these findings suggest that honokiol and catechin have the best protective effect on the normal cells, whereas curcumin and cinnamon act as radio- and light-sensitizers increasing the percentage of apoptotic cancer cells and DNA damage. The results constitute a significant step towards the investigation of possible strategies for the cell protection in space environment and provide new insights into the application of the examined compounds for the prevention and treatment of cancer. We believe that our research will remain valuable for resolving the difficulty of the human and biological material protection in space. Because of its relatively low costs, our approach remains economic alternative for simulated subcosmic conditions conducted in the laboratory, which requires far more expensive, specialized measurements.

6. Acknowledgments

The work was created as part of the activity of the Student Research Group "Biology of Cancer Cell" at the Wrocław Medical University (SKN No. K 148). The research was supported partially by the Statutory Funds of Wrocław Medical University no ST.E130.16.060 (manager prof. J. Zalewski), funds of Wrocław University of Science and Technology and by "Najlepsi z Najlepszyc 3.0" program of Polish Ministry of Science and Higher Education. We would like to thank Ms. E. Przydatek for help with the correction of the English language in the manuscript, Mr. J. Skórniak and Mr. W. Tarnowski for organization the balloon flight and transport of the biological samples.

7. References

1. Petrescu FI, Petrescu RV. The Aviation History. BoD–Books on Demand; 2012.
2. Jones WV. Evolution of scientific ballooning and its impact on astrophysics research. *Adv Sp Res*. 2014;53(10):1405–14.
3. Smith DJ, Sowa MB. Ballooning for Biologists: Mission Essentials for Flying Life Science Experiments to Near Space. *Gravitational Sp Res*. 2017;5(1).
4. Murgatroyd RJ. The physics and dynamics of the stratosphere and mesosphere. *Reports Prog Phys*. 1970;33(3):817–80.
5. Reitz G. Radiation Environment in the Stratosphere. *Radiat Prot Dosimetry* [Internet]. 1993 May 1 [cited 2018 Nov 3];48(1):5–20. Available from: <https://academic.oup.com/rpd/article/48/1/5/1611190/Radiation-Environment-in-the-Stratosphere>
6. Cadet J, Sage E, Douki T. Ultraviolet radiation-mediated damage to cellular DNA. *Mutat Res - Fundam Mol Mech Mutagen*. 2005;571(1–2 SPEC. ISS.):3–17.
7. Moan J, Peak MJ. Effects of UV radiation on cells. *J Photochem Photobiol B Biol*. 1989;4(1):21–34.
8. Desouky O, Ding N, Zhou G. Targeted and non-targeted effects of ionizing radiation. *J Radiat Res Appl Sci* [Internet]. 2015;8(2):247–54. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1687850715000333>
9. Anand AJ, Dzik WH, Imam A, Sadrzadeli SMH. Radiation-induced red cell damage: role of reactive oxygen species. *Transfusion*. 1997;37(2):160–5.
10. Cohen-Jonathan E, Bernhard EJ, McKenna WG. How does radiation kill cells? *Curr Opin Chem Biol*. 1999;3(1):77–83.
11. Narumi K, Sonoda JI, Shiotani K, Shigeru M, Shibata M, Kawachi A, et al. Simultaneous detection of green tea catechins and gallic acid in human serum after ingestion of green tea tablets using ion-pair high-performance liquid chromatography with electrochemical detection. *J Chromatogr B Anal Technol Biomed Life Sci* [Internet]. 2014;945–946:147–53. Available from: <http://dx.doi.org/10.1016/j.jchromb.2013.11.007>
12. Song Y, Sun H, Zhang A, Yan G, Han Y, Wang X. Plant-derived natural products as leads to anti-cancer drugs. *J Med Plant Herb Ther Res*. 2014;2:6–15.
13. Hosseini-mehr SJ, Rostamnejad M, Ghaffari-rad V. Epicatechin enhances anti-proliferative effect of bleomycin in ovarian cancer cell. *Res Mol Med*. 2013;1:24–7.
14. Vergne MJ, Patras A, Bhullar MS, Shade LM, Sasges M, Rakariyatham K, et al. UV-C Irradiation on the Quality of Green Tea : Effect on Catechins , Antioxidant Activity , and Cytotoxicity. 2018;0:1–7.
15. Li Q, Kakizaki M, Kuriyama S, Sone T, Yan H, Nakaya N, et al. Green tea consumption and lung cancer risk: the Ohsaki study. *Br J Cancer* [Internet]. 2008 Oct 2 [cited 2018 Mar 25];99(7):1179–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18766189>

16. Kang H, Rha SY, Oh KW, Nam CM. Green tea consumption and stomach cancer risk: a meta-analysis. *Epidemiol Health* [Internet]. 2010 Apr 26 [cited 2018 Mar 25];32:e2010001. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21191454>
17. Ni C-X, Gong H, Liu Y, Qi Y, Jiang C-L, Zhang J-P. Green Tea Consumption and the Risk of Liver Cancer: A Meta-Analysis. *Nutr Cancer* [Internet]. 2017 Feb 17 [cited 2018 Mar 25];69(2):211–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28095030>
18. Yang G, Zheng W, Xiang Y-B, Gao J, Li H-L, Zhang X, et al. Green tea consumption and colorectal cancer risk: a report from the Shanghai Men's Health Study. *Carcinogenesis* [Internet]. 2011 Nov [cited 2018 Mar 25];32(11):1684–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21856996>
19. Shrubsole MJ, Lu W, Chen Z, Shu XO, Zheng Y, Dai Q, et al. Drinking Green Tea Modestly Reduces Breast Cancer Risk. *J Nutr* [Internet]. 2009 Feb 1 [cited 2018 Mar 25];139(2):310–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19074205>
20. Kurahashi N, Sasazuki S, Iwasaki M, Inoue M, Tsugane S, JPHC Study Group. Green Tea Consumption and Prostate Cancer Risk in Japanese Men: A Prospective Study. *Am J Epidemiol* [Internet]. 2007 Nov 15 [cited 2018 Mar 25];167(1):71–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17906295>
21. Prasad R, Singh T, Katiyar SK. Honokiol inhibits ultraviolet radiation-induced immunosuppression through inhibition of ultraviolet-induced inflammation and DNA hypermethylation in mouse skin. *Sci Rep* [Internet]. 2017;7(1):1–14. Available from: <http://dx.doi.org/10.1038/s41598-017-01774-5>
22. Chilampalli S, Zhang X, Fahmy H, Kaushik RS, Zeman D, Hildreth MB, et al. Chemopreventive effects of honokiol on UVB-induced skin cancer development. *Anticancer Res* [Internet]. 2010 Mar [cited 2018 May 10];30(3):777–83. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20392996>
23. Vaid M, Sharma SD, Katiyar SK. Honokiol, a phytochemical from the Magnolia plant, inhibits photocarcinogenesis by targeting UVB-induced inflammatory mediators and cell cycle regulators: development of topical formulation. *Carcinogenesis* [Internet]. 2010 Nov 1 [cited 2018 May 10];31(11):2004–11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20823108>
24. Chilampalli C, Guillermo R, Zhang X, Kaushik RS, Young A, Zeman D, et al. Effects of magnolol on UVB-induced skin cancer development in mice and its possible mechanism of action. *BMC Cancer* [Internet]. 2011 Oct 20 [cited 2018 May 10];11:456. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22014088>
25. Ni H, Jin W, Zhu T, Wang J, Yuan B, Jiang J, et al. Curcumin modulates TLR4/NF-κB inflammatory signaling pathway following traumatic spinal cord injury in rats. *J Spinal Cord Med* [Internet]. 2015 Mar [cited 2018 Mar 26];38(2):199–206. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24621048>
26. Koosirirat C, Linpisarn S, Changsom D, Chawansuntati K, Wipasa J. Investigation of the anti-inflammatory effect of *Curcuma longa* in *Helicobacter pylori*-infected patients. *Int Immunopharmacol* [Internet]. 2010 Jul [cited 2018 Mar 26];10(7):815–8. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/20438867>

27. Moghadamtousi SZ, Kadir HA, Hassandarvish P, Tajik H, Abubakar S, Zandi K. A review on antibacterial, antiviral, and antifungal activity of curcumin. *Biomed Res Int* [Internet]. 2014 [cited 2018 Mar 26];2014:186864. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24877064>
28. Sinha D, Biswas J, Sung B, B. Aggarwal B, Bishayee A. Chemopreventive and Chemotherapeutic Potential of Curcumin in Breast Cancer. *Curr Drug Targets* [Internet]. 2012 Dec 31 [cited 2018 Mar 26];13(14):1799–819. Available from: <http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1389-4501&volume=13&issue=14&spage=1799>
29. Wang Y, Rishi AK, Wu W, Polin L, Sharma S, Levi E, et al. Curcumin suppresses growth of mesothelioma cells in vitro and in vivo, in part, by stimulating apoptosis. *Mol Cell Biochem* [Internet]. 2011 Nov [cited 2018 Mar 26];357(1–2):83–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21594647>
30. Darvesh AS, Carroll RT, Bishayee A, Novotny NA, Geldenhuys WJ, Van der Schyf CJ. Curcumin and neurodegenerative diseases: a perspective. *Expert Opin Investig Drugs* [Internet]. 2012 Aug 6 [cited 2018 Mar 26];21(8):1123–40. Available from: <http://www.tandfonline.com/doi/full/10.1517/13543784.2012.693479>
31. Khanna S, Park H-A, Sen CK, Golakoti T, Sengupta K, Venkateswarlu S, et al. Neuroprotective and antiinflammatory properties of a novel demethylated curcuminoid. *Antioxid Redox Signal* [Internet]. 2009 Mar [cited 2018 Mar 26];11(3):449–68. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18724833>
32. S. Darvesh A, B. Aggarwal B, Bishayee A. Curcumin and Liver Cancer: A Review. *Curr Pharm Biotechnol* [Internet]. 2012 Jan 1 [cited 2018 Mar 26];13(1):218–28. Available from: <http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1389-2010&volume=13&issue=1&spage=218>
33. Zhang J, Wang J, Xu J, Lu Y, Jiang J, Wang L, et al. Curcumin targets the TFEB-lysosome pathway for induction of autophagy. *Oncotarget* [Internet]. 2016 Nov 15 [cited 2018 Mar 26];7(46):75659–71. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27689333>
34. Hsieh Y-W, Huang C-Y, Yang S-Y, Peng Y-H, Yu C-P, Chao P-DL, et al. Oral intake of curcumin markedly activated CYP 3A4: in vivo and ex-vivo studies. *Sci Rep* [Internet]. 2014 Oct 10 [cited 2018 Mar 26];4:6587. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25300360>
35. Das L, Vinayak M. Long term effect of curcumin in restoration of tumour suppressor p53 and phase-II antioxidant enzymes via activation of Nrf2 signalling and modulation of inflammation in prevention of cancer. *PLoS One* [Internet]. 2015 [cited 2018 Mar 26];10(4):e0124000. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25860911>
36. Menon VP, Sudheer AR. Antioxidant and anti-inflammatory properties of curcumin. In: *The Molecular Targets and Therapeutic Uses of Curcumin in Health and Disease* [Internet]. Boston, MA: Springer US; 2007 [cited 2018 Mar 26]. p. 105–25. Available

from: <http://www.ncbi.nlm.nih.gov/pubmed/17569207>

37. Lin YG, Kunnumakkara AB, Nair A, Merritt WM, Han LY, Armaiz-Pena GN, et al. Curcumin Inhibits Tumor Growth and Angiogenesis in Ovarian Carcinoma by Targeting the Nuclear Factor- B Pathway. *Clin Cancer Res* [Internet]. 2007 Jun 1 [cited 2018 Mar 26];13(11):3423–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17545551>
38. Marchiani A, Rozzo C, Fadda A, Delogu G, Ruzza P. Curcumin and curcumin-like molecules: from spice to drugs. *Curr Med Chem* [Internet]. 2014 [cited 2018 Mar 26];21(2):204–22. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23590716>
39. Choi G-Y, Kim H-B, Hwang E-S, Lee S, Kim M-J, Choi J-Y, et al. Curcumin Alters Neural Plasticity and Viability of Intact Hippocampal Circuits and Attenuates Behavioral Despair and COX-2 Expression in Chronically Stressed Rats. *Mediators Inflamm* [Internet]. 2017 [cited 2018 Mar 26];6280925. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28167853>
40. Hewlings S, Kalman D. Curcumin: A Review of Its' Effects on Human Health. *Foods* [Internet]. 2017;6(10):92. Available from: <http://www.mdpi.com/2304-8158/6/10/92>
41. Lin YG, Kunnumakkara AB, Nair A, Merritt WM, Han LY, Armaiz-Pena GN, et al. Curcumin Inhibits Tumor Growth and Angiogenesis in Ovarian Carcinoma by Targeting the Nuclear Factor- B Pathway. *Clin Cancer Res*. 2007 Jun;13(11):3423–30.
42. Verwanger T, Krammer B, Bernardinelli E. Curcumin as a photosensitizer: Studies on different cell lines. *Photodiagnosis Photodyn Ther* [Internet]. 2011 Jun 1 [cited 2018 Mar 26];8(2):186. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1572100011002420>
43. Azab KS, Mostafa AHA, Ali EMM, Abdel-Aziz MAS. Cinnamon extract ameliorates ionizing radiation-induced cellular injury in rats. *Ecotoxicol Environ Saf* [Internet]. 2011;74(8):2324–9. Available from: <http://dx.doi.org/10.1016/j.ecoenv.2011.06.016>
44. Collins AR. The Comet Assay: Principles, Applications, and Limitations. In: *In Situ Detection of DNA Damage* [Internet]. New Jersey: Humana Press; 2002 [cited 2018 Jul 7]. p. 163–77. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12073440>
45. Nadin SB, Vargas-Roig LM, Ciocca DR. A Silver Staining Method for Single-cell Gel Assay. *J Histochem Cytochem* [Internet]. 2001 Sep 26 [cited 2018 Jul 7];49(9):1183–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11511687>
46. Fioletov V, Kerr JB, Fergusson A. The UV index: Definition, distribution and factors affecting it. *Can J Public Heal*. 2010;101(4):15–9.
47. J. C, T. B, Wiltberger M, Kent W, Grajewski B, Xu X. Atmospheric Ionizing Radiation from Galactic and Solar Cosmic Rays. *Curr Top Ioniz Radiat Res* [Internet]. 2012;(March 2017). Available from: <http://www.intechopen.com/books/current-topics-in-ionizing-radiation-research/atmospheric-ionizing-radiation-from-galactic-and-solar-cosmic-rays>
48. J. C, T. B, Wiltberger M, Kent W, Grajewski B, Xu X. Atmospheric Ionizing Radiation from Galactic and Solar Cosmic Rays. In: *Current Topics in Ionizing Radiation Research* [Internet]. InTech; 2012. Available from: <http://cdn.intechopen.com/pdfs-wm/32114.pdf>

49. Wasiak T. Promieniowanie kosmiczne i jego wpływ na organizm pilota. 2002.
50. Yamaguchi T. Relationship between survival period and dose of irradiation in rat thymocytes in vitro. *Int J Radiat Biol Relat Stud Phys Chem Med* [Internet]. 1967;12(3):235–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/5299143>
51. Yanagihara K, Nii M, Numoto M, Kamiya K, Tauchi H, Sawada S, et al. Radiation-induced apoptotic cell death in human gastric epithelial tumour cells; correlation between mitotic death and apoptosis. *Int J Radiat Biol* [Internet]. 1995 Jun;67(6):677–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7608631>
52. Mihara M, Erster S, Zaika A, Petrenko O, Chittenden T, Pancoska P, et al. p53 has a direct apoptogenic role at the mitochondria. *Mol Cell* [Internet]. 2003 Mar [cited 2018 Jun 20];11(3):577–90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12667443>
53. Hosoki A, Yonekura S-I, Zhao Q-L, Wei Z-L, Takasaki I, Tabuchi Y, et al. Mitochondria-targeted superoxide dismutase (SOD2) regulates radiation resistance and radiation stress response in HeLa cells. *J Radiat Res* [Internet]. 2012 [cited 2018 Jun 20];53(1):58–71. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22302046>
54. Garland JM, Rudin C. Cytochrome c induces caspase-dependent apoptosis in intact hematopoietic cells and overrides apoptosis suppression mediated by bcl-2, growth factor signaling, MAP-kinase-kinase, and malignant change. *Blood* [Internet]. 1998 Aug 15 [cited 2018 Jun 20];92(4):1235–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9694712>
55. Epperly MW, Bray JA, Esocobar P, Bigbee WL, Watkins S, Greenberger JS. Overexpression of the human manganese superoxide dismutase (MnSOD) transgene in subclones of murine hematopoietic progenitor cell line 32D cl 3 decreases irradiation-induced apoptosis but does not alter G2/M or G1/S phase cell cycle arrest. *Radiat Oncol Investig* [Internet]. 1999 [cited 2018 Jun 20];7(6):331–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10644056>
56. Hirose K, Longo DL, Oppenheim JJ, Matsushima K. Overexpression of mitochondrial manganese superoxide dismutase promotes the survival of tumor cells exposed to interleukin-1, tumor necrosis factor, selected anticancer drugs, and ionizing radiation. *FASEB J* [Internet]. 1993 Feb 1 [cited 2018 Jun 20];7(2):361–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8440412>
57. Candas D, Li JJ. MnSOD in oxidative stress response-potential regulation via mitochondrial protein influx. *Antioxid Redox Signal* [Internet]. 2014 Apr 1 [cited 2018 Jun 20];20(10):1599–617. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23581847>
58. Pearce LL, Epperly MW, Greenberger JS, Pitt BR, Peterson J. Identification of Respiratory Complexes I and III as Mitochondrial Sites of Damage Following Exposure to Ionizing Radiation and Nitric Oxide. *Nitric Oxide* [Internet]. 2001 Apr 1 [cited 2018 Jun 20];5(2):128–36. Available from: <https://www.sciencedirect.com/science/article/pii/S1089860301903386>
59. Epperly MW, Sikora CA, DeFilippi SJ, Gretton JA, Zhan Q, Kufe DW, et al. Manganese

- Superoxide Dismutase (SOD2) Inhibits Radiation-Induced Apoptosis by Stabilization of the Mitochondrial Membrane. [http://dx.doi.org/10.1667/0033-7587\(2002\)157\[0568:MSDSIR\]20CO;2](http://dx.doi.org/10.1667/0033-7587(2002)157[0568:MSDSIR]20CO;2) [Internet]. 2009 Jan 24 [cited 2018 Jun 20]; Available from: [http://www.bioone.org/doi/abs/10.1667/0033-7587\(2002\)157%5B0568:MSDSIR%5D2.0.CO;2](http://www.bioone.org/doi/abs/10.1667/0033-7587(2002)157%5B0568:MSDSIR%5D2.0.CO;2)
60. Wispé JR, Clark JC, Burhans MS, Kropp KE, Korfhagen TR, Whitsett JA. Synthesis and processing of the precursor for human manganese-superoxide dismutase. *Biochim Biophys Acta* [Internet]. 1989 Jan 19 [cited 2018 Jun 20];994(1):30–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2462451>
61. Wright G, Terada K, Yano M, Sergeev I, Mori M. Oxidative Stress Inhibits the Mitochondrial Import of Preproteins and Leads to Their Degradation. *Exp Cell Res* [Internet]. 2001 Feb 1 [cited 2018 Jun 20];263(1):107–17. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11161710>
62. Azzam EI, Jay-Gerin J-P, Pain D. Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Cancer Lett* [Internet]. 2012 Dec 31 [cited 2018 Jun 20];327(1–2):48–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22182453>
63. Sevgi K, Tepe B, Sarikurku C. Antioxidant and DNA damage protection potentials of selected phenolic acids. *Food Chem Toxicol* [Internet]. 2015 Mar 1 [cited 2018 Jun 20];77:12–21. Available from: <https://www.sciencedirect.com/science/article/pii/S0278691514005055>
64. Goel A, Aggarwal BB. Curcumin, the Golden Spice From Indian Saffron, Is a Chemosensitizer and Radiosensitizer for Tumors and Chemoprotector and Radioprotector for Normal Organs. *Nutr Cancer* [Internet]. 2010 Sep 23 [cited 2018 Jun 20];62(7):919–30. Available from: <http://www.tandfonline.com/doi/abs/10.1080/01635581.2010.509835>
65. Khristi V, Elias J, Dave N, Patel V. Gene Expression Analysis of Anti-oxidative Enzymes in Yeast against Oxidative Stress in Presence of *Cinnamomum zeylanicum*. *Res J Pharm , Biol Chem Sci*. 2014;5(1387):1387–96.
66. Evacuasiyany E, Ratnawati H, Liana LK, Widowati W, Maesaroh M, Mozef T, et al. Cytotoxic and antioxidant activities of catechins in inhibiting the malignancy of breast cancer. *Oxid Antioxid Med Sci* [Internet]. 2014 [cited 2018 Jun 20];3(2):141–6. Available from: <http://www.oamsjournal.com>
67. Huang L, Zhang K, Guo Y, Huang F, Yang K, Chen L, et al. Honokiol protects against doxorubicin cardiotoxicity via improving mitochondrial function in mouse hearts. *Sci Rep* [Internet]. 2017 Dec 20 [cited 2018 Jun 20];7(1):11989. Available from: <http://www.nature.com/articles/s41598-017-12095-y>
68. Zhai X, Qiao H, Guan W, Li Z, Cheng Y, Jia X, et al. Curcumin regulates peroxisome proliferator-activated receptor- γ coactivator-1 α expression by AMPK pathway in hepatic stellate cells in vitro. *Eur J Pharmacol* [Internet]. 2015 Jan 5 [cited 2018 Jun 20];746:56–62. Available from: <https://www.sciencedirect.com/science/article/pii/S0014299914007626>
69. Ducommun P, Ruffieux P-A, Kadouri A, von Stockar U, Marison IW. Monitoring of

- temperature effects on animal cell metabolism in a packed bed process. *Biotechnol Bioeng* [Internet]. 2002 Mar 30;77(7):838–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11835145>
70. Gao D, Critser AK. Mechanisms of Cryoinjury in Living Cells. *ILAR J* [Internet]. 2000;41(4):187–96. Available from: <https://academic.oup.com/ilarjournal/article/41/4/187/716700/Mechanisms-of-Cryoinjury-in-Living-Cells#36175702>
71. Karimi-busheri F, Rasouli-nia A, Weinfeld M. Biobanking and Cryopreservation of Stem Cells. 2016;951. Available from: <http://link.springer.com/10.1007/978-3-319-45457-3>
72. Savitskaya MA, Onishchenko GE. Apoptosis in Cryopreserved Eukaryotic Cells. *Biochemistry (Mosc)* [Internet]. 2016;81(5):445–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27297895>
73. Baust JG, Gao D, Baust JM. Cryopreservation: An emerging paradigm change. Vol. 5, *Organogenesis*. 2009. p. 90–6.
74. Cugia G, Centis F, Del Zotto G, Lucarini A, Argazzi E, Zini G, et al. High survival of frozen cells irradiated with gamma radiation. *Radiat Prot Dosimetry* [Internet]. 2011 Feb 1 [cited 2018 Jun 20];143(2–4):237–40. Available from: <https://academic.oup.com/rpd/article-lookup/doi/10.1093/rpd/ncq541>
75. Ashwood-Smith MJ, Friedmann GB. Lethal and chromosomal effects of freezing, thawing, storage time, and x-irradiation on mammalian cells preserved at -196° in dimethyl sulfoxide. *Cryobiology* [Internet]. 1979 Apr 1 [cited 2018 Jun 20];16(2):132–40. Available from: <https://www.sciencedirect.com/science/article/pii/0011224079900233?via%3Dihub>
76. Garman E. ‘Cool’ crystals: macromolecular cryocrystallography and radiation damage. *Curr Opin Struct Biol* [Internet]. 2003 Oct [cited 2018 Jun 20];13(5):545–51. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14568608>
77. Watase M, Awolesi MA, Ricotta J, Sumpio BE. Effect of pressure on cultured smooth muscle cells. *Life Sci*. 1997;61(10):987–96.
78. Lyon NC, Kraus SE, Kirsh RL, Parsons DF. Morphological Changes Induced in Cultured Mammalian Cells by Low Pressure Atmospheres. *Trans Am Microsc Soc* [Internet]. 1976 Jan [cited 2018 May 13];95(1):116. Available from: <https://www.jstor.org/stable/3225362?origin=crossref>
79. Tschopp A, Cogoli A. Hypergravity promotes cell proliferation. *Experientia*. 1983;39(12):1323–9.
80. Markin AA. Lipid peroxidation and the antioxidant protection system in humans exposed to hypergravity of varying intensity. *Hum Physiol* [Internet]. 2016 Sep 18 [cited 2018 May 10];42(5):546–9. Available from: <http://link.springer.com/10.1134/S0362119716050108>
81. Zhan H, Chen LM, Xin YM, Tang GX, Wen J. Effects of tea polyphenols on cerebral lipid peroxidation, liver and renal functions in rats after repeated +Gz stress. *Space Med Med Eng (Beijing)* [Internet]. 1999 Feb [cited 2018 May 10];12(1):1–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11765769>

82. Genchi GG, Rocca A, Marino A, Grillone A, Mattoli V, Ciofani G. Hypergravity as a Tool for Cell Stimulation: Implications in Biomedicine. *Front Astron Sp Sci* [Internet]. 2016;3(August):1–8. Available from: <http://journal.frontiersin.org/article/10.3389/fspas.2016.00026/abstract>

8. Author Disclosure Statement

D.P. was responsible for project administration. D.P., A.G., P.R. and W.B. wrote the original draft and performed most of investigations. O.M. helped with the formal analysis. J.R. performed flow cytometry (investigation). A.S. was responsible for CLSM (investigation). M.DZ. and P.K. helped with comet assay (investigation). J.G. was responsible for stratospheric conditions measurements (software). J.K. was the supervisor of the experiment and the main reviewer. All authors reviewed the manuscript. No competing financial interests exist.

9. Figures legend

Fig 1. Effect of the radiation on the DNA.

High-energy particles could affect DNA damage either by direct pathway or indirect pathway. Free radicals and by-products of the process of water radiolysis contribute to an increase of the oxidative stress in the cell affecting the DNA. Survival of the cell depends on the extent of the DNA damage (8–10). This figure was prepared using Servier Medical Art, available from www.servier.com/Powerpoint-image-bank.

Fig 2. The schematic representation of the procedure of balloon flight and the preparation of biological samples.

This figure was prepared using Servier Medical Art, available from www.servier.com/Powerpoint-image-bank.

Fig 3. The environmental analysis.

(A) The temperature fluctuations [°C] during the balloon flight outside the payload; the sensor was located at the top wall of gondola. (B) The pressure fluctuations [Pa] during the balloon flight, the sensor was located at the top wall of gondola. (C) UV radiation fluctuations [mV] during the balloon flight. The 1st UV sensor was located at the top wall of gondola, the 2nd UV sensor measured UV radiation at the side wall.

Fig 4. Cells viability verified by MTT assay after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Fig 5. Intracellular reactive oxygen species generation verified by DCF assay after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Fig 6 Cell membrane permeabilization determined by YO-PRO-1 fluorescence after the balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells protected by aluminum foil against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Fig 7. The percentage ratio of undamaged, necrotic, early and late apoptotic cells evaluated by Annexin V-APC/PI staining 24-hour after balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by the aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected by aluminum foil against radiation.

Fig 8. The comparison of early and late apoptotic CHO-K1 and SKOV-3 cells protected by aluminum foil and not protected against radiation, treated with various antioxidants, after 24-hour incubation after the balloon flight.

Results are represented as dot plots.

Fig 9. The representative microphotographs of the immunoreactivity of the manganese-dependent superoxide dismutase (SOD2) in CHO-K1 and SKOV-3 cells protected and not protected against radiation.

Fig 10. Number of colonies after 7-day incubation.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected by aluminum foil against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation (* $p \leq 0.05$). Data are presented as the mean percentage relative to control cells, which were not sent into the stratosphere.

Fig 11. Colony formation in clonogenic assay after 7 day incubation.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation, 1 – control cells, 2 – w/o antioxidant, 3 – catechin, 4 – honokiol, 5 – curcumin, 6 – cinnamon.

Fig 12. Different types of DNA damage visualized by comet assay.

(A) comet type 0 and 2; (B) comet type 0, 2 and 3; (C) comet type 0 and 1; (D) comet type 0, 1 and 3. Type 0 – size of the head the normal nucleus size, tail absent; type 1 – size of the head the

normal nucleus size, tail size less than normal nucleus size, type 2 – size of the head less than half size of normal nucleus, tail size more than 2 times the normal nucleus size, type 3 – size of the head the normal nucleus size, tail size about 2 times the normal nucleus size.

Fig 13 DNA damage evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, directly after balloon flight.

(A) CHO-K1 cells not protected against radiation, (B) CHO-K1 cells protected against radiation, (C) SKOV-3 cells not protected against radiation, (D) SKOV-3 cells protected against radiation (* $p \leq 0.05$). Results expressed as percentage of damaged DNA presented in tail of comet.

Fig 14 The representative photographs of the morphology of CHO-K1 and SKOV-3 cells protected and not protected against radiation stained with CellMask™ Deep Red (cell membranes) and DAPI (nuclei).

10. Tables

Tab. 1. Cell death evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, after 24-hour incubation after balloon flight. Results expressed as percentage of cells.

Tab. 2. The immunocytochemical reaction with SOD2 antibody in CHO-K1 and SKOV-3 cell lines after the balloon flight in the presence of various compounds, taking into account the exposure and the protection against radiation (% - percentage of stained cells, RI – Reaction Intensity).

Tab. 3. DNA damage evaluation of CHO-K1 and SKOV-3 cells protected and not protected against radiation, treated with various antioxidants, directly after balloon flight. Results expressed as percentage of cells.

BEFORE THE FLIGHT

CELL CULTURE



- SKOV-3
- CHO-K1

TREATMENT WITH ANTIOXIDANTS



24 h before the flight

- 1 - control
- 2 - w/o antioxidant
- 3 - catechin
- 4 - honokiol
- 5 - curcumin
- 6 - cinnamon

SAMPLE PREPARATION




1. Exposed to radiation
2. Protected against radiation
(by the aluminum foil)

BALLOON FLIGHT



- Time of flight: 2 h
- Max. altitude: 30 397 m
- T range: 20 °C - (-36 °C)
- The lowest pressure: 1 hPa
- Distance: 76 km

The biological samples



Box with the stratospheric sensors

AFTER THE FLIGHT

MTT assay

Viability

0h, 24h, 48h, 72h

DCF assay

Intracellular ROS generation

0h, 24h

YO-PRO-1 assay

Membrane permeabilization

0h, 24h



24h

Cell death assay

Apoptosis and necrosis

168h

0h

Immuno-staining

SOD2 expression

24h

Clonogenic assay

Long-term viability

Neutral comet assay

DNA damage

Figure 2




A**B****C**

Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

CHO-K1 cells

SKOV-3 cells

not protected against radiation

protected against radiation

not protected against radiation

protected against radiation

control


w/o antioxidant

catechin

honokiol

curcumin

cinnamon



bioRxiv preprint doi: <https://doi.org/10.1101/526376>; this version posted January 25, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 8

**CHO-K1 protected
against radiation**

**CHO-K1 not protected
against radiation**

**SKOV-3 protected
against radiation**

**SKOV-3 not protected
against radiation**

control

w/o antioxidant

catechin

honokiol

curcumin

cinnamon




Figure 9




Figure 10







Figure 11







Figure 12




Figure 13




Figure 1




Figure 14