Viral delivery of tissue nonspecific alkaline phosphatase

diminishes craniosynostosis in one of two FGFR2$^{C342Y/+}$ mouse

models of Crouzon syndrome.

Hwa Kyung Nam[1], Iva Vesela[1], Sara Dean Schutte[1], Nan E. Hatch[1*]

[1] Department of Orthodontics and Pediatric Dentistry, School of Dentistry, University of
Michigan, Ann Arbor, Michigan, United States of America
9

10 * Corresponding author
11 E-mail: nhatch@umich.edu
12

Abstract

Craniosynostosis is the premature fusion of cranial bones. The goal of this study was to determine if delivery of recombinant tissue nonspecific alkaline phosphatase (TNAP) could prevent or diminish the severity of craniosynostosis in a C57BL/6 FGFR2$^{C342Y/+}$ model of neonatal onset craniosynostosis or a BALB/c FGFR2$^{C342Y/+}$ model of postnatal onset craniosynostosis. Mice were injected with a lentivirus encoding a mineral targeted form of TNAP immediately after birth. Cranial bone fusion as well as cranial bone volume, mineral content and density were assessed by micro computed tomography. Craniofacial shape was measured with calipers. Alkaline phosphatase, alanine amino transferase (ALT) and aspartate amino transferase (AST) activity levels were measured in serum. Neonatal delivery of TNAP diminished craniosynostosis severity from 94% suture obliteration in vehicle treated mice to 67% suture obliteration in treated mice, $p<0.02$) and the incidence of malocclusion from 82.4% to 34.7% ($p<0.03$), with no effect on cranial bone in C57BL/6 FGFR2$^{C342Y/+}$ mice. In contrast, treatment with TNAP improved cranial bone volume ($p< 0.01$), density ($p< 0.01$) and mineral content ($p< 0.01$) but had no effect on craniosynostosis or malocclusion in BALB/c FGFR2$^{C342Y/+}$ mice. These results indicate that perinatal recombinant TNAP enzyme therapy diminishes craniosynostosis severity in the C57BL/6 FGFR2$^{C342Y/+}$ neonatal onset mouse model of Crouzon syndrome, and that effects of exogenous TNAP are genetic background dependent.

Introduction

Skull growth occurs via intramembranous bone deposition along the outer edge of each cranial vault bone, in coordination with endochondral growth of cranial base bones at the cranial base synchondroses (cartilaginous growth plates). Anterior-posterior growth of the skull is dependent upon both cranial vault and cranial base bone growth. Craniosynostosis (which, when syndromic also often includes premature fusion of cranial base bones) can lead to high intracranial pressure, abnormal skull and facial shapes, malocclusion, blindness, seizures and brain abnormalities [1-6]. Because the sole treatment is surgery, even with appropriately early diagnosis patients can suffer high morbidity [7-9]. A pharmaceutical treatment for craniosynostosis is not yet realized.

Activating mutations in Fibroblast Growth Factor Receptor 2 (*Fgfr2)* [10-13] and inactivating mutations in *Alpl*, the gene for tissue nonspecific alkaline phosphatase (TNAP) [14-17] can cause craniosynostosis. In this study we sought to determine if treatment with recombinant mineral-targeted TNAP could rescue craniosynostosis and associated craniofacial skeletal abnormalities in the FGFR2$^{C342Y/+}$ Crouzon mouse model of craniosynostosis, when delivered shortly post-natal with lentivirus. The FGFR2$^{C342Y/+}$ mutation was previously demonstrated to cause ligand independent signaling and is therefore widely considered to be an activating mutation leading to increased FGF signaling [18-21]. We pursued this investigation because we previously demonstrated that FGF signaling diminishes TNAP expression [22-24], and showed that TNAP deficiency in mice leads to a similar craniofacial phenotype to that seen in FGFR2$^{C342Y/+}$ Crouzon mice including coronal but not sagittal craniosynostosis, deficient growth of the cranial base with fusion of cranial base synchondroses, and brachycephalic/acrocephalic (wide/tall) head

shapes [11, 17, 25, 26]. Additionally, in a previous study using archival aliquots of lentivirus

expressing the mineral targeted recombinant form of TNAP that resulted in increases in serum

AP activity in only a small number of the treated mice, we found statistical differences in the

morphology of the inferior skull surface in treated vs. untreated BALB/c FGFR2$^{C342Y/+}$ mice [27].

Craniosynostosis onset in humans can occur pre-, peri- or postnatal, with earlier onset

forms leading to more severe outcomes and higher morbidity. We backcrossed FGFR2$^{C342Y/+}$

mice onto C57BL/6 and BALB/c strains, and found that C57BL/6 FGFR2$^{C342Y/+}$ mice exhibit

obliteration of the coronal cranial suture initiating shortly after birth, while BALB/c FGFR2$^{C342Y/+}$

mice exhibit point fusions across the coronal suture initiating at four weeks after birth [25].

Both FGFR2$^{C342Y/+}$ strains of mice also exhibit deficient cranial base growth with

brachycephalic/acrocephalic head shapes and tendency for a class III malocclusion (lower teeth

protruding anterior to upper teeth). The objective of this study was to determine if postnatal

delivery of recombinant TNAP could diminish the severity of craniosynostosis in FGFR2$^{C342Y/+}$

mice, a model of Crouzon craniosynostosis syndrome. Viral delivery of TNAP was tested in both

C57BL/6 FGFR2$^{C342Y/+}$ and BALB/c FGFR2$^{C342Y/+}$ mice to determine if efficacy is dependent upon

timing of craniosynostosis onset and/or severity of fusion.

Materials and methods

TNAP Lentivirus

Recombinant mineral-targeted TNAP lentivirus was generously provided by Dr. Jose Luis Millán

(Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA). Pseudotype of the virus is

VSV-G, for transduction into multiple cell types. The SJ1-based HIV-1 vector contains 0.25 kB

insulators derived from the chicken beta-globin locus to diminish insertional mutagenesis,

increase viral titer and increase protein expression [28]. The virus expresses a mineral-targeted

protein that is composed of soluble human TNAP enzyme fused to the constant region of

human IgG1 and a C-terminal deca-aspartate motif to confer targeting to hydroxyapatite. The

aspartate tag confers 30x higher affinity for hydroxyapatite than untagged enzyme [29].

Longevity of the virus for increasing AP levels was reported to last at least 60 days when

injected into neonatal mice and biodistribution of the virus was previously reported to be high

in the liver [30]. Treatment with this recombinant form of TNAP was previously shown to

increase serum alkaline phosphatase levels and rescue bone defects seen in hypophosphatasia

[14, 30-33]. Production and titer of the lentivirus for this study was performed by the University

of Michigan Vector Core.

Animal Procedures

Because severity of craniosynostosis and associated craniofacial shape defects are variable on

the mixed genetic background, FGFR2$^{C342Y/+}$mice were backcrossed with BALB/c and C57BL/6

mice (obtained from Charles River Laboratories) for at least fifteen generations prior to

experiments. BALB/c FGFR2$^{C342Y/+}$mice have a more moderate form of Crouzon syndrome with

craniosynostosis in the form of point fusions across the coronal suture first apparent between

three and four weeks after birth [25]. C57BL/6 mice have a more severe form of Crouzon

syndrome with craniosynostosis in the form of suture obliteration first apparent in neonatal mice

(**Fig. 1**). Both mice also exhibit abnormalities in the cranial base and deficient cranial base growth.

Genotyping was performed as previously described [11, 25]. Briefly, DNA from tail digests was

amplified by polymerase chain reaction using 5'-gagtaccatgctgactgcatgc-3'and 5'-

ggagaggcatctctgtttcaagacc-3' primers to yield a 200 base pair band for wild type FGFR2 and a 300

base pair band for mutant $FGFR2^{C342Y}$. Mice were fed ad libitum and housed under standard 12

hour dark/light cycles. Litters were randomly assigned to treatment/no treatment groups.

Treated mice were injected with 1.0 x 10^7 transforming units lentivirus or an equivalent volume

of phosphate buffered saline via the jugular vein after birth. BALB/c mice (n=12 $FGFR2^{+/+}$ vehicle

treated mice, n=14 $FGFR2^{C342Y/+}$ vehicle treated mice, n=16 $FGFR2^{C342Y/+}$ TNAP lentivirus treated

mice) were euthanized by CO_2 overdose at four weeks post-natal and C57BL/6 mice (n=14

$FGFR2^{+/+}$ vehicle treated mice, n=14 $FGFR2^{C342Y/+}$ vehicle treated mice, n=17 $FGFR2^{C342Y/+}$ TNAP

lentivirus treated mice) were euthanized by CO_2 overdose at three weeks post-natal for analyses.

BALB/c mice were sacrificed at a later age than C57BL/6 mice because craniosynostosis onset

occurs later in BALB/C than in C57BL/6 $FGFR2^{C342Y/+}$ mice. Blood was collected by aortic puncture

under surgical anesthesia. Mice were weighed, and body length was measured for each animal.

All animal procedures were prospectively approved of by the University of Michigan's University

Committee on Use and Care of Animals (UCUCA, protocol PRO00006815). All samples were de-

identified as to genotype and treatment group for analyses, then unblinded for statistical

comparison. The primary outcome assessment was craniosynostosis incidence. Secondary

outcome assessments included malocclusion incidence, cranial bone micro CT measurements,

craniofacial shape measurements, and cranial base bone lengths.

Serum Analyses

Mice were fasted for six hours prior to blood collection. Alkaline phosphatase activity (AP) in

serum was quantified using the colorimetric reagent 4-nitrophenyl-phosphate disodium

hexahydrate (Sigma-Aldrich), as compared to a standard curve using commercially available

alkaline phosphatase enzyme (Sigma-Aldrich). One unit of AP was defined as the amount of

enzyme needed to generate 1 umol of p-nitrophenol per minute. Inorganic phosphate

quantifications were performed using commercially available kits (Pointe Scientific), also as

compared to standard curves. The levels of alanine amino transferase (ALT) and Aspartate

amino transferase (AST) were determined using commercially available colorimetric assay kits

according to manufacturer instructions (Sigma-Aldrich) using serum from control and treated

BALB/c mice, as these mice yielded more serum per mouse. One unit of ALT was defined as the

as the amount of enzyme that would generate 1 nmol of pyruvate per minute. One unit of AST

was defined as the amount of enzyme that would generate 1 nmol of glutamate per minute.

Micro Computed Tomography

Whole skulls were scanned at an 18 μm isotropic voxel resolution using the eXplore Locus SP

micro computed tomography imaging system (GE Healthcare Pre-Clinical Imaging, London, ON,

Canada). Regions of interest (ROI's) for parietal and frontal bones were established as 0.5 mm

in length, 0.5 mm in width, and depth equivalent to thickness of bone, as previously described

[17, 25]. Density, volume and mineral content of cranial bones from mice were measured using

previously established methods using Microview version 2.2 software (GE Healthcare Pre-

Clinical Imaging, London, ON) and established algorithms [34, 35]. Linear measurements of

cranial base bones were measured on micro CT scans using Dolphin imaging software (Dolphin

Imaging & Management Solutions).

152 Cranial Suture and Cranial Base Synchondrosis Assessment

Fusions between frontal and parietal cranial bones (fusion of the coronal suture), fusion between

parietal bones (fusion of the sagittal suture) and fusion of the intersphenoidal synchondrosis (ISS)

and spheno-occipital synchondrosis (SOS) were identified on micro CT scans of mouse skulls.

Cranial sutures and synchondroses were viewed using the two-dimensional micro CT slices in an

orthogonal view across the entire length of the suture or synchondrosis, as previously described

[17, 25]. Synchondroses were identified as fused or not fused. To distinguish between more vs.

less severe forms of craniosynostosis, fusion of cranial suture was scored in the following

categories: 0) normal open suture, 1) diminished suture width with no fusion 2) diminished

suture width with point fusions, and 3) obliteration of the suture.

Reliability of suture fusion assessment was verified by both intra-operator and inter-

operator reliability statistics by calculating intraclass correlation coefficients (ICC). Intra-

operator reliability statistics was carried out by assessing suture fusion status of the coronal

and sagittal sutures as well as the ISS and SOS synchondroses on fifteen micro CT scans by one

investigator two times separated by a two-month period. Inter-operator reliability was carried

out by analyzing fifteen micro CT scans by a second investigator. The ICC for intraoperator

reliability for fusion assessment is .970 (p≤.0001) and the ICC for interoperator reliability is .972

(p≤.0001). Thus, there is high intraoperator and interoperator reliability for fusion assessment.

Linear Measurements

Craniofacial linear skeletal measurements were taken using digital calipers on dissected skulls. Linear measurements were calculated using previously reported craniofacial skeletal landmarks [25, 36, 37], including standard measurements currently in use by the Craniofacial Mutant Mouse Resource of Jackson Laboratory (Bar Harbor, ME). Linear measurements were normalized to total skull length (measured from nasale to opisthion) to account for size differences between FGFR2$^{+/+}$ and FGFR2$^{C342Y/+}$ mice. Measurements were performed twice and an average of the two measurements was utilized for statistical comparison by genotype and treatment. Cranial base anterior-posterior bone lengths were measured on micro CT scans using *Dolphin Imaging 11.0* software (Dolphin Imaging and Management Solutions, Chatsworth, CA), as previously described [26].

Statistics

Results from previous studies and this study demonstrate no sex difference for craniosynostosis in FGFR2$^{C342Y/+}$ mice, therefore sexes were combined for analyses [25]. Normality of data was evaluated by D'Agostino & Pearson Tests. Students t test was then used to compare groups of that were of normal distribution and the Mann Whitney test was utilized to compare groups of that were of non-normal distribution. Because serum AP levels varied in mice injected with the lentivirus, linear regressions were performed to determine if, and to what extent serum AP levels associated with changes body weight and serum liver enzyme levels. Incidences of

craniosynostosis severity category, synchondrosis fusion and malocclusion were analyzed by

the Fisher's exact test.

Results

Injection with the TNAP expression lentivirus significantly increased serum AP levels in all of the

treated mice (**Table 1**). BALB/c FGFR2$^{C342Y/+}$ mice injected with the lentivirus increased serum AP

levels by 1.2 U/mL when compared to BALB/c vehicle treated FGFR2$^{C342Y/+}$ mice (p<0.0001)

when measured at four weeks old. C57BL/6 FGFR2$^{C342Y/+}$ mice injected with the TNAP

expression lentivirus increased serum AP levels by 1.8 U/mL when compared to vehicle treated

C57BL/6 FGFR2$^{C342Y/+}$ mice (p<0.0001) when measured at three weeks old. No significant

difference in serum AP levels were seen between untreated FGFR2$^{C342Y/+}$ and FGFR2$^{+/+}$ mice on

the BALB/c or C57BL/6 backgrounds. Injection with the lentivirus did not alter serum inorganic

phosphate (P$_i$) levels.

Table 1. Serum Alkaline Phosphatase and Inorganic Phosphate Measurements
205   in vehicle vs. TNAP treated BALB/c and C57Bl/6 mice.

Strain	Genotype	Treatment	Serum AP Level (units/ml)	Serum Pi Level (mg/dl)
BALB/c	FGFR2$^{+/+}$	vehicle	0.03 +/- 0.01	10.9 +/- 0.8
BALB/c	FGFR2$^{C342Y/+}$	vehicle	0.03 +/- 0.78	9.9 +/- 1.4
BALB/c	FGFR2$^{C342Y/+}$	TNAP	1.30 +/- 0.54$^{\#}$	9.8 +/- 1.1
C57Bl/6	FGFR2$^{+/+}$	vehicle	0.01 +/- 0.01	9.5 +/- 1.0
C57Bl/6	FGFR2$^{C342Y/+}$	vehicle	0.02 +/- 0.01	8.7 +/- 1.1
C57Bl/6	FGFR2$^{C342Y/+}$	TNAP	1.93 +/- 0.77$^{\#}$	9.0 +/- 0.6

223                    $^{\#}$ p value < 0.01 between treatment groups

Statistical comparison of groups by Mann Whitney showed that FGFR2$^{C342Y/+}$ mice weigh less

and are shorter in body length than their wild type littermates, regardless of genetic

background (**Fig. 2**). Qualitative analysis of craniofacial skeletal shape suggested that

FGFR2$^{C342Y/+}$ mice differ in morphology from their wild type counterparts, and that delivery of

mineral-targeted TNAP via lentivirus did not impact skull morphology (**Fig. 3**).

Craniofacial skeletal linear measurements normalized to total skull length revealed many

differences between FGFR2$^{342Y/+}$ and FGFR2$^{+/+}$ mice on both congenic backgrounds (**Table 2**).

Table 2. Linear Craniofacial Skeletal Measurements reveals
no rescue of craniofacial shape in by TNAP treatment in FGFR2$^{C342Y/+}$ mice.

Strain	Measurement	FGFR2$^{+/+}$ vehicle	FGFR2$^{C342Y/+}$ vehicle	FGFR2$^{C342Y/+}$ TNAP
BALB/c	Cranial Height	.36 +/- .01*	.45 +/- .01	.45 +/- .01
BALB/c	Cranial Width	.55 +/- .01*	.63 +/- .01	.62 +/- .01
BALB/c	Inner Canthal Distance	.20 +/- .01*	.25 +/- .01	.26 +/- .01
BALB/c	Nose Length	.65+/- .01	.65 +/- .01	.65 +/- .01
BALB/c	Nasal Bone Length	.33 +/- .02*	.32 +/- .01	.32 +/- .03
BALB/c	Frontal Bone length	.33 +/- .02	.33 +/- .01	.34 +/- .03
BALB/c	Parietal Bone Length	.20 +/- .01*	.25 +/- .02	.26 +/- .01
BALB/c	Ratio Height to Width	.66 +/- .02*	.72 +/- .02	.72 +/- .02
C57Bl/6	Cranial Height	.38 +/- .01*	.52 +/- .01	.52 +/- .03
C57Bl/6	Cranial Width	.55 +/- .01*	.64 +/- .01	.64 +/- .02
C57Bl/6	Inner Canthal Distance	.23 +/- .01*	.29 +/- .01	.29 +/- .01
C57Bl/6	Nose Length	.65 +/- .01*	.62 +/- .01	.63 +/- .03
C57Bl/6	Nasal Bone Length	.32 +/- .01*	.23 +/- .02	.23 +/- .01
C57Bl/6	Frontal Bone length	.35 +/- .01*	.40 +/- .02	.41 +/- .03
C57Bl/6	Parietal Bone Length	.22 +/- .01*	.26 +/- .02	.27 +/- .02
C57Bl/6	Ratio Height to Width	.38 +/- .01*	.52 +/- .01	.52 +/- .03

233
234 Measures are reported as normalized to total skull length.
235 * p value < 0.01 between genotypes
236 No statistical differences between treatment groups were found.
237

BALB/c FGFR2$^{C342Y/+}$ mice had increased cranial height, cranial width, inner canthal distance,

parietal bone length and cranial height to width ratios, with decreased nasal bone length.

C57BL/6 FGFR2$^{C342Y/+}$ mice had increased cranial height, cranial width, inner canthal distance,

frontal bone length, parietal bone length and cranial height to width ratios, with decreased

nose and nasal bone lengths. Treatment with the TNAP lentivirus did not alter craniofacial

skeletal measurements in FGFR2$^{C342Y/+}$ mice on either genetic background.

Analysis of coronal suture fusions revealed diminished suture width (grade 1) in the

majority of BALB/c FGFR2$^{C342Y/+}$ vehicle treated mice with point fusions (grade 2) evident in

approximately 30% of the mice. In the more severe C57BL/6 FGFR2$^{C342Y/+}$ vehicle treated mice,

the majority of the mice had obliteration of the suture (grade 3) with point fusions (grade 2)

evident in the rest of the mice (**Fig. 4**). The sagittal suture was not fused in any of the mice, and

no cranial suture fusions were evident in wild type mice on either background. Treatment with

recombinant TNAP had no significant impact on the incidence of grade 1, 2 or 3 of the coronal

suture in the BALB/c FGFR2$^{C342Y/+}$ mice. While there was only a trend for decreased fusion in the

left coronal suture fusion upon treatment (control 94% suture obliteration vs. treated 78%

suture obliteration), there was a significant decrease in coronal suture fusion upon treatment in

the right coronal suture (control 94% suture obliteration vs. treated 57% suture obliteration,

$p < 0.03$). There was also a significant decrease in coronal suture fusion when combining both

right and left coronal sutures (control 94% suture obliteration vs. treated 67% suture

obliteration, $p < 0.02$). Consistent with rescue of coronal suture fusion in C57BL/6 but not BALB/c

mice, the incidence of malocclusion was also significantly decreased by treatment from 82.4%

to 34.7% in C57BL/6 but not BALB/c FGFR2$^{C342Y/+}$ mice (**Fig. 5**, $p < 0.03$).

Analysis of cranial base synchondrosis fusions revealed a 100% incidence of fusion of the

inter-sphenoidal synchondrosis (ISS) in both strains of FGFR2$^{C342Y/+}$ mice, with no fusions

evident in wild type mice. Comparison of vehicle vs. treated mice revealed no significant change

in the incidence of ISS fusion in FGFR2$^{C342Y/+}$ mice on either genetic background. Spheno-

occipital synchondronsis (SOS) fusion was seen in approximately half of C57BL/6 FGFR2$^{C342Y/+}$

mice but rarely in BALB/c FGFR2$^{C342Y/+}$ mice. Comparison of vehicle vs. treated mice revealed no

significant change in the incidence of SOS fusion in FGFR2$^{C342Y/+}$ mice on either genetic

background. Measurements of cranial base bone lengths demonstrated decreased length of the

basis-sphenoid and pre-sphenoid bones in both BALB/c and C57BL/6 vehicle treated

FGFR2$^{C342Y/+}$ as compared to FGFR2$^{+/+}$ mice (**Table 3**). Treatment increased length of the pre-

sphenoid bone in BALB/c and C57BL/6 FGFR2$^{C342Y/+}$ mice, but not to the equivalent of pre-

sphenoid bone length seen in FGFR2$^{+/+}$ mice.

Table 3. Cranial Base Bone Measurements in vehicle vs.
TNAP treated Balb/C and C57Bl/6 mice.

Strain	Genotype	Treatment	Basis Occipitus (mm)	Basis Sphenoid (mm)	Pre-Sphenoid (mm)
Balb/C	FGFR2$^{+/+}$	vehicle	3.1 +/- 0.1	2.9 +/- 0.1*	2.5 +/- 0.1*
Balb/C	FGFR2$^{C342Y/+}$	vehicle	2.9 +/- 0.3	2.4 +/- 0.3	1.8 +/- 0.1
Balb/C	FGFR2$^{C342Y/+}$	TNAP	2.9 +/- 0.2	2.5 +/- 0.3	1.9 +/- 0.1#
C57Bl/6	FGFR2$^{+/+}$	vehicle	2.8 +/- 0.2	2.9 +/- 0.2*	2.2 +/- 0.1*
C57Bl/6	FGFR2$^{C342Y/+}$	vehicle	2.8 +/- 0.1	2.7 +/- 0.1	1.7 +/- 0.1
C57Bl/6	FGFR2$^{C342Y/+}$	TNAP	3.0 +/- 0.2	2.6 +/-0.1	1.9 +/- 0.1#

* p value < 0.01 between genotypes
No significant differences between treatment groups were found.

Micro CT based analyses of cranial bones demonstrated significantly diminished bone

mineral density, tissue mineral content, tissue mineral density and bone volume fraction in

frontal bones, plus significantly diminished tissue mineral density and bone volume fraction in

parietal bones of vehicle treated FGFR2$^{C342Y/+}$ mice when compared to wild type littermates on

both BALB/c and C57BL/6 backgrounds (**Table 4**). Injection with the TNAP lentivirus

significantly increased frontal bone mineral density, tissue mineral content and bone volume

fraction, plus parietal bone volume fraction in FGFR2$^{C342Y/+}$ mice on the BALB/c background.

Injection with the TNAP lentivirus did not significantly impact any of the cranial bone

parameters in FGFR2$^{C342Y/+}$ mice on the C57BL/6 background.

302   **Table 4.** Cranial bone volume, density and mineral content
303   in vehicle vs. TNAP treated BALB/c and C57BL/6 mice.

Strain	Genotype	Treatment	Cranial Bone	Bone Mineral Content (mg)	Bone Mineral Density (mg/cc)	Tissue Mineral Content (mg)	Tissue Mineral Density (mg/cc)	Bone Volume Fraction
BALB/c	FGFR2$^{+/+}$	vehicle	Frontal	.035 +/- .004	405 +/- 14*	.028 +/- .007*	692 +/- 14*	0.41 +/- .03*
BALB/c	FGFR2$^{C342Y/+}$	vehicle	Frontal	.031 +/- .008	361 +/- 63	.020 +/- .006$^{\#\backslash}$	671 +/- 18	0.36 +/- .0
BALB/c	FGFR2$^{C342Y/+}$	TNAP	Frontal	.035 +/- .005	401 +/- 26$^{\#}$	.026 +/- .015$^{\#}$	683 +/- 29	0.42 +/- .06$^{\#}$
BALB/c	FGFR2$^{+/+}$	vehicle	Parietal	.034 +/- .004	405 +/- 12	.023 +/- .005	693 +/- 15*	0.43 +/- .03*
BALB/c	FGFR2$^{C342Y/+}$	vehicle	Parietal	.031 +/- .007	396 +/- 39	.020 +/- .005	669 +/- 24	0.36 +/- .06
BALB/c	FGFR2$^{C342Y/+}$	TNAP	Parietal	.034 +/- .006	403 +/- 27	.025 +/- .012	691 +/- 36	0.42 +/- .07$^{\#}$
C57BL/6	FGFR2$^{+/+}$	vehicle	Frontal	.017 +/- .003	245 +/- 25*	.006 +/- .001*	570 +/- 18*	0.12 +/- .01*
C57BL/6	FGFR2$^{C342Y/+}$	vehicle	Frontal	.013 +/- .002	209 +/- 25	.004 +/- .001	519 +/- 26	0.10 +/- .01
C57BL/6	FGFR2$^{C342Y/+}$	TNAP	Frontal	.016 +/- .004	225 +/- 30	.004 +/- .001	553 +/- 40	0.11 +/- .01
C57BL/6	FGFR2$^{+/+}$	vehicle	Parietal	.015 +/- .001	237 +/- 19	.005 +/- .001	590 +/- 13*	.121 +/- .023*
C57BL/6	FGFR2$^{C342Y/+}$	vehicle	Parietal	.012 +/- .003	217 +/- 18	.004 +/- .001	558 +/- 36	.098 +/- .006
C57BL/6	FGFR2$^{C342Y/+}$	TNAP	Parietal	.013 +/- .004	232 +/- 42	.004 +/- .001	575 +/- 32	.105 +/- .009

304
305   * p value < 0.01 between genotypes
306                         $^{\#}$ p value < 0.01 between treatment groups

Biodistribution of lentivirus containing the recombinant mineral-targeted form of TNAP was

previously shown to result in highest viral expression levels in the liver [30]. Because the

treated mice in this study received recombinant TNAP via lentivirus, while the control group

received no virus, we measured serum liver enzymes levels to determine if liver toxicity due to

lentivirus could account for a change in phenotype in the treated mice, and/or the increase in

serum AP levels seen in the treated mice. Alanine amino transferase (ALT) and Aspartate amino

transferase (AST) are liver enzymes that, when seen at high levels in serum, are indicative of

liver toxicity [38, 39]. Therefore, we tested serum ALT and AST levels in the mice. When

compared by genotypes, no differences were noted between FGFR2$^{+/+}$ and control FGFR2$^{C342Y/+}$

mice for either ALT or AST enzymes. When compared by treatment groups, FGFR2$^{C342Y/+}$ mice

that were treated had significantly lower ALT levels than their control counterparts, and

significantly higher levels of ALT than their control counterparts (**Fig. 6**). We next performed

correlation studies between ALT and AST serum levels with serum AP levels and body weight to

better understand if lentiviral delivery of TNAP led to liver toxicity. No significant correlations

were found between serum AP and AST levels for any of the groups. A significant reverse

correlation was found between serum AP and ALT levels only for the treated FGFR2$^{C342Y/+}$ group

(r = -.73; 95% confidence interval of -.898 to -.361, p < .005). No significant correlations were

found between serum AST or ALT levels and body weight for any of the groups. Together, this

data shows that the effect of lentiviral TNAP treatment decreased ALT and increased AST

enzymes in serum, but had no effect on body weight of the mice and was not likely causal for

the observed increased in serum AP levels seen in treated mice.

Discussion

Here we found that neonatal lentiviral delivery of recombinant TNAP improved cranial bone density, mineral content and volume fraction in the milder BALB/c FGFR2$^{C342Y/+}$ model of Crouzon syndrome but not in the more severe C57BL/6 FGFR2$^{C342Y/+}$ model of Crouzon syndrome. Improvement of cranial bone density, mineral content and volume fraction by TNAP treatment in BALB/c Crouzon mice is consistent with results showing that recombinant mineral targeted TNAP treatment rescues mineralization of craniofacial and long bones in in the *Alpl*$^{-/-}$ mouse model of infantile hypophosphatasia (HPP) and humans with infantile and childhood HPP [30, 31, 33, 40]. It is curious that delivery of mineral-targeted TNAP did not improve cranial bone parameters in the more severe C57BL/6 FGFR2$^{C342Y/+}$ mice. Wild type C57BL/6 mice have significantly less cranial bone density, mineral content and volume fraction than BALB/c wild type mice, which is diminished to an even greater extent in FGFR2$^{C342Y/+}$ C57BL/6 mice. While not quantified here, it is possible that the lack of rescue by TNAP is due to diminished bone matrix available to mineralize and/or a lack of cranial bone progenitor cells available to generate additional bone in the C57BL/6 mice, as TNAP is known to be essential for matrix mineralization [41] and for the formation of cranial bone progenitor cells [42].

Craniosynostosis severity was significantly diminished by treatment in the C57BL/6 FGFR2$^{C342Y/+}$ mice. The vast majority of these mice typically exhibit obliteration of the coronal suture within a few days after birth (**Fig. 1**). Suture obliteration went from 94% to 67% in the C57BL/6 FGFR2$^{C342Y/+}$ mice upon treatment with exogenous TNAP by three weeks after birth. This result is consistent with the rescue of craniosynostosis seen in *Alpl*$^{-/-}$ mice treated with mineral targeted recombinant TNAP protein [33]. While far from a complete rescue, the data

provided here does demonstrate the potential efficacy of TNAP for diminishing severity of

craniosynostosis in Crouzon syndrome. The data also supports the idea that convergence exists

between changes downstream of TNAP activity and FGFR2 signaling leading to coronal suture

fusion. We recently showed that TNAP regulates expression of FGFR2 and Erk1,2 activity [42].

While the current study was originally based upon the hypothesis that FGF signaling regulates

expression of TNAP, our more recent data suggests the alternative hypothesis that exogenous

delivery of TNAP rescues Crouzon craniosynostosis because TNAP diminishes the overactive FGF

and Erk1,2 signaling in $FGFR2^{C342Y/+}$ cells. We are currently working to confirm this latter

hypothesis and delineate how TNAP may mediate these changes.

This study was designed to include two mouse models of Crouzon syndrome: a severe form

(C57BL/6 $FGFR2^{C342Y/+}$ mice) that exhibited neonatal onset of coronal suture obliteration and a

more moderate form (BALB/c $FGFR2^{C342Y/+}$ mice) that exhibited small point fusions across the

coronal suture evident by four weeks after birth. We anticipated that neonatal delivery of TNAP

via lentivirus would have greater effects on the later onset BALB/c $FGFR2^{C342Y/+}$ mouse model,

as the drug would be delivered prior to onset of craniosynostosis. Results showed instead that

the treatment significantly diminished severity of coronal suture fusion in the neonatal onset

C5BL/6 $FGFR2^{C342Y/+}$ mouse with no effect on coronal suture fusion in the postnatal onset

BALB/c $FGFR2^{C342Y/+}$ mouse. This data suggests that exogenous TNAP may be able to prevent

cranial suture obliteration but not point fusions across the suture. This data also indicates that

cellular mechanisms leading to craniosynostosis in $FGFR2^{C342Y/+}$ mice are both independent and

dependent upon TNAP.

While lentiviral TNAP treatment did not rescue cranial base synchondrosis fusion in the

FGFR2$^{C342Y/+}$ mice, length of the pre-sphenoid bone was mildly increased in the treated mice.

This result is consistent with our previous study using archival lots of the lentivirus [27] which

suggested changes in inferior skull morphology and may indicate that TNAP can promote cranial

base growth in Crouzon syndrome. Delivery of recombinant TNAP also decreased the incidence

of class III malocclusion in the C57BL/6 but not BALB/c FGFR2$^{C342Y/+}$ mice. The malocclusion

rescue is likely due to changes in coronal suture fusion as opposed to changes in the cranial

base, as treatment with TNAP had minimal and similar changes on cranial base synchondroses

and cranial base bone lengths in both C57BL/6 and BALB/c FGFR2$^{C342Y/+}$ mice, with an effect on

coronal suture fusion only in C57BL/6 FGFR2$^{C342Y/+}$ mice.

A major limitation of this study is the method and timing of TNAP delivery. Lentiviral

delivery of TNAP was not ideal, as the control group did not receive lentivirus which can lead to

liver toxicity and increase serum AP levels. While we found that delivery of the TNAP lentivirus

decreased serum ALT and increased serum AST levels in the treated mice, there was no impact

on body weight of the mice. We interpret these findings to indicate that the treatment had

minimal significant impact on overall health of the mice. In addition, serum AP levels did not

correlate with serum AST levels and had a reverse correlation with serum ALT levels in treated

FGFR2$^{C342Y/+}$ mice. We interpret this latter data to indicate that while changes in the liver may

have occurred, liver toxicity was not the cause for increased serum AP levels in the treated

mice. Yet, we cannot be certain that the lentiviral vector itself had no impact on the craniofacial

skeletal phenotype of the treated FGFR2$^{C342Y/+}$ mice. A more ideal study design would

incorporate empty lentiviral vector delivery to the control mice, or delivery of a recombinant

TNAP protein to the mice. It would also be worthwhile to attempt prenatal delivery of TNAP, to

determine if earlier treatment would have a greater impact on rescue of coronal suture fusion

in C57BL/6 FGFR2$^{C342Y/+}$ mice.

Finally, it is also worth noting that earlier time points to assess changes in the coronal

suture were not performed. The decreased incidence of coronal suture obliteration in C57BL/6

Crouzon mice at 3 weeks postnatal suggests that onset occurred later in these mice. Earlier

time points assessing the coronal suture after treatment could have confirmed that this was

indeed the case. Earlier onset of craniosynostosis increases morbidity, as brain growth is limited

earlier and for a longer duration. If TNAP postpones fusion of the coronal suture in Crouzon

syndrome, treatment with TNAP could potentially diminish morbidity.

405 Acknowledgments

406   We thank Prof. José Luis Millán (La Jolla, CA) and Prof. Takashi Shimada (Tokyo, Japan) for

407   providing us with the lentiviral vector expressing mineral targeted TNAP, as reported in

408   Yamamoto et al., 2011 (30).

410 References

1. Flapper WJ, Anderson PJ, Roberts RM, David DJ. Intellectual outcomes following protocol
management in Crouzon, Pfeiffer, and Muenke syndromes. J Craniofac Surg. 2009;20(4):1252-5.
Epub 2009/07/25. doi: 10.1097/SCS.0b013e3181acdf9a. PubMed PMID: 19625842.
2. McCarthy JG, Warren SM, Bernstein J, Burnett W, Cunningham ML, Edmond JC, et al.
Parameters of care for craniosynostosis. Cleft Palate Craniofac J. 2012;49 Suppl:1S-24S. Epub
2011/08/19. doi: 10.1597/11-138. PubMed PMID: 21848431.

3.   Morriss-Kay GM, Wilkie AO. Growth of the normal skull vault and its alteration in
craniosynostosis: insights from human genetics and experimental studies. J Anat.
2005;207(5):637-53. Epub 2005/11/30. doi: 10.1111/j.1469-7580.2005.00475.x. PubMed PMID:
16313397; PubMed Central PMCID: PMCPMC1571561.
4.   Okajima K, Robinson LK, Hart MA, Abuelo DN, Cowan LS, Hasegawa T, et al. Ocular anterior
chamber dysgenesis in craniosynostosis syndromes with a fibroblast growth factor receptor 2
mutation. Am J Med Genet. 1999;85(2):160-70. Epub 1999/07/16. PubMed PMID: 10406670.
5.   Rasmussen SA, Yazdy MM, Frias JL, Honein MA. Priorities for public health research on
craniosynostosis: summary and recommendations from a Centers for Disease Control and
Prevention-sponsored meeting. Am J Med Genet A. 2008;146A(2):149-58. Epub 2007/12/15.
doi: 10.1002/ajmg.a.32106. PubMed PMID: 18080327.
6.   Seruya M, Oh AK, Boyajian MJ, Posnick JC, Keating RF. Treatment for delayed presentation
of sagittal synostosis: challenges pertaining to occult intracranial hypertension. J Neurosurg
Pediatr. 2011;8(1):40-8. Epub 2011/07/05. doi: 10.3171/2011.4.PEDS1160. PubMed PMID:
21721888.
7.   Abe H, Ikota T, Akino M, Kitami K, Tsuru M. Functional prognosis of surgical treatment of
craniosynostosis. Childs Nerv Syst. 1985;1(1):53-61. Epub 1985/01/01. PubMed PMID: 3986842.
8.   Baird LC, Gonda D, Cohen SR, Evers LH, LeFloch N, Levy ML, et al. Craniofacial reconstruction
as a treatment for elevated intracranial pressure. Childs Nerv Syst. 2012;28(3):411-8. Epub
2011/11/10. doi: 10.1007/s00381-011-1615-6. PubMed PMID: 22068642.
9.   Renier D, Lajeunie E, Arnaud E, Marchac D. Management of craniosynostoses. Childs Nerv
Syst. 2000;16(10-11):645-58. Epub 2001/01/11. doi: 10.1007/s003810000320. PubMed PMID:
11151714.
10. Chen L, Li D, Li C, Engel A, Deng CX. A Ser252Trp [corrected] substitution in mouse fibroblast
growth factor receptor 2 (Fgfr2) results in craniosynostosis. Bone. 2003;33(2):169-78. Epub
2003/09/23. PubMed PMID: 14499350.
11. Eswarakumar VP, Horowitz MC, Locklin R, Morriss-Kay GM, Lonai P. A gain-of-function
mutation of Fgfr2c demonstrates the roles of this receptor variant in osteogenesis. Proc Natl
Acad Sci U S A. 2004;101(34):12555-60. Epub 2004/08/19. doi: 10.1073/pnas.0405031101.
PubMed PMID: 15316116; PubMed Central PMCID: PMCPMC515096.
12. Reardon W, Winter RM, Rutland P, Pulleyn LJ, Jones BM, Malcolm S. Mutations in the
fibroblast growth factor receptor 2 gene cause Crouzon syndrome. Nat Genet. 1994;8(1):98-
103. Epub 1994/09/01. doi: 10.1038/ng0994-98. PubMed PMID: 7987400.
13. Wilkie AO, Slaney SF, Oldridge M, Poole MD, Ashworth GJ, Hockley AD, et al. Apert
syndrome results from localized mutations of FGFR2 and is allelic with Crouzon syndrome. Nat
Genet. 1995;9(2):165-72. Epub 1995/02/01. doi: 10.1038/ng0295-165. PubMed PMID:
7719344.
14. Whyte MP, Greenberg CR, Salman NJ, Bober MB, McAlister WH, Wenkert D, et al. Enzyme-
replacement therapy in life-threatening hypophosphatasia. N Engl J Med. 2012;366(10):904-13.
Epub 2012/03/09. doi: 10.1056/NEJMoa1106173. PubMed PMID: 22397652.
15. Collmann H, Mornet E, Gattenlohner S, Beck C, Girschick H. Neurosurgical aspects of
childhood hypophosphatasia. Childs Nerv Syst. 2009;25(2):217-23. Epub 2008/09/05. doi:
10.1007/s00381-008-0708-3. PubMed PMID: 18769927.

16. Fraser D. Hypophosphatasia. Am J Med. 1957;22(5):730-46. Epub 1957/05/01. PubMed
PMID: 13410963.
17. Liu J, Nam HK, Campbell C, Gasque KC, Millan JL, Hatch NE. Tissue-nonspecific alkaline
phosphatase deficiency causes abnormal craniofacial bone development in the Alpl(-/-) mouse
model of infantile hypophosphatasia. Bone. 2014;67:81-94. Epub 2014/07/12. doi:
10.1016/j.bone.2014.06.040. PubMed PMID: 25014884; PubMed Central PMCID:
PMCPMC4149826.
18. Galvin BD, Hart KC, Meyer AN, Webster MK, Donoghue DJ. Constitutive receptor activation
by Crouzon syndrome mutations in fibroblast growth factor receptor (FGFR)2 and FGFR2/Neu
chimeras. Proc Natl Acad Sci U S A. 1996;93(15):7894-9. Epub 1996/07/23. PubMed PMID:
8755573; PubMed Central PMCID: PMCPMC38845.
19. Neilson KM, Friesel RE. Constitutive activation of fibroblast growth factor receptor-2 by a
point mutation associated with Crouzon syndrome. J Biol Chem. 1995;270(44):26037-40. Epub
1995/11/03. PubMed PMID: 7592798.
20. Robertson SC, Meyer AN, Hart KC, Galvin BD, Webster MK, Donoghue DJ. Activating
mutations in the extracellular domain of the fibroblast growth factor receptor 2 function by
disruption of the disulfide bond in the third immunoglobulin-like domain. Proc Natl Acad Sci U S
A. 1998;95(8):4567-72. Epub 1998/05/16. PubMed PMID: 9539778; PubMed Central PMCID:
PMCPMC22530.
21. Pfaff MJ, Xue K, Li L, Horowitz MC, Steinbacher DM, Eswarakumar JVP. FGFR2c-mediated
ERK-MAPK activity regulates coronal suture development. Dev Biol. 2016;415(2):242-50. Epub
2016/04/02. doi: 10.1016/j.ydbio.2016.03.026. PubMed PMID: 27034231; PubMed Central
PMCID: PMCPMC5580088.
22. Hatch NE, Li Y, Franceschi RT. FGF2 stimulation of the pyrophosphate-generating enzyme,
PC-1, in pre-osteoblast cells is mediated by RUNX2. J Bone Miner Res. 2009;24(4):652-62. Epub
2008/12/04. doi: 10.1359/jbmr.081213. PubMed PMID: 19049325; PubMed Central PMCID:
PMCPMC2659512.
23. Hatch NE, Nociti F, Swanson E, Bothwell M, Somerman M. FGF2 alters expression of the
pyrophosphate/phosphate regulating proteins, PC-1, ANK and TNAP, in the calvarial
osteoblastic cell line, MC3T3E1(C4). Connect Tissue Res. 2005;46(4-5):184-92. Epub
2006/03/21. doi: 10.1080/03008200500237203. PubMed PMID: 16546821.
24. Liu J, Kwon TG, Nam HK, Hatch NE. Craniosynostosis-associated Fgfr2(C342Y) mutant bone
marrow stromal cells exhibit cell autonomous abnormalities in osteoblast differentiation and
bone formation. Biomed Res Int. 2013;2013:292506. Epub 2013/06/14. doi:
10.1155/2013/292506. PubMed PMID: 23762837; PubMed Central PMCID: PMCPMC3665166.
25. Liu J, Nam HK, Wang E, Hatch NE. Further analysis of the Crouzon mouse: effects of the
FGFR2(C342Y) mutation are cranial bone-dependent. Calcif Tissue Int. 2013;92(5):451-66. Epub
2013/01/30. doi: 10.1007/s00223-013-9701-2. PubMed PMID: 23358860; PubMed Central
PMCID: PMCPMC3631296.
26. Nam HK, Sharma M, Liu J, Hatch NE. Tissue Nonspecific Alkaline Phosphatase (TNAP)
Regulates Cranial Base Growth and Synchondrosis Maturation. Front Physiol. 2017;8:161. Epub
2017/04/06. doi: 10.3389/fphys.2017.00161. PubMed PMID: 28377728; PubMed Central
PMCID: PMCPMC5359511.

27. Wang E, Nam HK, Liu J, Hatch NE. The effects of tissue-non-specific alkaline phosphatase
gene therapy on craniosynostosis and craniofacial morphology in the FGFR2C342Y/+ mouse
model of Crouzon craniosynostosis. Orthod Craniofac Res. 2015;18 Suppl 1:196-206. Epub
2015/04/14. doi: 10.1111/ocr.12080. PubMed PMID: 25865549; PubMed Central PMCID:
PMCPMC4396711.
28. Hanawa H, Yamamoto M, Zhao H, Shimada T, Persons DA. Optimized lentiviral vector design
improves titer and transgene expression of vectors containing the chicken beta-globin locus
HS4 insulator element. Mol Ther. 2009;17(4):667-74. Epub 2009/02/19. doi:
10.1038/mt.2009.1. PubMed PMID: 19223867; PubMed Central PMCID: PMCPMC2835111.
29. Nishioka T, Tomatsu S, Gutierrez MA, Miyamoto K, Trandafirescu GG, Lopez PL, et al.
Enhancement of drug delivery to bone: characterization of human tissue-nonspecific alkaline
phosphatase tagged with an acidic oligopeptide. Mol Genet Metab. 2006;88(3):244-55. Epub
2006/04/18. doi: 10.1016/j.ymgme.2006.02.012. PubMed PMID: 16616566; PubMed Central
PMCID: PMCPMC2587042.
30. Yamamoto S, Orimo H, Matsumoto T, Iijima O, Narisawa S, Maeda T, et al. Prolonged
survival and phenotypic correction of Akp2(-/-) hypophosphatasia mice by lentiviral gene
therapy. J Bone Miner Res. 2011;26(1):135-42. Epub 2010/08/06. doi: 10.1002/jbmr.201.
PubMed PMID: 20687159; PubMed Central PMCID: PMCPMC3179312.
31. Millan JL, Narisawa S, Lemire I, Loisel TP, Boileau G, Leonard P, et al. Enzyme replacement
therapy for murine hypophosphatasia. J Bone Miner Res. 2008;23(6):777-87. Epub 2007/12/19.
doi: 10.1359/jbmr.071213. PubMed PMID: 18086009; PubMed Central PMCID:
PMCPMC2652241.
32. Yadav MC, Lemire I, Leonard P, Boileau G, Blond L, Beliveau M, et al. Dose response of
bone-targeted enzyme replacement for murine hypophosphatasia. Bone. 2011;49(2):250-6.
Epub 2011/04/05. doi: 10.1016/j.bone.2011.03.770. PubMed PMID: 21458605; PubMed Central
PMCID: PMCPMC3117961.
33. Liu J, Campbell C, Nam HK, Caron A, Yadav MC, Millan JL, et al. Enzyme replacement for
craniofacial skeletal defects and craniosynostosis in murine hypophosphatasia. Bone.
2015;78:203-11. Epub 2015/05/12. doi: 10.1016/j.bone.2015.05.005. PubMed PMID:
25959417; PubMed Central PMCID: PMCPMC4466206.
34. Meganck JA, Kozloff KM, Thornton MM, Broski SM, Goldstein SA. Beam hardening artifacts
in micro-computed tomography scanning can be reduced by X-ray beam filtration and the
resulting images can be used to accurately measure BMD. Bone. 2009;45(6):1104-16. Epub
2009/08/05. doi: 10.1016/j.bone.2009.07.078. PubMed PMID: 19651256; PubMed Central
PMCID: PMCPMC2783193.
35. Umoh JU, Sampaio AV, Welch I, Pitelka V, Goldberg HA, Underhill TM, et al. In vivo micro-CT
analysis of bone remodeling in a rat calvarial defect model. Phys Med Biol. 2009;54(7):2147-61.
Epub 2009/03/17. doi: 10.1088/0031-9155/54/7/020. PubMed PMID: 19287088.
36. Perlyn CA, DeLeon VB, Babbs C, Govier D, Burell L, Darvann T, et al. The craniofacial
phenotype of the Crouzon mouse: analysis of a model for syndromic craniosynostosis using
three-dimensional MicroCT. Cleft Palate Craniofac J. 2006;43(6):740-8. Epub 2006/11/16. doi:
10.1597/05-212. PubMed PMID: 17105336.
37. Richtsmeier JT, Baxter LL, Reeves RH. Parallels of craniofacial maldevelopment in Down
syndrome and Ts65Dn mice. Dev Dyn. 2000;217(2):137-45. Epub 2000/03/08. doi:

10.1002/(SICI)1097-0177(200002)217:2<137::AID-DVDY1>3.0.CO;2-N. PubMed PMID:
10706138.
38. Franca RF, Zucoloto S, da Fonseca BA. A BALB/c mouse model shows that liver involvement
in dengue disease is immune-mediated. Exp Mol Pathol. 2010;89(3):321-6. Epub 2010/08/03.
doi: 10.1016/j.yexmp.2010.07.007. PubMed PMID: 20673760.
39. Huang F, Zhang W, Gong G, Yuan C, Yan Y, Yang S, et al. Experimental infection of Balb/c
nude mice with Hepatitis E virus. BMC Infect Dis. 2009;9:93. Epub 2009/06/16. doi:
10.1186/1471-2334-9-93. PubMed PMID: 19523236; PubMed Central PMCID:
PMCPMC2717970.
40. Whyte MP, Simmons JH, Moseley S, Fujita KP, Bishop N, Salman NJ, et al. Asfotase alfa for
infants and young children with hypophosphatasia: 7 year outcomes of a single-arm, open-
label, phase 2 extension trial. Lancet Diabetes Endocrinol. 2018. Epub 2018/12/19. doi:
10.1016/S2213-8587(18)30307-3. PubMed PMID: 30558909.
41. Murshed M, Harmey D, Millan JL, McKee MD, Karsenty G. Unique coexpression in
osteoblasts of broadly expressed genes accounts for the spatial restriction of ECM
mineralization to bone. Genes Dev. 2005;19(9):1093-104. Epub 2005/04/19. doi:
10.1101/gad.1276205. PubMed PMID: 15833911; PubMed Central PMCID: PMCPMC1091743.
42. Nam HK, Vesela I, Siismets E, Hatch NE. Tissue nonspecific alkaline phosphatase promotes
calvarial progenitor cell cycle progression and cytokinesis via Erk1,2. Bone. 2018;120:125-36.
Epub 2018/10/21. doi: 10.1016/j.bone.2018.10.013. PubMed PMID: 30342227.
567

Figure Legends:

Figure 1. Neonatal Coronal Suture Fusion in C57BL/6 FGFR2$^{C342Y/+}$ Mice. Representative images

of dissected FGFR2$^{+/+}$ (A) and FGFR2$^{C342Y/+}$ (B) day three mouse skulls are shown. Arrows point

to loss of the coronal suture in C57BL/6 FGFR2$^{C342Y/+}$ mice.

Figure 2. Body Weight and Length of BALB/c and C57BL/6 Mice. Body weight and length of

vehicle vs. TNAP treated mice are shown. FGFR2$^{C342Y/+}$ mice are lighter and smaller than

FGFR2$^{+/+}$ mice, regardless of strain. Treatment does not alter body weight or body length.

*p<0.01 between genotypes.

Figure 3. Micro CT Images of Vehicle and TNAP Treated FGFR2$^{C342Y/+}$ Mice. Micro CT isosurface

axial images of day 28 BALB/c and day 21 C57BL/6 mouse skulls are shown. Denser bone tissues

are lighter in color. Comparison of BALB/c FGFR2$^{+/+}$ (WT) mice (A), BALB/c FGFR2$^{C342Y/+}$ (CZ)

vehicle treated mice (C), and BALB/c FGFR2$^{C342Y/+}$ (CZ) TNAP treated mice (E) indicates

morphologic differences between genotypes but not treatment groups. Comparison of C57BL/6

FGFR2$^{+/+}$ (WT) mice (B), C57BL/6 /c FGFR2$^{C342Y/+}$ (CZ) vehicle treated mice (D), and C57BL/6

FGFR2$^{C342Y/+}$ (CZ) TNAP treated mice (F) indicates morphologic differences between genotypes

but not treatment groups.

Figure 4. Coronal Suture Fusion in Vehicle and TNAP Treated FGFR2$^{C342Y/+}$ Mice. Percentage of

mice with fusion of the coronal suture are shown. Fusion was scored in the following

categories: 0) normal open suture, 1) diminished suture width with no fusion, 2) diminished

suture width with point fusions across the suture, and 3) obliteration of the suture. Results

show diminished suture obliteration in C57BL/6 FGFR2$^{C342Y/+}$ (Crouzon) mice with no changes

noted in BALB/c FGFR2$^{C342Y/+}$ (Crouzon) mice. *p<0.03 between treatment groups.

Figure 5. Incidence of Malocclusion in Vehicle and TNAP Treated FGFR2$^{C342Y/+}$ Mice.

Percentage of FGFR2$^{C342Y/+}$ (Crouzon) mice with a class III malocclusion are shown. Treatment

with TNAP lentivirus significantly diminished malocclusion in C57BL/6 FGFR2$^{C342Y/+}$ (Crouzon)

but not BALB/c FGFR2$^{C342Y/+}$ (Crouzon) mice. *p<0.03.

Figure 6. Liver Toxicity Enzyme Tests in Vehicle and TNAP Treated FGFR2$^{C342Y/+}$ Mice. Alanine

amino transferase (ALT) and aspartate amino transferase (AST) liver enzymes were measured in

serum of vehicle and treated mice. When compared by groups, no differences were seen

between FGFR2$^{+/+}$ and FGFR2$^{C342Y/+}$ vehicle treated mice. Treatment decreased serum ALT (A)

and increased serum AST (B) in FGFR2$^{C342Y/+}$ mice. Linear regression analyses showed a

significant reverse correlation (r = -.73; 95% CI of -.898 to -.361, p < .005) between AP and ALT

enzyme levels in FGFR2$^{C342Y/+}$ mice (C). No correlation was found between AP and AST enzyme

levels in FGFR2$^{C342Y/+}$ mice (D). No correlations were found between ALT (E) or AST (F) enzyme

levels and body weight.

FIGURES

Fig. 1. Neonatal Suture Fusion in C57BL/6 Crouzon Mice.

Fig. 2. Body Weight and Length of BALB/c and C57BL/6 Mice.

Fig. 3. Isosurface Images of BALB/c and C57BL/6 Mice.

Fig. 4 Coronal Suture Fusion in BALB/c and C57BL/6 Mice.

Fig. 5. Class III Malocclusion in BALB/c and C57BL/6 Mice.

Fig. 6. Liver Enzyme Tests in Control and Treated Mice