Vibrio cholerae **MARTX toxin multifunctionality silences inflammatory response to**

toxin-mediated cytoskeletal collapse

Patrick J. Woida[1], **Karla J. F. Satchell**[1,2*]

[1]Department of Microbiology-Immunology, Northwestern University Feinberg School of

Medicine, Chicago, IL 60611, USA

[2]Corresponding author

*Correspondence. Email k-satchell@northwestern.edu

ABSTRACT

Multifunctional autoprocessing repeats-in-toxin (MARTX) toxins are pore-forming toxins that

translocate multiple functionally independent effector domains into a target eukaryotic cell.

Vibrio cholerae colonizes intestinal epithelial cells (IECs) and utilizes a MARTX toxin with three

effector domains — the actin cross-linking domain (ACD), the Rho inactivation domain (RID),

and the α/β hydrolase domain (ABH) — to regulate innate immunity and enhance colonization.

Whether these multiple catalytic enzymes delivered from a single toxin have coordinated

function has not been explored. Using cultured IECs, we demonstrate ACD-induced cytoskeletal

collapse activates a robust proinflammatory response that is blocked by the action of co-

delivered RID and ABH. Thus, MARTX toxins utilize multiple enzymatic activities on a single

toxin to silence the host response to both bacterial factors and effector function. Further, these

data explain that *V. cholerae* utilizes the MARTX toxin to suppress intestinal inflammation and

contribute to cholera being classically defined as non-inflammatory diarrheal disease.

Keywords: *Vibrio cholerae*, MARTX toxin, interleukin-8, CXCL8, RNA-seq, MAP kinases,

signaling

GRAPHICAL ABSTRACT
30
31

INTRODUCTION

Multifunctional autoprocessing repeats-in-toxin (MARTX) toxins utilize multiple enzymatic functions to promote virulence of various *Vibrio* species. MARTX toxins are secreted as single 3500 – 5300 amino acid polypeptides that contain conserved glycine-rich repeats at the N- and C-termini that flank multiple arrayed effector domains and an autoprocessing cysteine protease domain (CPD) (Satchell, 2015). The glycine-rich repeats are proposed to form a pore in the plasma membrane of eukaryotic cells to translocate the arrayed effectors and the CPD into the target cell (Dolores et al., 2015; Kim et al., 2015; Kim et al., 2008). In the cytoplasm, CPD is activated by binding host inositol hexakisphosphate ($InsP_6$) and then auto-cleaves to free the effector domains from the large holotoxin. The individual effectors then can traffic through the cell to identify targets and to perform their catalytic functions (Egerer and Satchell, 2010; Prochazkova et al., 2009; Shen et al., 2009) (Figure 1A). Due to this enzymatic multifunctionality, MARTX toxins have been described as bacterial "cluster bombs" that release multiple cytotoxic bomblets into host cells from a single toxin warhead. While the biochemical function of many of the effector domains is known (Kim, 2018), the additive or synergistic benefit of having all these enzymatic functions delivered on a single toxin has yet to be identified (Woida and Satchell, 2018).

 Vibrio cholerae is the causative agent of the severe diarrheal disease cholera (Ali et al., 2012). In addition to its primary virulence factor, the ADP-ribosylating cholera toxin, pandemic *V. cholerae* El Tor O1 strains secrete a 4,545 amino acid (a.a) MARTX$_{Vc}$ toxin that contributes to enhanced bacterial colonization of the small intestine by protecting the pathogen from neutrophil-mediated clearance during the earliest stages of infection (Olivier et al., 2007; Olivier et al., 2009; Queen and Satchell, 2012; Weill et al., 2017). The early timing of these events suggests the inhibition of neutrophils and other innate immune cells does not reflect destruction of the cells by the MARTX$_{Vc}$ toxin, but rather a failure of neutrophils to be recruited to the site of

infection. Therefore, the MARTX$_{Vc}$ toxin might function to limit host signaling that results in

innate immune cell recruitment.

Although MARTX toxins can have highly variable effector domain organizations in some

species, the MARTX$_{Vc}$ toxin of nearly all *V. cholerae* strains have the same three effector

domains (Dolores and Satchell, 2013) (Figure 1A). The first effector domain is the actin cross-

linking domain (ACD) that covalently cross-links monomeric actin to depolymerize actin (Fullner

and Mekalanos, 2000; Kudryashov et al., 2008; Sheahan et al., 2004). The cross-linked actin

oligomers also have high binding affinity for formins and other actin binding proteins to further

disrupt the cytoskeleton (Heisler et al., 2015). The second domain is the Rho inactivation

domain (RID) that inactivates Rac1 and other Rho family GTPases by transferring fatty acids

onto lysine residues in the C-terminal polybasic region of the GTPases. The acylation of the

Rho GTPases blocks them from interacting with downstream effectors. The action of both ACD

and RID results in destruction of the actin cytoskeleton and loss of epithelial cell junction

integrity (Dolores et al., 2015; Sheahan and Satchell, 2007; Zhou et al., 2017). The final effector

domain is the α/β hydrolase domain (ABH), a highly specific phospholipase A1 that cleaves only

phosphatidylinositol 3-phosphate (PI3P) to inhibit autophagy and endocytic trafficking in HeLa

cells (Agarwal et al., 2015). All three effector domains can also independently disable

macrophages, suggesting a mechanism by which the MARTX$_{Vc}$ toxin protects against clearance

of *V. cholerae* from the intestine. However, while ACD potently inhibits phagocytosis, inhibition

by RID and ABH is far less robust (Dolores et al., 2015). Therefore, in the context of co-

introduction into cells in parallel with ACD, phagocytosis is not likely a biologically significant

outcome of RID and ABH intoxication. Further, since the MARTX$_{Vc}$ toxin promotes colonization

possibly even well before onset of inflammation, inhibition of phagocytic clearance may not be a

primary function of the MARTX$_{Vc}$ toxin.

Intestinal epithelial cells (IECs) act as a barrier to contain bacteria to the lumen, and also

as sensors to detect microbial-associated molecular patterns and release chemokines that

recruit immune cells to the site of infection (Kagnoff and Eckmann, 1997; Peterson and Artis,

2014). While *V. cholerae* is typically considered a secretory, non-inflammatory diarrheal

disease, these bacteria do elicit mild inflammation in human intestines (Bishop et al., 2014;

Qadri et al., 2002). In addition, *V. cholerae* is known to stimulate release of the neutrophil-

recruiting chemokine interleukin-8 (IL-8 or CXCL8) from human colonic intestinal cells in

response to pathogen-associated molecular patterns (PAMPs) including purified flagellin and

lipopolysaccharide (LPS) (Harrison et al., 2008; Rodriguez et al., 2001; Soriani et al., 2002;

Zhou et al., 2004). However, the extent to which *V. cholerae* induces IL-8 secretion *in vitro* in

response to live bacteria varies dramatically based on the strain isolate (Rodriguez et al., 2001;

Zhou et al., 2004). Since *V. cholerae* routinely induces inflammation in response to purified

PAMPs, we hypothesized that the variable response to live bacteria may be attributed to strain-

dependent variation in secreted exotoxins impacting initiation of proinflammatory signaling in

IECs.

In this study, we show that *V. cholerae* globally suppresses proinflammatory gene

expression in IECs and this signaling inhibition is mediated by the catalytic action of the

secreted MARTX$_{Vc}$ toxin, and not by other accessory toxins. These results explain observed

differences in the inflammatory response to various *V. cholerae* strains and why infections with

MARTX$_{Vc}^{+}$ *V. cholerae* induces a highly secretory, but non-inflammatory, diarrhea. Further, we

find that the cytoskeletal destruction initiated by the ACD conversely functions as a damage-

associated molecular pattern (DAMP) to potently induce proinflammatory gene expression

through the mitogen-activated protein kinase (MAPK) pathway; and, this response is more

robust even than that induced by bacterial PAMPs. However, in the context of the MARTX$_{Vc}$

holotoxin, co-delivery on the same toxin of RID and ABH silences pathways that would normally

transduce the signals to induce proinflammatory gene expression in response to both DAMPs

and PAMPs. These data reveal that simultaneous delivery of all three effector domains on a

single multifunctional toxin is advantageous as it promotes the ACD-mediated destruction of the

actin cytoskeleton without detection of the associated damage by the infected host. Thus,

multifunctional toxins can coordinate their multiple enzymatic activities to fine-tune the host

response to infection.

RESULTS

The MARTX$_{Vc}$, and not HlyA or HapA, inhibits IL-8 secretion from IECs

V. cholerae secretes three accessory toxins in addition to cholera toxin: the MARTX$_{Vc}$ toxin

encoded by the gene *rtxA*, α-hemolysin encoded by *hlyA*, and hemagglutinin/protease encoded

by *hapA*. To evaluate whether accessory toxins modulate IL-8 secretion in IECs, cultured IECs

were treated with live *V. cholerae* strain N16961 or previously generated derivatives with

deletions in *rtxA*, *hlyA*, or *hapA* (Fullner et al., 2002). N16961 stimulated IL-8 secretion at

concentrations similar to untreated control cells (Figure 1B), while N16961Δ*rtxA* induced

significantly more IL-8 secretion (Figure 1B). No difference was observed due to loss of HlyA or

HapA, or both toxins (*ΔhlyAΔhapA*, heretofor referred to as KFV119) (Figure 1B). These results

reveal that N16961 has the potential to stimulate release of IL-8 from IECs, but the MARTX$_{Vc}$

toxin actively suppresses the response.

The atypical El Tor O1 strain 2010EL-1786 from the 2010 Haiti outbreak is known to be

more proinflammatory (Satchell et al., 2016). This strain has a premature stop codon in *rtxA* that

prevents secretion of the MARTX$_{Vc}$ toxin (Dolores and Satchell, 2013). Consistent with results

using N16961, 2010EL-1786 induced a significant increase in IL-8 secretion from IECs, but this

response was suppressed when the stop codon in *rtxA* was restored to a Trp codon (Figure

1C). These results show that *V. cholerae* activates proinflammatory chemokine secretion, but

this is suppressed in *rtxA$^+$* *V. cholerae* strains.

MARTX$_{Vc}$ toxin effector domains, and not the pore, inhibit IL-8 secretion in IECs

The MARTX$_{Vc}$ toxin has four putative cytotoxic functions, the formation of a pore and the

biochemical activities of its three effector domains (Woida and Satchell, 2018). In previous

studies (Dolores et al., 2015), a derivative of KFV119 was generated in which the *rtxA* gene was

modified to replace sequences that encode the effector domains with an in-frame sequence for

β-lactamase (Bla). This strain produces a toxin that forms a functional pore and is able to

translocate, but lacks all effector activity (*rtxA::bla*). The individual effector domains ACD

(*acd::bla*), RID (*rid::bla*), and ABH (*abh::bla*) were then individually restored into the parent

rtxA::bla strain to create effector gain of function toxins (Figure 2A-E). Catalytically inactive

effector derivatives of these toxins and a Δ*rtx* negative control strain were also generated and

characterized.

The *rtxA::bla* strain induced similar levels of IL-8 secretion from intestinal cells as the

Δ*rtx* strain, but significantly higher than KFV119 (Figure 2F). This result indicates that the

MARTX$_{Vc}$ pore alone does not suppress IL-8 secretion. Surprisingly, the *acd::bla* strain induced

significantly more IL-8 secretion compared to both the Δ*rtx* and *rtxA::bla* strains (Figure 2F). The

rid::bla strain induced IL-8 secretion equivalent to KFV119 (Figure 2F). These results indicate

that RID is sufficient to suppress *V. cholerae*-induced IL-8 secretion, while ACD actually

exacerbates IL-8 secretion beyond stimulation by live bacteria. The *abh::bla* strain had no effect

on *V. cholerae*-induced IL-8 secretion (Figure 2F). In fact, when intestinal cells were

simultaneously co-inoculated with both the *acd::bla* and the *rid::bla* strains, IL-8 secretion was

also suppressed (Figure 2G). This was due to the acylation activity of RID, since the *rid-*

H2782A*::bla* catalytically inactive strain did not suppress IL-8. In addition, although less robust,

co-inoculation of *acd::bla* with *abh::bla* also attenuated IL-8 secretion (Figure 2G). These results

reveal that RID, and to a lesser extent ABH, inhibits both bacterial- and ACD-induced IL-8

secretion.

The MARTX$_{Vc}$ toxin inhibits expression of the *CXCL8* gene

A block to IL-8 secretion can occur by inhibition of IL-8 protein secretion or inhibition of

IL-8 (*CXCL8*) gene expression. At two hours post-inoculation, there was no significant change in

transcription of the *CXCL8* gene in intestinal cells inoculated with KFV119 or Δ*rtx* compared to

mock-treated cells (Figure 2H). There was also no significant change due to RID or ABH. By

contrast, a significant 32-fold increase was induced by ACD (Figure 2H). These results indicate

that MARTX$_{Vc}$ effector domain modulation of IL-8 secretion occurred primarily due to blocking of

CXCL8 gene transcription.

The absence of a response to Δ*rtx* was unexpected since *V. cholerae* alone induced IL-8

secretion (Figure 1B, 2F) (Zhou et al., 2004). Since chemokine secretion from cells by ELISA

requires a long incubation period following bacterial challenge, we hypothesized that the IEC

response to *V. cholerae* occurs after the 120 minutes (min) bacterial challenge in which *CXCL8*

expression was assayed. Therefore, changes in *CXCL8* expression were measured overtime

following bacterial challenge. Confirming previous results, no change in *CXCL8* expression was

observed by KFV119 at any timepoint measured (Figure 2I). However, while Δ*rtx* did not induce

changes in expression at 120 min post-inoculation, there was a significant 10-fold increase in

expression at six hours post-inoculation that eventually returned to baseline between 12 and 22

hours (Figure 2I). As observed previously, *acd::bla* induced a significant 42-fold change in

CXCL8 expression at two hours. While this response diminished overtime, *CXCL8* mRNA levels

remained 10- to 18-fold higher than mock-treated cells (Figure 2I). These data further show that

ACD-mediated *CXCL8* expression is stimulated as much as four hours before *V. cholerae*-

induced gene expression. Therefore, in addition to suppressing the host response to effector

activity, *V. cholerae* may utilize the MARTX$_{Vc}$ toxin to disable host inflammatory responses prior

to detection of bacterial PAMPs, such as flagellin and LPS (Harrison et al., 2008; Soriani et al.,

2002).

ACD, but not the MARTX$_{Vc}$ holotoxin, induces proinflammatory gene expression in IECs

To determine the breadth of the proinflammatory immune response stimulated by ACD, we

utilized a whole transcriptome RNA-sequencing (RNA-seq) approach. Cell were treated with

various bacterial strains for two hours prior to transcriptional profiling. Statistically significant

results were given a $-1 \leq 0 \leq 1$ log$_2$ fold change cut off to adjust for biologically significant

changes (Figure 3A-B). A single gene *JUN* was upregulated by KFV119 compared to

uninoculated cells, but this upregulation was not consistent when validated by qPCR (Figure

3C). Thus, there was no significant upregulation of any host gene after two hours treatment of

IECs with MARTX$_{Vc}$+ *V. cholerae*.

Similarly, *rid::bla, abh::bla,* and *Δrtx* also did not induce significant changes in IEC gene

expression by two hours. By contrast, the *acd::bla* strain induced differential regulation of over

200 genes. Many of the significantly upregulated genes by ACD were identified as regulators of

inflammation. These include genes for chemokines CXCL8 (IL-8) and CXCL3 (IL-3), cytokine

TNF, and for proinflammatory transcription factors JUN, FOS*,* and EGR1/2/3 (Fig 3B-G).

Upregulation of select proinflammatory genes was validated by qPCR (Figure 2H, 3C-G). These

data suggest that when delivered independently, ACD induces proinflammatory gene

expression more rapidly than by *V. cholerae* alone. However, when ACD is delivered on the

complete MARTX$_{Vc}$ toxin, the response is abolished.

RID and ABH suppress host MAPK signaling to modulate the ACD proinflammatory

response

Bioinformatic analysis of the *acd::bla* differentially regulated genes revealed that many

were associated with the ERK, p38, JNK MAPK and NF-κB pathways (Figure 3H). These

pathways have also been associated with induction of IL-8 by purified *V. cholerae* flagellin

(Harrison et al., 2008). The phosphorylation of MAPK in bacterial-treated cells was monitored

after two hours. While KFV119-treated cells showed no ERK phosphorylation, *Δrtx* activated

ERK signaling (Figure 3I). Further, *acd::bla* stimulated phospho-ERK levels above that of *Δrtx*

alone (Figure 3I). Both the *rid::bla* and *abh::bla* strains suppressed phospho-ERK, while *rid*-

H2782A*::bla* and *abh*-H3369A::*bla* strains did not (Figure 3I). KFV119-treated cells also showed

no phosphorylation of p38 or JNK. However, these MAPK pathways were activated by *acd::bla*

(Figure 3J and K). While previous studies suggest *V. cholerae* PAMPs may also activate these

pathways (Harrison et al., 2008), the observed limited or undetectable p38 and JNK MAPK

phosphorylation by *Δrtx* alone is likely a result of timing, since *V. cholerae* does not induce

expression of proinflammatory genes until six hours post-inoculation (Figure 2I and 3A). Finally,

none of the effector domains inhibited NF-κB, as measured by IκBα degradation (Figure 3L).

The lack of MAPK activation in MARTX$_{Vc}$$^+$ KFV119-treated cells suggests there is active

suppression of ACD-induced MAPK signaling due to interplay between the MARTX$_{Vc}$ effector

domains. However, these experiments rely on comparing strains with only one effector to those

with either all or no effectors. To study interactions between effector domains requires MARTX$_{Vc}$

toxin strains that simultaneously deliver two active effectors into a cell on a single toxin. Six new

strains with introduced codon changes in *rtxA* were generated to produce MARTX$_{Vc}$ toxins that

retain natural processing and delivery of all effectors, but with one, two, or all three of the

effectors carrying point mutations in essential catalytic site residues (Figure 4A and Figure S1).

Cells treated with these strains confirmed results above that ACD-induced IL-8 secretion is

suppressed by both RID and ABH (Figure 4B). However, unlike *abh:bla*, the ABH Active Only

strain also significantly reduced bacterial secretion of IL-8 (Figure 4B). Therefore, when

delivered on the complete holotoxin, ABH can suppress *V. cholerae* induction of IL-8.

To determine if both RID and ABH can suppress ACD upregulation of MAPK signaling,

activation of these pathways were examined in cells treated with the newly constructed strains.

The single active ACD strain stimulated ERK activation beyond that of *V. cholerae* alone, while

both the single active RID and ABH strains suppressed bacterial activation of ERK (Figure 4C).

Additionally, both the Active ACD/RID and Active ACD/ABH strain showed attenuation of ACD-

induced phospho-ERK, phospho-JNK and phospho-p38 compared to the ACD Active only strain

(Figure 4C-E).

Further, cells treated with PD98059, an inhibitor of ERK, showed no attenuation of the

IL-8 response (Figure 4F). However, inhibition of the p38 MAPK pathway by SB202190 did

inhibit IL-8 secretion (Figure 4G), while inhibition of the JNK pathway by SP600125 had no

effect on IL-8 secretion (Figure 4H).

In total, these data show that suppression of MAPK signaling pathways, particularly the

p38 pathway, can modulate the ACD-induced proinflammatory response, and this response is

attenuated by the action of RID and ABH.

Cytoskeletal collapse can activate proinflammatory signaling in ACD-treated cells, but

the signaling is not detected due to prior action of RID and ABH

ACD has two putative cytotoxic functions. The first is the inhibitor action of toxic actin oligomers

(dimers and trimers) formed when only 2-4% of total actin has been cross-linked (Heisler et al.,

2015; Kudryashova et al., 2018). The second is ACD will eventually cross-link nearly 100% of

cellular actin into higher order oligomers (10 to 15-mers) to sequester bulk actin and induce

cytoskeletal collapse (Fullner and Mekalanos, 2000). Kinetic experiments comparing actin

cross-linking abundance to MAPK activation were performed to determine which of these

functions activate MAPK signaling. The ACD Active Only strain induced a maximum of 4% of

total actin cross-linked between 5 and 15 min after addition of *V. cholerae*, corresponding to the

formation of early toxic actin oligomers (Figures 5A-B and S2). Cross-linking increased around

60 min (Figure 5A-B) corresponded to ACD sequestering bulk actin to induce actin

depolymerization and cytoskeletal collapse. In the same assayed samples, MAPK signaling was

activated between 60 and 90 min (Figure 5C-H), when over 50% total monomeric actin was

cross-linked. These results indicate that a significant portion of actin needs to be cross-linked to

activate proinflammatory signaling. While the Triple* strain showed stochastic activation of

MAPK signaling between 5 and 15 min, it induced significantly less ERK phosphorylation than

the ACD Active Only strain and there was no detection of phospho-p38 or phospho-JNK

between 30 and 120 min (Figure S3C-H). These data suggest that it is ACD sequestration of

bulk actin and cytoskeletal destruction that leads to activation of proinflammatory signaling in

IECs.

To completely block the intestinal inflammatory response, RID and ABH would have to

inactivate MAPK signaling prior to host detection of ACD-induced cytoskeletal collapse. This

would indicate that instead of reversing the activation state of MAPK pathways after 60-90 min,

RID and ABH inactive these pathways ahead of ACD-induced activation. Cells treated with

KFV119 failed to induce any changes in MAPK activation following significant actin cross-linking

(Figure 6A-F). Further, both the ACD/RID Active strain and the ACD/ABH Active strain showed

reduced phosphorylation of ERK, p38, and JNK, even following significant actin cross-linking at

60 min (Figure S4A-L). Additionally, MAPK activation induced by the Active ACD/RID and Active

ACD/ABH strains was still significantly less than activation by the ACD Active only strain at 120

min (Figure 6D-F). These data would suggest that RID and ABH are each sufficient to attenuate

ACD induction of MAPK signaling. However, modest, yet significant, activation of the JNK

pathway by the Active ACD/RID strain was still observed at 120 min (Figure S4A-F). The Active

ACD/ABH strain also induced slight, yet significant, activation of the p38 and JNK pathways at

120 min (Figure S4G-L). Since both RID and ABH alone do not completely abolish activation of

these pathways, both effectors may be required to completely abolish the global inflammatory

response observed in the RNA-seq analysis (Figure 3A). The inhibition of MAPK signaling was

not due to RID and ABH directly inhibiting ACD actin cross-linking activity. In fact, the ACD/ABH

strain showed slight, yet significant, increase in actin cross-linking at 30 and 60 min before

returning to similar abundances at 90 and 120 min (Figure S4M). Therefore, both RID and ABH

independently silence signal transduction pathways prior to host detection of ACD induced

collapse of the actin cytoskeleton without modulating ACD activity.

RID can inhibit IL-8 secretion due to actin destruction by ACD and by latrunculin A,

independent of live bacteria or formin inhibition.

To determine if the action of RID is specific to MARTX toxins or more broadly applicable,

cells were treated with purified recombinant ACD fused to the N-terminal of the anthrax lethal

toxin (LF$_N$ACD) in combination with the anthrax protective antigen (PA). This previously

characterized system allows for the delivery of MARTX effector domains into cells in the

absence of bacteria and MARTX toxin delivery (Figure 7A) (Cordero et al., 2006). Only cells

treated with LF$_N$ACD and PA induced IL-8 secretion, demonstrating that ACD is sufficient for

induction of a proinflammatory immune response in the absence of a bacterium. This response

was inhibited by co-treating cells with both LF$_N$ACD and LF$_N$RID in the presence of PA (Figure

7B). Chemical depolymerization of actin also can induce IL-8 secretion (Bobo et al., 2013).

Indeed, cells treated with latrunculin A, a sponge toxin that binds and sequesters G-actin, also

showed IL-8 secretion. Showing that the mechanisms are conserved, treatment of cells with

LF$_N$RID in the presence of PA suppressed latrunculin A induced IL-8 secretion (Figure 7C).

However, inhibiting formins with SMIFH2 failed to elicit an IL-8 response (Figure 7D). These

data support that cytoskeletal collapse stimulates the inflammatory pathways and, regardless of

the inducer, these pathways are inhibited by RID.

DISCUSSION

MARTX toxins are unique hybrids of independently secreted toxins and multifunctional effector delivery systems (Gavin and Satchell, 2015; Kim, 2018). This study demonstrates that the *V. cholerae* MARTX toxin utilizes this multifunctionality to "self-regulate" and silence the host response both to its own cytotoxic activity and detection of bacterial PAMPs. Following translocation, ACD begins cross-linking actin to produce early toxic oligomers that sequester actin binding proteins and disrupt intestinal tight junctions. While these toxic oligomers are being formed, RID and ABH block MAPK signaling. Once ACD begins to sequester bulk actin to induce cytoskeletal collapse, the cell is unable to trigger MAPK activation. Therefore, the MARTX$_{Vc}$ toxin blocks IECs from activating an innate immune response (Figure 7E). While previous studies have suggested MARTX$^+$ *V. cholerae* strains induce IL-8 secretion, those studies found this proinflammatory response to be growth phase dependent in which stationary phase cultures contain more IL-8 than log phase cultures (Zhou et al., 2004). Secretion of the MARTX$_{Vc}$ toxin is also growth phase dependent and is expressed and secreted during log phase and then toxin present in the supernatant fluids is degraded by proteases during stationary phase (Boardman et al., 2007). Therefore, no toxin would be present to suppress IL-8 secretion. Our findings support a new model in which MARTX$_{Vc}$ toxin produced during the early active growth stage in the intestines acts to suppress the IEC inflammatory response. These data indicate that MARTX$_{Vc}$ toxin suppression of intestinal innate immunity prevents host recruitment of immune cells to protect the bacteria from neutrophil-mediated clearance. Additionally, MARTX$_{Vc}$ toxin immunomodulatory activities may contribute to the differences in inflammation observed between various *V. cholerae* strains (Rodriguez et al., 2001; Zhou et al., 2004).

The current predominant circulating *V. cholerae* strains responsible for disease are altered El Tor isolate that are hypervirulent and exhibit increased clinical severity of diarrhea

(Alam et al., 2011; Son et al., 2011). This is in part due to the acquisition of mutations in H-NS

and VieA resulting in increased production of cholera toxin and HlyA, increased motility, and

inflammasome activation suggesting increased inflammation (Russell et al., 2018; Satchell et

al., 2016; Son et al., 2011). In addition, these strains naturally lack the MARTX$_{Vc}$ toxin due to a

stop codon in *rtxA*. Restoring toxin secretion in altered El Tor isolate 2010EL-1786 attenuated

IL-8 induction comparable to the parent strain (Figure 1C). Thus, loss of the MARTX$_{Vc}$ toxin

immunomodulatory activities in these altered strains may increase intestinal inflammation, which

could exacerbate disease severity and contribute to its hypervirulence.

The effector domains, and not the MARTX$_{Vc}$ pore, were identified to modulate innate

immune signaling in IECs (Figure 2, 4). These data support a previously established model in

which the MARTX pore functions primarily as an effector delivery platform and not a direct

virulence mechanism (Gavin et al., 2017). How each delivered effector contributes to

pathogenesis is still debated. While other studies have connected RID and ABH to inhibition of

macrophage phagocytosis *in vitro* (Chen et al., 2017; Zhou et al., 2017), these conclusions

contradicted previous findings that 5% inhibition by these effectors is biologically insignificant

compared to the 90% inhibition by ACD (Dolores et al., 2015). Our study validates that RID and

ABH most likely function primarily to abolish host detection of cytoskeletal damage and bacterial

PAMPs. RID-mediated inactivation of MAPK pathways most likely occurs through its direct

inactivation of Rho family GTPases. Rho GTPases regulate MAPK and other cell signaling

pathways and bacterial toxin inactivation of Rho GTPase can inhibit these pathways in specific

cell types (Schwartz, 2004; Woolery et al., 2014). However, toxin inactivation of Rho GTPases

through covalent modifications, such as ampylation or glycosylation, induces IL-8 secretion in

IECs and activates the pyrin inflammasome in macrophages (Mahida et al., 1996; Xu et al.,

2014). However, RID does not activate the inflammasome (Xu et al., 2014). Therefore, RID

acylation of the C-terminal polybasic region of Rho family GTPases allows for the inactivation of

Rho GTPases to suppress proinflammatory signaling pathways in IECs, while also evading host

detection of Rho inactivation through the inflammasome.

How cleavage of PI3P by ABH blocks proinflammatory pathways is still unknown. PI3P

is required for the formation of autophagosomes and ABH-mediated cleavage of PI3P inhibits

formation of autophagosomes and endocytic trafficking (Agarwal et al., 2015). We postulate that

ABH depletion of PI3P could block formation of scaffolding complexes on autophagosomes and

endosomes that promote recruitment and activation of MAPK signaling molecules

(Fehrenbacher et al., 2009; Martinez-Lopez et al., 2013). The loss of the scaffolds would then

prevent activation of the MAPK pathway.

ACD production of toxic actin oligomers was recently characterized as the primary

cytotoxic mechanism of ACD based on the hypothesis that the kinetics of ACD actin cross-

linking prevent it from cross-linking the majority of cellular actin (Heisler et al., 2015). Our study

reveals ACD stimulation of MAPK signaling pathways occur following cross-linking of over 50%

of cellular actin (Figure 6, 7). This supports previous findings in which type VI secreted ACD

cross-links significant quantities of actin and induce intestinal inflammation mice (Ma and

Mekalanos, 2010). While the toxic oligomers may still contribute to cytoskeletal collapse, our

data indicate that it is the destruction of the cytoskeleton, and not the direct formation of toxic

actin oligomers inhibition of formins, that activate proinflammatory signaling.

RID and ABH silencing the global response to ACD activity provide evidence that

different MARTX effector domain combinations could impact the overall host response to co-

delivered effectors. These data suggest that an effector's contribution to virulence and disease

could be enhanced or attenuated depending on which other effector domains it is delivered with.

V. vulnificus with different natural effector combinations have varying virulence potential and

promote altered host responses (Gavin and Satchell, 2018; Kwak et al., 2011; Murciano et al.,

2017). Therefore, MARTX toxin multifunctionality, in combination with variability in the effector

domain repertoire, allows for variety of effector interplay combinations which could either

enhance or attenuate MARTX toxin associated virulence.

ACKNOWLEDGEMENTS

We would like to thank the Northwestern University Center for Genetic Medicine NUSeq core

facility, especially Xinkun Wang and Matthew Schipma, for technical assistance, bioinformatic

analysis of the RNA-sequencing experiments and DNA sequencing. We would like to thank

members of the Satchell lab for their valuable input and technical support and Dr. Nicholas

Cianciotto and Dr. Gail Hecht for review of the manuscript. This work was supported by the NIH

Ruth L. Kirschstein Institutional National Research Service Award Training Grant in Immunology

and Microbial Pathogenesis T32AI007476 (to P.J.W.) and NIH grants R01AI092825 and

R01AI098369 (to K.J.F.S.).

AUTHOR CONTRIBUTIONS

P.J.W. conceptualized, designed, and conducted all experiments. K.J.F.S advised on all

experiments. P.J.W. wrote the original draft of the manuscript and both P.J.W. and K.J.F.S.

reviewed and edited the manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Agarwal, S., Kim, H., Chan, R.B., Agarwal, S., Williamson, R., Cho, W., Paolo, G.D., and

Satchell, K.J. (2015). Autophagy and endosomal trafficking inhibition by *Vibrio cholerae* MARTX

toxin phosphatidylinositol-3-phosphate-specific phospholipase A1 activity. Nat Commun. *6*,

8745.

Ahrens, S., Geissler, B., and Satchell, K.J. (2013). Identification of a His-Asp-Cys catalytic triad

essential for function of the Rho inactivation domain (RID) of *Vibrio cholerae* MARTX toxin. J

Biol Chem. *288*, 1397-1408.

Alam, M., Islam, A., Bhuiyan, N.A., Rahim, N., Hossain, A., Khan, G.Y., Ahmed, D., Watanabe,

H., Izumiya, H., Faruque, A.S., et al. (2011). Clonal transmission, dual peak, and off-season

cholera in Bangladesh. Infect Ecol Epidemiol. *1*.

Ali, M., Lopez, A.L., You, Y.A., Kim, Y.E., Sah, B., Maskery, B., and Clemens, J. (2012). The

global burden of cholera. Bull World Health Organ. *90*, 209-218A.

Anders, S., Pyl, P.T., and Huber, W. (2015). HTSeq--a Python framework to work with high-

throughput sequencing data. Bioinformatics. *31*, 166-169.

Bishop, A.L., Patimalla, B., and Camilli, A. (2014). *Vibrio cholerae*-induced inflammation in the

neonatal mouse cholera model. Infect Immun. *82*, 2434-2447.

Boardman, B.K., Meehan, B.M., and Fullner Satchell, K.J. (2007). Growth phase regulation of

Vibrio cholerae RTX toxin export. J Bacteriol. *189*, 1827-1835.

Bobo, L.D., El Feghaly, R.E., Chen, Y.S., Dubberke, E.R., Han, Z., Baker, A.H., Li, J., Burnham,

C.A., and Haslam, D.B. (2013). MAPK-activated protein kinase 2 contributes to Clostridium

difficile-associated inflammation. Infect Immun. *81*, 713-722.

Carmona-Saez, P., Chagoyen, M., Tirado, F., Carazo, J.M., and Pascual-Montano, A. (2007).

GENECODIS: a web-based tool for finding significant concurrent annotations in gene lists.

Genome Biol. *8*, R3.

Chen, C.L., Chien, S.C., Leu, T.H., Harn, H.I., Tang, M.J., and Hor, L.I. (2017). *Vibrio vulnificus*

MARTX cytotoxin causes inactivation of phagocytosis-related signaling molecules in

macrophages. J Biomed Sci. *24*, 58.

Cordero, C.L., Kudryashov, D.S., Reisler, E., and Satchell, K.J. (2006). The Actin cross-linking

domain of the *Vibrio cholerae* RTX toxin directly catalyzes the covalent cross-linking of actin. J

Biol Chem. *281*, 32366-32374.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics. *29*,

15-21.

Dolores, J., and Satchell, K.J. (2013). Analysis of *Vibrio cholerae* genome sequences reveals

unique rtxA variants in environmental strains and an rtxA-null mutation in recent altered El Tor

isolates. MBio. *4*, e00624.

Dolores, J.S., Agarwal, S., Egerer, M., and Satchell, K.J. (2015). *Vibrio cholerae* MARTX toxin

heterologous translocation of beta-lactamase and roles of individual effector domains on

cytoskeleton dynamics. Mol Microbiol. *95*, 590-604.

Egerer, M., and Satchell, K.J. (2010). Inositol hexakisphosphate-induced autoprocessing of

large bacterial protein toxins. PLoS Pathog. *6*, e1000942.

Fehrenbacher, N., Bar-Sagi, D., and Philips, M. (2009). Ras/MAPK signaling from

endomembranes. Mol Oncol. *3*, 297-307.

Fullner, K.J., Boucher, J.C., Hanes, M.A., Haines, G.K., 3rd, Meehan, B.M., Walchle, C.,

Sansonetti, P.J., and Mekalanos, J.J. (2002). The contribution of accessory toxins of *Vibrio*

cholerae O1 El Tor to the proinflammatory response in a murine pulmonary cholera model. J

Exp Med. *195*, 1455-1462.

Fullner, K.J., and Mekalanos, J.J. (1999). Genetic characterization of a new type IV-A pilus

gene cluster found in both classical and El Tor biotypes of *Vibrio cholerae*. Infect Immun. *67*,

1393-1404.

Fullner, K.J., and Mekalanos, J.J. (2000). In vivo covalent cross-linking of cellular actin by the

Vibrio cholerae RTX toxin. EMBO J. *19*, 5315-5323.

Gavin, H.E., Beubier, N.T., and Satchell, K.J. (2017). The Effector Domain Region of the *Vibrio*

vulnificus MARTX Toxin Confers Biphasic Epithelial Barrier Disruption and Is Essential for

Systemic Spread from the Intestine. PLoS Pathog. *13*, e1006119.

Gavin, H.E., and Satchell, K.J. (2015). MARTX toxins as effector delivery platforms. Pathog Dis.

73, ftv092.

Gavin, H.E., and Satchell, K.J.F. (2018). RRSP and RID Effector Domains Dominate Virulence

Impact of *Vibrio vulnificus* MARTX Toxin. J Infect Dis. jiy590.

Harrison, L.M., Rallabhandi, P., Michalski, J., Zhou, X., Steyert, S.R., Vogel, S.N., and Kaper,

J.B. (2008). *Vibrio cholerae* flagellins induce Toll-like receptor 5-mediated interleukin-8

production through mitogen-activated protein kinase and NF-kappaB activation. Infect Immun.

76, 5524-5534.

Heisler, D.B., Kudryashova, E., Grinevich, D.O., Suarez, C., Winkelman, J.D., Birukov, K.G.,

Kotha, S.R., Parinandi, N.L., Vavylonis, D., Kovar, D.R., et al. (2015). ACD toxin-produced actin

oligomers poison formin-controlled actin polymerization. Science. *349*, 535-539.

Kagnoff, M.F., and Eckmann, L. (1997). Epithelial cells as sensors for microbial infection. J Clin

Invest. *100*, 6-10.

Kim, B.S. (2018). The Modes of Action of MARTX Toxin Effector Domains. Toxins (Basel). *10*.

Kim, B.S., Gavin, H.E., and Satchell, K.J. (2015). Distinct roles of the repeat-containing regions

and effector domains of the *Vibrio vulnificus* multifunctional-autoprocessing repeats-in-toxin

(MARTX) toxin. MBio. *6*, e00324.

Kim, Y.R., Lee, S.E., Kook, H., Yeom, J.A., Na, H.S., Kim, S.Y., Chung, S.S., Choy, H.E., and

Rhee, J.H. (2008). *Vibrio vulnificus* RTX toxin kills host cells only after contact of the bacteria

with host cells. Cell Microbiol. *10*, 848-862.

Kudryashov, D.S., Cordero, C.L., Reisler, E., and Satchell, K.J. (2008). Characterization of the

enzymatic activity of the actin cross-linking domain from the *Vibrio cholerae* MARTX$_{Vc}$ toxin. J

Biol Chem. *283*, 445-452.

Kudryashova, E., Heisler, D.B., Williams, B., Harker, A.J., Shafer, K., Quinlan, M.E., Kovar,

D.R., Vavylonis, D., and Kudryashov, D.S. (2018). Actin Cross-Linking Toxin Is a Universal

Inhibitor of Tandem-Organized and Oligomeric G-Actin Binding Proteins. Curr Biol. *28*, 1536-

1547 e1539.

Kwak, J.S., Jeong, H.G., and Satchell, K.J. (2011). *Vibrio vulnificus* rtxA1 gene recombination

generates toxin variants with altered potency during intestinal infection. Proc Natl Acad Sci U S

A. *108*, 1645-1650.

Lin, W., Fullner, K.J., Clayton, R., Sexton, J.A., Rogers, M.B., Calia, K.E., Calderwood, S.B.,

Fraser, C., and Mekalanos, J.J. (1999). Identification of a *Vibrio cholerae* RTX toxin gene cluster

that is tightly linked to the cholera toxin prophage. Proc Natl Acad Sci U S A. *96*, 1071-1076.

Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-

time quantitative PCR and the 2(-$\Delta\Delta$C(T)) Method. Methods. *25*, 402-408.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Ma, A.T., and Mekalanos, J.J. (2010). In vivo actin cross-linking induced by *Vibrio cholerae* type

VI secretion system is associated with intestinal inflammation. Proc Natl Acad Sci U S A. *107*,

4365-4370.

Mahida, Y.R., Makh, S., Hyde, S., Gray, T., and Borriello, S.P. (1996). Effect of *Clostridium*

difficile toxin A on human intestinal epithelial cells: induction of interleukin 8 production and

apoptosis after cell detachment. Gut. *38*, 337-347.

Martinez-Lopez, N., Athonvarangkul, D., Mishall, P., Sahu, S., and Singh, R. (2013). Autophagy

proteins regulate ERK phosphorylation. Nat Commun. *4*, 2799.

Murciano, C., Lee, C.T., Fernandez-Bravo, A., Hsieh, T.H., Fouz, B., Hor, L.I., and Amaro, C.

(2017). MARTX Toxin in the Zoonotic Serovar of *Vibrio vulnificus* Triggers an Early Cytokine

Storm in Mice. Front Cell Infect Microbiol. *7*, 332.

Nogales-Cadenas, R., Carmona-Saez, P., Vazquez, M., Vicente, C., Yang, X., Tirado, F.,

Carazo, J.M., and Pascual-Montano, A. (2009). GeneCodis: interpreting gene lists through

enrichment analysis and integration of diverse biological information. Nucleic Acids Res. *37*,

W317-322.

Olivier, V., Haines, G.K., 3rd, Tan, Y., and Satchell, K.J. (2007). Hemolysin and the

multifunctional autoprocessing RTX toxin are virulence factors during intestinal infection of mice

with *Vibrio cholerae* El Tor O1 strains. Infect Immun. *75*, 5035-5042.

Olivier, V., Queen, J., and Satchell, K.J. (2009). Successful small intestine colonization of adult

mice by *Vibrio cholerae* requires ketamine anesthesia and accessory toxins. PLoS One. *4*,

e7352.

Peterson, L.W., and Artis, D. (2014). Intestinal epithelial cells: regulators of barrier function and

immune homeostasis. Nat Rev Immunol. *14*, 141-153.

Philippe, N., Alcaraz, J.P., Coursange, E., Geiselmann, J., and Schneider, D. (2004).

Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria. Plasmid. *51*,

246-255.

Prochazkova, K., Shuvalova, L.A., Minasov, G., Voburka, Z., Anderson, W.F., and Satchell, K.J.

(2009). Structural and molecular mechanism for autoprocessing of MARTX toxin of Vibrio

cholerae at multiple sites. J Biol Chem. *284*, 26557-26568.

Qadri, F., Raqib, R., Ahmed, F., Rahman, T., Wenneras, C., Das, S.K., Alam, N.H., Mathan,

M.M., and Svennerholm, A.M. (2002). Increased levels of inflammatory mediators in children

and adults infected with *Vibrio cholerae* O1 and O139. Clin Diagn Lab Immunol. *9*, 221-229.

Queen, J., and Satchell, K.J. (2012). Neutrophils are essential for containment of *Vibrio*

cholerae to the intestine during the proinflammatory phase of infection. Infect Immun. *80*, 2905-

2913.

Rodriguez, B.L., Rojas, A., Campos, J., Ledon, T., Valle, E., Toledo, W., and Fando, R. (2001).

Differential interleukin-8 response of intestinal epithelial cell line to reactogenic and

nonreactogenic candidate vaccine strains of *Vibrio cholerae.* Infect Immun. *69*, 613-616.

Russell, R., Wang, H., Benitez, J.A., and Silva, A.J. (2018). Deletion of gene encoding the

nucleoid-associated protein H-NS unmasks hidden regulatory connections in El Tor biotype

Vibrio cholerae. Microbiology. *164*, 998-1003.

Satchell, K.J. (2015). Multifunctional-autoprocessing repeats-in-toxin (MARTX) Toxins of

Vibrios. Microbiol Spectr. *3*, VE-0002-2014.

Satchell, K.J., Jones, C.J., Wong, J., Queen, J., Agarwal, S., and Yildiz, F.H. (2016). Phenotypic

Analysis Reveals that the 2010 Haiti Cholera Epidemic Is Linked to a Hypervirulent Strain. Infect

Immun. *84*, 2473-2481.

Schwartz, M. (2004). Rho signalling at a glance. J Cell Sci. *117*, 5457-5458.

Sheahan, K.L., Cordero, C.L., and Satchell, K.J. (2004). Identification of a domain within the

multifunctional *Vibrio cholerae* RTX toxin that covalently cross-links actin. Proc Natl Acad Sci U

S A. *101*, 9798-9803.

Sheahan, K.L., and Satchell, K.J. (2007). Inactivation of small Rho GTPases by the

multifunctional RTX toxin from *Vibrio cholerae*. Cell Microbiol. *9*, 1324-1335.

Shen, A., Lupardus, P.J., Albrow, V.E., Guzzetta, A., Powers, J.C., Garcia, K.C., and Bogyo, M.

(2009). Mechanistic and structural insights into the proteolytic activation of *Vibrio cholerae*

MARTX toxin. Nat Chem Biol. *5*, 469-478.

Son, M.S., Megli, C.J., Kovacikova, G., Qadri, F., and Taylor, R.K. (2011). Characterization of

Vibrio cholerae O1 El Tor biotype variant clinical isolates from Bangladesh and Haiti, including a

molecular genetic analysis of virulence genes. J Clin Microbiol. *49*, 3739-3749.

Soriani, M., Bailey, L., and Hirst, T.R. (2002). Contribution of the ADP-ribosylating and receptor-

binding properties of cholera-like enterotoxins in modulating cytokine secretion by human

intestinal epithelial cells. Microbiology. *148*, 667-676.

Tabas-Madrid, D., Nogales-Cadenas, R., and Pascual-Montano, A. (2012). GeneCodis3: a non-

redundant and modular enrichment analysis tool for functional genomics. Nucleic Acids Res. *40*,

W478-483.

Weill, F.X., Domman, D., Njamkepo, E., Tarr, C., Rauzier, J., Fawal, N., Keddy, K.H., Salje, H.,

Moore, S., Mukhopadhyay, A.K., et al. (2017). Genomic history of the seventh pandemic of

cholera in Africa. Science. *358*, 785-789.

Woida, P.J., and Satchell, K.J.F. (2018). Coordinated delivery and function of bacterial MARTX

toxin effectors. Mol Microbiol. *107*, 133-141.

Woolery, A.R., Yu, X., LaBaer, J., and Orth, K. (2014). AMPylation of Rho GTPases subverts

multiple host signaling processes. J Biol Chem. *289*, 32977-32988.

Xu, H., Yang, J., Gao, W., Li, L., Li, P., Zhang, L., Gong, Y.N., Peng, X., Xi, J.J., Chen, S., et al.

(2014). Innate immune sensing of bacterial modifications of Rho GTPases by the Pyrin

inflammasome. Nature. *513*, 237-241.

Zhou, X., Gao, D.Q., Michalski, J., Benitez, J.A., and Kaper, J.B. (2004). Induction of interleukin-

8 in T84 cells by *Vibrio cholerae*. Infect Immun. *72*, 389-397.

Zhou, Y., Huang, C., Yin, L., Wan, M., Wang, X., Li, L., Liu, Y., Wang, Z., Fu, P., Zhang, N., et

al. (2017). N(epsilon)-Fatty acylation of Rho GTPases by a MARTX toxin effector. Science. *358*,

528-531.

FIGURES AND FIGURE LEGENDS

Figure 1. The MARTX$_{Vc}$ toxin, and not other *V. cholerae* accessory toxins, suppress IL-8

in IECs

(A) Schematic of MARTX$_{Vc}$ toxin activity in IECs. The MARTX$_{Vc}$ toxin utilizes N- and C-terminal

regions to form a pore in a target eukaryotic membrane and translocate the central effector

domains and CPD. CPD binds to InsP$_6$, which activates the domain autoproteolytic activity to

separate and release the three effector domains from the holotoxin.

(B, C) IL-8 secretion to the media from T84 intestinal cells inoculated with *V. cholerae* strains as

indicated Data are reported as means ± standard deviation (s.d.) (*n*=3-6) (** p<0.01,

****p<0.0001, One-way ANOVA with multiple comparison's test).

594

595 **Figure 2. The MARTX_{Vc} toxin effector domains, and not the pore, differentially regulate**

596 **IL-8 secretion in IECs**

597 (A-E) Schematics of MARTX$_{Vc}$ toxin effector arrangements in strains modified to translocate β-

598 lactamase (Bla) and single effector gain-of-function strains (Dolores et al., 2015).

599 (F, G) IL-8 secretion measured from T84 intestinal cells inoculated with one or two *V. cholerae*

600 effector-free Bla and single effector gain of function strains as indicated. Data are reported as

601 means ± s.d. (*p<0.05, ** p<0.01, ****p<0.0001, One-way ANOVA with multiple comparisons).

602 (H, I) qPCR detection of *CXCL8* expression from T84 cells at (H) 120 min post-inoculation or (I)

603 over 22 hours with *V. cholerae* strains as indicated. Data are reported as means ± s.d. (**

604 p<0.01, ***p<0.001, ****p<0.0001, One-way ANOVA with multiple comparisons. For panel I,

605 multiple comparisons between *acd::bla* and PBS were set as a normalized fold change of 1. ##

606 p<0.001, One-way ANOVA with multiple comparisons between Δ*rtx* and PBS set as a

607 normalized fold change of 1).

608

Figure 3. ACD induces a global proinflammatory response that is silenced when

delivered with RID and ABH on the wild-type toxin

(A, B) Heat maps of differentially expressed genes identified in RNA-sequencing analysis of T84

cells inoculated with *V. cholerae* strains as indicated. Heat maps represent statistically

significant differentially expressed genes with a $-1 \leq 0 \leq 1$ $\log_2$ fold change cut off.

(C-G) qPCR validation of select differentially expressed genes. Data are reported as the mean $\pm$

s.d (** p<0.01, ***p<0.001, ****p<0.0001, One-way ANOVA with multiple comparisons between

acd::bla and indicated strain. No other strains induced a statistically different response

compared to the normalized PBS control or another strain).

(H) ACD significantly upregulated genes involved in regulation or downstream of the MAPK and

NF-κB signaling pathways.

(I-L) Western blot analysis of (I) phospho-ERK, (J) phospho-p38, (K) phospho-JNK, and (L)

IκBα degradation from T84 cells inoculated with *V. cholerae* strain as indicated. Blots are

representative of three independent experiments.

Figure 4. Both RID and ABH suppress ACD induced proinflammatory response

(A) Schematic of MARTX$_{Vc}$ toxin identifying catalytic residues mutated in single, double, and triple catalytically inactive MARTX$_{Vc}$ toxin effector strains.

(B) IL-8 secretion measured from T84 cells inoculated with *V. cholerae* strains as indicated. Data are reported as means ± s.d (*p<0.05, ** p<0.01, ****p<0.0001, One-way ANOVA with multiple comparisons).

(C-E) Western blot analysis of (C) phospho-ERK, (D) phospho-p38, and (E) phospho-JNK from

T84 cells inoculated with strains as indicated. Blots are representative of three independent

experiments.

(F-H) IL-8 secretion measured from T84 cells pre-treated with (F) ERK inhibitor PD98059, (G)

p38 inhibitor SB202190, and (F) JNK inhibitor SP600125. Data are reported as means ±

s.d.($*p<0.05$, ** $p<0.01$, ****$p<0.0001$, Student's t-test).

Figure 5. ACD activates proinflammatory MAPK signaling following sequestration of bulk actin

(A, B) Representative western blot and quantification (*n*=3) of actin cross-linking from T84 cells inoculated with the ACD only strain of *V. cholerae*. Actin oligomers start forming around 5 min to sequester actin binding proteins while significant cross-linking and sequestration of bulk actin and cytoskeleton destruction start forming at 60 min. Data are reported as means ± s.d.(** p<0.01, ***p<0.001, One-way ANOVA with multiple comparisons between indicated timepoints and 0% actin cross-linking from 0 min control).

(C - E) Fold change of (C) phospho-ERK (*n*=4), (D) phospho-p38 (*n*=2), and (E) phospho-JNK (*n*=3) compared to 0 min control quantified from western blot band densities from T84 cells

inoculated with the ACD Active only strain plotted with quantification of percent actin cross-

linked over the 120 min incubation. Data are reported as means ± s.d (** $p<0.01$, ***$p<0.001$,

One-way ANOVA with multiple comparisons between indicated timepoints and 0% actin cross-

linking from 0 min control from Figure 5B. ## $p<0.01$, ### $p<0.001$, #### $p<0.0001$, One-way

ANOVA with multiple comparisons between phospho-ERK/JNK at indicated timepoints and a 0

min control with a normalized fold change of 1). A third biological replicate was performed for

phospho-p38 quantification but the ACD Active only strain induced a stochastically high 273-fold

change compared to 0 min control (Figure S3A-B).

(F-H) Representative western blots (*n*=3-4) of (F) phospho-ERK, (G) phospho-p38, and (H)

phospho-JNK activation during 120 min bacterial challenge.

Figure 6. RID and ABH block MAPK signaling prior to host detection to ACD

sequestration of bulk actin

(A–C) Fold change of (A) phospho-ERK, (B) phospho-p38, and (C) phospho-JNK compared to 0

min control quantified from western blot band densities from T84 cells inoculated with the WT

KFV119 *V. cholerae* strain plotted with quantification of percent actin cross-linked over the 120

min incubation. Data are reported as the mean ± s.d.(****p<0.0001, One-way ANOVA with

multiple comparisons between indicated timepoints and 0% actin cross-linking from 0 min

control).

(D-F) Representative western blots of (D) phospho-ERK, (E) phospho-p38, and (F) phospho-

JNK activation during 120 min bacterial challenge.

(G-I) Quantification of fold change in (G) phospho-ERK, (H) phospho-p38, and (I) phospho-JNK

compared to the Active ACD only strain at 120 minutes post-inoculation. Data are reported as

the mean $\pm$ s.d (*$p<0.05$, **$p<0.01$, ****$p<0.0005$, Student's t-tests compared to Active ACD only

strain). No statistically significant difference was observed between KFV119, ACD/RID Active,

or ACD/ABH Active strains compared to each other.

Figure 7. Cytoskeletal collapse, and not formation of toxic actin oligomers, activates

proinflammatory response in IECs

(A) Schematic of LF_N- Effector intoxication. (1) PA binds to the anthrax toxin receptor. (2) PA is

processed to its 63 kDa active form (PA_{63}). (3) PA_{63} oligomerizes to a heptamer complex. (4)

The LF_N domain of the LF_N-MARTX effector fusion binds to the PA_{63} heptamer. (5) The LF_N-

Effector + PA63 complex enters the cell through receptor mediated endocytosis. (6) Acidification

of the vacuole promotes the PA_{63} heptamer to form a pore in the vacuole membrane to release

LF_N-Effector fusion protein into the cytosol.

(B) IL-8 secretion measured from T84 intestinal cells treated with LF_N alone, LF_NACD, or

LF_NRID in the presence or absence of PA. Data are reported as means ± s.d.(****$p < 0.0001$,

One-way ANOVA with multiple comparisons).

(C, D) IL-8 measured from T84 cells treated with (C) latrunculin A alone or (D) SMIFH2 in

combination with LF_NRID in the presence of PA where indicated. Data are reported as means ±

s.d.(*** $p < 0.001$, ****$p < 0.0001$, One-way ANOVA with multiple comparisons).

(E) Model of interplay between the $MARTX_{Vc}$ toxin multiple functions. (Left) $MARTX_{Vc}$ toxin

effector domains are separated from the holotoxin. (Middle) During the early stages of effector

activity, ACD actin cross-linking forms toxic actin oligomers to disrupt tight junctions. RID and

ABH inactive MAPK signaling pathways. (Right). Inactivation of MAPK pathways blocks the

inflammatory response to ACD destruction of the cytoskeleton and host detection of bacterial

PAMPs.

Materials and Methods

Antibodies and chemical reagents

Antibodies used in this study include phospho-p44/p42 MAPK (ERK1/2) (Cell Signaling Technology, #4337S), p44/p42 MAPK (ERK1/2) (Cell Signaling Technology, #4965S), phospho-p38 MAPK (Cell Signaling Technology, #9211S), p38 MAPK (Cell Signaling Technology, #9212S), phospho-SAPK/JNK (Cell Signaling Technology, #9255S), SAPK/JNK (Cell Signaling Technology, #9252S), Vinculin (Cell Signaling Technology, #13901), IKBα (Cell Signaling Technology, #9242S), Tubulin (Cell Signaling Technology, #2144), GAPDH (Santa Cruz, #sc-25779), Actin (Sigma, #A2066), and LICOR IRDye 800CW/680LT secondary antibodies (LICOR, #926-3221, #926-3211, #926-68070, #926-68071). Chemical reagents used in this study were ERK MAPK inhibitor PD98059 (Cell Signaling Technology, #9900L), p38 MAPK inhibitor SB202190 (Cell Signaling Technology, #S7067), JNK MAPK inhibitor SP600125 (Abcam, #120065), Latrunculin A (Millipore, #428021), and SMIFH2 (Fisher Scientific, #440110).

Cell culture

T84 male colorectal carcinoma cells acquired from the American Type Culture Collection (ATCC, #CCL-248) were cultured in Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12 (DMEM/F12), GlutaMAX supplement (ThermoFisher Gibco, #10565018) with 10% (w/v) fetal bovine serum (FBS, Gemini Bio-products, #900-108) and 0.1% penicillin/streptomycin (ThermoFisher Gibco). Cells were maintained at 37°C in the presence of 5% CO_2.

Bacterial growth medium

Bacterial strains and plasmids used in this study are listed in Table S1. All *V. cholerae* strains used in this study are derived from spontaneous streptomycin-resistant derivatives of

clinical isolates N16961 or 2010EL-1796. *V. cholerae* and *Escherichia coli* were grown on Luria-

Bertani (LB) broth or agar. *V. cholerae* medium was supplemented with 100 μg mL^{-1}

streptomycin, 2 μg mL^{-1} chloramphenicol, 100 μg mL^{-1} ampicillin, or 5% (w/v) sucrose as

needed. *E. coli* growth medium was supplemented with 10 μg mL^{-1} chloramphenicol, 100 μg mL^{-1}

^{1} ampicillin, or 50 μg mL^{-1} kanamycin as needed.

<u>Treatment of cells with live bacteria</u>

V. cholerae strains were grown at 30°C in LB medium supplemented with 100 μg mL^{-1}

streptomycin. Overnight cultures were diluted 1:100 and grown at 30°C with shaking until

exponential phase (OD$_{600}$≈0.40-0.60). Bacteria from 1 mL were pelleted by centrifugation and

resuspended in phosphate buffered saline (PBS) to a final concentration of 5 x 10^{8} bacterial

cells mL^{-1}. T84 cells (seeded the day before into cell culture-treated plates) were twice washed

with PBS and then media changed to antibiotic- and FBS-free DMEM/F12. Resuspended *V.*

cholerae was added to media over cells (multiplicity of infection = 5). Inoculations were

synchronized by centrifugation at 500 x*g* for 3 min. T84 cells were subsequently incubated at

37°C in the presence of 5% CO_2 until processed for downstream applications.

<u>LF$_N$Effector intoxication of T84 cells</u>

LF$_N$ fusion proteins were expressed from pTCO24 (LF$_N$ACD) and pKS119 (LF$_N$RID) and

purified as previously described (Cordero et al., 2006; Sheahan and Satchell, 2007). In brief, *E.*

coli containing BL21(DE3)(pMagic) with overexpression plasmids were grown in Terrific broth

and expression of protein induced with 1 mM isopropyl β-D-thiogalactopyranoside at 25°C

overnight. Bacteria were harvested in Buffer A (10mM Tris, 500 mM NaCl pH 8.3), lysed by

sonication, and lysate clarified by centrifugation at 16,0000 x*g* for 30 min. The 6xHis-tagged

recombinant proteins were purified using Ni-NTA HisTrap column followed by size-exclusion

chromatography using a Superdex 75 column in Buffer A with 5 mM β-mercaptoethanol using

the ÄKTA protein purification system (GE Healthcare). Proteins were stored in 10% glycerol at

-80°C.

For intoxication of cells, media was replaced over 10^5 cells previously seeded in a 12-well

plate. 31.7 nM PA alone (List Labs, #171E) or in the presence of 13.6nM LF_N, LF_N-ACD, or LF_N-

RID was added to media and cells were incubated for 20 hr at 37°C in the presence of 5% CO_2.

For co-intoxication of 13.6 nM LF_NACD and LF_N-RID, 3x excess PA was used to ensure equal

translocation of both effectors.

Construction of pDS132 *sacB*-counterselectable plasmids

Fragments (gBlocks) containing either the ACD E1990A mutation or the ABH H3369A

mutation along with 500 base pairs up and downstream from the mutations were commercially

synthesized. Each fragment also has additional secondary silent mutations to introduce novel

MfeI and PstI restriction sites near the E1990A and H3369A mutations, respectively. Sequences

are listed in Table S2. Fragments were cloned into pDS132 digested with SphI (New England

Biolabs, #R3182S) using Gibson Assembly (New England Biolabs, #E2611S) according to

manufacturer's instructions. Plasmids were recovered and propagated in DH5αλ*pir* on LB

supplemented with chloramphenicol and confirmed by sequencing.

Transfer of E1990A, H2782A, and H3369A mutations to the *V. cholerae* chromosome

The ACD E1990A and ABH H3369A mutations on the above pDS132-based plasmids

were transferred to SM10λ*pir*. These mutations or the RID H2782A mutation from pSA129 were

transferred to the *V. cholerae* chromosome by conjugation followed by *sacB*-dependent

counterselection for double homologous recombination as previously described (Ahrens et al.,

2013; Dolores et al., 2015). To confirm recombinants gained the desired mutations, regions

corresponding to the mutations were amplified by PCR, digested with the introduced novel

restriction site, and products were separated on agarose gel. Presence of introduced mutations

were also confirmed by sequencing.

IL-8 enzyme-linked immunosorbent assay (ELISA)

1- to 2 x 10^5 T84 cells seeded in a 12-well tissue culture treated plate were treated as

described above for 2 hr. Media from inoculated cells was removed, cells were washed once

with warm PBS and media changed to DMEM12 GlutaMAX media supplemented with serum,

pen-strep, and 100 μg mL^{-1} gentamicin. Cells were incubated for an additional 20 hr at 37°C in

the presence of 5% CO_2. For IL-8 chemical inhibitor studies, 10 μM of MAPK inhibitor or 0.33%

DMSO control were added to T84 cells one hour prior to bacterial challenge. Inhibitors were

reapplied following 2 hr bacterial challenge. Media from the T84 cells was harvested and spun

down at 20,000xg for 1 min at 4°C and supernatant was collected. Concentration of IL-8 in cell

media was measured using the IL-8 Human Matched Antibody Pair ELISA kit (ThermoFisher,

#CHC1303) following manufacturer's instructions.

Western blot analysis of actin crosslinking and cell signaling pathways

1 x 10^6 T84 cells in a 6-well tissue culture dish were treated with various *V. cholerae*

strains as described above for 2 hr. Cells were washed once with cold PBS and removed from

the plate in 150 μL lysis buffer (150 mM NaCl, 20 mM TRIS pH 7.5, 1% Triton X-100, and Pierce

Protease and Phosphatase inhibitor added prior to use) using a cell scrapper. Lysates were

incubated on ice for 15 min and then clarified by centrifugation at 20,000xg at 4°C for 10 min.

Concentration of protein in the collected supernatant fluid determined using the bicinchoninic

(BCA) assay (ThermoFisher, #23227). Normalized samples were boiled for 5 min at 95°C in

SDS loading buffer and protein separated on either a 10% or 15% SDS-polyacrylamide gel.

Proteins were transferred to nitrocellulose membranes and blocked in TBS (10 mM Tris pH=7.4,

150 mM NaCl) with 5% (w/v) milk for 1 hour. Membranes were washed with TBS and then

incubated in indicated primary antibodies 1:1000 in TBS with 5% (w/v) bovine serum album

(BSA, Fisher Bioreagents) overnight at 4°C. Membranes were washed with TBS and probed in

1:10,000 IRDye 800CW/680LT secondary antibody for 1 hour before being washed again and

imaged using the LI-COR Bioscience Odyssey imaging system. Quantification of band density

was conducted using Fiji/ImageJ.

RNA-seq

The stranded mRNA-seq was conducted in the Northwestern University NUSeq Core

Facility. Briefly, total RNA examples were checked for quality using RINs generated from Agilent

Bioanalyzer 2100. RNA quantity was determined with Qubit fluorometer. The Illumina TruSeq

Stranded mRNA Library Preparation Kit was used to prepare sequencing libraries from 750 ng

of high-quality RNA samples (RIN=10). The Kit procedure was performed without modifications.

This procedure includes mRNA purification and fragmentation, cDNA synthesis, 3' end

adenylation, Illumina adapter ligation, library PCR amplification and validation. Illumina NextSeq

500 Sequencer was used to sequence the libraries with the production of single-end, 75 bp

reads.

The quality of DNA reads, in fastq format, was evaluated using FastQC. Adapters were

trimmed, and reads of poor quality or aligning to rRNA sequences were filtered. The cleaned

reads were aligned to the human reference genome using STAR (Dobin et al., 2013). Read

counts for each gene were calculated using htseq-count (Anders et al., 2015). Normalization

and differential expression were determined using DESeq2 (Love et al., 2014). The cutoff for

determining significantly differentially expressed genes was an FDR-adjusted p-value less than

0.05. A pathway analysis was performed on both gene lists using GeneCoDis (Carmona-Saez

et al., 2007; Nogales-Cadenas et al., 2009; Tabas-Madrid et al., 2012) to identify pathways that

are enriched with genes that are upregulated and downregulated.

<u>Quantitative RT-PCR</u>

1×10^6 T84 cells in a 6-well tissue culture treated plate were treated with *V. cholerae* as

described above. At indicated time points, mRNA was harvested using the Qiagen RNeasy kit

(Qiagen, #74014) following manufacturer's instructions. RNA isolated was measured using a

Nano-drop 100 spectrophotometer. Reverse transcription was performed using random

hexamers (Roche) and Superscript III Reverse Transcriptase (Invitrogen, #18080044) in the

presence of RNase OUT (Invitrogen, #10777019) or RNasin (Promega, #N2611) under the

following conditions: 25°C for 5 min, then 55°C for 60 min, 95°C for 5 min. Remaining RNA was

hydrolyzed using 1 N NaOH. Quantitative PCR was performed using iQ SYBR Green supermix

(Bio-Rad, #1708880) on the iQ5 Multicolor RealTime PCR Detection System using gene

specific primers indicated in Table S3. Relative change in gene expression compared to PBS

control was determined the $\Delta\Delta$CT method (Livak and Schmittgen, 2001).

<u>Quantification and statistical analysis</u>

All experiments were done at least in triplicate and quantitative results are reported as the

mean $\pm$ standard deviation (s.d.). Statistical analysis was performed using GraphPad Prism v6.0

as detailed in the figure legends. Statistical differences in ELISA and qPCR results were

determined by one-way ANOVA followed by multiple comparison's test. Statistical difference in

results comparing suppression of ERK, p38, and JNK signaling by various *V. cholerae* strains

compared to the ACD Active only strain were determined using Student's t-tests.

Figure S1. Schematic of single and double catalytically active MARTX$_{Vc}$ toxin effector strains and triple inactive MARTX$_{Vc}$ toxin effector strain.

(A) Wild type toxin. (B) RID Active only strain with RID and ABH catalytically inactive. (C) RID Active only strain with ACD and ABH catalytically inactive. (D) ABH Active only strain with ACD and RID catalytically inactive. (E) ACD/RID active strain with only ABH catalytically inactive. (F) ACD/ABH Active strain with only RID catalytically inactive. (G) RID/ABH Active strain with only ACD catalytically inactive. (H) Triple inactive strain (Triple*) with all three effectors catalytically inactive.

Figure S2. Cells harvested in Triton X-100 lysis buffer for assessment of MAPK phosphorylation underestimates the rate of actin cross-linking further supporting that extensive cross-linking occurs ahead of inflammation.

(A) Quantification of actin cross-linking from T84 cells treated with KFV119 or the ACD Active Only strain with washed cells being lysed and collected in Triton X-100 lysis buffer followed by centrifugation and protein quantification or by direct resuspension and boiling in 2x SDS loading buffer. Data reported as mean ± s.d.

(B) Representative western blot of actin cross-linking from T84 cells inoculated with KFV119 and samples collected in SDS loading buffer.

(C) Representative western blot of actin cross-linking form T84 cells inoculated with the

ACD Active only strain and samples collected in SDS loading buffer.

Triton X-100 lysis underestimates actin cross-linking at early time points further

supporting that extensive cross-linking activate the inflammatory response. Specifically,

there is a statistically significant increase in actin cross-linking observed when KFV119

treated cells were lysed using SDS loading buffer compared to Triton X-100 at 30

minutes (*$p<0.05$, Two-way ANOVA with multiple comparisons). However, no significant

difference was observed at any time point between the two methods when cells were

treated with the ACD Active only strain. No significant difference was observed

comparing samples treated with either KFV119 and the ACD Active only strain when

using either method.

Figure S3. MAPK activation over 120 min challenge with the triple inactive (Triple*)

MARTX$_{Vc}$ toxin effector strain.

(A) Western blot analysis of p-38 activation by the ACD only strain 273-fold times higher at 120

min compared to the 0 min control. Non-specific bands are indicated as n.s.

(B) Quantification of p-38 activation from western blot presented in panel A.

(C-E) Fold change of (C) phospho-ERK, (D) phospho-p38, and (E) phospho-JNK compared to 0

min control quantified from western blot band densities from T84 cells inoculated with the Triple*

V. cholerae strain plotted with quantification of percent actin cross-linked over the two hour

incubation.

(F-H) Representative western blots of (G) phospho-ERK, (H) phospho-p38, and (I) phospho-

JNK activation during 120 min bacterial challenge.

Figure S4. RID and ABH block MAPK signaling prior to host detection of sequestration of

bulk actin without modulating ACD activity.

(A-C) Fold change of (A) phospho-ERK, (B) phospho-p38, and (C) phospho-JNK compared to 0

min control quantified from western blot band densities from T84 cells inoculated with the

ACD/RID active *V. cholerae* strain plotted with quantification of percent actin cross-linked over

the two hour incubation (**$p<0.01$, ****$p<0.0001$, One-way ANOVA with multiple comparisons

between indicated timepoints and 0% actin cross-linking from 0 min control. # $p<0.05$, One-way

ANOVA with multiple comparisons between phospho-JNK at indicated timepoint and a 0 min

control with a normalized fold change of 1).

(D-F) Representative western blots of (D) phospho-ERK, (E) phospho-p38, and (F) phospho-

JNK activation during 120 min bacterial challenge.

(G-I) Fold change of (G) phospho-ERK, (H) phospho-p38, and (I) phospho-JNK compared to 0

min control quantified from western blot band densities from T84 cells inoculated with the

ACD/ABH active *V. cholerae* strain plotted with quantification of percent actin cross-linked over

the 120 min incubation. (****$p<0.0001$, One-way ANOVA with multiple comparisons between

indicated timepoints and 0% actin cross-linking from 0 min control. # $p<0.05$, One-way ANOVA

with multiple comparisons between phospho-p38/JNK at indicated timepoints and a 0 min

control with a normalized fold change of 1).

(J-L) Representative western blots of (J) phospho-ERK, (K) phospho-p38, and (L) phospho-JNK

activation during 120 min bacterial challenge.

(M) Comparison of percent actin cross-linking during 120 min bacterial challenge of ACD

positive *V. cholerae* strains (* $p<0.05$, ** $p<0.01$ Two-way ANOVA with multiple comparisons

between KFV119 and Active ACD/ABH, #$p<0.05$, Two-way ANOVA with multiple comparisons

between Active ACD only and Active ACD/ABH).

Table S1. Bacterial strains and plasmid used in study.

Designation	Relevant Description	Source
E. coli		
BL21(DE3)(pMagic)	Protein overexpression, Km^R	A. Joachimiak (Argonne National Laboratory)
DH5α λ*pir*	Plasmid cloning	Lab stock
SM10λ*pir*	Conjugation to *V. cholerae,* Sm^R	Lab stock
V. cholerae		
N16961Sm	El Tor O1, Wild-type, Sm^R	(Lin et al., 1999)
KFV43	N16961Δ*hapA*	(Fullner and Mekalanos, 1999)
VOV21	N16961Δ*hlyA*	(Olivier et al., 2007)
CW123	N16961Δ*rtxA*	(Sheahan et al., 2004)
KFV119	N16961Δ*hlyA*Δ*hapA*	(Dolores et al., 2015)
JD23	KFV119Δ*rtxABCD*	(Dolores et al., 2015)
JD1	KFV119 *rtxA::bla*	(Dolores et al., 2015)
JD20	KFV119 *acd::bla*	(Dolores et al., 2015)
JD19	KFV119 *rid::Bla*	(Dolores et al., 2015)
JD21	KFV119 *rid-H2782A::bla*	(Dolores et al., 2015)
JD2	KFV119 *abh::bla*	(Dolores et al., 2015)
JD15	KFV119 *abh-H3369A::bla*	(Dolores et al., 2015)
2010EL-1786	Atypical El Tor O1, Wild-type, Sm^R	ATCC #BAA-2163
JD16	2010EL-1786Ω pJD22 (2010EL-1786 *rtxA+*)	(Dolores and Satchell, 2013)
PJWV1	KFV119 *rtxA-E1990A* (RID/ABH Active)	This Study
PJWV2	KFV119 *rtxA-H2782A* (ACD/ABH Active)	This Study
PJWV3	KFV119 *rtxA-H3369A* (ACD/RID Active)	This Study
PJWV4	KFV119 *rtxA-E1990A-H2782A* (ABH Active Only)	This Study
PJWV5	KFV119 *rtxA-E1990A-H3369A* (RID Active Only)	This Study
PJWV6	KFV119 *rtxA-H2782A-H3369A (ACD Active Only)*	This Study
PJWV7	KFV119 *rtxA-E1990A-H2782A-H3369A* (Triple*)	This Study
Plasmids		
pTCO24	Overexpression of LF$_N$-ACD, Amp^R	(Cordero et al., 2006)

pKS119	Overexpression of LF$_N$-RID, Amp^R	(Sheahan and Satchell, 2007)
pSA129	*sacB* counterselection cloning vector pWM91with *rtxA* with fragment of H2782A codon change, Amp^R	(Ahrens et al., 2013)
pDS132	*sacB* counterselection cloning vector, oriR6K, oriT, Chl^R	(Philippe et al., 2004)
pPJW4	pDS132 with fragment of *rtxA* with E1990A codon change, Chl^R	This Study
pPJW5	pDS132 with fragment of *rtxA* with H3369A codon change, Chl^R	This Study

924
925

Table S2. Sequences of gBlocks used to clone into pDS132 and modify ACD (E1990A)

and ABH (H3369A) effector domains of *rtxA* on the *V. cholerae* genome.

Mutation	Sequence (5'-3') (introduce codon changes highlighted)
ACD E1990A	CCCAAGCTTCTTCTAGAGGTACCGCATGCCAAGCACAAGCCGATGCTC AAGGTGCTAAACAAAACGAAGGTGATCGTCCTGATCGTCAAGGCGTGA CTGGTAGTGGCCTTTCGGGTAATGCTCATAGTGTGGAAGGCGCTGGCG AAACAGACAGTCATGTCAACACCGACAGCCAAACCAACGCCGATGGCC GATTCAGTGAAGGTTTAACCGAACAAGAGCAAGAAGCGCTAGAAGGTG CGACCAACGCAGTGAACCGTTTGCAAATTAACGCAGGTATTCGAGCGA AAAACAGCGTTAGCAGTATGACTTCTATGTTCTCTGAAACAAATAGCAAG AGCATTGTTGTTCCTACCAAAGTCTCGCCTGAACCAGAGCGCCAAGAAG TGACTCGTAGAGACGTCCGTATCTCAGGGGTGAACCTCGAAAGTCTAA GTGCGGTACAGGGAAGTCAACCAACGGGTCAACTGGCTTCGAAAAGTG TCCCCGGATTTAAAAGCCATTTCGCATCGACATCAATTGGTATA**GCA**AA TGAGTTATCCGGTCTGGTGGTGGTTTTACCGAAAAACTCAGCGCAGACT TTTGGCTATGTGCATGATTCACAAGGTAACCCATTGTTCATGCTAACCAA GGATATGAATCAAGGTGGTTATAGCAACCCAGTGGGTATCAATGATATT CAAGGGGTGAACAACTGGCAGACGCATACGATTGAACTGGTTACATATC CTAGTGAAATCAGTGATACAGCAGCGGTTGAAAGTCGTAAAGAGGCAAT GCTATGGCTTGCGAAAGAGTTTACCGATCATATCAATCAGTCTAACCAC CAAAGCTTACCTCATTTAGTGAGTGATGACGGTCGTTTCACTCTGGTTAT ATCGAACTCTAAGCATCTTATTGCGGCGGGTAACGGAACCTCTATTGAT GCACAAGGCAAGACCATAGGAATGACCCCTAGTGGCCAACAAGCAACA ATGGCGATCAGTGCGAAAGAATTTGGTACAAGCTCGTCGCCGGAAGTC AGACTGCTGCATGCGATATCGAGCTCTCCCGGG
ABH H3369A	CCCAAGCTTCTTCTAGAGGTACCGCATGCAATCACTACCAGAAGCAAGG TATCGATATGCTCGCAGTCAACCTGCGTGGCTATGGTGAAAGCGACGG TGGACCAAGCGAAAAAGGCTTGTACCAAGATGCTCGCACCATGTTCAAC TACCTAGTGAATGATAAGGGTATTGACCCAAGCAACATCATCATTCACG GCTACTCAATGGGCGGTCCAATTGCCGCAGATTTAGCACGTTATGCCG CGCAAAACGGCCAAGCGGTGTCTGGCTTATTGCTTGACCGTCCTATGC CAAGCATGACCAAAGCAATCACCGCTCACGAAGTGGCGAATCCAGCGG GCATTGTGGGGGGCTATCGCGAAAGCGGTTAACGGCCAGTTCTCTGTAG AGAAAAATCTCGAAGGTTTGCCAAAAGAGACATCCATTCTGCTGTTGAC CGATAACGAAGGTTTGGGTAACGAAGGTGAGAAACTTCGTACCAAACTC ACTGCCTCTGGTTACAACGTCACTGGCGAGCAGACATTCTATGGT**GCT**G AAGCAAGCAACCGTTTGATGAGTCAATATGCGGATCAAATTGTCTCCGG TTTGTCCAGCAGTGCAAGTGTAGATGAAGACCTAGATCAACAAGGGTTG GATACCACATCAACCAAGGATCAAGGTATCTCAAATAAGAATGATCATCT GCAGGTGGTGGATAGTAAAGAAGCATTAGCGGATGGAAAAATACTCCAT AATCAAAATGTTAATAGCTGGGGCCCGATTACGGTTACACCAACGACAG ATGGTGGTGAAACCCGCTTCGACGGTCAAATCATCGTTCAAATGGAAAA CGACCCGGTAGTAGCAAAAGCGGCAGCCAATTTAGCAGGTAAACATGC TGAAAGCAGTGTGGTGGTGCAGCTCGATTCAGACGGCAACTATCGCGT GGTGTATGGCGATCCGTCAAAACTGGATGGAAAGCTACGTTGGCAGTT GGTGGGGCATGGTCGCGACCACTCAGAAACTAACAATACTCGCTTAAG TGGTTACAGTGCCGCATGCGATATCGAGCTCTCCCGGG

929 **Table S3 Primers used in study.**

Primer	Sequence	Function
IL-8_5'	AGCACTCCTTGGCAAAACTG	CXCL8 qPCR
IL-8_3'	CAAGAGCCAGGAAGAAACCA	CXCL8 qPCR
qTNF_5'	GCCAGAGGGCTGATTAGAGA	TNF qPCR
qTNF_3'	TCAGCCTCTTCTCCTTCCTG	TNF qPCR
qJUN_5'	GTCCTTCTTCTCTTGCGTGG	JUN qPCR
qJUN_3'	GGAGACAAGTGGCAGAGTCC	JUN qPCR
qFOS_5'	GGGGCAAGGTGGAACAGTTAT	FOS qPCR
qFOS_3'	CCGCTTGGAGTGTATCAGTCA	FOS qPCR
qCXCL3_5'	CGCCCAAACCGAAGTCATAG	CXCL3 qPCR
qCXCL3_3'	GCTCCCCTTGTTCAGTATCTTTT	CXCL3 qPCR
qEGR1_5'	AAAGCGGCCAGTATAGGTGA	EGR1 qPCR
qEGR1_3'	AGCCCTACGAGCACCTGAC	EGR1 qPCR
GAPDH_2_5' FWD	TTGAGGTCAATGAAGGGGTC	GAPDH qPCR
GAPDH_2_3' REV	GAAGGTGAAGGTCGGAGTCA	GAPDH qPCR
ACDVc_cat_FWD	GCACAAGCCGATGCTCAAGGTGCTAAACAAAC	*acd* E1990A PCR
ACDVc_cat_REV	GTCTGACTTCCGGCGACGAGCTTGTACCAAATT	*acd* E1990A PCR
RIDVc_cat_FWD	GGCAAAGGTAATCTTGCCAATATCGATCTGCTAGG	*rid* H2782A PCR
RIDVc_cat_REV	CTCGATAAAGCGTTTCAGCTTAATGTCGCCATC	*rid* H2782A PCR
ABHVc_cat_FWD	CACTACCAGAAGCAAGGTATCGATATGTCGCAG	*abh* H3369A PCR
ABHVc_cat_REV	CCACTTAAGCGAGTATTGTTAGTTTCTGAG	*abh* H3369A PCR