Adaptation and the Parliament of Genes

Thomas W. Scott[1]* & Stuart A. West[1]

[1]Department of Zoology, University of Oxford, Oxford, United Kingdom

*Corresponding author: E-mail: thomas.scott@zoo.ox.ac.uk

Abstract

Fields such as behavioural and evolutionary ecology are built on the assumption that natural selection leads to organisms that behave as if they are trying to maximise their fitness. However, there is considerable evidence for selfish genetic elements that change the behaviour of individuals to increase their own transmission. How can we reconcile these contradictions? We found theoretically that, when selfish genetic elements have a greater impact at the individual level, they are more likely to be suppressed, and suppression spreads more quickly. Consequently, selfish genetic elements will either have a minor impact at the individual level, or tend to be suppressed. In addition, we found that selection on selfish genetic elements favours higher levels of distortion. Consequently, selfish genetic elements will tend to evolve to make themselves more likely to be suppressed. Overall, our results suggest that even when there is the potential for considerable genetic conflict, this will often have negligible impact at the individual level.

Introduction

There is a clear contradiction between major branches of modern evolutionary biology. On the one hand, fields such as behavioural and evolutionary ecology are based on the assumption that organisms will behave as if they are trying to maximise their fitness[1-4]. Models based on fitness maximisation are used to make predictions about the selective forces (reasons) for adaptation, and these are then tested empirically[5,6]. This approach has been phenomenally successful, explaining many aspects of behaviour, life history and morphology. For example, research on: foraging, resource competition, sexual selection, parental care, sex allocation, signalling and cooperation[7-12].

However, on the other hand, there is considerable evidence for selfish genetic elements, which increase their own contribution to future generations at the expense of other genes in the same organism[13-16]. These 'selfish genes' distort traits away from the values that would maximise individual fitness, to increase their own transmission[14,17-21]. Evidence for such genetic conflict has been found across the tree of life, from simple prokaryotes to complex animals. The contradiction is that selfish genetic elements mess up individual fitness maximisation, and appear to be common. But individual fitness maximisation still appears to occur[22,23]. For example, research on sex allocation has provided phenomenal support for both the individual fitness maximisation approach, and the existence of selfish sex ratio distorters[9,14,24-26]. How can we resolve this contradiction?

Leigh[27] provided a potential solution to this contradiction by suggesting that selfish

genetic elements would be suppressed by the 'parliament of genes'. Leigh's

argument was that, because selfish genetic elements reduce the fitness of the other

genes in the organism, the rest of the genome will have a united interest in

suppressing selfish genetic elements. Furthermore, because those other genes are

far more numerous, they will be likely to win the conflict. Consequently, even when

there is considerable potential for conflict within individuals, we would still expect

fitness maximisation at the individual level[28-32]. Leigh[27] demonstrated the plausibility

of his argument by showing theoretically how a suppressor of a meiotic drive gene

could be favoured.

However, there are a number of potential problems with the parliament of genes

hypothesis. First, even if a parliament of genes could easily generate suppressors,

whether a suppressor spreads can depend upon biological details such as any cost

associated with the suppressor, the extent to which a selfish genetic element is

distorting a trait, and the population frequency of that selfish genetic element[14,33-38].

Second, even if suppressors can spread, prolonged non-equilibrium trait distortion is

possible if the spread of suppressors through populations is slow. Third, selfish

genetic elements are themselves also under evolutionary pressure to reach a level

of distortion that would maximise their transmission to the next generation – how will

this influence the likelihood that they are suppressed?[31] Fourth, segregation at

suppressor loci might expose previously suppressed selfish genetic elements[34].

We investigated the parliament of genes hypothesis theoretically. Our aim was to

investigate the extent to which genetic conflict distorts traits away from the value that

would maximise individual fitness. We found that: (i) the greater the level of distortion

caused by a selfish genetic element, the more likely and the quicker it will be

suppressed; (ii) selection on selfish genetic elements leads towards greater

distortion, making them more likely to be suppressed. We found the same patterns

with an illustrative model, and when examining three specific examples: selfish

distortion of the sex ratio by an X chromosome driver; an altruistic helping behaviour

encoded by an imprinted gene; and production of a cooperative public good encoded

on a horizontally transmitted bacterial plasmid. Furthermore, we found close

agreement when analysing scenarios with population genetic analyses and

individual based simulations. Our results suggest that even when there is potential

for considerable genetic conflict, it will have relatively little impact on traits at the

individual level.

82　Results

A selfish genetic element may be able to gain a propagation advantage through trait

distortion ('*distorter*'). Any part of the genome that does not gain the propagation

advantage from the trait distortion will be selected to suppress the distorter. This

collection of genes will comprise most of the genome, and so will constitute the

majority within the parliament of genes[20]. We take into account the large size of this

collection of genes by assuming that it is highly likely that a potential suppressor of a

distorter can arise by mutation (high mutational accessibility). Consequently, we

focus our analyses on when a distorter and its suppressor can spread. Prolonged

trait distortion may be possible if a suppressor is hard to evolve, but our assumption

of suppressor availability allows us to focus on the evolutionary dynamics that

succeed the acquisition of a suppressor by mutation[31].

Our overall aim is to assess, given the potential for suppression, the extent to which

a distorter can distort the organism trait away from the optimum for individuals (the

largest "coreplicon"[21]). In order to elucidate the selective forces in operation, we ask

four questions in a step-wise manner, with increasing complexity:

(1) In the absence of a suppressor, when can a distorter invade?

(2) When can a costly suppressor of the distorter invade?

(3) What are the overall consequences of distorter-suppressor dynamics for trait

102   values, at the individual and population level, at evolutionary equilibrium and

103   before equilibrium has been reached?

104   (4) If the extent to which the distorter manipulates the organism trait can evolve,

105   how will this influence the likelihood that it is suppressed, and hence the

106   individual and population trait values?

Illustrative Model

We assume an arbitrary trait that influences organism fitness. In the absence of

distorters, all individuals have the trait value that maximises their individual fitness.

The distorter manipulates the trait away from the individual optimum, to increase

their own transmission to offspring. We assume a large population of diploid,

randomly mating individuals. The aim of this model is to establish key aspects of the

population genetics governing distorters and their suppressors, in an abstract

setting. We will subsequently address analogous issues in three specific biological

scenarios.

(1) Spread of a Distorter

We consider a distorter, which we denote by y_1, that is dominant and distorts an

organism trait value by some positive amount, denoted by k ($k>0$). This distortion

increases the transmission of the distorter to offspring. Specifically, the distorter (y_1)

drives at meiosis, in heterozygotes, against a non-distorter (y_0), being passed into

the proportion $(1+t(k))/2$ of offspring. $t(k)$ denotes the transmission bias ($0 \leq t(k) \leq 1$)

and is a monotonically increasing function of trait distortion $\left(\frac{dt}{dk} \geq 0\right)$.

Trait distortion leads to a fitness (viability) cost ($c_{drive}(k)$) at the individual level,

reducing an individual's number of offspring from 1 to $1\text{-}c_{drive}(k)$ ($0 \leq c_{drive}(k) \leq 1$).

Owing to distorter dominance, the fitness cost of trait distortion is borne by

heterozygous as well as distorter-homozygous individuals. The fitness cost is a

monotonically increasing function of trait distortion $\left(\frac{dc_{drive}}{dk} \geq 0\right)$. We assume that $t(k)$

and $c_{drive}(k)$ do not change with population allele frequencies, but relax this

assumption in our specific models.

We first ask what frequency the distorter will reach in the population in the absence

of suppression. If we take p and p' as the population frequency of the distorter in two

consecutive generations, then the population frequency of the distorter in the latter

generation is:

$$\bar{w}\, p' = (1 - c_{drive}(k))\,(p^2 + (1-p)\,p\,(t(k)+1)),$$ (1)

where $\bar{w}$ is the average fitness of individuals in the population in the current

generation, and can be written in full as: $\bar{w} = (1 - c_{drive}(k))\big(p^2 + 2p(1-p)\big) +$

$(1-p)^2$. In Appendix 1 we show, with a population genetic analysis of equation (1),

that the distorter will spread from rarity and reach fixation when $c_{drive}(k) < t(k)(1-$

$c_{drive}(k))$. This shows that distortion will evolve when the number of offspring that the

distorter gains as a result of distortion ($t(k)(1-c_{drive}(k))$) is greater that the number of

offspring bearing the distorter that are lost as a result of reduced individual fitness

($c_{drive}(k)$).

(2) Spread of an autosomal suppressor

We assume that the distorter (y_1) can be suppressed by an unlinked autosomal allele

(suppressor), denoted by *sup*. This suppressor (*sup*) is dominant and only

expressed in the presence of the distorter (facultative), and its expression may lead

to a fitness cost to the individual, c_{sup} ($0 \leq c_{sup} \leq 1$)[39,40]. This cost can arise for multiple

reasons, including energy expenditure, or errors relating to the use of gene silencing

machinery, and is likely to be relatively low[41]. Gene silencing generally precedes the

translation of the targeted gene, and so we assume that the costs of suppression

(c_{sup}) is independent of the amount of distortion caused by the distorter (k).

We can write recursions detailing the generational change in the frequencies of the

four possible gametes, y_0/+, y_0/*sup*, y_1/+, y_1/*sup*, with the respective frequencies in

the current generation denoted by x_1, x_2, x_3 and x_4, and the frequencies in the

subsequent generation denoted by an appended dash ('):

$\bar{w} \, x_1' = x_1^2+x_1x_2+(1\text{-}t)(1\text{-}c_{drive})x_1x_3+((1\text{-}c_{sup})/2)x_1x_4+((1\text{-}c_{sup})/2)x_2x_3$ $\qquad$ (2)

$\bar{w} \, x_2' = x_1x_2+((1\text{-}c_{sup})/2)x_1x_4+x_2^2+((1\text{-}c_{sup})/2)x_2x_3+(1\text{-}c_{sup})x_2x_4$

$\bar{w} \, x_3' = (1\text{+}t)(1\text{-}c_{drive})x_1x_3+((1\text{-}c_{sup})/2)x_1x_4+((1\text{-}c_{sup})/2)x_2x_3+(1\text{-}c_{drive})x_3^2+(1\text{-}c_{sup})x_3x_4$

$\bar{w} \, x_4' = ((1\text{-}c_{sup})/2)x_1x_4+((1\text{-}c_{sup})/2)x_2x_3+(1\text{-}c_{sup})x_2x_4+(1\text{-}c_{sup})x_3x_4+(1\text{-}c_{sup})x_4^2.$

$\bar{w}$ is the average fitness of individuals in the current generation, and equals the sum

of the equations' right-hand sides. In Appendix 2, we show, with a population genetic

analysis of our system of equations (2), that a suppressor will spread from rarity if

distortion, k, is greater than some threshold value at which the cost of suppression

(c_{sup}) is less than the cost of being subjected to trait distortion, $c_{sup}<c_{drive}(k)$. A

threshold with respect to the level of distortion (k) arises because the cost of trait

distortion ($c_{drive}(k)$) increases with greater distortion, but the cost of suppression

(c_{sup}) is constant.

(3) Consequences for organism trait values

The extent of trait distortion at the individual level shows a discontinuous relationship

with the strength of the distorter. When distortion is low, a suppressor will not spread

($c_{sup}>c_{drive}(k)$) and so the level of trait distortion at the individual level will increase

with the level of trait distortion induced by the distorter (k). However, once a

threshold is reached ($c_{sup}<c_{drive}(k)$), the suppressor spreads. We show in Appendix 3

that the spread of the suppressor causes the distorter (y_1) to lose its selective

Figure 1. Distorter-suppressor dynamics and consequences for the organism.
The trait distorter (y_1) and its suppressor (*sup*) are introduced from rarity. In part (a), the resulting average trait distortion ($x_3 k$) is plotted at equilibrium, as the solid line, against the extent to which the distorter causes trait values to deviate from the individual optimum (k). Below a certain threshold strength ($c_{sup} > c_{drive}(k)$), to the left of the dashed line, the suppressor does not invade, and so the resulting trait distortion increases with the strength of the distorter (k). However, above this threshold, the suppressor invades, and the distorter is purged, restoring the trait to the individual optimum. In part (b), the number of generations for distorters that are purged at equilibrium (having been suppressed), which lie to the right of the dashed line, is plotted on a $\log_{10}$ scale. Stronger distorters are purged more quickly than weaker distorters. The numerical solutions displayed graphically assume that the cost of suppression, the transmission benefit of distortion, and the individual fitness cost of distortion, are respectively given by: $c_{sup}=0.15$; $t=0.87k$ and $c_{drive}=0.9k^{1.5}$.

advantage and be eliminated from the population, leading to an absence of distortion

at the individual level.

Overall, these results suggest that, given a relatively low cost of suppression (c_{sup}),

the level of distortion observed at the individual level will either be low or absent.

When a distorter is weak (low k), it will not be suppressed, but it will only have a

small influence at the level of the individual. When a distorter is strong (high k), it will

be suppressed and so there will be no influence at the level of the individual (Fig.

1a).

We also found that stronger distorters are suppressed more quickly (Fig. 1b). In

Appendix 4, we numerically iterated our recursions to determine how many

generations it takes for suppressors to reach equilibrium. As long as trait distortion

continues to reduce individual fitness non-negligibly after suppression is favoured

(such that $\frac{dt}{dk} / \frac{dc_{drive}}{dk}$ is not excessively high after $c_{sup} < c_{drive}(k)$), stronger distorters

(higher k) are suppressed and purged more rapidly than weaker distorters, limiting

the potential for non-equilibrium trait distortion (Fig. 1b).

(4) Evolution of trait distortion

We then considered the consequence of allowing the level of trait distortion (k) to

evolve. We assume a distorter (y_1) that distorts by k, and then introduce a rare

mutant (y_2) that distorts by a different amount $\hat{k}$ ($\hat{k} \neq k$). This mutant (y_2) is propagated

into the proportion $(1+t(\hat{k})-t(k))/2$ of the offspring of y_2y_1 heterozygotes, and into the

proportion $(1+t(\hat{k}))/2$ of the offspring of y_2y_0 heterozygotes. We assume that the

stronger of the two distorters is dominant, but found similar results when assuming

additivity (Appendix 5). We assume that the similarity in coding sequence and

regulatory control means that the original distorter and the mutant are both

Figure 2. Evolution of trait distortion. In part (a), a distorter and suppressor are introduced in our
agent-based simulation model (Appendix 7), and the following parameter regime is chosen: c_{sup}=0.1,
t=k and c_{drive}=max(k_a,k_b)/2. The population average distorter and suppressor strengths over 100
simulation runs are plotted for successive generations. Initially, both distorter and suppressor strength
increases. Eventually, a threshold is passed, after which, distorters are purged from the population,
meaning the trait is undistorted at equilibrium. In part (b), each location on the graph corresponds to a
different parameter regime. Along the *x* axis, we vary the rate of increase in the marginal individual
cost of trait distortion $\left(\frac{d^2 c_{drive}}{dk^2}\right)$ relative to the rate of decrease in the marginal transmission benefit
$\left(-\frac{d^2 t}{dk^2}\right)$, which determines the target level of trait distortion (k_{target}) (Supplementary Information 1).
Along the *y* axis, we vary the cost of suppression (c_{sup}). We see that, when there is a low target level
of distortion (left hand side of the heat map), suppressors fail to invade, leading to a low level of trait
distortion at equilibrium. When there is a high target level of distortion and a relatively low cost of
suppression (bottom right hand side), suppressors invade, leading to no trait distortion at equilibrium.
When there is a high target level of distortion and a relatively high cost of suppression (top right hand
side), suppressors do not invade, leading to appreciable trait distortion at equilibrium.
241

suppressed by the same suppressor allele, at the same cost (c_{sup})[49]. In Appendix 5,

we write the recursions that detail the generational frequency changes in the

different possible gametes (y_0/+, y_0/sup, y_1/+, y_1/sup, y_2/+, y_2/sup).

We found that stronger mutant distorters ($\hat{k}>k$) will invade from rarity when the

marginal increase in offspring they are propagated into exceeds the marginal

increase in offspring they are lost from as a result of reduced fitness ($\Delta t(1-$

$c_{drive}(\hat{k}))>\Delta c_{drive}$, where Δ denotes marginal change ($\Delta t=t(\hat{k})-t(k)$; $\Delta c_{drive}=c_{drive}(\hat{k})-$

$c_{drive}(k)$)). Consequently, if distortion is initially low, and successive mutant distorters

are introduced, each deviating only slightly from the distorters from which they are

derived ("δ-weak selection"[42]), invading distorters will approach a 'target' strength,

denoted by k_{target}. The target strength is that at which the marginal benefit of

transmission is exactly counterbalanced by the marginal individual cost of reduced

offspring, which occurs when $\frac{dt}{dk}(1-c_{drive})=\frac{dc_{drive}}{dk}$. The target strength of distortion

(k_{target}) will therefore be greater if increased trait distortion (k) leads to a low rate of

decrease in marginal transmission benefit $\left(-\frac{d^2t}{dk^2}\right)$ relative to the rate of increase in

marginal individual cost $\left(\frac{d^2c_{drive}}{dk^2}\right)$ (Fig. 2b). If mutations are larger (strong selection),

invading distorters may overshoot the target strength of distortion ($\hat{k}>k_{target}$). Weaker

mutant distorters ($\hat{k}<k$) are recessive so cannot invade from rarity.

As evolution on the distorter increases the level of distortion, it makes it more likely

that the distorter reaches the critical level of distortion where suppression will be

favoured. When this is the case ($c_{sup}<c_{drive}(k_{target})$), the distorter spreads to high

frequency, which then causes the suppressor to increase in frequency, reversing the

direction of selection on the distorter, towards non-distortion (y_0), resulting in zero

trait distortion at equilibrium ($k^*=0$) (Fig. 2a; Appendix 6). Suppression only fails to

spread if the individual fitness cost associated with suppression is greater than the

individual fitness cost associated with the target trait distortion ($c_{sup}>c_{drive}(k_{target})$; Fig.

2a). Given that the individual fitness cost of pre-translational suppression at a single

locus is likely to be low, then any non-negligible distorter is likely to be suppressed.

Overall, our results suggest that selection on distorters will tend to drive the eventual

suppression of those distorters. In Appendix 7 we developed an agent-based

simulation, which allowed us to continuously vary the level of both distortion and

suppression, and obtained results in close agreement (Fig. 2a; Supplementary

Information 3, Fig. S2).

Specific Models

We then tested the robustness of our above conclusions by developing models for

three different biological scenarios: a sex ratio distorter on an X chromosome (X

driver); an imprinted gene that is only expressed when maternally inherited; and a

gene for the production of a public good by bacteria, which is encoded on a mobile

genetic element. We examined these cases because they are different types of

distortion, involving different selection pressures, in very different organisms. We

obtained qualitatively similar results in all three cases.

In all of our specific models, we assume that the suppressor: is dominant; is only

expressed in the presence of the distorter (facultative); completely suppresses the

distorter; and may incur a fitness (viability) cost to the individual when it is

expressed, independent of distorter strength, denoted by c_{sup} ($0 \leq c_{sup} \leq 1$)[43,44]. These

assumptions fit well to a molecularly characterised suppressor ("*nmy*") of a sex ratio

distorter ("*Dox*")[39,40]; and more generally to suppressors that act pre-

translationally[45,46]. We also relax a simplifying assumption of our illustrative model,

by allowing the transmission benefit and individual fitness cost of trait distortion to

vary with the population frequency of the distorter.

Sex Ratio Distortion

We examined sex ratio evolution in a diploid species, in a large outbreeding

(panmictic) population, with non-overlapping generations, and where males and

females are equally costly to produce. Fisher[1] and many others have shown that, in

this scenario, individuals would be selected to invest equally in male and female

offspring[9,24]. We assumed genetic sex determination, with males as XY and females

as XX, and that females mate with λ mates per generation. The distorter (y_1) that we

considered is an X driving chromosome, which acts in males, killing Y-bearing

sperm, and causing the male's mating partners to produce a higher proportion of

female (XX) offspring. The proportion is given by $(1+k)/2$, where k denotes the

proportion of Y-bearing sperm that are killed ($0 < k \leq 1$). We assumed that the sex ratio

distorter can be suppressed by a costly autosomal suppressor (*sup*). This biology

corresponds to sex ratio distortion in flies[25].

Our sex ratio model showed very similar results to our illustrative model. In

Supplementary Information 4 we show with population genetic analyses that when

distortion is weak (low k), suppressors are not favoured. In contrast, if distortion is

strong (high k), then suppression is favoured, and so there is no net influence on the

individual trait value. Consequently, the extent to which the sex ratio deviates from a

50:50 investment will be small or zero[47] (Fig. 3ai). In addition to being more likely to

be suppressed, the stronger distortion is, the quicker suppressors spread (Fig. S5).

Finally, when we allowed the X chromosome driver to evolve, it evolved to high

levels of sex ratio distortion (high k_{target}), increasing the likelihood of suppression.

We only obtained appreciable and detectable levels of sex ratio distortion (>60%

females) if the cost of suppression exceeded a 15-35% viability reduction (Fig. S8).

This is a much greater cost than what we would expect from natural gene

suppression pathways[48]. A suppressor will only fail to spread when the individual

cost of sex ratio distortion is less than the cost of suppressing the distorter (Equation

S3). Given that the cost of suppression is likely to be low, we would only expect

distorters that have relatively little impact at the individual level to evade

suppression. We tested the robustness of our population genetic analysis with an

agent-based simulation, and a game theory model, and found close agreement (Fig.

S6).

Our predictions are consistent with the available data on X drivers in *Drosophila*. As

predicted by our model: (1) Across natural populations of *D. simulans*, there is a

positive correlation between the extent of sex ratio distortion and the extent of

suppression[49]. (2) In both *D. mediopunctata* and *D. simulans* the presence of an X

linked driver led to the experimental evolution of suppression[50,51]. In addition,

consistent with our model: (3) In natural populations of *D. simulans* the prevalence of

an X driver has been shown to sometimes decrease under complete suppression[52].

(4) Crossing different species of *Drosophila* has been shown to lead to appreciable

sex ratio deviation, by unlinking distorters from their suppressors, and hence

revealing previously hidden distorters[53]. Work on other sex ratio distorters has also

shown that suppressors can spread extremely quickly from rarity, reaching fixation in

as little as ~5 generations[54].

362   Genomic Imprinting and Altruism

Genomic imprinting occurs at a minority of genes in mammals and flowering plants.

An imprinted allele has different epigenetic marks, and corresponding expression

levels, when maternally and paternally inherited[55]. We examined the evolution of an

altruistic helping behaviour in a population capable of genomic imprinting. A

behaviour is altruistic if it incurs a cost (c) to perform, by the actor, and provides a

benefit (b) to another individual, the recipient. Altruism is favoured if the genetic

relatedness (R) between the actor and recipient is sufficiently high, such that $Rb>c$[3].

An individual may be more closely related to their social partners via their maternal

or paternal genes[14,15,18]. For example, if a female mates two males, then on average

her offspring would be related by $R_m=1/2$ at maternal genes and $R_p=1/4$ at paternal

genes. If genes can 'gain information' about where they came from, by imprinting,

then they could be selected to adjust traits accordingly. Assume that relatedness to

social partners is R_p and R_m at paternal and maternal genes respectively. In this

case, altruistic helping would be favoured at: maternally imprinted genes when

$R_mb>c$; paternally imprinted genes when $R_pb>c$; and unimprinted genes when

$((R_p+R_m)/2)b>c$[18,56,57]. Consequently, if $R_mb>c>((R_p+R_m)/2)b$, then altruistic helping is

favoured at maternally imprinted genes, when it is disfavoured at unimprinted genes

(selfish trait distortion).

We modelled the evolution of altruism in a large population of diploid, sexually

reproducing individuals. The distorter (y_1) increases altruistic investment by some

amount (k), at a fitness cost to the individual ($0 \leq c_{GI}(k) \leq 1$) and benefit to the social

partner ($b_{GI}(k) > c_{GI}(k)$) that are both monotonically increasing functions of investment

$\left(\frac{d\{b_{GI}, c_{GI}\}}{dk} \geq 0 \right)$. The distorter ($y_1$) is only expressed when maternally inherited,

whereas its potential suppressor (*sup*) is unimprinted[58]. Every generation, individuals

associate in pairs with kin that they are maximally related to via their maternally

inherited genes ($R_m = 1$) but minimally related to via their paternally inherited genes

($R_p = 0$). Individuals then have the opportunity to be altruistic to their partner, before

mating at random in proportion to their fitness (fecundity), reproducing, then dying

(non-overlapping generations).

In Supplementary Information 5, we showed with a population genetic analysis that

our imprinting model produces very similar results to our illustrative model. When

distortion is weak (low k), such that the cost of trait distortion is less than the cost of

suppression, suppressors are not favoured (Equation S8). Given that the distorter

suppression cost is likely to be low, these distorters that evade suppression will have

relatively little impact at the individual level. When distortion is strong (high k), then

suppression is favoured, and so there is no influence on the individual trait value[58].

Consequently, the extent to which altruistic investment deviates from the individual

optimum of zero investment will be small or zero (Fig. 3bi). Finally, when we allowed

the imprinted gene to evolve, we found that cooperative distortion increased until it

reached the value (k_{target}) at which the marginal benefit of cooperation is cancelled

out by the marginal cost of cooperation ($\frac{\partial b_{GI}}{\partial k} = \frac{\partial c_{GI}}{\partial k}$; Equation S10), increasing the

likelihood of suppression (Fig. 3bii).

Although there have been no direct tests, our predictions are consistent with data on

imprinted genes. There is no evidence that traits influenced by imprinted genes

deviate significantly from individual level optima under normal development[14].

Significant deviation is only observed when imprinted genes are deleted, implying

that imprinted trait distorters are either suppressed, or counterbalanced by

oppositely imprinted genes pulling the trait in the opposite direction[32,58]. Furthermore,

although many different parties (coreplicons) have vested interests in genomic

imprinting, our analysis suggests why the unimprinted majority could win control[59].

This could help explain both why imprinting appears to be relatively rare within the

genome[18,55,60], and why imprints are removed and re-added every generation in

mice, handing control of genomic patterns of imprinting to unimprinted genes[18,59,61].

Horizontal Gene Transfer and Public Goods

Bacteria produce and excrete many extracellular factors that provide a benefit to the

local population of cells and so can be thought of as public goods[62]. We modelled

the evolution of investment in a public good in a large, clonally reproducing

population. We assume a public good that costs c to produce, and provides a benefit

b to the group. We assume a well-mixed population, meaning genetic relatedness at

vertically inherited genes is zero ($R_{vertical}$=0), and so public good production is

disfavoured at the individual level ($R_{vertical}b$=0<c)[3,63].

We consider a distorter (y_1) of public goods production that is on a mobile gene,

such as a plasmid. Mobile genes can spread within groups, increasing genetic

relatedness at the mobile locus ($R_{horizontal}$>0), potentially favouring public goods

production[64,65]. We assume that the distorter increases public goods investment by

some amount (k), at a fitness cost to the individual (0≤$c_{HGT}(k)$≤1) and benefit shared

within the group ($b_{HGT}(k)$>$c_{HGT}(k)$), that are both monotonically increasing functions

of investment $\left(\frac{d\{b_{HGT}, c_{HGT}\}}{dk} \geq 0 \right)$. We also assume a potential suppressor (*sup*) that is

immobile[46,66,67]. Each generation, individuals randomly aggregate into groups, and

one allele at the mobile locus (y_0, y_1, y_2) spreads horizontally within each group, each

with equal likelihood, increasing relatedness at the mobile locus. Public goods may

then be produced and shared within groups, before individuals reproduce and die

(non-overlapping generations).

In Supplementary Information 6, we show with a population genetic analysis that our

plasmid model produces similar results to our illustrative model. When distortion is

weak (low k), suppressors are not favoured, but the distorter has relatively little

impact at the individual level (Fig. 3ci). When distortion is strong (high k), then

suppression is favoured, and so there is no influence on the individual trait value[67].

Consequently, the extent to which public goods investment deviates from the

individual optimum of zero investment will be relatively small or zero (Fig. 3ci).

Finally, when we allowed the mobile distorter to evolve, we found that higher levels

of public goods investment (high k_{target}) are favoured, leading to selection for

suppression (Fig. 3cii). We lack empirical data that would allow us to test our model

of mobile public goods genes.

Discussion

We have found that the individual level consequences of selfish genetic elements

('*distorters*') will be either small or non-existent. If distorters lead to only small

distortions of traits, then they will not be suppressed, but they will only have small

effects on traits (Figs. 1a & 3i). Specifically, distorters will only remain unsuppressed

if trait distortion compromises individual fitness less than suppression of the trait

distorter does ($c_{drive}(k) < c_{sup}$). Given that the individual fitness cost of pre-translational

suppression at a single locus is likely to be low, we can say that trait distortion

conferred by unsuppressed distorters is likely to be relatively negligible.

However, if distorters lead to large distortions of traits then this selects for their

suppression, and so they will have no net effect on traits at the individual level (Figs.

1a & 3i). Stronger distorters will also be suppressed more quickly (Fig. 1b).

Furthermore, selection on distorters favours higher levels of distortion, which will

render them more likely to be ultimately suppressed. Consequently, the evolution of

distorters will often drive their own demise (Figs. 2, 3ii & 4). These results suggest

that even if there is substantial potential for genetic conflict, distorters will have

relatively little influence at the individual level, in support of Leigh's[27] parliament of

genes hypothesis.

Trait 1

Trait 2

Figure 4: Selfish genetic elements evolve to be suppressed by the parliament of genes. The cross represents the position in phenotype space, here defined with respect to two traits, *1* and *2*, that maximises the fitness of an individual. The circle surrounding the cross represents the phenotype space where suppression of selfish genetic elements, that have distorted traits 1 or 2, would not be selected for. The surrounding area represents the phenotype space in which the parliament of genes is selected to suppress selfish genetic elements. The three dots represent three possible individuals, which, owing to weakly selfish genetic elements, are not maximising individual fitness (the dots do not lie exactly on the cross), although they are approximately (e.g. within the bounds of experimental error in measurement). Because these deviations from individual fitness maximisation are only slight, costly suppression of the weakly selfish genetic elements does not evolve. However, the selfish genetic elements will evolve to become more distorting (solid arrows), bringing individuals into the non-tolerated area of phenotype space, where they will be suppressed and individual fitness maximisation (the black cross) is regained (dashed arrows).

We have assumed that there will be a much greater number of genes where

suppression is favoured, and so it is relatively easy for a suppressor to be reached

by mutation[31]. We have therefore examined, given the potential for suppression,

what direction would we expect natural selection to take on average. We are not

claiming that appreciable trait distortion will never evolve, and there are clearly cases

where it has[14]. A number of biological details will matter for different systems,

including: (i) whether a cost of suppression is incurred when the distorter is not

present; (ii) whether distorters can outstrip suppressors in coevolutionary arms

races, leading to distorters going unsuppressed for extended time periods[31,54,68]; (iii)

the rate at which trait-distorting selfish genetic elements arise by mutation, relative to

the rate at which their suppressors arise by mutation. Our results are supported by

cases where appreciable distortion is only revealed in hybrid crosses, implying that

they are generally suppressed[53,69].

We emphasise that when the assumption of individual fitness maximisation is made

in behavioural and evolutionary ecology, it is not being assumed that natural

selection produces perfect fitness maximisers[5]. Many factors could constrain

adaptation, such as genetic architecture, mutation and phylogenetic constraints[70,71].

Instead, the assumption of fitness maximisation is used as a basis to investigate the

selective forces that have favoured particular traits (adaptations). The aim is not to

test if organisms maximise fitness, or behave 'optimally', but rather to try to

understand the selective forces favouring particular traits or behaviours[2].

To conclude, debate over the validity of assuming individual level fitness

maximisation has revolved around whether selfish genetic elements are common or

rare[4,19,20,23,72]. We have shown that that even if selfish genetic elements are

common, they will tend to be either weak and negligible, or suppressed. This

suggests that even if there is the potential for appreciable genetic conflict, individual

level fitness maximisation will still often be a reasonable assumption. This allows us

to explain why certain traits, especially the sex ratio, have been able to provide such

clear support for both individual level fitness maximisation and genetic conflict.

Appendix

Appendix 1: Distorter population frequency

We ask when a rare distorter (y_1) can invade a population fixed for the non-distorter

(y_0). We take Equation (1), set $p'=p=p^*$, and solve to find two possible equilibria:

$p^*=0$ (non-distorter fixation) and $p^*=1$ (distorter fixation). The distorter (y_1) can invade

from rarity when the $p^*=0$ equilibrium is unstable, which occurs when the differential

of p' with respect to p, at $p^*=0$, is greater than one. The distorter invasion criterion is

therefore $c_{drive}(k)<t(k)(1-c_{drive}(k))$.

We now ask what frequency the distorter (y_1) will reach after invasion. The distorter

(y_1) can spread to fixation if the $p^*=1$ equilibrium is stable, which requires that the

differential of p' with respect to p, at $p^*=1$, is less than one. This requirement always

holds true, demonstrating that there is no negative frequency dependence on the

distorter, and that it will always spread to fixation after its initial invasion.

Appendix 2: Suppressor invasion condition

We ask when the suppressor (*sup*) can spread from rarity in a population in which

the distorter (y_1) and non-suppressor (+) are fixed at equilibrium. We derive the

Jacobian stability matrix for this equilibrium, which is a matrix of each genotype

frequency (x_1', x_2', x_3', x_4') differentiated by each genotype frequency in the prior

generation (x_1, x_2, x_3, x_4), at the equilibrium position given by $x_1^*=0$, $x_2^*=0$, $x_3^*=1$,

$x_4^*=0$:

$$J = \begin{pmatrix} 1-t & \frac{1-c_{sup}}{2(1-c_{drive})} & 0 & 0 \\ 0 & \frac{1-c_{sup}}{2(1-c_{drive})} & 0 & 0 \\ t-1 & \frac{-3(1-c_{sup})}{2(1-c_{drive})} & 0 & \frac{-(1-c_{sup})}{1-c_{drive}} \\ 0 & \frac{1-c_{sup}}{2(1-c_{drive})} & 0 & \frac{1-c_{sup}}{1-c_{drive}} \end{pmatrix}.$$

(A1)

The suppressor can invade when the equilibrium is unstable, which occurs when the

leading eigenvalue is greater than one. The leading eigenvalue is $(1-c_{sup})/(1-c_{drive})$,

meaning the suppressor invasion criterion is $c_{drive} > c_{sup}$.

Appendix 3: Equilibrium distorter and suppressor frequencies

We ask what frequency the distorter (y_1) and suppressor (*sup*) will reach after initial

suppressor (*sup*) invasion. We assume that the suppressor is introduced from rarity

when the distorter has reached the population frequency given by f ($x_1 \rightarrow f$, $x_3 \rightarrow 1-f$,

$\{x_2, x_4\} \rightarrow 0$). We numerically iterate Equations (2), over successive generations, until

equilibrium has been reached. At equilibrium, for all parameter combinations

(f, t, c_{sup}, c_{drive}), the suppressor reaches an internal equilibrium and the distorter is lost

from the population ($x_1^* + x_2^* = 1$, $x_3^* = 0$, $x_4^* = 0$). This equilibrium arises because

distorter-presence gives the suppressor (*sup*) a selective advantage, leading to high

suppressor frequency, which in turn reverses the selective advantage of the distorter

(y_1), leading to distorter loss and suppressor equilibration.

Appendix 4: Non-equilibrium trait distortion

We consider a distorter that is suppressed and therefore purged at equilibrium

($c_{drive} > c_{sup}$), and ask to what extent it can contribute to individual trait distortion in the

period after its initial invasion but before its eventual loss (non-equilibrium). We

introduce the distorter (y_1) and suppressor (*sup*) from rarity and numerically iterate

our recursions until the distorter has been purged from the population (or a cap of

20,000,000 generations has been reached). We vary parameters between $0 \leq t \leq 1$,

$c_{sup} < c_{drive} \leq 1$, $0 \leq c_{sup} \leq 1$.

We find that a higher cost of trait distortion (c_{drive}) relative to suppression (c_{sup}) leads

to shorter non-equilibrium maintenance of the distorter in the population. This is

because the cost of trait distortion relative to suppression mediates selection on the

suppressor (Appendix 2). We find that a higher transmission bias (t) leads to longer

non-equilibrium maintenance of the distorter in the population, but this effect is

diluted as the cost of trait distortion (c_{drive}) is increased relative to suppression (c_{sup})

(Supplementary Information 2, Fig. S1). Stronger distorters (with higher k, leading to

higher c_{drive} and t) are therefore generally suppressed and purged more rapidly than

weaker distorters (Fig. 1b). Exceptions are distorters that reduce individual fitness

relatively negligibly after the point (k) at which suppression is favoured, such that

$\frac{dt}{dk} / \frac{dc_{drive}}{dk}$ is very high for values of k satisfying $c_{sup} < c_{drive}(k)$.

Appendix 5: Invasion of a mutant distorter

We ask when a mutant distorter (y_2) will invade against a resident distorter (y_1) that

is unsuppressed and at fixation ($k \neq \hat{k}$). We write recursions detailing the generational

frequency changes in the six possible gametes, y_0/+, y_0/*sup*, y_1/+, y_1/*sup*, y_2/+,

y_2/*sup*, with current generation frequencies denoted respectively by x_1, x_2, x_3, x_4, x_5,

x_6, and next generation frequencies denoted with an appended dash ('):

$\bar{w}\,x_1' = x_1x_1 + x_1x_2 + (1-t(k))(1-c_{drive}(k))x_1x_3 + ((1-c_{sup})/2)x_1x_4 + (1-t(\hat{k}))(1-c_{drive}(\hat{k}))$ (A2)

$\quad x_1x_5 + ((1-c_{sup})/2)x_1x_6 + ((1-c_{sup})/2)x_2x_3 + ((1-c_{sup})/2)x_2x_5$

$\bar{w}\,x_2' = x_1x_2 + ((1-c_{sup})/2)x_1x_4 + ((1-c_{sup})/2)x_1x_6 + x_2x_2 + ((1-c_{sup})/2)x_2x_3 + (1-c_{sup})x_2x_4$

$\quad + ((1-c_{sup})/2)x_2x_5 + (1-c_{sup})x_2x_6$

$\bar{w}\,x_3' = (1+t(k))(1-c_{drive}(k))x_1x_3 + ((1-c_{sup})/2)x_1x_4 + ((1-c_{sup})/2)x_2x_3 + (1-c_{drive}(k))$

$\quad x_3x_3 + (1-c_{sup})x_3x_4 + (1+t(k)-t(\hat{k}))(1-c_{drive}(\max(k,\hat{k})))x_3x_5 + ((1-c_{sup})/2)x_3x_6$

$\quad + ((1-c_{sup})/2)x_4x_5$

$\bar{w}\,x_4' = ((1-c_{sup})/2)x_1x_4 + ((1-c_{sup})/2)x_2x_3 + (1-c_{sup})x_2x_4 + (1-c_{sup})x_3x_4 + ((1-c_{sup})/2)x_3x_6$

$\quad + (1-c_{sup})x_4x_4 + ((1-c_{sup})/2)x_4x_5 + (1-c_{sup})x_4x_6$

$\bar{w}\,x_5' = (1+t(\hat{k}))(1-c_{drive}(\hat{k}))x_1x_5 + ((1-c_{sup})/2)x_1x_6 + ((1-c_{sup})/2)x_2x_5 + (1-t(k)+t(\hat{k}))$

$\quad (1-c_{drive}(\max(k,\hat{k})))x_3x_5 + ((1-c_{sup})/2)x_3x_6 + ((1-c_{sup})/2)x_4x_5 + (1-c_{drive}(\hat{k})))$

$\quad x_5x_5 + (1-c_{sup})x_5x_6$

$\bar{w}\,x_6' = ((1-c_{sup})/2)x_1x_6 + ((1-c_{sup})/2)x_2x_5 + (1-c_{sup})x_2x_6 + ((1-c_{sup})/2)x_3x_6 + ((1-c_{sup})/2)$

$\quad x_4x_5 + (1-c_{sup})x_4x_6 + (1-c_{sup})x_5x_6 + (1-c_{sup})x_6x_6.$

$\bar{w}$ is the average fitness of individuals in the current generation, and equals the sum

of the right-hand side of the system of equations. The mutant distorter can invade

when the equilibrium given by $x_1^* = 0$, $x_2^* = 0$, $x_3^* = 1$, $x_4^* = 0$, $x_5^* = 0$, $x_6^* = 0$ is unstable,

which occurs when the leading eigenvalue of the Jacobian stability matrix for this

equilibrium is greater than one. Testing for stability in this way, we find that, if the

mutant distorter is weaker than the resident, it can never invade. If the mutant

distorter is stronger than the resident, it invades from rarity when $\Delta t(1-$

$c_{drive}(\hat{k})) > \Delta c_{drive}$, where $\Delta t = t(\hat{k}) - t(k)$, $\Delta c_{drive} = c_{drive}(\hat{k}) - c_{drive}(k)$.

The implication is that, if trait distortion is initially low, and mutant distorters are

successively introduced, each deviating only very slightly from the resident distorter

from which they are derived, such that $\hat{k}=k\pm\delta$, where δ is very small ("δ-weak

selection"[42]), then distorters will approach a 'target' strength at which $\frac{dt}{dk}(1-$

$c_{drive}) = \frac{dc_{drive}}{dk}$. In the absence of suppression, this target (k_{target}) is the equilibrium

level of distortion ($k^*=k_{target}$). However, if mutant distorters (y_2) are allowed to deviate

appreciably from residents (y_1) (strong selection), then distorters may invade even if

they overshoot the target ($\hat{k}>k_{target}$). In the absence of suppression, k_{target} is then not

the equilibrium level of distortion, but rather, the minimum equilibrium level of

distortion ($k^*>k_{target}$) (Supplementary Information 3, Fig. S2b).

We could alternatively have assumed that an individual's trait is distorted according

to the average strength of its alleles (additive gene interactions), rather than

according to the stronger (higher-k) allele (dominance). Such an assumption leads to

a single invasion criterion for a mutant distorter, regardless of whether the mutant

distorter is stronger or weaker than the resident distorter, given by:

$\Delta t(2 - c_{drive}(k) - c_{drive}(\hat{k})) > \Delta c_{drive}$. In the absence of suppression, this leads to an

equilibrium level of distortion (k^*), that holds even under strong selection, that

satisfies $2\frac{dt}{dk}(1 - c_{drive}) = \frac{dc_{drive}}{dk}$.

Appendix 6: Equilibrium distorter and suppressor frequencies (long term

evolution)

We ask what equilibrium state will arise after the invasion of a mutant distorter. We

assume that the mutant distorter (y_2) is introduced from rarity when the resident

distorter (y_1) has reached the population frequency given by q. We numerically

iterate Equations (A2), over successive generations, until equilibrium has been

reached. At equilibrium, for all parameter combinations (q, $t(k)$, $t(\hat{k})$, c_{sup}, $c_{drive}(k)$,

$c_{drive}(\hat{k})$), the resident distorter (y_1) is lost from the population ($x_3,x_4=0$), with either

the mutant distorter (y_2) and non-suppressor (+) at fixation ($x_5^*=1$), or the non-

distorter at fixation alongside the suppressor at an internal equilibrium ($x_1^*+x_2^*=1$).

The latter scenario arises if the mutant distorter triggers suppressor invasion

($c_{sup}<c_{drive}(\hat{k})$). This equilibrium arises because mutant distorter-presence gives the

suppressor (*sup*) a selective advantage, leading to high suppressor frequency, which

in turn reverses the selective advantage of distortion, leading to distorter (y_1,y_2) loss

and suppressor equilibration.

Appendix 7: Agent-based simulation

We construct an agent-based simulation to ask what level of trait distortion evolves

when continuous variation is permitted at distorter and suppressor loci. We model a

population of N=2000 individuals and track evolution at two autosomal loci: a

distorter locus (L1) and a suppressor locus (L2). Each individual has two alleles at

the distorter locus, with strengths denoted by k_a and k_b, and two alleles at the

suppressor locus, with strengths denoted by m_a and m_b (diploid). Strengths can take

any continuous value between zero and one. We assume that, for both loci, the

strongest (highest value) allele within an individual is dominant. The absolute fitness

of an individual with at least one active meiotic driver ($\max(k_a,k_b)>0$) is: 1-

$c_{drive}(\max(k_a,k_b))(1- \max(m_a,m_b))-c_{sup}\max(m_a,m_b)$, and the absolute fitness of an

individual lacking an active distorter ($\max(k_a,k_b)=0$) is 1. The function

$c_{drive}(\max(k_a,k_b))$ is given an explicit form in simulations (Supplementary Information

3, Fig. S2).

Each generation, there are N breeding pairs. To fill each position in each breeding

pair, individuals are drawn from the population, with replacement, with probabilities

given by their fitness (hermaphrodites). Breeding pairs then reproduce to produce

one offspring, before dying (non-overlapping generations). Alleles at the suppressor

locus (L2) are inherited in Mendelian fashion. Alleles at the distorter locus may drive,

meaning the parental allele of strength k_a is inherited, rather than the allele of

strength k_b, with the probability $(1+(t(k_a)-t(k_b))(1-\max(m_a,m_b)))/2$. The transmission

bias function, t, is given an explicit form in simulations (Supplementary Information 3,

Fig. S2). Each generation, distorter and suppressor alleles have a 0.01 chance of

mutating to a new value, which is drawn from a normal distribution centred around

the pre-mutation value, with variance 0.2, and truncated between 0 and 1. We track

the population average distorter strength, denoted by $E[k]$, and suppressor strength,

denoted by $E[m]$, over 20,000 generations. We see that, allowing for continuous

variation at the distorter and suppressor loci, if the cost of suppression (c_{sup}) is not

excessively high, trait distortion at equilibrium is either low or nothing (Fig. 2a; Fig.

S2b).

References

1.   Fisher, R. A. *The Genetical Theory of Natural Selection*. (Oxford University

Press, 1930).

2.   Gardner, A. Adaptation as organism design. *Biology Letters* **5,** 861–864

(2009).

3.   Hamilton, W. D. The genetical evolution of social behaviour. I & II. *Journal of*

Theoretical Biology **7** (1964).

4.   Grafen, A. The formal darwinism project in outline. *Biol Philos* **29,** 155–174

(2014).

5.   Parker, G. A. & Maynard Smith, J. Optimality theory in evolutionary biology.

Nature Publishing Group **348,** 27–33 (1990).

6.   Frank, S. A. *Foundations of Social Evolution*. (Princeton University Press,

1998). doi:10.1038/sj.hdy.6885351

7.   Stephens, D. W. & Krebs, J. R. *Foraging Theory*. (Princeton University Press,

1986).

8.   Stearns, S. C. *The Evolution of Life Histories*. (Oxford University Press on

Demand, 1992).

9.   West, S. *Sex Allocation*. (Princeton University Press, 2009).

doi:10.1515/9781400832019

10.   Davies, N. B., Krebs, J. R. & West, S. A. *An Introduction to Behavioural*

Ecology. (John Wiley & Sons, 2012).

11.   Westneat, D. & Fox, C. W. *Evolutionary Behavioral Ecology*. (Oxford

University Press on Demand, 2010).

12.   Bourke, A. F. G. *Principles of Social Evolution*. (OUP Oxford, 2011).

13. Werren, J. H., NUR, U. & Wu, C. I. Selfish Genetic Elements. *Trends in*

Ecology & Evolution **3,** 297–302 (1988).

14. Burt, A. & Trivers, R. *Genes in Conflict.* (Harvard University Press, 2006).

doi:10.4159/9780674029118

15. Gardner, A. & Úbeda, F. The meaning of intragenomic conflict. *Nature Ecology*

& Evolution **1,** 1–9 (2017).

16. Ågren, J. A. & Clark, A. G. Selfish genetic elements. *PLoS Genet* **14,**

e1007700–20 (2018).

17. Hamilton, W. D. Extraordinary sex ratios. A sex-ratio theory for sex linkage and

inbreeding has new implications in cytogenetics and entomology. *Science* **156,**

477–488 (1967).

18. Haig, D. *Genomic Imprinting and Kinship.* (Rutgers University Press, 2002).

19. Haig, D. Genetic dissent and individual compromise. *Biol Philos* **29,** 233–239

(2014).

20. Bourke, A. F. G. The gene's-eye view, major transitions and the formal

darwinism project. *Biol Philos* **29,** 241–248 (2014).

21. Cosmides, L. M. & Tooby, J. Cytoplasmic inheritance and intragenomic

conflict. *Journal of Theoretical Biology* **89,** 83–129 (1981).

22. West, S. A. & Gardner, A. Adaptation and Inclusive Fitness. *Current Biology*

23, R577–R584 (2013).

23. Ågren, J. A. Selfish genetic elements and the gene's-eye view of evolution.

Curr Zool **62,** 659–665 (2016).

24. Charnov, E. L. *The Theory of Sex Allocation.* (Princeton University Press,

1982).

25. Jaenike, J. Sex Chromosome Meiotic Drive. *Annual Review of Ecology and*

Systematics **32,** 25–49 (2001).

26. Leigh, E. G., Herre, E. A. & Fischer, E. A. Sex allocation in animals.

Experientia **41,** 1265–1276 (1985).

27. Leigh, E. G. *Adaptation and diversity*. (1971).

28. Leigh, E. G. When does the good of the group override the advantage of the

individual? *Proc Natl Acad Sci USA* **80,** 2985–2989 (1983).

29. Leigh, E. G. How does selection reconcile individual advantage with the good

of the group? *Proc Natl Acad Sci USA* **74,** 4542–4546 (1977).

30. Strassmann, J. E. & Queller, D. C. The Social Organism: Congresses, Parties,

and Committees. *Evolution* **64,** 605–616 (2010).

31. Queller, D. C. & Strassmann, J. E. Evolutionary Conflict. *Annu. Rev. Ecol.*

Evol. Syst. **49,** annurev–ecolsys–110617–062527–21 (2018).

32. Gardner, A. & Ross, L. Mating ecology explains patterns of genome

elimination. *Ecol Lett* **17,** 1602–1612 (2014).

33. Crow, J. F. Why is mendelian segregation so exact? *Bioessays* **13,** 305–312

(1991).

34. Ridley, M. *Mendel's Demon*. 176–177 (Weidenfeld & Nicolson, 2000).

35. Prout, T., Bundgaard, J. & Bryant, S. Population genetics of modifiers of

meiotic drive I. The solution of a special case and some general implications.

Theoretical Population Biology **4,** 446–465 (1973).

36. Caubet, Hatcher, MocquardRigaud. Genetic conflict and changes in

heterogametic mechanisms of sex determination. *Journal of Evolutionary*

Biology **13,** 766–777 (2000).

37. Randerson, J. P. The evolutionary dynamics of male-killers and their hosts.

Heredity **84,** 1–9 (2000).

38. Haig, D. Intragenomic politics. *Cytogenet Genome Res* **113,** 68–74 (2006).

39. Tao, Y., Masly, J. P., Araripe, L., Ke, Y. & Hartl, D. L. A sex-ratio meiotic drive

system in Drosophila simulans. I: an autosomal suppressor. *PLoS Biol* **5,** e292

(2007).

40. Tao, Y. *et al.* A sex-ratio meiotic drive system in Drosophila simulans. II: an X-

linked distorter. *PLoS Biol* **5,** e293 (2007).

41. Qiu, S. A computational study of off-target effects of RNA interference. *Nucleic*

Acids Research **33,** 1834–1847 (2005).

42. Wild, G. & Traulsen, A. The different limits of weak selection and the

evolutionary dynamics of finite populations. *Journal of Theoretical Biology* **247,**

382–390 (2007).

43. Vaz, S. C. & Carvalho, A. B. Evolution of autosomal suppression of the sex-

ratio trait in Drosophila. *Genetics* **166,** 265–277 (2004).

44. Hall, D. W. Meiotic drive and sex chromosome cycling. *Evolution* **58,** 925–931

(2004).

45. Aravin, A. A., Hannon, G. J. & Brennecke, J. The Piwi-piRNA pathway

provides an adaptive defense in the transposon arms race. *Science* **318,** 761–

764 (2007).

46. Doron, S. *et al.* Systematic discovery of antiphage defense systems in the

microbial pangenome. *Science* **359,** eaar4120–13 (2018).

47. Rood, E. S. & Freedberg, S. Intragenomic conflict produces sex ratio dynamics

that favor maternal sex ratio distorters. *Ecol Evol* **6,** 8085–8093 (2016).

48. Unckless, R. L., Larracuente, A. M. & Clark, A. G. Sex-ratio meiotic drive and

Y-linked resistance in Drosophila affinis. *Genetics* **199,** 831–840 (2015).

49. Atlan, A., Mercot, H., Landre, C. & Montchamp-Moreau, C. The Sex-Ratio Trait

in Drosophila simulans: Geographical Distribution of Distortion and

Resistance. *Evolution* **51,** 1886 (1997).

50. Carvalho, A. B., Sampaio, M. C., Varandas, F. R. & Klaczko, L. B. An

experimental demonstration of Fisher's principle: Evolution of sexual

proportion by natural selection. *Genetics* **148,** 719–731 (1998).

51. Capillon, C. & Atlan, A. Evolution of Driving X Chromosomes and Resistance

Factors in Experimental Populations of Drosophila simulans. *Evolution* **53,** 506

(1999).

52. Bastide, H. *et al.* Rapid Rise and Fall of Selfish Sex-Ratio X Chromosomes in

Drosophila simulans: Spatiotemporal Analysis of Phenotypic and Molecular

Data. *Molecular Biology and Evolution* **28,** 2461–2470 (2011).

53. Blows, M. W., Berrigan, D. & Gilchrist, G. W. Rapid evolution towards equal

sex ratios in a system with heterogamety. *Evol Ecol Res* 277–283 (1999).

54. Hornett, E. A. *et al.* Evolution of Male-Killer Suppression in a Natural

Population. *PLoS Biol* **4,** e283–6 (2006).

55. Peters, J. The role of genomic imprinting in biology and disease: an expanding

view. *Nature Publishing Group* **15,** 517–530 (2014).

56. Úbeda, F. & Gardner, A. A model for genomic imprinting in the social brain:

juveniles. *Evolution* **64,** 2587–2600 (2010).

57. Queller, D. C. Theory of genomic imprinting conflict in social insects. *BMC*

Evol Biol **3,** 15 (2003).

58.   Wilkins, J. F. & Haig, D. Genomic imprinting of two antagonistic loci.

803   *Proceedings of the Royal Society B: Biological Sciences* **268,** 1861–1867

804   (2001).

805   59.   Burt, A. & Trivers, R. Genetic conflicts in genomic imprinting. *Proceedings of*

806   *the Royal Society B: Biological Sciences* **265,** 2393–2397 (1998).

807   60.   Gregg, C. *et al.* High-Resolution Analysis of Parent-of-Origin Allelic Expression

808   in the Mouse Brain. *Science* **329,** 643–648 (2010).

809   61.   Kafri, T., Gao, X. H. & Razin, A. Mechanistic Aspects of Genome-Wide

810   Demethylation in the Preimplantation Mouse Embryo. *Proc Natl Acad Sci USA*

811   **90,** 10558–10562 (1993).

812   62.   West, S. A., Diggle, S. P., Buckling, A., Gardner, A. & Griffin, A. S. The Social

813   Lives of Microbes. *Annu. Rev. Ecol. Evol. Syst.* **38,** 53–77 (2007).

814   63.   Frank, S. A. A general model of the public goods dilemma. *Journal of*

815   *Evolutionary Biology* **23,** 1245–1250 (2010).

816   64.   Nogueira, T. *et al.* Horizontal Gene Transfer of the Secretome Drives the

817   Evolution of Bacterial Cooperation and Virulence. *Current Biology* **19,** 1683–

818   1691 (2009).

819   65.   Smith, J. The social evolution of bacterial pathogenesis. *Proceedings of the*

820   *Royal Society B: Biological Sciences* **268,** 61–69 (2001).

821   66.   Johnson, L. J. The Genome Strikes Back: The Evolutionary Importance of

822   Defence Against Mobile Elements. *Evolutionary Biology* **34,** 121–129 (2007).

823   67.   Mc Ginty, S. É. & Rankin, D. J. The evolution of conflict resolution between

824   plasmids and their bacterial hosts. *Evolution* **66,** 1662–1670 (2012).

68. Dyson, E. A. & Hurst, G. Persistence of an extreme sex-ratio bias in a natural

population. *Proc Natl Acad Sci USA* **101,** 6520–6523 (2004).

69. Corbett-Detig, R., Medina, P., Frérot, H., Blassiau, C. & Castric, V. Bulk pollen

sequencing reveals rapid evolution of segregation distortion in the male

germline of Arabidopsis hybrids. *Evolution Letters* **129,** 1393–11 (2019).

70. Maynard Smith, J. *et al.* Developmental Constraints and Evolution: A

Perspective from the Mountain Lake Conference on Development and

Evolution. *The Quarterly Review of Biology* **60,** 265–287 (1985).

71. Lehmann, L., Alger, I. & Weibull, J. Does evolution lead to maximizing

behavior? *Evolution* **69,** 1858–1873 (2015).

72. Gardner, A. & Grafen, A. Capturing the superorganism: a formal theory of

group adaptation. *Journal of Evolutionary Biology* **22,** 659–671 (2009).

Acknowledgements

839 We thank Geoff Wild, Alan Grafen, Egbert Giles Leigh, Jr., Guy Cooper, Sam Levin,

840 Asher Leeks and Matishalin Patel for helpful comments and discussion.

Author Contributions

843 TWS and SAW designed the study and wrote the paper. TWS carried out

844 mathematical analysis.

Competing Interests

847 The authors declare no competing interests.

Data Accessibility

850 The data that support the findings of this study are available from the corresponding

851 author upon reasonable request.