Krüppel-like factor gene function in the ctenophore _Mnemiopsis_ suggests an ancient role in

promoting cell proliferation in metazoan stem cell niches

Jason S Presnell[1,2], William E Browne[1,*]

[1]Department of Biology, University of Miami, Cox Science Center, 1301 Memorial Drive,

Miami, FL 33146, USA

[2]Present Address: Department of Integrative Biology, Oregon State University, 3029 Cordley

Hall, Corvallis, OR 97331, USA

*Correspondence: w.browne@miami.edu

Abstract

All animals possess pluripotent stem cells that during an organism's lifespan give rise to

differentiated cell types and are capable of self-renewal. To date studies assessing the function of

explicit genes driving the regulation of stem cell renewal and pluripotency have been

characterized in bilaterians and a small number of cnidarians. Transcription factor genes critical

for regulation of stem cell self-renewal and pluripotency include *Oct4*, *Sox2*, *Klf4, and c-myc*

(OSKM) [1–3]. In particular, the role of *Klfs* in stem cell renewal and pluripotency has been

thoroughly investigated [1,2,4–9]. In mouse embryonic stem cells the three *Krüppel-like factor* (*Klf*)

genes *Klf2*, *Klf4*, and *Klf5* participate in an auto-regulatory activation loop and maintain a

transcriptional circuit with *Oct4*, *Sox2*, *c-myc*, and *Nanog* regulating stem cell self-renewal [4].

While OSKM gene homologs have been found in the genomes of many metazoans [3,10,11], their

function is unknown in non-bilaterians. Ctenophora, an ancient non-bilaterian lineage [12–14],

possess putative stem cells [11,15,16] and have prodigious regenerative capacity [17,18], making them

informative models for investigating the evolution and genetic underpinnings of stem cell

maintenance in Metazoa. Ctenophores possess homologs of the OSKM genes [11,19–21], and while

the expression patterns of ctenophore *Sox* paralogs are associated with high rates of cell

proliferation in putative stem cell niches [11,19], little is known about *Pou*, *Klf*, or *c-myc* expression

or function. Here we examine the expression of *Klf* homologs in the ctenophore *Mnemiopsis*

leidyi and disrupt zygotic *Klf* gene function during embryogenesis using both morpholino

oligonucleotides and CRISPR-Cas9. We find that zygotic *MleKlf5a* and *MleKlf5b* play a role in

patterning the gastrovascular cavity, pharynx, apical organ, and tentacle bulbs in *M. leidyi*.

Simultaneous zygotic knockdown of both *MleKlf5a* and *MleKlf5b* results in the diminution or

loss of cell proliferation in endodermal stem cell niches and aberrant endodermal patterning. Our

results highlight an ancestral functional role for *Klf* transcription factors in the regulation of rapid

cell proliferation in metazoan stem cell niches.

Main

Ctenophores are marine, invertebrate predators characterized by rows of giant cilia fused

into 'ctenes' used for locomotion, an aboral and neuronal-rich gravity-sensing (apical) organ, and

muscular tentacles lined with sticky colloblast cells used for prey capture [22–27]. *M. leidyi* embryos

undergo a ctenophore-specific early cleavage program [28,29], gastrulation takes place ~3-5 hours

post fertilization (hpf) followed by tissue organization and organogenesis over the course of the

next several hours before hatching as cydippids ~18-22 hpf [30].

We used EdU incorporation assays to identify dividing cell populations during

embryogenesis. EdU incorporation identified several regions during mid to late embryogenesis

with rapid rates of cell proliferation (Fig. 1a-h). Four pairs of ctene rows, one pair in each

quadrant, are typically among the first ectodermal structures to form [29]. Each ctene plate is made

up of polster cells bearing fused giant cilia [31]. Beginning at 13 hpf, we observed EdU

incorporation in a set of ectodermal cells surrounding each ctene row pair (Fig. 1b, f). In the

pharynx, cell proliferation was initially widespread aborally during pharyngeal development and

elongation (Fig. 1d). EdU incorporation in the pharynx subsequently becomes spatially restricted

to two domains, one orally near the stomodeal opening, and the other aborally that includes the

pharyngeal region forming the junction with the gastrovascular cavity (Fig. 1h). In the apical

organ, cell proliferation was localized to the developing polar fields and a small subset of cells

located deep within the epithelial floor along tentacular axis (Fig. 1a, e). Due to their proximity

to where new lithocytes are formed, these deeper proliferative cells likely represent lithocyte

progenitor cells [32].

Flanking the apical organ along the tentacular axis, ectodermal invaginations and internal

medial endodermal cell bands form the tentacle bulb organ [33–35] contributing to both the

ectodermal tentacular lateral ridges and the endodermal tentacular median ridges respectively,

each harboring progenitor cells that give rise to colloblasts and tentacle muscle [33–35]. During

tentacle bulb development EdU incorporation was initially observed in the median ridges (Fig.

1c-d). Later EdU incorporation was detected along the length of the median ridges and in the

lateral ridges (Fig. 1g-h). An Edu pulse followed by a 24 hour chase revealed that these

proliferative cells gave rise preferentially to tentacular muscle cells (Extended Data Fig. 1).

Previous cell proliferation studies in another ctenophore species, *Pleurobrachia pileus*, identified

gene expression indicative of 'stemness' such as *Piwi*, *Vasa*, *Nanos*, and *Sox* coincident with

EdU-positive cells in proliferative stem cell niches [15,19] similar to those observed during *M. leidyi*

embryogenesis.

The genome of *M. leidyi* contains three *Klf* genes: *MleKlf5a*, *MleKlf5b*, and *MleKlfX* [20].

MleKlf5a and *MleKlf5b* are the result of a lineage-specific duplication within the Ctenophora.

MleKlfX is highly derived with no clear orthology to any known metazoan *Klf* clade. *MleKlf5a*

and *MleKlf5b* transcripts are maternally loaded [21] and transcripts are detected in all embryonic

cells through gastrulation (Fig. 1i-l). Maternal loading has been observed in a number of *Klf*

genes among bilaterians [36–38]. Following gastrulation, *MleKlf5a* and *MleKlf5b* zygotic expression

is spatially localized to regions undergoing high rates of cell proliferation coincident with EdU

incorporation (Fig1 a-h, m-x). *MleKlf5a* and *MleKlf5b* expression overlap extensively in

endodermal domains during embryonic development (Fig. 1m-x). Interestingly a single

ectodermal expression domain unique to *MleKlf5b* is detected in proliferative cells flanking

developing ctene row polster cells (Fig. 1b, f, Extended Data Fig. 2a, b).

Within the pharynx, *MleKlf5a* and *MleKlf5b* expression is confined to the interior-most

cell layer. Expression in the aboral-most portion of the pharynx is associated with cells that form

the junction with the central gastrovascular cavity (infundibulum) and also throughout the

epithelial cell layer lining the developing infundibulum (Fig. 1n, p, r, t, v, x). During the initial

formation of the aboral apical organ, *MleKlf5a* and *MleKlf5b* expression is detected in the

epithelial floor. Subsequently, *MleKlf5a* and *MleKlf5b* expression in the apical organ becomes

restricted to cells that positionally correlate with putative lithocyte progenitors (Fig. 1r, t, v, x).

Within the developing tentacle bulbs, expression patterns of *MleKlf5a* and *MleKlf5b* mirror EdU

incorporation in regions of rapid cell proliferation (Fig. 1c-e, g-h, m-x). *MleKlf5a* and *MleKlf5b*

are highly expressed in the median ridge of developing tentacle bulbs (Fig. 1m-x), suggesting

that *MleKlf5a* and *MleKlf5b* may play a role in the maintenance of the median ridge tentacular

stem cell niche. In summary, *MleKlf5a* and *MleKlf5b* paralog expression extensively overlaps in

embryonic domains with high rates of cell proliferation. These regions express additional genes

associated with stemness such as *Piwi*, *Vasa*, *Nanos*, and *Sox* [11,15,16,19]. *MleKlf5a* and *MleKlf5b*

expression identifies putative stem cell niches within the aboral region of the pharynx, coincides

with putative lithocyte progenitor cells located in epithelial floor of the apical organ and

progenitor cells located in the median ridge of the tentacle bulb.

In contrast to *MleKlf5a* and *MleKlf5b* expression patterns, *MleKlfX* expression is

restricted to late stage embryos, first appearing ~16 hpf and does not appear to be associated with

EdU-positive cells (Extended Data Fig. 2c, d). *MleKlfX* expression is restricted to a small

number of cells within the apical organ. A deep set of *MleKlfX* expressing cells are located

centrally in the epithelial floor of the developing apical organ. A second shallower set is located

in each quadrant of the developing apical organ, just medial of the ciliated grooves. The two cell

clusters located along the tentacular plane are perpendicular to two additional cell clusters,

forming a cross-shaped pattern. These *MleKlfX* expressing cells positionally correspond with the

apical organ lamellate bodies, putative photoreceptor cells [39,40], suggesting that *MleKlfX* may

play a role in differentiation of light sensitive neuronal cell types in the apical organ. In

summary, *MleKlfX* a highly divergent gene in *M. leidyi*, appears to be expressed in a set of

differentiated neuronal cells, possibly involved with light sensing.

To characterize zygotic *Klf* gene function in *M. leidyi* we used both splice-blocking

morpholinos (sbMOs) and CRISPR-Cas9 genome editing to abrogate gene function during

embryonic development (Fig. 2). Knockdown of *MleKlfX* failed to produce any obvious

morphological phenotypes, so we focus here on the *MleKlf5a* and *MleKlf5b* results. It has been

shown that co-expressed KLFs bind to shared downstream regulatory targets resulting in

complex functional outcomes. For example, KLF2, KLF4, and KLF5 have redundant roles in the

downstream regulation of *Nanog* [4], whereas competition between KLF1 and KLF3 for binding

sites can result in differential functional outcomes [41]. For this reason we sought to maximize the

efficiency of generating an observable phenotype by performing simultaneous *MleKlf5a* and

MleKlf5b knockdown with both sbMO and CRISPR-Cas9 genome editing experiments. We

injected single-cell embryos with either *MleKlf5a+MleKlf5b* sbMOs or *MleKlf5a-*

sgRNA+MleKlf5b-sgRNA. Microinjected embryos were incubated to ~20 hpf and stained with

vital dyes (Fig. 3). Morphant and crispant specimens were imaged and compared to equivalent

late stage wildtype embryos from the same spawning events. A selection of morphants and

crispants were then subjected to single embryo RNA or DNA extractions respectively for gene

knockdown validation. Two sbMOs were designed for each gene (Fig. 2a, e). These produced

splice-blocking errors in morphants through exon skipping and/or intron retention (Fig. 2b, f).

An initial set of 4 sgRNAs were designed for *MleKlf5a* and *MleKlf5b* (Table 1) based on the *M.*

leidyi reference genome [24, 49]. A single sgRNA for each gene, *MleKlf5a-sgRNA4* and *MleKlf5b-*

sgRNA3, proved efficient at mediating Cas9 double-stranded break activity at the target loci (Fig.

2c-d, g-h). Sanger sequencing followed by ICE analysis revealed clear degradation of signal in

the crispants at the target loci compared to control embryos (Fig. 2c,g), indicating *MleKlf5a-*

sgRNA4 and *MleKlf5b-sgRNA3* targeted Cas9 exonuclease activity produced indels and

frameshift mutations (Extended Data Fig. 3b). Control injections with a standard control MO,

Cas9 protein alone, and sgRNAs alone had no detectable effect on embryonic development

(Extended Data Fig. 3a). No evidence of off-target Cas9 activity was observed in tested loci

(Table 2) thus phenotypes generated via both double gene abrogation approaches in our study

were due to the disruption of *MleKlf5a* and *MleKlf5b* activity.

Both *MleKlf5a*+*MleKlf5b* morphants and crispants generated an allelic series of

phenotypes (Fig. 3a-p). While morphants and crispants phenocopied one another, a higher

proportion of severe phenotypes were observed among cripsants as compared to morphants as

expected (Fig. 3s) reflecting the effects of Cas9 mediated genome editing versus titration of

functional mRNAs by sbMOs. In contrast to the observation of primarily severe phenotypes in

the crispant *MleKlf5a-sgRNA4*+*MleKlf5b-sgRNA3* double gene knockdowns (Fig. 3o-p; Fig. 4e-

f, l-m), single gene knockdown with either *MleKlf5a-sgRNA4* or *MleKlf5b-sgRNA3* generated

predominantly mild phenotypes (Extended Data Fig. 4). This suggests that *MleKlf5a* and

MleKlf5b likely have functionally redundant regulatory roles in maintaining proliferative

endodermal stem cell niches during embryonic development. Further investigation will be

needed to determine whether *MleKlf5a* and *MleKlf5b* are acting on the same direct downstream

targets, or acting indirectly on complementary non-overlapping sets of genes regulating cell

proliferation. For example, in mouse embryonic stem cells, pluripotency and proliferation are

maintained through differential inhibition of endoderm and mesoderm by *Klf4* and *Klf5*

respectively [42]. With regard to the unique ectodermal expression domain of *MleKlf5b*, no

ectodermal or ctene row phenotype was observed in *MleKlf5b-sgRNA3* or *MleKlf5a-sgRNA4*

crispants nor double gene abrogation experiments. Ctenes are initially established by maternal

factors [29], new polster cells and ctene row expansion begin post-hatching in the cydippid stage [43].

Thus the observed EdU- and *MleKlf5b*-positive ectodermal cells (Fig. 1b, f: Extended Data Fig.

2a-b) likely contribute to ctene row expansion post-hatching and/or perhaps also play a role in

the growth and maintenance of surrounding ectodermal epithelia post-hatching.

Embryos with mild phenotypes underwent pharyngeal elongation concomitant with

mesogleal deposition and an increase in size; however, they displayed disorganized endodermal

patterning associated with the aboral end of the pharynx and infundibular gastrovascular cavity

(Fig. 3e-f, m-n). In some cases we observed instances of pharyngeal bifurcation at the junction

with the gastrovascular cavity (Fig. 3 f,n; Extended Data Fig. 4b; Extended Data Fig. 5). In

contrast, embryos classified as having severe phenotypes (Fig. 3g-h, o-p) failed to increase in

size, likely reflecting a lack of ECM extrusion into the mesogleal space, resulting in internal

embryonic spaces being disproportionately occupied by gastrovascular endoderm and a failure of

normal pharyngeal elongation. Assessment of cell proliferation in *MleKlf5a-sgRNA4+MleKlf5b-*

sgRNA3 crispants revealed a dramatic reduction of EdU incorporation at the aboral end of the

pharynx (Fig. 3q, r). These results suggest that zygotic *MleKlf5a* and *MleKlf5b* play a

coordinated role in the growth and patterning of the gastrovascular cavity and pharynx,

particularly at the pharyngeal gastrovascular junction, presumably through maintenance of cell

proliferation in putative endodermal stem cell niches in these tissues.

Across both morphants and crispants, additional patterning defects were observed in the

apical organ epithelial floor and in the tentacle bulbs (Fig. 3; Fig. 4). *MleKlf5a* and *MleKlf5b*

knockdown resulted in a significant reduction of apical organ lithocytes as compared to control

embryos (Fig. 4a-g). By 20 hpf, control embryo statocysts contain ~7 lithocytes (Fig. 4a-b, g).

MleKlf5a+MleKlf5b morphants had an average ~4 lithocytes (Fig. 4g), with three lacking

lithocytes entirely (Fig. 4c, d). *MleKlf5a-sgRNA4+MleKlf5b-sgRNA3* crispants had an average of

~2 lithocytes (Fig. 4g), with five completely lacking lithocytes (Fig. 4e, f). In embryos lacking

lithocytes, both balancer cilia and dome cilia were present (Fig. 4d, f), reflecting that lithocytes

are derived from endoderm while the balancer and dome cilia are derived from ectoderm [34]. This

suggests that *MleKlf5a* and *MleKlf5b* play a role in maintenance of endodermally derived

lithocyte progenitor cells in the apical organ.

Simultaneous abrogation of *MleKlf5a* and *MleKlf5b* also resulted in a dramatic reduction

in tentacle bulb size, particularly notable in the tentacular median ridge (Fig. 4h-n). We

measured the length of the tentacular median ridge and found significant differences between

control and injected embryos (Fig. 4n; Extended Data Fig. 6). In control embryos the tentacular

median ridge was, on average, 23 µm in length, whereas in *MleKlf5a+MleKlf5b* morphants and

MleKlf5a-sgRNA4+MleKlf5b-sgRNA3 crispants, the average tentacular median ridge length was

~18 µm and ~9 µm respectively (Fig. 4n). Furthermore, we observed that 15% of the

MleKlf5a+MleKlf5b morphants and 29% of the *MleKlf5a-sgRNA4+MleKlf5b-sgRNA3* crispants

lacked tentacular median ridges altogether (Fig. 4j-m). Assessment of cell proliferation in

MleKlf5a-sgRNA4+MleKlf5b-sgRNA3 crispants revealed a dramatic reduction in EdU

incorporation in the tentacular median ridges, but not in the tentacular lateral ridges (Fig. 3q, r).

This indicates that proliferation in the tentacular lateral ridge stem cell niches were largely

unaffected in both morphants and crispants. Lateral ridge stem cells give rise to tentacle epithelia

and colloblasts [15,35], which were present in embryos with reduced median ridges (Fig. 4j-m). Our

results support a requirement for *MleKlf5a* and *MleKlf5b* activity for normal tentacle bulb

development and specifically for production of tentacle root muscle through maintenance of cell

proliferation in the endodermally-derived tentacular median ridge stem cell niche.

In this study, we used both sbMOs and Cas9-mediated genome editing to coordinately

abrogate zygotic *MleKlf5a* and *MleKlf5b* gene function during *M. leidyi* embryonic development.

We show that Cas9-mediated genome editing is an effective method for performing

combinatorial gene function assays in F0 ctenophore embryos. Along with recent RNA-seq data

highlighting candidate genes associated with zygotic gene activation and specific cell types in

ctenophores [21,35,44], sbMO and CRISPR-Cas9 based gene function assays in *M. leidyi* will

provide much needed insight into the genetic underpinnings of ctenophore biology and further

our understanding of early metazoan evolution. Our results highlight an essential role for zygotic

MleKlf5a and *MleKlf5b* activity in promoting cell proliferation in tissue and organ specific stem

cell niches that remain active throughout the lifespan of *M. leidyi* that produce a number of

terminally differentiated cell types unique to the Ctenophora. Our results further suggest that the

functional activity of *Klfs* in promoting cell proliferation within stem cell niches represents a

conserved ancestral function of this gene family across the Metazoa.

Fig. 1: *MleKlf5a* and *MleKlf5b* are expressed in regions of rapid cell proliferation during

embryogenesis in *Mnemiopsis leidyi.* Schematic depiction of tentacular axis (TA) and

pharyngeal axis (PA) orientation are located in panel upper right. Aboral view in **a, c, e, g, j, l,**

m, o, q, s, u, w. Lateral view, oral up, in **b, d, f, h, k, n, p, r, t, v, x**. Scale bars are 50 μm. **(a-h)**

25 min EdU pulse in 13 hours post fertilization (hpf) (**a-d**) and 16 hpf (**e-h**) embryos. EdU

incorporation is localized to two cell clusters in the apical organ (**a, e**), epithelial cells flanking

the developing ctene rows (bounded by dashed lines) (**b, f**), epithelial cells lining the

gastrovascular cavity (gvc) (**d, h**) in the developing tentacular median ridges (arrowheads), and

the oral (oph) and aboral (aph) regions of the pharynx (**d**). (**g, h**) Later in development, EdU

incorporation is found in the tentacular median ridge (arrowheads) and lateral ridges (lr) of the

tentacle bulbs and in both oral (oph) and aboral (aph) regions of the pharynx. (**i-x**) mRNA

expression patterns for *MleKlf5a* and *MleKlf5b* during embryogenesis. Maternal transcripts for

both *MleKlf5a* (**i, j**) and *MleKlf5b* (**k, l**) are found ubiquitously in zygotes (**i**), early cleavage

stages (**k**) and gastrulae (**j, l**). (**m-p**) Zygotic *MleKlf5a* and *MleKlf5b* transcripts exhibit

overlapping endodermal expression domains. At 9 hpf expression of *MleKlf5a* and *MleKlf5b*

transcripts are localized to the forming tentacular median ridges (arrowheads) and the developing

pharynx (oph + aph). (**q-t**) At 14 hpf *MleKlf5a* and *MleKlf5b* transcript expression is also found

in the apical organ (arrows) and epithelia of the newly formed gastrovascular cavity (gvc). (**u-x**)

At 20 hpf *MleKlf5a* and *MleKlf5b* transcripts are found in the tentacular median ridge

(arrowheads) and lateral ridge (lr), on either side of the apical organ floor (arrows), localized

towards the aboral end of the pharynx (aph), and the gastrovascular cavity epithelium (gvc).

Fig. 2: Two independent methods used to abrogate *MleKlf5a* and *MleKlf5b* gene function.

MleKlf5a (**a**) and *MleKlf5b* (**e**) exon-intron schematics show the location of sbMO targets (blue

boxes) and sgRNA targets (black triangle) used in this study. (**b,f**) Electrophoretic gels of PCR

products obtained using *MleKlf5a* and *MleKlf5b* sbMO RT-PCR primers (arrows in predicted

spliceform schematics to the right) on cDNA obtained from an individual morphant exemplar (*n*

= 21 morphants), 2-log DNA ladder was used for band size reference. Wildtype (wt) and mis-

spliced transcripts due to exon-skipping (-E) and/or intron retention (+I) were present in

morphants. (**c-d, g-h**) Crispant discordance plots and corresponding Sanger sequence traces.

sgRNA target site is underlined, Cas9 cut site is represented by dashed line and shows clearly

elevated sequence discordance relative to genome sequence reference and control sanger

sequencing. Sanger sequencing signal degradation is caused by the introduction of indels in an

individual crispant exemplar ($n = 17$ crispants; also see **EDF3b**).

Fig. 3: *MleKlf5a* and *MleKlf5b* double gene knockdown via sbMO and *sgRNA*-Cas9 genome

editing. (a-p) Live imaging showing morphology of 20 hpf cydippids. Aboral view in **a, c, e, g,**

i, k, m, o. Lateral view, oral up, in **b, d, f, h, j, l, n, p, q, r**. Schematic depiction of tentacular

axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Scale bars are 50

μm. Hoechst (blue) marks nuclei. Mitotracker (red) preferentially marks position of ctene row

polster cells, one pair per embryonic quadrant. Lysotracker (yellow) preferentially stains

epithelial cells lining the gastrovascular cavity (gvc). Tentacular median ridges (arrowheads) are

positioned medially along the tentacular axis and contacted by gvc epithelial cells. The pharynx

(ph) is positioned centrally and joins with the gvc aborally. The apical organ (arrow) is located at

the aboral pole of the embryo. Un-injected wildtype embryo (**a, b**). Morphology is unaffected in

embryos sham injected with control morpholino (MO) (**c, d**), *sgRNA* only (**i, j**) or Cas9 protein

only (**k, l**). In contrast, mild phenotypes in double gene knockdown morphants (**e, f**) and double

gene edited crispants (**m, n**) display aberrant distributions of gvc epithelial cells (Lysotracker

signal), aberrant patterning of the pharynx (ph) including aboral bifurcations (**f**, **n**, **EDF4**,

EDF5), aberrant patterning of the tentacle bulb and tentacular median ridges (arrowheads), and

atypical apical organ (arrow) morphology. Severe phenotypes in double gene knockdown

morphants (**g, h**) and double gene edited crispants (**o, p**) are reduced in size due to lack of

mesogleal ECM extrusion, display collapsed pharynx with gvc junction defects, significantly

reduced tentacle bulbs and tentacular median ridges (arrowheads), and apical organ defects

(arrow). (**q**) 25 min EdU pulse in wildtype 16 hpf embryo. Strong EdU incorporation signal

(orange) is found in the oral (oph) and aboral (aph) regions of the developing pharynx, epithelial

cells lining the gastrovascular cavity (gvc), the developing tentacular median ridges (arrowheads)

and lateral ridges (lr) of the tentacle bulbs. (**r**) 25 min EdU pulse in 16 hpf crispant embryo. EdU

incorporation signal (orange) is significantly diminished in regions that also express *MleKlf5a*

and *MleKlf5b* (refer to **Fig. 1**) including the pharynx, gvc and tentacular median ridges. Note that

lr cell proliferation is not affected. (**s**) Comparison of phenotypic proportions in morphants (*n* =

45) (51% no phenotype, 29% mild phenotype, 20% severe phenotype) vs crispants (*n* = 16) (6%

no phenotype, 19% mild phenotype, 75% severe phenotype).

Fig. 4: *MleKlf5a* and *MleKlf5b* double gene knockdown negatively affects apical organ

lithocyte production and negatively affects tentacular median ridge size. Live images of

embryos at 20 hpf. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA)

orientation are located in panel upper right. Aboral view in **a, c, e, h, j, l**. Lateral view, oral up, in

b, d, f, i, k, m. Scale bars are 50 μm. (**a, b**) lithocytes (arrow), dome cilia (arrowhead) and apical

organ (ao) in wildtype embryo. (**c, d**) double gene knockdown morphant lacking lithocytes. (**e, f**)

double gene edited crispant lacking lithocytes. Dome cilia (arrowheads) and balancer cilia are

present in both morphants and crispants. (**g**) *MleKlf5a* and *MleKlf5b* double gene knockdown

significantly reduces lithocyte production. Box plot center lines show medians, box limit at 25th

and 75th percentiles, whiskers extend 1.5 times interquartile range from the 25th and 75th

percentiles, data points are plotted as open circles. * = two-tailed *t*-test, $t = 3.47$, $P<0.005$. ** =

two-tailed *t*-test, $t = 6.52$, $P<0.00001$. Wildtype ($n = 14$), morphant ($n = 33$), crispant ($n = 15$).

(**h, i**) Wildtype tentacular median ridge (arrowhead) and lateral ridge (lr). (**j, k**) double gene

knockdown morphant with dramatically reduced tentacular median ridge. (**l, m**) double gene

edited crispant with dramatically reduced tentacular median ridge. The tentacle bulb lateral ridge

remains present in both morphants and crispants. (**n**) *MleKlf5a* and *MleKlf5b* double gene

knockdown significantly reduces tentacular median ridge length. Box plot center lines show the

median, box limit at 25th and 75th percentiles, whiskers extend 1.5 times interquartile range

from the 25th and 75th percentiles, data points are plotted as open circles. * = two-tailed *t*-test, *t*

$= 4.01$, $P<0.0005$. ** = two-tailed *t*-test, $t = 8.32$, $P<0.00001$. Each data point represents a

single tentacular median ridge length, with a maximum of 2 data points (i.e., 2 tentacular median

ridges) from each embryo. Data value of 0 indicate the absence of a tentacular median ridge

and/or tentacle bulb. Wildtype ($n = 28$), morphant ($n = 61$), crispant ($n = 34$).

Extended Data Fig. 1: Proliferating cells in the tentacular median ridge give rise to tentacle

muscle in *M. leidyi*. Aboral view and schematic depiction of tentacular axis (TA) and

pharyngeal axis (PA) orientation are located in panel upper right. Scale bars are 50 μm. EdU

pulse:chase immunohistochemistry on tentacle bulbs (**a**) after 25 min pulse:0 min chase and (**b**)

after 25 min pulse:24 hr chase. (**a**) 25 min pulse EdU marks rapidly proliferating cells in the

tentacular median ridge (arrow) and lateral ridges (lr) of the tentacle bulb. (**b**) After 25 min

pulse:24 hr chase, EdU incorporation is primarily detected in emergent tentacle muscle cells.

Extended Data Fig. 2: Additional *MleKlf5b* and *MleKlfX* transcript expression domains.

Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in

panel upper right. Scale bars are 50 μm. (**a, b**) 9 hpf and 14 hpf embryos, lateral view, oral up.

Dashed lines bound *MleKlf5b* ectodermal expression (magenta) in rapidly dividing cells

(compare with **Fig. 1b, f**) that flank developing ctene row pairs (visible as aligned columns of

DAPI stained nuclei bounded by *MleKlf5b* expression). (**c, d**) Magnified aboral views of

MleKlfX expression (magenta) in epithelial floor cells of the apical organ. (**c**) Initially *MleKlfX*

expression is detected at 16 hpf in four small clusters of cells at the boundary of each developing

embryonic quadrant converging in the center of the apical organ. (**d**) Expression of *MleKlfX*

resolves into several cell clusters by 18 hpf in both the tentacular (TA) and pharyngeal (PA)

axes.

Extended Data Fig. 3: ICE analysis of Crispants. (**a**) Sham injection has no effect on

embryonic development. Bar graph comparison of percentage of normal development in

wildtype $n = 161$ (24 abnormal); devitellinized but uninjected $n = 217$ (42 abnormal), two-tailed

Fisher's exact test $P=0.2761$; control MO injected $n = 49$ (10 abnormal), two-tailed Fisher's

exact test $P=0.379$; Cas9 only injected $n = 7$ (1 abnormal), two-tailed Fisher's exact test $P=1$;

sgRNA only injected $n = 4$ (1 abnormal), two-tailed Fisher's exact test $P=0.4851$. (**b**)

Distribution of ICE scores (estimate of indel proportion in signal trace from genomic DNA of

individual crispants) and KO scores (estimate of indel proportion that result in frameshift

mutation) for *MleKlf5a-sgRNA4* ($n = 14$) and *MleKlf5b-sgRNA3* ($n = 17$). Cas9 cut sites were

PCR amplified from genomic DNA prepared from individual crispants and subsequently Sanger

sequenced. Each cut site data point represents Sanger trace analyses from an individual embryo.

Box plot center lines show the median, box limit at 25th and 75th percentiles, whiskers extend

1.5 times interquartile range from the 25th and 75th percentiles, data points are plotted as open

circles.

Extended Data Fig. 4: Cas9 mediated genome editing of either *MleKlf5a* or *MleKlf5b*

resulted in reduced phenotype penetrance. Schematic depiction of tentacular axis (TA)

orientation located in panel upper right. Lateral view, oral up, scale bars are 50 μm. Single gene

editing produced primarily mild phenotypes for both *MleKlf5a* crispants (**a**) and *MleKlf5b*

crispants (**b**). (**c**) Bar graph comparison of distribution of mild (gray) vs severe (black)

phenotypes in *MleKlf5a* crispants ($n = 5$, 80% mild) and *MleKlf5b* crispants ($n = 5$, 80% mild).

Extended Data Fig. 5: Aberrant pharyngeal patterning. Schematic depiction of tentacular

axis (TA) and pharyngeal axis (PA) orientation located in panel upper right. Lateral views, oral

up, scale bars are 50 μm. (**a, b**) 20 hpf double gene morphant embryo with bifurcated (arrow)

pharynx (ph). (**c, d**) Same individual morphant 3 days post fertilization (dph) showing

persistence of pharyngeal bifurcation patterning defect just anterior of the pharyngeal-

gastrovascular junction resulting in a deletion of the pharyngeal folds.

Extended Data Fig. 6: Schematic of cydippid tentacle bulb. Aboral top, lateral bottom.

Tentacular median ridge colored light gray. Tentacular lateral ridges colored darker gray.

Tentacle (not shown) is rooted at the aboral medial base of the tentacle bulb. Location of

measurement taken for the tentacular median ridge length (refer to **Fig. 4n**) is denoted with red

dashed line.

Methods

Cloning and *in situ* hybridization

RNA was extracted using Trizol (Thermo Fisher Scientific) from *Mnemiopsis* embryos

collected at different stages of development and used to generate cDNA libraries (SMARTer kit,

Clontech). The coding sequences of *MleKlf5a*, *MleKlf5b*, and *MleKlfX* were amplified from

cDNA (Table 1) and cloned into pGEM-T Easy vector (Promega). The cloned fragments were

used as templates for *in vitro* transcription (MEGAscript, Ambion) of antisense digoxigenin-

labeled (Digoxigenin-11-UTP, Roche) riboprobes.

In situ hybridization followed ([46]). Riboprobes were used at a final concentration of ~0.5

ng/µl and hybridized with embryos for 24 hours. After color development, nuclei were labeled

with either DAPI (Molecular Probes) or Hoechst 33342 (Molecular Probes) in 1x PBS. Embryos

were immediately imaged or stored at -20°C in 70% glycerol in 1x PBS. Images were acquired

using a Zeiss Axio Imager.Z2, Zeiss AxioCam MRm Rev3 camera, and Zeiss Zen Blue software.

Fluorescent Z-stacks were deconvolved, post-processed for brightness and contrast and

assembled in Adobe Photoshop. Monochrome brightfield images were inverted, pseudocolored,

and overlaid onto fluorescent images of labeled nuclei. Samples were stored at -20°C in 70%

glycerol in 1x PBS.

Vital Dye staining

Live embryos were incubated in filtered seawater (FSW) containing a final concentration

of 100 nM MitoTracker (Deep Red FM, Molecular Probes), 100 nM LysoTracker (Red DND-99,

Molecular Probes), and 10 ng/µl Hoechst 33342 for one hour at room temperature. Live embryos

were then placed on glass microscope slides in a drop of FSW and relaxed with a drop of 6.5%

$MgCl_2$ (in dH_2O) on a coverslip positioned with clay feet. Images were acquired using a Zeiss

Axio Imager.Z2, Zeiss AxioCam MRm Rev3 camera, and Zeiss Zen Blue software. Fluorescent

Z-stacks were deconvolved, post-processed for brightness and contrast, and assembled in Adobe

Photoshop.

EdU labeling

Click-iT® EdU Alexa Fluor® 647 Imaging Kit (ThermoFisher Scientific) was used for

identification of proliferating cells. Embryos were collected at different developmental stages

and pulse incubated for 25 minutes with 100 µM EdU in a solution of a 1:1 volumetric ratio of

FSW to 6.5% $MgCl_2$ (dissolved in dH_2O) at room temperature. The EdU solution was washed

out and embryos were either fixed immediately or allowed to continue to develop during a 24

hour chase and subsequently fixed. Embryos were fixed with 4% PFA:FSW for 30 minutes at

room temperature, washed with 3% BSA:1xPBS), and incubated with 0.5% Triton X-100:1x

PBS for 20 minutes at room temperature. Fixed embryos were washed with 3% BSA:1x PBS and

stored at 4°C until used for EdU detection as per manufacturer protocol. Embryos were

subsequently washed with 1x PBS and mounted on glass microscope slides. Images were

acquired using a Zeiss Axio Imager.Z2, Zeiss AxioCam MRm Rev3 camera, and Zeiss Zen Blue

software. Fluorescent Z-stacks were deconvolved, post-processed for brightness and contrast,

and assembled in Adobe Photoshop or FIJI [47].

Microinjection

Laboratory cultures of adult *Mnemiopsis leidyi* on a ~12 hr:12 hr light:dark cycle were

spawned ~4 hours post darkness (hpd). At ~3.5 hpd individual adult *M. leidyi* were placed into 8-

inch glass bowls and screened for mature sperm and eggs. Freshly fertilized eggs were collected

by pipette.

Microinjection needles were pulled with a Brown micropipette puller (P-1000, Sutter

Instrument Company) using aluminosilicate glass capillaries (AF100-64-10, Sutter Instrument

Company). Pulled capillaries were beveled using a microelectrode beveler (BV-10, Sutter

Instrument Company). Beveled capillary needles were loaded via backfilling with injection

cocktails mixed with fluorescently-conjugated dextran (Invitrogen) (used to assess injection

success). Loaded capillary needles were mounted to a Xenoworks microinjection system (Sutter

Instrument Company) paired to a Zeiss Discovery V8 epiflourescent stereomicroscope.

Fertilized *M. leidyi* eggs were collected and passed sequentially through a 500 μm and a

400 μm cell strainer (pluriSelect Life Science) to remove excess mucus and egg jelly. Embryos

were then washed with 1x Penicillin/Streptomycin:FSW (PS:FSW). The embryonic vitelline

membranes were removed in gelatin coated dishes with acid-sharpened tungsten needles.

Embryos were then carefully transferred to an injection dish. After microinjection, embryos were

kept at room temperature until reaching the desired development stage for further analyses.

To determine if removing the vitelline membrane had an effect on embryogenesis, we

scored the percentage of normal development in embryos that were removed from the vitelline

membrane but not injected. There was no significant difference in the percentage of normal

development between embryos kept in their vitelline membrane (85%, n=161) and those that

were devitellinized but uninjected (80%, n=217; χ^2=1.5272, p=0.217; Extended Data Fig. 3a).

Morpholino oligonucleotides

Splice-blocking morpholino oligonucleotides (sbMOs, Gene Tools) were designed for

both *MleKlf5a* (*ML00922a*) and *MleKlf5b* (*ML25776a*). *MleKlf5a* sbMO #1 and sbMO #2

targeted intron 2-exon 3 and intron 3-exon 4 boundaries, respectively. *MleKlf5b* sbMO #1 and

sbMO #2 targeted exon 6-intron 6 and exon 7-intron 7 boundaries, respectively. A standard

control MO was used as a negative control. Sequences of sbMOs are listed in Table 1. Stock

solutions of sbMO in dH$_2$O were stored at room temperature. sbMO injection cocktail solutions

consisted of a final sbMO concentration of ~333 nM and ~0.5 mg/ml fluorescent dextran

(rhodamine or Alexa-Fluor 488, 10,000 MW, Invitrogen) in 35% glycerol. After phenotypic

analyses RNA was extracted from individual embryos (Arcturus PicoPure, ThermoFisher) and

cDNA prepared. Gene-specific primers were used on cDNA (OneTaq One-Step RT-PCR, New

England Biolabs) to evaluate aberrant transcript splicing via gel electrophoresis.

CRISPR/Cas9

We followed a cloning-free method to generate sgRNAs [48,49]. PCR amplified templates

were generated by annealing a 20-nt universal tracrRNA oligo to a sgRNA-specific oligo that

consisted of a T7 promoter, followed by the sgRNA target sequence, and a complementary

sequence to the tracrRNA oligo (Table 1). These templates were then *in vitro* transcribed

(MEGAscript, Ambion) to generate sgRNAs. The CasOT program [50] and *M. leidyi* reference

genome [51] were used to identify sgRNA target sites for *MleKlf5a* (*ML00922a*), *MleKlf5b*

(*ML25776a*), and *MleKlfX* (*ML20061a*). We selected sgRNAs that had no fewer than four

mismatches to alternative genomic sites to minimize potential off-target site (OTS) activity

(Table 1). Recombinant Cas9 protein (PNA Bio) and sgRNAs were injected at concentrations of

400 ng/µl of Cas9 protein and 100 ng/µl for each sgRNA. After phenotypic analysis, genomic

DNA was extracted from individual embryos (QIAamp DNA Micro, Qiagen) and each sgRNA

target site was amplified and Sanger sequenced. The ICE analysis tool (Synthego,

https://ice.synthego.com/#/) was used to determine Cas9 efficiency for each sgRNA. ICE

analysis gives two scores: an ICE score which reflects the percentage of indels found and a KO

score which reflects the percentage of indels that produce a frameshift mutation.

References

1. Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic

and adult fibroblast cultures by defined factors. *Cell* **126,** 663–676 (2006).

2. Takahashi, K. *et al.* Induction of pluripotent stem cells from adult human fibroblasts by

defined factors. *Cell* **131,** 861–872 (2007).

3. Rosselló, R. A. *et al.* Mammalian genes induce partially reprogrammed pluripotent stem

cells in non-mammalian vertebrate and invertebrate species. *eLIFE* **2,** e00036 (2013).

4. Jiang, J. *et al.* A core Klf circuitry regulates self-renewal of embryonic stem cells. *Nat. Cell*

Biol. **10,** 353–360 (2008).

5. Parisi, S. *et al.* Direct targets of Klf5 transcription factor contribute to the maintenance of

mouse embryonic stem cell undifferentiated state. *BMC Biol.* **8,** 128 (2010).

6. McConnell, B. B. & Yang, V. W. Mammalian Krüppel-like factors in health and diseases.

Physiol. Rev. **90,** 1337–1381 (2010).

7. Parisi, S. *et al.* Klf5 is involved in self-renewal of mouse embryonic stem cells. *J. Cell Sci.*

121, 2629–2634 (2008).

8. Nandan, M. O. & Yang, V. W. The role of Krüppel-like factors in the reprogramming of

somatic cells to induced pluripotent stem cells. *Histol. Histopathol.* **24,** 1343–1355 (2009).

9.   Nandan, M. O., Ghaleb, A. M., Bialkowska, A. B. & Yang, V. W. Krüppel-like factor 5 is

essential for proliferation and survival of mouse intestinal epithelial stem cells. *Stem Cell*

Res. **14,** 10–19 (2015).

10.   Steele, R. E., David, C. N. & Technau, U. A genomic view of 500 million years of cnidarian

evolution. *Trends Genet.* **27,** 7–13 (2011).

11.   Schnitzler, C. E., Simmons, D. K., Pang, K., Martindale, M. Q. & Baxevanis, A. D.

Expression of multiple *Sox* genes through embryonic development in the ctenophore

Mnemiopsis leidyi is spatially restricted to zones of cell proliferation. *Evodevo* **5,** 15 (2014).

12.   Dunn, C. W. *et al.* Broad phylogenomic sampling improves resolution of the animal tree of

life. *Nature* **452,** 745–749 (2008).

13.   Hejnol, A. *et al.* Assessing the root of bilaterian animals with scalable phylogenomic

methods. *Proc. Biol. Sci.* **276,** 4261–4270 (2009).

14.   Whelan, N. V. *et al.* Ctenophore relationships and their placement as the sister group to all

other animals. *Nat Ecol Evol* (2017). doi:10.1038/s41559-017-0331-3

15.   Alié, A. *et al.* Somatic stem cells express *Piwi* and *Vasa* genes in an adult ctenophore:

Ancient association of 'germline genes' with stemness. *Dev. Biol.* **350,** 183–197 (2011).

16.   Reitzel, A. M., Pang, K. & Martindale, M. Q. Developmental expression of 'germline'- and

'sex determination'-related genes in the ctenophore *Mnemiopsis leidyi*. *Evodevo* **7,** 17

(2016).

17.   Martindale, M. Q. The ontogeny and maintenance of adult symmetry properties in the

ctenophore, *Mnemiopsis mccradyi*. *Dev. Biol.* **118,** 556–576 (1986).

18.   Martindale, M. Q. The onset of regenerative properties in ctenophores. *Curr. Opin. Genet.*

Dev. **40,** 113–119 (2016).

19. Jager, M., Quéinnec, E., Chiori, R., Le Guyader, H. & Manuel, M. Insights into the early

evolution of SOX genes from expression analyses in a ctenophore. *J. Exp. Zool. B Mol.*

Dev. Evol. **310,** 650–667 (2008).

20. Presnell, J. S., Schnitzler, C. E. & Browne, W. E. KLF/SP Transcription Factor Family

Evolution: Expansion, Diversification, and Innovation in Eukaryotes. *Genome Biol. Evol.* **7,**

2289–2309 (2015).

21. Davidson, P. L. *et al.* The maternal-zygotic transition and zygotic activation of the

Mnemiopsis leidyi genome occurs within the first three cleavage cycles. *Mol. Reprod. Dev.*

(2017). doi:10.1002/mrd.22926

22. Chun, C. *Die ctenophoren des Golfes von Neapel und der angrenzenden meeres-abschnitte.*

(W. Engelmann, 1880).

23. von Byern, J., Mills, C. E. & Flammang, P. Bonding tactics in ctenophores—morphology

and function of the colloblast system. in *Biological Adhesive Systems: From Nature to*

Technical and Medical Application (eds. von Byern, J. & Grunwald, I.) 29–40 (Springer,

2010).

24. Ryan, J. F. *et al.* The genome of the ctenophore *Mnemiopsis leidyi* and its implications for

cell type evolution. *Science* **342,** 1242592 (2013).

25. Moroz, L. L. *et al.* The ctenophore genome and the evolutionary origins of neural systems.

Nature **510,** 109–114 (2014).

26. Tamm, S. L. Cilia and the life of ctenophores. *Invertebr. Biol.* **133,** 1–46 (2014).

27. Pang, K. & Martindale, M. Q. Ctenophores. *Curr. Biol.* **18,** R1119–20 (2008).

28. Freeman, G. The effects of altering the position of cleavage planes on the process of

localization of developmental potential in ctenophores. *Dev. Biol.* **51,** 332–337 (1976).

29. Fischer, A. H., Pang, K., Henry, J. Q. & Martindale, M. Q. A cleavage clock regulates

features of lineage-specific differentiation in the development of a basal branching

metazoan, the ctenophore *Mnemiopsis leidyi*. *Evodevo* **5,** 4 (2014).

30. Pang, K. & Martindale, M. Q. Comb Jellies (Ctenophora): A Model for Basal Metazoan

Evolution and Development. *CSH Protoc.* **2008,** db.emo106 (2008).

31. Tamm, S. L. Mechanisms of Ciliary Co-Ordination in Ctenophores. *J. Exp. Biol.* **59,** 231–

245 (1973).

32. Tamm, S. L. Formation of the statolith in the ctenophore *Mnemiopsis leidyi*. *Biol. Bull.* **227,**

7–18 (2014).

33. Martindale, M. Q. & Henry, J. J. Experimental Analysis of Tentacle Formation in the

Ctenophore *Mnemiopsis leidyi*. *Biol. Bull.* **193,** 245–247 (1997).

34. Martindale, M. Q. & Henry, J. Q. Intracellular fate mapping in a basal metazoan, the

ctenophore *Mnemiopsis leidyi*, reveals the origins of mesoderm and the existence of

indeterminate cell lineages. *Dev. Biol.* **214,** 243–257 (1999).

35. Babonis, L. S. *et al.* Integrating embryonic development and evolutionary history to

characterize tentacle-specific cell types in a ctenophore. *Mol. Biol. Evol.* (2018).

doi:10.1093/molbev/msy171

36. De Graeve, F. *et al.* Identification of the *Drosophila* progenitor of mammalian Krüppel-like

factors 6 and 7 and a determinant of fly development. *Gene* **314,** 55–62 (2003).

37. Weber, U., Rodriguez, E., Martignetti, J. & Mlodzik, M. Luna, a Drosophila KLF6/KLF7,

is maternally required for synchronized nuclear and centrosome cycles in the preblastoderm

embryo. *PLoS One* **9,** e96933 (2014).

38. Blakeley, P. *et al.* Defining the three cell lineages of the human blastocyst by single-cell

RNA-seq. *Development* **142,** 3613 (2015).

39. Horridge, G. A. Presumed photoreceptive cilia in a ctenophore. *Quart J Micr Sci* **105,** 311–

317 (1964).

40. Schnitzler, C. E. *et al.* Genomic organization, evolution, and expression of photoprotein and

opsin genes in *Mnemiopsis leidyi*: a new view of ctenophore photocytes. *BMC Biol.* **10,** 107

(2012).

41. Ilsley, M. D. *et al.* Krüppel-like factors compete for promoters and enhancers to fine-tune

transcription. *Nucleic Acids Res.* **45,** 6572–6588 (2017).

42. Aksoy, I. *et al.* Klf4 and Klf5 differentially inhibit mesoderm and endoderm differentiation

in embryonic stem cells. *Nat. Commun.* **5,** 3719 (2014).

43. Tamm, S. L. Patterns of comb row development in young and adult stages of the

ctenophores *Mnemiopsis leidyi* and *Pleurobrachia pileus*. *J. Morphol.* **273,** 1050–1063

(2012).

44. Sebé-Pedrós, A. *et al.* Early metazoan cell type diversity and the evolution of multicellular

gene regulation. *Nat Ecol Evol* **2,** 1176–1188 (2018).

45. Harbison, G. R. & Miller, R. L. Not all ctenophores are hermaphrodites. Studies on the

systematics, distribution, sexuality and development of two species of *Ocyropsis*. *Mar. Biol.*

90, 413–424 (1986).

46. Pang, K. & Martindale, M. Q. Ctenophore whole-mount in situ hybridization. *CSH Protoc.*

2008, db.prot5087 (2008).

47. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*

Methods **9,** 676–682 (2012).

48.   Kistler, K. E., Vosshall, L. B. & Matthews, B. J. Genome engineering with CRISPR-Cas9

in the mosquito Aedes aegypti. *Cell Rep.* **11,** 51–60 (2015).

49.   Varshney, G. K. *et al.* High-throughput gene targeting and phenotyping in zebrafish using

CRISPR/Cas9. *Genome Res.* **25,** 1030–1042 (2015).

50.   Xiao, A. *et al.* CasOT: a genome-wide Cas9/gRNA off-target searching tool. *Bioinformatics*

30, 1180–1182 (2014).

51.   Moreland, R. T. *et al.* A customized Web portal for the genome of the ctenophore

Mnemiopsis leidyi. *BMC Genomics* **15,** 316 (2014).

ACKNOWLEDGEMENTS

577   This work was supported in part by startup funds from the University of Miami College of Arts

578   and Sciences to WEB. JSP was supported by the University of Miami College of Arts and

579   Sciences. We thank Ricardo Cepeda for additional animal support.

AUTHOR CONTRIBUTIONS

582   WEB originally conceived the study and designed the research. JSP and WEB performed

583   experiments, collected and analyzed data, and wrote the manuscript. JSP and WEB read and

584   approved the final manuscript.

AUTHOR INFORMATION

587   Correspondence and requests for materials should be addressed to WEB.

588   (w.browne@miami.edu)

590   **Competing financial interests:** The authors declare no competing financial interests.

Table 1 | Primers and Oligonucleotides used in this study. All sequences are oriented 5'-3'. Blue nucleotide sequences correspond to T7 promoter. Black bold italicized nucleotide sequences correspond to genomic *MleKlf* targets and include addition of two 5' *G* residues to aid T7 polymerase binding. Red nucleotide sequences denote region of complementary between templated primers and Universal tracrRNA primer, which are annealed to the form the sgRNA transcription template.

Name	Use	Forward	Reverse
MleKlf5a	ISH probe	ATGAGTGCTATGACATG	AAACGTGTTCAAATGCCTCTT
MleKlf5b	ISH probe	ATGGACGTTTCCACGC	AGACGAGCTAGGGGGAACG
MleKlfX	ISH probe	GGCAGTTTAGTTCGATCGG	TGCAGTGAGTGGTAGGTT
MleKlf5a sbMO#1	sbMO	TCTCGTGTCTGAAACAATTTTAAGT	n/a
MleKlf5a sbMO#2	sbMO	GTCTACCACCTGCAAGATTTTAAGT	n/a
MleKlf5b sbMO#1	sbMO	CAGTTGATTTCTCACCTGCCAAGAA	n/a
MleKlf5b sbMO#2	sbMO	CAAACAGACTTACCTTCAAATGTGA	n/a
Standard Control	sbMO	CCTCTTACCTCAGTTACAATTTATA	n/a
MleKlf5a sbMO RT-PCR	sbMO validation	CCCTTGTAAACTTGAGCA	TCTTCGTGTAAACCTTCG
MleKlf5b sbMO RT-PCR	sbMO validation	GACAAGTTCCAAAGACTAAC	TACAGTAGATGAGGAGGTTT
Universal tracrRNA	sgRNA synthesis	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC	n/a
sgRNA template	sgRNA synthesis	GAAATTAATACGACTCACTATA**GG[N$_{XX}$]**GTTTTAGAGCTAGAAATAGC	n/a
MleKlf5a-gRNA1	sgRNA synthesis	AGCAACGGGTCCGTCCGT	n/a

Table1

MleKlf5a-gRNA2	sgRNA synthesis	TTGAGGGACGCGGGAGCAA	n/a
MleKlf5a-gRNA3	sgRNA synthesis	ACGGAGGGAATCGGCGAT	n/a
MleKlf5a-gRNA4	sgRNA synthesis	TTAGACTCTGTCGCGGGG	n/a
MleKlf5b-gRNA1	sgRNA synthesis	TGGTGATATACCAGGCG	n/a
MleKlf5b-gRNA2	sgRNA synthesis	ATCTTTCACGCTTAGGGGC	n/a
MleKlf5b-gRNA3	sgRNA synthesis	CGCTTGGAGGGAACCTAA	n/a
MleKlf5b-gRNA4	sgRNA synthesis	CTGAAACACCGGTCGCAG	n/a
MleKlf5a-sgRNA4 sequencing	sgRNA/Cas9 cut site validation	AAGACGTCCGATATTCTCTC	GGTGATCACTCCTACTGAAA
MleKlf5b-sgRNA3 sequencing	sgRNA/Cas9 cut site validation	GGTGTTCATACCTAGACGAT	TGTGTCTGTGTATAGTCGAG

Table 2 | Off-target CRISPR/Cas9 loci with mismatches to either *MleKlf5a* or *MleKl5b* target sequence and primers used for Sanger sequencing. All sequences are oriented 5'-3'.

Locus ID	# Mismatches	Forward	Reverse
ML090223a	4 (*MleKlf5a-sgRNA4)*	AAATTGTTTGTGTTCACACT	AGTTTTCTTTGTTTTCAGGG
ML021138a	5 (*MleKlf5a-sgRNA4)*	CAGCTTCATTGTAAAGAGTC	TTAGTTCTTATGTTTTCGCG
ML200217a	5 (*MleKlf5b-sgRNA3)*	GGATACTAGTTCCATAGCAG	CTCCTTGTTGATATTCTGGA
ML00363a	6 (*MleKlf5b-sgRNA3)*	TATGATTCTTGTTACCAGGG	ACATACGTCATCCTATTAGC
ML02979a	7 (*MleKlf5b-sgRNA3)*	CTGCATGATTACAAAGGTTT	ATGCTAAGAAGGATGCAATA

Figure1

Figure 1. *MleKlf5a* and *MleKlf5b* **are expressed in regions of rapid cell proliferation during embryogenesis in** *Mnemiopsis leidyi.* Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Aboral view in **a, c, e, g, j, l, m, o, q, s, u, w**. Lateral view, oral up, in **b, d, f, h, k, n, p, r, t, v, x**. Scale bars are 50 μm. **(a-h)** 25 min EdU pulse in 13 hours post fertilization (hpf) **(a-d)** and 16 hpf **(e-h)** embryos. EdU incorporation is localized to two cell clusters in the apical organ **(a, e)**, epithelial cells flanking the developing ctene rows (bounded by dashed lines) **(b, f)**, epithelial cells lining the gastrovascular cavity (gvc) **(d, h)** in the developing tentacular median ridges (arrowheads), and the oral (oph) and aboral (aph) regions of the pharynx **(d)**. **(g, h)** Later in development, EdU incorporation is found in the tentacular median ridge (arrowheads) and lateral ridges (lr) of the tentacle bulbs and in both oral (oph) and aboral (aph) regions of the pharynx. **(i-x)** mRNA expression patterns for *MleKlf5a* and *MleKlf5b* during embryogenesis. Maternal transcripts for both *MleKlf5a* **(i, j)** and *MleKlf5b* **(k, l)** are found ubiquitously in zygotes **(i)**, early cleavage stages **(k)** and gastrulae **(j, l)**. **(m-p)** Zygotic *MleKlf5a* and *MleKlf5b* transcripts exhibit overlapping endodermal expression domains. At 9 hpf expression of *MleKlf5a* and *MleKlf5b* transcripts are localized to the forming tentacular median ridges (arrowheads) and the developing pharynx (oph + aph). **(q-t)** At 14 hpf *MleKlf5a* and *MleKlf5b* transcript expression is also found in the apical organ (arrows) and epithelia of the newly formed gastrovascular cavity (gvc). **(u-x)** At 20 hpf *MleKlf5a* and *MleKlf5b* transcripts are found in the tentacular median ridge (arrowheads) and lateral ridge (lr), on either side of the apical organ floor (arrows), localized towards the aboral end of the pharynx (aph), and the gastrovascular cavity epithelium (gvc).

Figure2

Figure 2. Two independent methods used to abrogate *MleKlf5a* and *MleKlf5b* gene function. *MleKlf5a* (**a**) and *MleKlf5b* (**e**) exon-intron schematics show the location of sbMO targets (blue boxes) and sgRNA targets (black triangle) used in this study. (**b,f**) Electrophoretic gels of PCR products obtained using *MleKlf5a* and *MleKlf5b* sbMO RT-PCR primers (arrows in predicted spliceform schematics to the right) on cDNA obtained from an individual morphant exemplar ($n = 21$ morphants), 2-log DNA ladder was used for band size reference. Wildtype (wt) and mis-spliced transcripts due to exon-skipping (-E) and/or intron retention (+I) were present in morphants. (**c-d, g-h**) Crispant discordance plots and corresponding Sanger sequence traces. sgRNA target site is underlined, Cas9 cut site is represented by dashed line and shows clearly elevated sequence discordance relative to genome sequence reference and control sanger sequencing. Sanger sequencing signal degradation is caused by the introduction of indels in an individual crispant exemplar ($n = 17$ crispants; also see **EDF3b**).

Fig3

Figure 3. *MleKlf5a* and *MleKlf5b* double gene knockdown via sbMO and *sgRNA*-Cas9 genome editing. (**a-p**) Live imaging showing morphology of 20 hpf cydippids. Aboral view in **a, c, e, g, i, k, m, o**. Lateral view, oral up, in **b, d, f, h, j, l, n, p, q, r**. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Scale bars are 50 μm. Hoechst (blue) marks nuclei. Mitotracker (red) preferentially marks position of ctene row polster cells, one pair per embryonic quadrant. Lysotracker (yellow) preferentially stains epithelial cells lining the gastrovascular cavity (gvc). Tentacular median ridges (arrowheads) are positioned medially along the tentacular axis and contacted by gvc epithelial cells. The pharynx (ph) is positioned centrally and joins with the gvc aborally. The apical organ (arrow) is located at the aboral pole of the embryo. Un-injected wildtype embryo (**a, b**). Morphology is unaffected in embryos sham injected with control morpholino (MO) (**c, d**), *sgRNA* only (**i, j**) or Cas9 protein only (**k, l**). In contrast, mild phenotypes in double gene knockdown morphants (**e, f**) and double gene edited crispants (**m, n**) display aberrant distributions of gvc epithelial cells (Lysotracker signal), aberrant patterning of the pharynx (ph) including aboral bifurcations (**f**, **n**, **EDF4**, **EDF5**), aberrant patterning of the tentacle bulb and tentacular median ridges (arrowheads), and atypical apical organ (arrow) morphology. Severe phenotypes in double gene knockdown morphants (**g, h**) and double gene edited crispants (**o, p**) are reduced in size due to lack of mesogleal ECM extrusion, display collapsed pharynx with gvc junction defects, significantly reduced tentacle bulbs and tentacular median ridges (arrowheads), and apical organ defects (arrow). (**q**) 25 min EdU pulse in wildtype 16 hpf embryo. Strong EdU incorporation signal (orange) is found in the oral (oph) and aboral (aph) regions of the developing pharynx, epithelial cells lining the gastrovascular cavity (gvc), the developing tentacular median ridges (arrowheads) and lateral ridges (lr) of the tentacle bulbs. (**r**) 25 min EdU pulse in 16 hpf crispant embryo. EdU incorporation signal (orange) is significantly diminished in regions that also express *MleKlf5a* and *MleKlf5b* (refer to **Fig. 1**) including the pharynx, gvc and tentacular median ridges. Note that lr cell proliferation is not affected. (**s**) Comparison of phenotypic proportions in morphants (*n* = 45) (51% no phenotype, 29% mild phenotype, 20% severe phenotype) vs crispants (*n* = 16) (6% no phenotype, 19% mild phenotype, 75% severe phenotype).

Fig4

Figure 4. *MleKlf5a* and *MleKlf5b* double gene knockdown negatively affects apical organ lithocyte production and negatively affects tentacular median ridge size. Live images of embryos at 20 hpf. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Aboral view in **a, c, e, h, j, l**. Lateral view, oral up, in **b, d, f, i, k, m**. Scale bars are 50 μm. (**a, b**) lithocytes (arrow), dome cilia (arrowhead) and apical organ (ao) in wildtype embryo. (**c, d**) double gene knockdown morphant lacking lithocytes. (**e, f**) double gene edited crispant lacking lithocytes. Dome cilia (arrowheads) and balancer cilia are present in both morphants and crispants. (**g**) *MleKlf5a* and *MleKlf5b* double gene knockdown significantly reduces lithocyte production. Box plot center lines show medians, box limit at 25th and 75th percentiles, whiskers extend 1.5 times interquartile range from the 25th and 75th percentiles, data points are plotted as open circles. * = two-tailed *t*-test, = 3.47, <0.005. ** = two-tailed *t*-test, = 6.52, <0.00001. Wildtype (*n* = 14), morphant (*n* = 33), crispant (*n* = 15). (**h, i**) Wildtype tentacular median ridge (arrowhead) and lateral ridge (lr). (**j, k**) double gene knockdown morphant with dramatically reduced tentacular median ridge. (**l, m**) double gene edited crispant with dramatically reduced tentacular median ridge. The tentacle bulb lateral ridge remains present in both morphants and crispants. (**n**) *MleKlf5a* and *MleKlf5b* double gene knockdown significantly reduces tentacular median ridge length. Box plot center lines show the median, box limit at 25th and 75th percentiles, whiskers extend 1.5 times interquartile range from the 25th and 75th percentiles, data points are plotted as open circles. * = two-tailed *t*-test, = 4.01, <0.0005. ** = two-tailed *t*-test, = 8.32, <0.00001. Each data point represents a single tentacular median ridge length, with a maximum of 2 data points (i.e., 2 tentacular median ridges) from each embryo. Data value of 0 indicate the absence of a tentacular median ridge and/or tentacle bulb. Wildtype (*n* = 28), morphant (*n* = 61), crispant (*n* = 34).

EDF1

20hpf + no chase

a

TA ← → PA

ten

lr

EdU
Dapi

20hpf + 24 hr chase

b

TA ← → PA

ten

lr

EdU
Dapi

c

TA ← → PA

ten

lr

EdU

Extended Data Figure 1. Proliferating cells in the tentacular median ridge give rise to tentacle muscle in *M. leidyi.* Aboral view and schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Scale bars are 50 µm. EdU pulse:chase immunohistochemistry on tentacle bulbs (**a**) after 25 min pulse:0 min chase and (**b**) after 25 min pulse:24 hr chase. (**a**) 25 min pulse EdU marks rapidly proliferating cells in the tentacular median ridge (arrow) and lateral ridges (lr) of the tentacle bulb. (**b**) After 25 min pulse:24 hr chase, EdU incorporation is primarily detected in emergent tentacle muscle cells.

EDF2

9 hpf

a

TA

14 hpf

b

MleKlf5b

MleKlf5b
Dapi

16 hpf

c

TA ← → PA

18 hpf

d

MleKlfx

MleKlfx
Dapi

Extended Data Figure 2. Additional *MleKlf5b* and *MleKlfX* transcript expression domains. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. Scale bars are 50 µm. (**a, b**) 9 hpf and 14 hpf embryos, lateral view, oral up. Dashed lines bound *MleKlf5b* ectodermal expression (magenta) in rapidly dividing cells (compare with **Fig. 1b, f**) that flank developing ctene row pairs (visible as aligned columns of DAPI stained nuclei bounded by *MleKlf5b* expression). (**c, d**) Magnified aboral views of *MleKlfX* expression (magenta) in epithelial floor cells of the apical organ. (**c**) Initially *MleKlfX* expression is detected at 16 hpf in four small clusters of cells at the boundary of each developing embryonic quadrant converging in the center of the apical organ. (**d**) Expression of *MleKlfX* resolves into several cell clusters by 18 hpf in both the tentacular (TA) and pharyngeal (PA) axes.

EDF3

Extended Data Figure 3. ICE analysis of Crispants. (**a**) Sham injection has no effect on embryonic development. Bar graph comparison of percentage of normal development in wildtype *n* = 161 (24 abnormal); devitellinized but uninjected *n* = 217 (42 abnormal), two-tailed Fisher's exact test *P*=0.2761; control MO injected *n* = 49 (10 abnormal), two-tailed Fisher's exact test *P*=0.379; Cas9 only injected *n* = 7 (1 abnormal), two-tailed Fisher's exact test *P*=1; sgRNA only injected *n* = 4 (1 abnormal), two-tailed Fisher's exact test *P*=0.4851. (**b**) Distribution of ICE scores (estimate of indel proportion in signal trace from genomic DNA of individual crispants) and KO scores (estimate of indel proportion that result in frameshift mutation) for *MleKlf5a-sgRNA4* (*n* = 14) and *MleKlf5b-sgRNA3* (*n* = 17). Cas9 cut sites were PCR amplified from genomic DNA prepared from individual crispants and subsequently Sanger sequenced. Each cut site data point represents Sanger trace analyses from an individual embryo. Box plot center lines show the median, box limit at 25th and 75th percentiles, whiskers extend 1.5 times interquartile range from the 25th and 75th percentiles, data points are plotted as open circles.

EDF4

Extended Data Figure 4. Cas9 mediated genome editing of either *MleKlf5a* or *MleKlf5b* resulted in reduced phenotype penetrance. Schematic depiction of tentacular axis (TA) orientation located in panel upper right. Lateral view, oral up, scale bars are 50 µm. Single gene editing produced primarily mild phenotypes for both *MleKlf5a* crispants (**a**) and *MleKlf5b* crispants (**b**). (**c**) Bar graph comparison of distribution of mild (gray) vs severe (black) phenotypes in *MleKlf5a* crispants (*n* = 5, 80% mild) and *MleKlf5b* crispants (*n* = 5, 80% mild).

EDF5

MleKlf5a + MleKlf5b sbMO
20 hpf   3 dph

Extended Data Figure 5. Aberrant pharyngeal patterning. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation located in panel upper right. Lateral views, oral up, scale bars are 50 μm. (**a, b**) 20 hpf double gene morphant embryo with bifurcated (arrow) pharynx (ph). (**c, d**) Same individual morphant 3 days post fertilization (dph) showing persistence of pharyngeal bifurcation patterning defect just anterior of the pharyngeal-gastrovascular junction resulting in a deletion of the pharyngeal folds.

EDF6

Extended Data Figure 6. Cydippid tentacle bulb schematic. Aboral top, lateral bottom. Tentacular median ridge colored light gray. Tentacular lateral ridges colored darker gray. Tentacle (not shown) is rooted at the aboral medial base of the tentacle bulb. Location of measurement taken for the tentacular median ridge length (refer to **Fig. 4n**) is denoted with red dashed line.