***Krüppel-like factor* gene function in the ctenophore *Mnemiopsis leidyi* assessed by**

CRISPR/Cas9-mediated genome editing

Jason S Presnell[1,2], William E Browne[1,*]

[1]Department of Biology, University of Miami, Cox Science Center, 1301 Memorial Drive,

Miami, FL 33146, USA

[2]Present Address: Department of Human Genetics, University of Utah, 15 N 2030 E, Salt Lake

City, UT 84112, USA

ORCID ID: 0000-0002-7225-6692

*Corresponding Author:

William E. Browne, Department of Biology, University of Miami, Cox Science Center, 1301

Memorial Drive, Miami, FL 33146, USA

305-284-3319

w.browne@miami.edu

ORCID ID: 0000-0001-8200-6489

Keywords: Ctenophore, Krüppel-like factor, KLF5, endoderm, evolution, lithocyte

Abstract

The *Krüppel-like factor* (*Klf*) gene family encodes for transcription factors that play an important role in the regulation of stem cell proliferation, cell differentiation, and development in bilaterians. While *Klf* genes have been shown to be expressed in various cell types in non-bilaterian animals, their functional role in early diverging animal lineages has not been assessed. Thus, the ancestral activity of these transcription factors in animal development is not well understood. The ctenophore *Mnemiopsis leidyi* has emerged as an important non-bilaterian model system for understanding early animal evolution. Here we characterize the expression and functional role of *Klf* genes during *M. leidyi* embryogenesis. Zygotic *Klf* gene function was assessed with both CRISPR/Cas9-mediated genome editing knockout and splice-blocking morpholino oligonucleotide knockdown approaches. Abrogation of zygotic *Klf* expression during *M. leidyi* embryogenesis results in the irregular development of several organs including the pharynx, tentacle bulbs, and apical organ. Our data suggest an ancient role for *Klf* genes in regulating endodermal patterning.

Summary Statement (~15-30 words)

Using morpholino oligonucleotide knockdown and CRISPR/Cas9 genome editing, this study shows that in the ctenophore *Mnemiopsis leidyi,* tissues derived from the endoderm are dependent upon *Klf5* ortholog expression for proper development and patterning.

Introduction

Members of the *Krüppel-like factor (Klf)* gene family encode transcription factors with a

characteristic DNA binding domain composed of three C-terminal C2H2-zinc fingers

(McConnell and Yang, 2010; Presnell et al., 2015). During metazoan diversification, the *Klf*

transcription factor gene family expanded via duplication and domain shuffling events (Presnell

et al., 2015). *Klf* transcription factors are expressed in a variety of cells and tissues and have roles

in many biological processes, including proliferation of stem and progenitor cells, embryonic

development, germ layer differentiation, neuronal growth and regeneration, immune system

regulation and metabolic regulation (Bialkowska et al., 2017; McConnell and Yang, 2010;

Moore et al., 2009; Nagai et al., 2009; Oishi and Manabe, 2018; Pearson et al., 2008; Sweet et

al., 2018).

While KLF functional studies have been restricted to bilaterians, *Klf* genes are found in

the genomes of all metazoans (Presnell et al., 2015). Within Cnidaria, *Klfs* are expressed in

multipotent interstitial stem cells and their various downstream lineages, as well as in ectodermal

epithelial stem cells in *Hydra vulgaris* (Hemmrich et al., 2012; Siebert et al., 2019). In

Hydractinia symbiolongicarpus, *Klf* genes are upregulated in male sexual polyp bodies vs female

sexual polyp bodies (DuBuc et al., 2020). In a single-cell RNA-seq dataset for *Nematostella*

vectensis, *Klfs* were also shown to be expressed in various cell types, including the gastrodermis

and neuronal cells (Sebé-Pedrós et al., 2018a). Within Porifera, *Klfs* are upregulated in the stem-

cell like archaeocytes, epithelial pinacocytes, and mesenchymal cells in both *Spongilla lacustris*

and *Amphimedon queenslandica* (Musser et al., 2019; Sebé-Pedrós et al., 2018b). Single-cell

RNA-seq data for the Placozoan *Trichoplax adhaerens* revealed a single *Klf* gene upregulated in

epithelial cells (Sebé-Pedrós et al., 2018b). In ctenophores, three *Klf* genes have been identified

in two distantly related species, *Pleurobrachia bachei* and *Mnemiopsis leidyi* (Presnell et al.,

2015). The genome of *M. leidyi* contains *MleKlf5a*, *MleKlf5b*, and *MleKlfX* (Presnell et al.,

2015). *MleKlf5a* and *MleKlf5b* are the result of a lineage-specific duplication within the

Ctenophora, while *MleKlfX* is highly derived with no clear orthology to any known metazoan *Klf*

clade (Presnell et al., 2015). To date, single-cell and tissue-specific RNA-seq studies in *M. leidyi*

have not established differential expression signatures for *Klf* genes (Babonis et al., 2018; Sebé-

Pedrós et al., 2018b).

M. leidyi is a species of the non-bilaterian phylum Ctenophora, one of the earliest-
diverging extant metazoan lineages (Dunn et al., 2008; Hejnol et al., 2009; Whelan et al., 2017).
M. leidyi has been used extensively as a model for investigating early metazoan developmental
patterning, regeneration, and the evolution of animal traits (Babonis et al., 2018; Fischer et al.,
2014; Martindale and Henry, 1999; Presnell et al., 2016; Reitzel et al., 2016; Salinas-Saavedra
and Martindale, 2020; Schnitzler et al., 2014; Yamada et al., 2010). *M. leidyi* embryos undergo a
ctenophore-specific early cleavage program with gastrulation taking place ~3-5 hours post-
fertilization (hpf) followed by tissue organization and organogenesis over the next several hours
(Fischer et al., 2014; Freeman, 1976)(Fig. 1A). Four pairs of ctene rows, one pair in each
quadrant, are typically among the first differentiated ectodermal structures to appear (Fischer et
al., 2014). Each ctene plate is made up of polster cells bearing fused giant cilia (Tamm, 1973).
While initial ctene plate development is established by maternal factors (Fischer et al., 2014),
new ctene row expansion begins post-hatching during the juvenile cydippid stage (Tamm, 2012).
After the formation of the initial ctene rows, the developing embryo rapidly increases in size.
This period of rapid growth is accompanied by pharynx elongation along the aboral/oral axis, the
development of tentacle bulbs, and deposition of the first lithocytes onto the balancer cilia of the
apical organ (Martindale and Henry, 2015). Flanking the apical organ along the tentacular axis, a
pair of ectodermal invaginations and internal endodermal cells form the developing tentacle bulb
organs and the tentacular lateral and median ridges, respectively (Martindale and Henry, 1997b;
Martindale and Henry, 1999). Embryonic and adult tentacle bulb organs contain populations of
highly proliferative cells in the lateral ridge and median ridge tissues that give rise to
differentiated colloblast and tentacle muscle cells, respectively (Alié et al., 2011; Babonis et al.,
2018; Jager et al., 2008; Schnitzler et al., 2014). Moreover, genes associated with germline
development and stemness, including *Piwi*, *Vasa*, *Nanos*, and *Sox* homologs are highly expressed
in both the lateral and median ridges of the tentacle bulb, as well as in proliferative cell
populations in the developing apical organ and ctene rows, suggestive of the presence of
undifferentiated progenitor cells with stem-cell like properties in these tissues. In adult animals,
these undifferentiated progenitor cells presumably play a role in the continuous replacement of
lost cells (Alié et al., 2011; Jager et al., 2008; Reitzel et al., 2016; Schnitzler et al., 2014). After
~18-20 hpf, the fully developed cydippid hatches and maintains a feeding, pelagic lifestyle

before transitioning into the adult lobate body plan ~20 days post hatching (Martindale and

Henry, 2015).

Here we address *Klf* gene function in ctenophores by examining the expression of the

three known *Klf* homologs in *M. leidyi* during embryogenesis. We show results disrupting the

zygotic function for two *Klf* genes, *MleKlf5a* and *MleKlf5b,* using both morpholino

oligonucleotides and CRISPR/Cas9. Knockdown of *Klf* gene expression is associated with the

abnormal development of various organs during ctenophore embryogenesis and our data

provides additional insight into the function of the *Klf* gene family in the animal stem lineage.

Results

***MleKlf5a*, *MleKlf5b*, and *MleKlfX* expression**

MleKlf5a and *MleKlf5b* transcripts are maternally loaded in *M. leidyi* (Davidson et al.,

2017) similar to maternal loading of a number of *Klf* genes in other metazoans (Blakeley et al.,

2015; De Graeve et al., 2003; Weber et al., 2014). *MleKlf5a* and *MleKlf5b* transcripts are

detected in all embryonic cells through gastrulation (Fig. 1B-E). Post-gastrulation, transcripts for

both *MleKlf5a* and *MleKlf5b* become spatially restricted to cell populations associated with the

developing pharynx, gastrovascular system, tentacle bulb median ridges, and within the

developing apical organ (Fig. 1F-Q).

Within the developing pharynx, *MleKlf5a* and *MleKlf5b* expression are initially

widespread (Fig. 1G,I). As the pharynx elongates, *MleKlf5a* and *MleKlf5b* expression become

restricted to the interior-most cell layers of the medial and aboral pharyngeal regions (Fig.

1K,M,O,Q). The aboral-most region of the pharynx includes cells that form the junction with the

central gastrovascular cavity, or infundibulum. *MleKlf5a* and *MleKlf5b* expression is found

throughout the endodermal epithelial lining of the presumptive gastrodermis (Fig. 1J-Q). During

the initial development of the aboral apical organ, *MleKlf5a* and *MleKlf5b* expression is detected

in the apical organ floor epithelia. As the apical organ develops, *MleKlf5a* and *MleKlf5b*

expression becomes progressively restricted to cells located along the tentacular axis that are

positionally correlated with sites of lithocyte formation (Tamm, 2014; Fig. 1K,O,M,Q). Within

the developing tentacle bulbs, both *MleKlf5a* and *MleKlf5b* are expressed in the tentacular

median ridge (Fig. 1F-Q). An additional unique *MleKlf5b* expression domain is detected in a

narrow band of epidermal cells surrounding newly formed ctene row polster cells (Fig. 1H,L,P;

Fig. S1A,B).

In contrast to both *MleKlf5a* and *MleKlf5b*, *MleKlfX* expression is restricted to late

embryogenesis, first appearing ~16 hpf. Expression of *MleKlfX* transcripts are localized to a

small number of cells within the apical organ (Fig. S1C,D). One group of *MleKlfX* expressing

cells is located deep within the central epithelial floor of the developing apical organ. These cells

are located along the tentacular axis and pharyngeal axis forming a cross-shaped pattern (Fig.

S1C). A second shallower group of *MleKlfX* expressing cells is located within each quadrant just

medial of the ciliated grooves in the developing apical organ (Fig. S1D). These *MleKlfX*

expressing cells correspond positionally with the apical organ lamellate bodies, which may

represent putative photoreceptor cells (Horridge, 1964; Schnitzler et al., 2012), suggesting that

MleKlfX expression may be associated with light sensing neuronal cell types in the apical organ.

Knockdown of zygotic *MleKlf5a* and *MleKlf5b* expression

To characterize zygotic *Klf* gene function in *M. leidyi,* we used both splice-blocking

morpholinos (sbMOs) and CRISPR/Cas9-mediated genome editing to knockdown *Klf* gene

expression during embryonic development (Fig. 2). *MleKlfX* gene knockdown experiments failed

to produce any obvious morphological phenotypes, so we focus here on *MleKlf5a* and *MleKlf5b*

knockdown experiments. It has been shown that co-expressed KLFs bind to shared downstream

regulatory targets resulting in complex functional outcomes. For example, KLF2, KLF4, and

KLF5 have redundant roles in the downstream regulation of *Nanog* (Jiang et al., 2008), whereas

competition between KLF1 and KLF3 for binding sites can result in disparate functional

outcomes (Ilsley et al., 2017). For this reason, we sought to maximize the efficiency of

generating an observable phenotype by performing simultaneous *MleKlf5a* and *MleKlf5b*

knockdown with both sbMO and CRISPR/Cas9 genome editing experiments.

We injected single-cell embryos with either *MleKlf5a+MleKlf5b* sbMOs (KLF-MO

embryos) or *MleKlf5a-sgRNA+MleKlf5b-sgRNA* (KLF-Cas9 embryos). Microinjected embryos

were allowed to develop to ~20 hpf, stained with vital dyes, live-imaged, and compared to

equivalent late-stage wildtype embryos from the same spawns. Microinjections with standard

control MO, Cas9 protein alone, or sgRNAs alone had no detectable effect on embryonic

development (Fig. S2A). A selection of KLF-MO, KLF-Cas9, and wildtype embryos were

subjected to individual-embryo RNA or DNA analyses, respectively, to validate gene expression

knockdown. For both *MleKlf5a* and *MleKlf5b*, sbMOs produced mRNA splicing errors in KLF-

MO embryos via exon skipping and/or intron retention (Fig. 2A,B,E,F). An initial set of 4 single

guide RNAs (sgRNAs) were designed for *MleKlf5a* and *MleKlf5b* (Table 1) based on the *M.*

leidyi reference genome (Moreland et al., 2014; Moreland et al., 2020; Varshney et al., 2015).

For each gene, a single sgRNA, *MleKlf5a-sgRNA4* and *MleKlf5b-sgRNA3* (Fig. 2A,E), proved

efficient at mediating Cas9 double-stranded break activity at the target loci (Fig. 2C,D,G,H).

Sanger sequencing followed by ICE analysis (Hsiau et al., 2019) revealed a clear degradation of

sequence trace signal at the target loci in KLF-Cas9 embryos as compared to control embryos

(Fig. 2C,G), indicating the presence of indels and putative frameshift mutations generated by

sgRNA targeted Cas9 exonuclease activity (Fig. S2B). No evidence of off-target Cas9

exonuclease activity was observed (Table 2). Thus phenotypes generated by both gene

abrogation approaches in our study were due to the simultaneous disruption of *MleKlf5a* and

MleKlf5b gene expression.

179   KLF-MO and KLF-Cas9 embryos phenocopied one another and displayed phenotypes of

varying penetrance (Fig. 3A-P). A higher proportion of severe phenotypes were observed among

KLF-Cas9 embryos as compared to KLF-MO embryos (Fig. 3S), reflecting the effects of Cas9-

mediated genome editing versus titration of functional mRNAs by sbMOs. In contrast to the

observation of predominantly severe phenotypes in KLF-Cas9 embryos injected with *MleKlf5a-*

sgRNA4+MleKlf5b-sgRNA3 (Fig. 3O-P; Fig. 4E-F, L-M), single gene knockdown using either

MleKlf5a-sgRNA4 or *MleKlf5b-sgRNA3* primarily generated mild phenotypes (Fig. S3).

186   KLF-MO and KLF-Cas9 embryos with mild phenotypes underwent pharyngeal

elongation simultaneous with both mesoglea extrusion and a concomitant increase in size similar

to that observed in control embryos; however, experimental embryos displayed disorganized

patterning at the aboral end of the pharynx and the infundibular gastrovascular cavity (Fig. 3E-F,

M-N). Occasionally, we observed pharyngeal bifurcation at the junction of the pharynx with the

infundibular gastrovascular cavity (Fig. 3F,N; Fig. S3B; Fig. S4). In contrast, embryos having

severe phenotypes (Fig. 3G-H, O-P) failed to increase in size, reflecting a lack of extracellular

matrix (ECM) extrusion into the mesoglea space. Thus in severely affected embryos, the internal

embryonic space typically occupied by mesoglea is absent and the interior volume is completely

occupied by gastrovascular endoderm and abnormally elongated pharyngeal tissue. Both the

stomodeum and oral regions of the pharynx were still visible in severe mutant embryos,

indicating that the entire pharyngeal structure was not lost. However, it is unclear whether the

observed abnormal pharyngeal elongation is caused by *Klf* gene abrogation directly or is a spatial

effect due to the absence of mesogleal ECM.

The extrusion of ECM correlates with increased squamousity of epidermal cells (Fig.

4O). In severely affected animals we observed reduced epidermal squamousity and significantly

closer spacing of epidermal cells relative to normally developing animals (Fig. 4O-T). This

suggests reduced lateral tension forces on ectodermal cells in animals lacking mesogeal ECM.

Thus despite a decreased total body size, the ectodermal cell contribution to the epidermis

appears to be largely unaffected.

Among both KLF-MO and KLF-Cas9 embryos, patterning defects were also observed in

the apical organ (Fig. 3; Fig. 4A-G). *MleKlf5a* and *MleKlf5b* knockdown resulted in a significant

reduction of apical organ lithocytes as compared to control embryos (Fig. 4A-G). By 20 hpf,

control embryo statocysts contained an average of ~7 lithocytes (Fig. 4A-B, G). KLF-MO

embryos had an average ~4 lithocytes, with three embryos lacking lithocytes entirely (Fig.

4C,D,G). KLF-Cas9 embryos had an average of ~2 lithocytes, with five embryos completely

lacking lithocytes (Fig. 4E,F,G). Notably both KLF-MO and KLF-Cas9 embryos lacking

lithocytes still possessed phenotypically normal balancer cilia and dome cilia derived from

ectoderm (Fig. 4D,F).

The simultaneous abrogation of *MleKlf5a* and *MleKlf5b* also resulted in a dramatic

reduction in tentacle bulb size, particularly in the tentacular median ridge (Fig. 3; Fig. 4H-N; Fig.

S6K). We measured the tentacular median ridge width and found significant differences between

control and injected embryos (Fig. 4N; Fig. S5). The control embryo average median ridge width

was ~23 µm. KLF-MO and KLF-Cas9 embryo average median ridge width was ~18 µm and ~9

µm, respectively (Fig. 4N). Moreover, we observed that 15% of KLF-MO embryos and 29% of

KLF-Cas9 embryos lacked tentacular median ridges altogether (Fig. 4J-N).

The median ridge in adult *Pleurobrachia pileus* and juvenile *M. leidyi* cydippids has

previously been shown to contain populations of proliferative cells (Alié et al., 2011; Reitzel et

al., 2016; Schnitzler et al., 2014). In our *MleKlf5a* and *MleKlf5b* knockdown experiments the

relative size of the median ridge was consistently reduced, therefore we decided to perform EdU

incorporation assays during mid-late embryogenesis to assess cell proliferation. We observed

dramatically reduced EdU incorporation in areas affected by the knockdown of *MleKlf5a* and

MleKlf5b, including the median ridge and pharynx, suggesting that reduced cell proliferation

rates are associated with the attenuation of zygotic *Mle*KLF5a and *Mle*KLF5b activity (Fig.

3Q,R; Fig. S6).

Discussion

Our expression analyses of *MleKlf5a* and *MleKlf5b* in *M. leidyi* show that transcripts of

both genes are maternally loaded and ubiquitously distributed through gastrulation, corroborating

previous RNA-seq results which detected abundant transcripts for both *MleKlf5a* and *MleKlf5b*

during *M. leidyi* early embryonic cleavage stages (Davidson et al., 2017). Knockdown of zygotic

MleKlf5a and *MleKlf5b* expression does not appear to impact early embryonic development. The

zygotic expression of *MleKlf5a* and *MleKlf5b* display localized spatio-temporal patterns in post-

gastrulation embryos. Both *MleKlf5a* and *MleKlf5b* are expressed in the developing pharynx,

gastrodermis, tentacle bulbs and apical organ. These similar expression patterns could be due to

functionally redundant roles (Lynch and Conery, 2000). In contrast, the expression of *MleKlfX* is

restricted to late stages of development in a subset of apical organ epithelial cells. The *M. leidyi*

KlfX gene sequence is highly divergent relative to other metazoan *Klf* genes (Presnell et al.,

2015), suggestive of a *Mnemiopsis*-specific functional role for *MleKlfX*.

The *Klf* gene complement in *M. leidyi* is reduced compared to other non-bilaterian

lineages (Presnell et al., 2015), a trend observed in other gene families (Moroz et al., 2014; Ryan

et al., 2013). *Klf5-like* genes are found in all metazoans (McCulloch and Koenig, 2020; Presnell

et al., 2015). Among the non-bilaterian phyla, a *Klf5* ortholog in the cnidarian *Nematostella*

vectensis genome was shown to be expressed in a cluster of cells associated with digestive

filaments and the gastrodermis (Sebé-Pedrós et al., 2018a). In sponges, a *Klf5* ortholog was

found to be expressed in stem-cell like archaeocytes in the marine sponge *Amphimedon*

queenslandica (Sebé-Pedrós et al., 2018b), and in the digestive choanocytes and peptidocytes of

the freshwater sponge *Spongilla lacustris* (Musser et al., 2019). In vertebrates, *Klf5* orthologs are

required for the maintenance of intestinal crypt epithelia in the gut (Gao et al., 2015; Kuruvilla et

al., 2015; McConnell et al., 2011; Nandan et al., 2015). While less is known about *Klf5* orthologs

from invertebrate bilaterians, *Klf5* is expressed in several cephalopod embryonic tissues

including yolk cells and the developing mouth (McCulloch and Koenig, 2020). In our previous

phylogenetic study, it was unclear whether the few identified invertebrate sequences were either

Klf4 or *Klf5*, which share high sequence similarity (Presnell et al., 2015). One of these

sequences, *Drosophila melanogaster dar1*, shares sequence similarity to human *Klf5* but has a

functional role more similar to human *Klf4*, and was shown to play a role in regulation of gut

proliferation (Wu et al., 2018). Based on our expression analysis of *MleKlf5a* and *MleKlf5b* and

the observed dysregulation of gastrodermal patterning in *MleKlf5a+MleKlf5b* knockdown

embryos, our data suggest an evolutionarily conserved role for *Klf5-like* orthologs in the

regulation and maintenance of gut epithelia among metazoans.

M. leidyi endodermal cell lineages stem from early cleavage stage E and M oral

macromeres while ectodermal lineages originate from the aboral micromeres. Fate mapping

experiments show that the ectodermal micromeres contribute to the epidermis, ctene rows,

tentacle epithelia and colloblasts, balancer cilia and the epithelial floor of the apical organ, while

the endodermal macromeres give rise to the gastrodermis and associated endodermal canal

system, muscle, the tentacular median ridge, and the apical organ lithocytes (Henry and

Martindale, 2001; Martindale and Henry, 1997a; Martindale and Henry, 1999). Dysregulation of

MleKlf5a and *MleKlf5b* show consistent abnormal phenotypes associated with the development

of the apical organ and tentacle bulbs. In the apical organ of *MleKlf5a* and *MleKlf5b*

dysregulated embryos, the development of endodermally derived lithocytes is reduced or absent

while the ectodermally derived epithelial floor, balancer cilia, and dome cilia appear normal (Fig.

4A-G). Similarly in the developing tentacle bulb, abrogation of *MleKlf5a* and *MleKlf5b* activity

resulted in the absence or reduction in size of the endodermally derived median ridge (Fig. 4H-

N). Remaining tentacle tissue likely represents ectodermal contributions to tentacle epithelia and

colloblasts. The development of other ectodermally derived structures including the stomodeum

and epidermal cells were unaffected (Fig. 3; Fig. 4O-T). These results suggest that *MleKlf5a* and

MleKlf5b play a functional role in the development and maintenance of endodermally derived

tissues during *M. leidyi* embryogenesis.

With regard to the unique ectodermal expression domain of *MleKlf5b* (Fig. S1A-B), no

ectodermal or ctene row patterning phenotypes were observed in KLF-Cas9 embryos. However,

ctene rows showed gross spatial disorganization in phenotypically severe KLF-MO and KLF-

Cas9 embryos, possibly reflecting a requirement for coordinated contact between ectoderm and

underlying endoderm for precise ctene row alignment (Fig. 3G,O). For example, in

phenotypically mild KLF-MO and KLF-Cas9 embryos, ctene row morphogenesis did not occur

in 'quadrants' in which endodermal tissue failed to contact ectodermal tissue (Fig. 3E,M). This

result corroborates prior analyses indicating that ctene row development is at least partially

regulated through inductive interactions between endodermal and ectodermal cell lineages

(Fischer et al., 2014; Henry and Martindale, 2001; Henry and Martindale, 2004; Martindale and

Henry, 1997a). One possible explanation for the observed *MleKlf5b* expression pattern could be

that *MleKlf5b* is expressed in developing light producing photocytes derived from endodermal

2M macromeres that run subjacent to the ctene rows (Anctil, 1985; Fischer et al., 2014; Freeman

and Reynolds, 1973; Martindale and Henry, 1999; Schnitzler et al., 2012). An EdU-positive ring

of proliferative cells is situated around the ctene rows (Fig. S6C,G). These proliferative,

MleKlf5b positive cells may represent photocyte progenitor cells, as photocytes differentiate

relatively early during development (Fischer et al., 2014). Notably, the initial development of

differentiated polster cells/ctenes is specified by maternal factors, with additional ctenes

generated post embryonically. Therefore zygotic *MleKlf5b* would not directly impact the

specification of the initial ctenes during the stages observed in our study. An alternative

explanation is that these *MleKlf5b-* and EdU-positive ectodermal cells represent progenitor cells

that will give rise to new polster cells post-hatching and thus contribute to ctene row expansion.

In mammalian lineages *Klf5* orthologs help maintain stem cell renewal and promote

proliferation in the intestinal crypt and in pluripotent embryonic stem cells (Jiang et al., 2008;

Kuruvilla et al., 2015; Nandan et al., 2015; Parisi et al., 2008; Parisi et al., 2010). However, a

recent study suggests that mammalian pluripotency factors are not necessarily conserved in all

animals, and the ancestral metazoan stem cell toolkit primarily consists of genes associated with

the germline multipotency program (Alié et al., 2015; Juliano et al., 2010). Germline genes,

including *Piwi, Bruno, Pl-10*, have been shown to be expressed in putative progenitor cell

populations in the tentacle bulb, ctene rows, and apical organ of adult *Pleurobrachia* (Alié et al.,

2011). In *M. leidyi* cydippids, *Piwi, Vasa,* as well as *Sox* pluripotency factors are expressed in

these same tissues, suggesting that progenitor cell populations in these tissues express both

pluripotency factors as well as germline factors (Reitzel et al., 2016; Schnitzler et al., 2014). Our

EdU-staining recapitulates earlier work identifying zones of cell proliferation associated with the

developing pharynx, gastrodermis, areas around the ctene rows, and in the apical organ epithelial

floor (Reitzel et al., 2016; Schnitzler et al., 2014). These areas of cell proliferation correlate with

the zygotic transcript expression domains of both *MleKlf5a* and *MleKlf5b* (Fig. 1; Fig. 3Q; Fig.

S6).

Notably, sponge orthologs to *Klf5*, *Piwi, Bruno* and *Pl-10* are expressed in archaeocyte

and choanocyte cell types recognized as sponge equivalents to pluripotent and multipotent stem

cells respectively (Alié et al., 2015; Musser et al., 2019; Sebé-Pedrós et al., 2018b). Although we

were unable to perform quantitative analyses, our qualitative assessments show a diminution/loss

of EdU-positive cells in the median ridge and apical organ in *MleKlf5a+MleKlf5b* knockdown

embryos (Fig. 3Q-R; Fig. S6K). One interpretation of our results is that *MleKlf5a* and *MleKlf5b*

are expressed in proliferative cells and play a functional role in the maintenance of multipotent

endodermal progenitor cell populations.

To resolve whether *MleKlf5a* or *MleKlf5b* expressing cells are both proliferative and

multipotent will require additional experimentation. Future experiments involving the

knockdown of pluripotency and germline determination genes, such as *Piwi* and *Vasa*, along

with EdU assays may reveal further aspects of cellular proliferation and specification associated

with *Klf* activity. Alternatively, the observed phenotypes may be due to proliferation-

independent mechanisms establishing terminal cell identity. For example *MleKlf5a* and *MleKlf5b*

may regulate the terminal specification of lithocyte and tentacle muscle cell types. Based on this

work, while the explicit regulatory role of *MleKlf5a* and *MleKlf5b* remains unclear, our results

show that *MleKlf5a* and *MleKlf5b* are functionally associated with the formation, developmental

patterning and maintenance of endodermally derived structures in *M. leidyi* including the

gastrodermis, the tentacular median ridge, tentacle muscle, and apical organ lithocytes.

Conclusions

In this study, we used two independent methods, splice blocking morpholinos and Cas9-

mediated genome editing, to singly and coordinately abrogate zygotic *MleKlf5a* and *MleKlf5b*

gene function during *M. leidyi* embryonic development. We show that Cas9-mediated genome

editing is an effective method for performing combinatorial gene function assays in F0

ctenophore embryos. Along with recent RNA-seq data highlighting candidate genes associated

with zygotic gene activation and patterning of specific cell types in ctenophores (Babonis et al.,

2018; Davidson et al., 2017; Sebé-Pedrós et al., 2018b), both sbMO and CRISPR/Cas9 based

gene function assays in *M. leidyi* will provide much needed insight into the genetic mechanisms

underlying ctenophore biology and further our understanding of early metazoan evolution

(Jokura et al., 2019; Yamada et al., 2010). Our results begin to lay the groundwork for assessing

gene function essential for the embryonic development of *M. leidyi* and thus inform

developmental mechanisms unique to Ctenophora for the specification of terminally

differentiated cell types (e.g., lithocytes). Furthermore, our functional data suggest that *Klf*

activity is required for the proper development and maintenance of several endodermally derived

structures, possibly through the maintenance of multipotent progenitor cell proliferation. This

likely represents a conserved ancestral function for this transcription factor gene family in the

animal stem lineage.

Materials and methods

Cloning and *in situ* hybridization

RNA was extracted using Trizol (Thermo Fisher Scientific) from *Mnemiopsis* embryos

collected at different developmental stages and used to generate cDNA libraries (SMARTer kit,

Clontech). The coding sequences of *MleKlf5a*, *MleKlf5b*, and *MleKlfX* were amplified from

cDNA (Table 1) and cloned into pGEM-T Easy vector (Promega). The cloned fragments were

used as templates for *in vitro* transcription (MEGAscript, Ambion) of antisense digoxigenin-

labeled (Digoxigenin-11-UTP, Roche) riboprobes.

In situ hybridization followed (Pang and Martindale, 2008). Riboprobes were used at a

final concentration of ~0.5 ng/µl and hybridized with embryos for 24 hours. After color

development, nuclei were labeled with either DAPI (Molecular Probes) or Hoechst 33342

(Molecular Probes) in 1x PBS. Embryos were immediately imaged or stored at -20°C in 70%

glycerol in 1x PBS. Images were acquired using a Zeiss Axio Imager.Z2, Zeiss AxioCam MRm

Rev3 camera, and Zeiss Zen Blue software. Fluorescent Z-stacks were deconvolved, post-

processed for brightness and contrast and assembled in Adobe Photoshop. Monochrome

brightfield images were inverted, pseudo colored and overlaid onto fluorescent images of labeled

nuclei.

Vital Dye staining

Live embryos were incubated in filtered seawater (FSW) containing a final concentration

of 100 nM MitoTracker (Deep Red FM, Molecular Probes), 100 nM LysoTracker (Red DND-99,

Molecular Probes), and 10 ng/µl Hoechst 33342 for one hour at room temperature. Live embryos

were then placed on glass slides in a drop of FSW and relaxed with a drop of 6.5% $MgCl_2$ (in

dH_2O) on a coverslip positioned with clay feet. Images were acquired using a Zeiss Axio

Imager.Z2, Zeiss AxioCam MRm Rev3 camera, and Zeiss Zen Blue software. Fluorescent Z-

stacks were deconvolved, post-processed for brightness and contrast, and assembled in Adobe

Photoshop.

EdU labeling

Click-iT® EdU Alexa Fluor® 647 Imaging Kit (ThermoFisher Scientific) was used for

identification of proliferating cells. Embryos were collected at different developmental stages

and pulse incubated for 25 minutes with 100 µM EdU in a solution of a 1:1 volumetric ratio of

FSW to 6.5% $MgCl_2$ (dissolved in dH_2O) at room temperature. The EdU solution was washed

out and embryos were either fixed immediately or allowed to continue to develop during a 24-

hour chase and subsequently fixed. Embryos were fixed with 4% PFA in FSW for 30 minutes at

room temperature, washed with 3% BSA in 1x PBS, and incubated with 0.5% Triton X-100 in

1x PBS for 20 minutes at room temperature. Fixed embryos were washed with 3% BSA in 1x

PBS and stored at 4°C until used for EdU detection as per manufacturer protocol. Embryos were

subsequently washed with 1x PBS and mounted on glass microscope slides. Images were

acquired using a Zeiss Axio Imager.Z2, Zeiss AxioCam MRm Rev3 camera, and Zeiss Zen Blue

software. Fluorescent Z-stacks were deconvolved, post-processed for brightness and contrast,

and assembled in Adobe Photoshop or FIJI (Schindelin et al., 2012).

Microinjection

Laboratory cultures of adult *Mnemiopsis leidyi* on a ~12 hr:12 hr light:dark cycle were

spawned ~4 hours post darkness (hpd). At ~3.5 hpd individual adult *M. leidyi* were placed into 8-

inch glass bowls and screened for mature sperm and eggs. Freshly fertilized eggs were collected

by pipette.

Microinjection needles were pulled with a Brown micropipette puller (P-1000, Sutter

Instrument Company) using aluminosilicate glass capillaries (AF100-64-10, Sutter Instrument

Company). Pulled capillaries were beveled using a microelectrode beveler (BV-10, Sutter

Instrument Company). Beveled capillary needles were loaded via backfilling with injection

cocktails mixed with fluorescently-conjugated dextran (Invitrogen) (used to assess injection

success). Loaded capillary needles were mounted to a Xenoworks microinjection system (Sutter

Instrument Company) paired to a Zeiss Discovery V8 epifluorescence stereomicroscope.

Fertilized *M. leidyi* eggs were collected and passed sequentially through a 500 µm and a

400 µm cell strainer (pluriSelect Life Science) to remove excess mucus and egg jelly. Embryos

were then washed with 1x Penicillin/Streptomycin:FSW (PS:FSW). The embryonic vitelline

membranes were removed in gelatin-coated dishes with acid-sharpened tungsten needles.

Embryos were then carefully transferred to an injection dish. After microinjection, embryos were

kept at room temperature until reaching the desired development stage for further analyses.

To determine if removing the vitelline membrane had an effect on embryogenesis, we

scored the percentage of normal development in embryos that were removed from the vitelline

membrane but not injected. There was no significant difference in the percentage of normal

development between embryos kept in their vitelline membrane (85%, *n*=161) and those that

were devitellinized but un-injected (80%, *n*=217; χ^2 =1.5272, *p*=0.217; S3a).

Morpholino oligonucleotides

Splice-blocking morpholino oligonucleotides (sbMOs, Gene Tools) were designed for

both *MleKlf5a* (*ML00922a*) and *MleKlf5b* (*ML25776a*). *MleKlf5a* sbMO #1 and sbMO #2

targeted intron 2-exon 3 and intron 3-exon 4 boundaries, respectively. *MleKlf5b* sbMO #1 and

sbMO #2 targeted exon 6-intron 6 and exon 7-intron 7 boundaries, respectively. A standard

control MO was used as a negative control. Sequences of sbMOs are listed in Table 1. Stock

solutions of sbMO in dH$_2$O were stored at room temperature. sbMO injection cocktail solutions

consisted of a final sbMO concentration of ~333 nM and ~0.5 mg/ml fluorescent dextran

(rhodamine or Alexa-Fluor 488, 10,000 MW, Invitrogen) in 35% glycerol. After phenotypic

analyses RNA was extracted from individual embryos (Arcturus PicoPure, ThermoFisher) and

cDNA prepared. Gene-specific primers were used on cDNA (OneTaq One-Step RT-PCR, New

England Biolabs) to evaluate aberrant transcript splicing via gel electrophoresis.

CRISPR/Cas9

We followed a cloning-free method to generate sgRNAs (Kistler et al., 2015; Varshney et

al., 2015). PCR amplified templates were generated by annealing a 20-nt universal tracrRNA

oligo to a sgRNA-specific oligo that consisted of a T7 promoter, followed by the sgRNA target

sequence, and a complementary sequence to the tracrRNA oligo (Table 1). These templates were

then *in vitro* transcribed (MEGAscript, Ambion) to generate sgRNAs. The CasOT program

(Xiao et al., 2014) and *M. leidyi* reference genome (Moreland et al., 2014; Moreland et al., 2020)

were used to identify sgRNA target sites for *MleKlf5a* (*ML00922a*), *MleKlf5b* (*ML25776a*), and

MleKlfX (*ML20061a*). We selected sgRNAs that had no fewer than four mismatches to

alternative genomic sites to minimize potential off-target site (OTS) activity (Table 1; Table 2).

Recombinant Cas9 protein (PNA Bio) and sgRNAs were injected at concentrations of 400 ng/µl

of Cas9 protein and 100 ng/µl for each sgRNA. After phenotypic analysis, genomic DNA was

extracted from individual embryos (QIAamp DNA Micro, Qiagen) and each sgRNA target site

was amplified and Sanger sequenced. The ICE analysis tool (Hsiau et al., 2019) was used to

determine Cas9 efficiency for each sgRNA. ICE analysis gives two scores: an ICE score which

reflects the percentage of indels found and a KO score which reflects the percentage of indels

that produce a frameshift mutation.

Epidermal nuclei counts

Live images of wildtype, KLF-MO, and KLF-Cas9 embryos were used. Individual Z-

sections from Hoechst channels were focused on the outer epidermal layer for each embryo

oriented along the tentacular axis (TA). A 100 μm^2 region of interest (roi) was positioned

medially and oral of the ctene rows. Nuclei within the roi were manually counted. Nuclei counts

were quantified and plotted using R (http://shiny.chemgrid.org/boxplotr/).

ACKNOWLEDGEMENTS

This work was supported in part by startup funds from the University of Miami College of Arts

and Sciences to WEB. JSP was supported by the University of Miami College of Arts and

Sciences. We thank Ricardo Cepeda for additional animal support.

AUTHOR CONTRIBUTIONS

WEB originally conceived the study and designed the research. JSP and WEB performed

experiments, collected and analyzed data, and wrote the manuscript. JSP and WEB read and

approved the final manuscript.

AUTHOR INFORMATION

Correspondence and requests for materials should be addressed to WEB.

(w.browne@miami.edu)

Competing financial interests: The authors declare no competing financial interests.

References

Alié, A., Leclère, L., Jager, M., Dayraud, C., Chang, P., Le Guyader, H., Quéinnec, E. and Manuel,
M. (2011). Somatic stem cells express *Piwi* and *Vasa* genes in an adult ctenophore: Ancient
association of "germline genes" with stemness. *Dev. Biol.* **350**, 183–197.

Alié, A., Hayashi, T., Sugimura, I., Manuel, M., Sugano, W., Mano, A., Satoh, N., Agata, K. and
Funayama, N. (2015). The ancestral gene repertoire of animal stem cells. *Proc. Natl. Acad. Sci. U.*
S. A. **112**, E7093–100.

Anctil, M. (1985). Ultrastructure of the luminescent system of the ctenophore Mnemiopsis leidyi. *Cell*
Tissue Res. **242**, 333–340.

Babonis, L. S., DeBiasse, M. B., Francis, W. R., Christianson, L. M., Moss, A. G., Haddock, S. H.
D., Martindale, M. Q. and Ryan, J. F. (2018). Integrating embryonic development and
evolutionary history to characterize tentacle-specific cell types in a ctenophore. *Mol. Biol. Evol.*

Bialkowska, A. B., Yang, V. W. and Mallipattu, S. K. (2017). Krüppel-like factors in mammalian stem
cells and development. *Development* **144**, 737–754.

Blakeley, P., Fogarty, N. M. E., Del Valle, I., Wamaitha, S. E., Hu, T. X., Elder, K., Snell, P.,
Christie, L., Robson, P. and Niakan, K. K. (2015). Defining the three cell lineages of the human
blastocyst by single-cell RNA-seq. *Development* **142**, 3613.

Davidson, P. L., Koch, B. J., Schnitzler, C. E., Henry, J. Q., Martindale, M. Q., Baxevanis, A. D.
and Browne, W. E. (2017). The maternal-zygotic transition and zygotic activation of the
Mnemiopsis leidyi genome occurs within the first three cleavage cycles. *Mol. Reprod. Dev.*

De Graeve, F., Smaldone, S., Laub, F., Mlodzik, M., Bhat, M. and Ramirez, F. (2003). Identification
of the *Drosophila* progenitor of mammalian Krüppel-like factors 6 and 7 and a determinant of fly
development. *Gene* **314**, 55–62.

DuBuc, T. Q., Schnitzler, C. E., Chrysostomou, E., McMahon, E. T., Febrimarsa, Gahan, J. M.,
Buggie, T., Gornik, S. G., Hanley, S., Barreira, S. N., et al. (2020). Transcription factor AP2
controls cnidarian germ cell induction. *Science* **367**, 757–762.

Dunn, C. W., Hejnol, A., Matus, D. Q., Pang, K., Browne, W. E., Smith, S. A., Seaver, E., Rouse, G.
W., Obst, M., Edgecombe, G. D., et al. (2008). Broad phylogenomic sampling improves resolution

of the animal tree of life. *Nature* **452**, 745–749.

Fischer, A. H., Pang, K., Henry, J. Q. and Martindale, M. Q. (2014). A cleavage clock regulates
features of lineage-specific differentiation in the development of a basal branching metazoan, the
ctenophore *Mnemiopsis leidyi*. *Evodevo* **5**, 4.

Freeman, G. (1976). The effects of altering the position of cleavage planes on the process of localization
of developmental potential in ctenophores. *Dev. Biol.* **51**, 332–337.

Freeman, G. and Reynolds, G. T. (1973). The development of bioluminescence in the ctenophore
Mnemiopsis leidyi. *Dev. Biol.* **31**, 61–100.

Gao, Y., Cao, Q., Lu, L., Zhang, X., Zhang, Z., Dong, X., Jia, W. and Cao, Y. (2015). Kruppel-like
factor family genes are expressed during Xenopus embryogenesis and involved in germ layer
formation and body axis patterning. *Dev. Dyn.* **244**, 1328–1346.

Hejnol, A., Obst, M., Stamatakis, A., Ott, M., Rouse, G. W., Edgecombe, G. D., Martinez, P.,
Baguñà, J., Bailly, X., Jondelius, U., et al. (2009). Assessing the root of bilaterian animals with
scalable phylogenomic methods. *Proc. Biol. Sci.* **276**, 4261–4270.

Hemmrich, G., Khalturin, K., Boehm, A.-M., Puchert, M., Anton-Erxleben, F., Wittlieb, J.,
Klostermeier, U. C., Rosenstiel, P., Oberg, H.-H., Domazet-Loso, T., et al. (2012). Molecular
signatures of the three stem cell lineages in hydra and the emergence of stem cell function at the base
of multicellularity. *Mol. Biol. Evol.* **29**, 3267–3280.

Henry, J. Q. and Martindale, M. Q. (2001). Multiple inductive signals are involved in the development
of the ctenophore *Mnemiopsis leidyi*. *Dev. Biol.* **238**, 40–46.

Henry, J. Q. and Martindale, M. Q. (2004). Inductive interactions and embryonic equivalence groups in
a basal metazoan, the ctenophore *Mnemiopsis leidyi*. *Evol. Dev.* **6**, 17–24.

Horridge, G. A. (1964). Presumed photoreceptive cilia in a ctenophore. *Quart J Micr Sci* **105**, 311–317.

Hsiau, T., Conant, D., Rossi, N., Maures, T., Waite, K., Yang, J., Joshi, S., Kelso, R., Holden, K.,
Enzmann, B. L., et al. (2019). Inference of CRISPR Edits from Sanger Trace Data. *bioRxiv* 251082.

Ilsley, M. D., Gillinder, K. R., Magor, G. W., Huang, S., Bailey, T. L., Crossley, M. and Perkins, A.
C. (2017). Krüppel-like factors compete for promoters and enhancers to fine-tune transcription.
Nucleic Acids Res. **45**, 6572–6588.

Jager, M., Quéinnec, E., Chiori, R., Le Guyader, H. and Manuel, M. (2008). Insights into the early
evolution of SOX genes from expression analyses in a ctenophore. *J. Exp. Zool. B Mol. Dev. Evol.*
310, 650–667.

Jiang, J., Chan, Y.-S., Loh, Y.-H., Cai, J., Tong, G.-Q., Lim, C.-A., Robson, P., Zhong, S. and Ng,
H.-H. (2008). A core Klf circuitry regulates self-renewal of embryonic stem cells. *Nat. Cell Biol.* **10**,
353–360.

Jokura, K., Shibata, D., Yamaguchi, K., Shiba, K., Makino, Y., Shigenobu, S. and Inaba, K. (2019).
CTENO64 Is Required for Coordinated Paddling of Ciliary Comb Plate in Ctenophores. *Curr. Biol.*
29, 3510–3516.e4.

Juliano, C. E., Swartz, S. Z. and Wessel, G. M. (2010). A conserved germline multipotency program.
Development **137**, 4113–4126.

Kistler, K. E., Vosshall, L. B. and Matthews, B. J. (2015). Genome engineering with CRISPR-Cas9 in
the mosquito Aedes aegypti. *Cell Rep.* **11**, 51–60.

Kuruvilla, J. G., Ghaleb, A. M., Bialkowska, A. B., Nandan, M. O. and Yang, V. W. (2015). Role of
Krüppel-like factor 5 in the maintenance of the stem cell niche in the intestinal crypt. *Stem Cell*
Transl Investig **2**,.

Lynch, M. and Conery, J. S. (2000). The evolutionary fate and consequences of duplicate genes. *Science*
290, 1151–1155.

Martindale, M. Q. and Henry, J. Q. (1997a). Reassessing embryogenesis in the Ctenophora: the
inductive role of e1 micromeres in organizing ctene row formation in the "mosaic" embryo,
Mnemiopsis leidyi. *Development* **124**, 1999–2006.

Martindale, M. Q. and Henry, J. J. (1997b). Experimental Analysis of Tentacle Formation in the
Ctenophore *Mnemiopsis leidyi*. *Biol. Bull.* **193**, 245–247.

Martindale, M. Q. and Henry, J. Q. (1999). Intracellular fate mapping in a basal metazoan, the
ctenophore *Mnemiopsis leidyi*, reveals the origins of mesoderm and the existence of indeterminate
cell lineages. *Dev. Biol.* **214**, 243–257.

Martindale, M. Q. and Henry, J. Q. (2015). Ctenophora. In *Evolutionary Developmental Biology of*
Invertebrates 1: Introduction, Non-Bilateria, Acoelomorpha, Xenoturbellida, Chaetognatha (ed.

Wanninger, A.), pp. 179–201. Vienna: Springer Vienna.

McConnell, B. B. and Yang, V. W. (2010). Mammalian Krüppel-like factors in health and diseases.
Physiol. Rev. **90**, 1337–1381.

McConnell, B. B., Kim, S. S., Yu, K., Ghaleb, A. M., Takeda, N., Manabe, I., Nusrat, A., Nagai, R.
and Yang, V. W. (2011). Krüppel-like factor 5 is important for maintenance of crypt architecture
and barrier function in mouse intestine. *Gastroenterology* **141**, 1302–13, 1313.e1–6.

McCulloch, K. J. and Koenig, K. M. (2020). Krüppel-like factor/specificity protein evolution in the
Spiralia and the implications for cephalopod visual system novelties. *Proc. Biol. Sci.* **287**, 20202055.

Moore, D. L., Blackmore, M. G., Hu, Y., Kaestner, K. H., Bixby, J. L., Lemmon, V. P. and
Goldberg, J. L. (2009). KLF family members regulate intrinsic axon regeneration ability. *Science*
326, 298–301.

Moreland, R. T., Nguyen, A.-D., Ryan, J. F., Schnitzler, C. E., Koch, B. J., Siewert, K., Wolfsberg,
T. G. and Baxevanis, A. D. (2014). A customized Web portal for the genome of the ctenophore
Mnemiopsis leidyi. *BMC Genomics* **15**, 316.

Moreland, R. T., Nguyen, A.-D., Ryan, J. F. and Baxevanis, A. D. (2020). The *Mnemiopsis* Genome
Project Portal: integrating new gene expression resources and improving data visualization.
Database **2020**,.

Moroz, L. L., Kocot, K. M., Citarella, M. R., Dosung, S., Norekian, T. P., Povolotskaya, I. S.,
Grigorenko, A. P., Dailey, C., Berezikov, E., Buckley, K. M., et al. (2014). The ctenophore
genome and the evolutionary origins of neural systems. *Nature* **510**, 109–114.

Musser, J. M., Schippers, K. J., Nickel, M. and Mizzon, G. (2019). Profiling cellular diversity in
sponges informs animal cell type and nervous system evolution. *BioRxiv*.

Nagai, R., Friedman, S. L. and Kasuga, M. eds. (2009). *The Biology of Krüppel-like Factors*. Springer,
Tokyo.

Nandan, M. O., Ghaleb, A. M., Bialkowska, A. B. and Yang, V. W. (2015). Krüppel-like factor 5 is
essential for proliferation and survival of mouse intestinal epithelial stem cells. *Stem Cell Res.* **14**,
10–19.

Oishi, Y. and Manabe, I. (2018). Krüppel-Like Factors in Metabolic Homeostasis and Cardiometabolic

593   Disease. *Front Cardiovasc Med* **5**, 69.

594  **Pang, K. and Martindale, M. Q.** (2008). Ctenophore whole-mount in situ hybridization. *CSH Protoc.*
595   **2008**, db.prot5087.

596  **Parisi, S., Passaro, F., Aloia, L., Manabe, I., Nagai, R., Pastore, L. and Russo, T.** (2008). Klf5 is
597   involved in self-renewal of mouse embryonic stem cells. *J. Cell Sci.* **121**, 2629–2634.

598  **Parisi, S., Cozzuto, L., Tarantino, C., Passaro, F., Ciriello, S., Aloia, L., Antonini, D., De Simone, V.,**
599   **Pastore, L. and Russo, T.** (2010). Direct targets of Klf5 transcription factor contribute to the
600   maintenance of mouse embryonic stem cell undifferentiated state. *BMC Biol.* **8**, 128.

601  **Pearson, R., Fleetwood, J., Eaton, S., Crossley, M. and Bao, S.** (2008). Krüppel-like transcription
602   factors: a functional family. *Int. J. Biochem. Cell Biol.* **40**, 1996–2001.

603  **Presnell, J. S., Schnitzler, C. E. and Browne, W. E.** (2015). KLF/SP Transcription Factor Family
604   Evolution: Expansion, Diversification, and Innovation in Eukaryotes. *Genome Biol. Evol.* **7**, 2289–
605   2309.

606  **Presnell, J. S., Vandepas, L. E., Warren, K. J., Swalla, B. J., Amemiya, C. T. and Browne, W. E.**
607   (2016). The Presence of a Functionally Tripartite Through-Gut in Ctenophora Has Implications for
608   Metazoan Character Trait Evolution. *Curr. Biol.* **26**, 2814–2820.

609  **Reitzel, A. M., Pang, K. and Martindale, M. Q.** (2016). Developmental expression of "germline"- and
610   "sex determination"-related genes in the ctenophore *Mnemiopsis leidyi*. *Evodevo* **7**, 17.

611  **Ryan, J. F., Pang, K., Schnitzler, C. E., Nguyen, A.-D., Moreland, R. T., Simmons, D. K., Koch, B.**
612   **J., Francis, W. R., Havlak, P., NISC Comparative Sequencing Program, et al.** (2013). The
613   genome of the ctenophore *Mnemiopsis leidyi* and its implications for cell type evolution. *Science*
614   **342**, 1242592.

615  **Salinas-Saavedra, M. and Martindale, M. Q.** (2020). Par protein localization during the early
616   development of Mnemiopsis leidyi suggests different modes of epithelial organization in the
617   metazoa. *Elife* **9**,.

618  **Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,**
619   **Rueden, C., Saalfeld, S., Schmid, B., et al.** (2012). Fiji: an open-source platform for biological-
620   image analysis. *Nature Methods* **9**, 676–682.

Schnitzler, C. E., Pang, K., Powers, M. L., Reitzel, A. M., Ryan, J. F., Simmons, D., Tada, T., Park,
M., Gupta, J., Brooks, S. Y., et al. (2012). Genomic organization, evolution, and expression of
photoprotein and opsin genes in *Mnemiopsis leidyi*: a new view of ctenophore photocytes. *BMC*
Biol. **10**, 107.

Schnitzler, C. E., Simmons, D. K., Pang, K., Martindale, M. Q. and Baxevanis, A. D. (2014).
Expression of multiple *Sox* genes through embryonic development in the ctenophore *Mnemiopsis*
leidyi is spatially restricted to zones of cell proliferation. *Evodevo* **5**, 15.

Sebé-Pedrós, A., Saudemont, B., Chomsky, E., Plessier, F., Mailhé, M.-P., Renno, J., Loe-Mie, Y.,
Lifshitz, A., Mukamel, Z., Schmutz, S., et al. (2018a). Cnidarian Cell Type Diversity and
Regulation Revealed by Whole-Organism Single-Cell RNA-Seq. *Cell* **173**, 1520–1534.e20.

Sebé-Pedrós, A., Chomsky, E., Pang, K., Lara-Astiaso, D., Gaiti, F., Mukamel, Z., Amit, I., Hejnol,
A., Degnan, B. M. and Tanay, A. (2018b). Early metazoan cell type diversity and the evolution of
multicellular gene regulation. *Nat Ecol Evol* **2**, 1176–1188.

Siebert, S., Farrell, J. A., Cazet, J. F., Abeykoon, Y., Primack, A. S., Schnitzler, C. E. and Juliano,
C. E. (2019). Stem cell differentiation trajectories in Hydra resolved at single-cell resolution.
Science **365**,.

Sweet, D. R., Fan, L., Hsieh, P. N. and Jain, M. K. (2018). Krüppel-Like Factors in Vascular
Inflammation: Mechanistic Insights and Therapeutic Potential. *Front Cardiovasc Med* **5**, 6.

Tamm, S. L. (1973). Mechanisms of Ciliary Co-Ordination in Ctenophores. *J. Exp. Biol.* **59**, 231–245.

Tamm, S. L. (2012). Patterns of comb row development in young and adult stages of the ctenophores
Mnemiopsis leidyi and *Pleurobrachia pileus*. *J. Morphol.* **273**, 1050–1063.

Tamm, S. L. (2014). Formation of the statolith in the ctenophore *Mnemiopsis leidyi*. *Biol. Bull.* **227**, 7–
18.

Varshney, G. K., Pei, W., LaFave, M. C., Idol, J., Xu, L., Gallardo, V., Carrington, B., Bishop, K.,
Jones, M., Li, M., et al. (2015). High-throughput gene targeting and phenotyping in zebrafish using
CRISPR/Cas9. *Genome Res.* **25**, 1030–1042.

Weber, U., Rodriguez, E., Martignetti, J. and Mlodzik, M. (2014). Luna, a Drosophila KLF6/KLF7, is
maternally required for synchronized nuclear and centrosome cycles in the preblastoderm embryo.

PLoS One **9**, e96933.

Whelan, N. V., Kocot, K. M., Moroz, T. P., Mukherjee, K., Williams, P., Paulay, G., Moroz, L. L.
and Halanych, K. M. (2017). Ctenophore relationships and their placement as the sister group to all
other animals. *Nat Ecol Evol*.

Wu, X., Chen, Z., Gao, Y., Wang, L., Sun, X., Jin, Y. and Liu, W. (2018). The krüppel-like factor
Dar1 restricts the proliferation of Drosophila intestinal stem cells. *FEBS J.* **285**, 3945–3958.

Xiao, A., Cheng, Z., Kong, L., Zhu, Z., Lin, S., Gao, G. and Zhang, B. (2014). CasOT: a genome-wide
Cas9/gRNA off-target searching tool. *Bioinformatics* **30**, 1180–1182.

Yamada, A., Martindale, M. Q., Fukui, A. and Tochinai, S. (2010). Highly conserved functions of the
Brachyury gene on morphogenetic movements: insight from the early-diverging phylum Ctenophora.
Dev. Biol. **339**, 212–222.

Figure Legends

Fig 1. Zygotic *MleKlf5a* and *MleKlf5b* are primarily expressed in endodermally derived

tissues during embryogenesis in *Mnemiopsis leidyi*. A) Schematics highlighting major

morphological landmarks (e.g., ctene rows, pharynx, tentacle bulbs, apical organ) during *M.*

leidyi embryogenesis. Gastrulation typically occurs within 6 hours post-fertilization (hpf),

followed by rapid tissue remodeling and organogenesis over the next several hours. By 24 hpf

embryos are ready to hatch as cydippid larvae and have fully developed organ systems. The top

row is an aboral view. The bottom row is a lateral view with oral up and aboral down. (**B-Q**)

Whole-mount *in situ* hybridization for *MleKlf5a* and *MleKlf5b* during embryogenesis.

Orientation follows schematics from **A**. Aboral views in **C, E, F, H, J, L, N, P**. Lateral views in

D, G, I, K, M, O, Q. (**B-E**) Maternal transcripts for both *MleKlf5a* and *MleKlf5b* are found

ubiquitously in zygotes (**B**), early cleavage stages (**D**) and gastrulae (**C,E**). (**F-Q**) Zygotic

MleKlf5a and *MleKlf5b* transcript expression domains with corresponding schematics. (**F-I**)

Initially, expression of *MleKlf5a* and *MleKlf5b* zygotic transcripts are localized to the forming

tentacular median ridges (arrowheads) and the developing pharynx (oph + aph). (**J-M**) Later in

development, *MleKlf5a* and *MleKlf5b* transcript expression are also found in the developing

apical organ (arrows) and epithelia of the newly formed gastrovascular cavity (gvc). (**N-Q**) In

cydippids, *MleKlf5a* and *MleKlf5b* transcripts are found in the tentacular median ridge

(arrowheads) and lateral ridge (lr), on either side of the apical organ floor (arrows), localized

towards the aboral end of the pharynx (aph), and throughout the gastrovascular cavity epithelium

(gvc). (**L, P**) *MleKlf5b* transcripts are also expressed in an additional domain around the ctene

rows (asterisks). See also Fig. S1a-b. Scale bars: 50 µm. oph, oral end of the pharynx; TA,

tentacular axis; PA, pharyngeal axis.

Fig 2. Validation of independent methods used to abrogate *MleKlf5a* and *MleKlf5b* gene

function. *MleKlf5a* (**A**) and *MleKlf5b* (**E**) exon-intron schematics show the location of splice-

blocking morpholino oligonucleotide (sbMO) targets (blue boxes) and single-guide RNA

(sgRNA) targets (black triangles) used in this study. The orange bars indicate the location of the

DNA binding domain. (**B,F**) Electrophoretic gels of PCR products obtained using *MleKlf5a* and

MleKlf5b sbMO RT-PCR primers (arrows in predicted splice form schematics to the right) on

cDNA obtained from a whole individual KLF-MO embryo exemplar (*n* = 21 KLF-MO embryos)

and control embryo exemplar. A 2-log DNA ladder was used for band size reference. Wildtype

(wt) and mis-spliced transcripts due to exon-skipping (-E) and/or intron retention (+I) were

present in KLF-MO embryos. (**C-G**) Discordance plots produced using ICE software show

elevated sequence discordance downstream of the Cas9 cut site relative to control genomic

sequence. (**D,H**) Corresponding Sanger sequence traces from genomic DNA extracted from a

whole individual KLF-Cas9 embryo exemplar show signal degradation downstream of the Cas9

cut site as compared to a wildtype embryo. Sanger sequencing signal degradation is caused by

the introduction of indels in KLF-Cas9 embryos ($n = 17$ KLF-Cas9 embryos; also see S2b). The

sgRNA target site is underlined, the position of the Cas9 cut site is represented by a vertical

dashed line.

Fig 3. *MleKlf5a* and *MleKlf5b* double gene knockdown via sbMO and *sgRNA*-Cas9 genome

editing. (**A-P**) Live imaging showing morphology of 20 hpf cydippids. Aboral view in **A, C, E,**

G, I, K, M, O. Lateral view, oral up, in **B, D, F, H, J, L, N, P, Q, R**. Schematic depiction of

tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper right. (**A, B**)

Un-injected wildtype embryo. Hoechst (blue) marks nuclei. Mitotracker (red) preferentially

marks the position of ctene row polster cells, one pair per embryonic quadrant. Lysotracker

(yellow) preferentially stains epithelial cells lining the gastrovascular cavity (gvc). Tentacular

median ridges (arrowheads) are positioned medially along the tentacular axis and contacted by

gvc epithelial cells. The pharynx (ph) is positioned centrally and joins with the gvc aborally. The

apical organ (arrow) is located at the aboral pole of the embryo. Morphology is unaffected in

embryos sham injected with control morpholino (MO) (**C, D**), *sgRNA* only (**I, J**) or Cas9 protein

only (**K, L**). In contrast, mild phenotypes in double gene knockdown KLF-MO embryos (**E, F**)

and double gene edited KLF-Cas9 embryos (**M, N**) display aberrant distributions of gvc

epithelial cells (Lysotracker signal), aberrant patterning of the pharynx (ph) including aboral

bifurcations (**F**, **N**; refer to Fig. S4, Fig. S5), aberrant patterning of the tentacle bulb and

tentacular median ridges (arrowheads), and atypical apical organ (arrow) morphology. Severe

phenotypes in double gene knockdown KLF-MO embryos (**G, H**) and double gene edited KLF-

Cas9 embryos (**O, P**) are reduced in size due to lack of mesoglea ECM extrusion, display

collapsed pharynx with gvc junction defects, significantly reduced tentacle bulbs and tentacular

median ridges (arrowheads), and apical organ defects (arrow). (**Q**) 25 min EdU pulse in wildtype

16-20 hpf embryo. Strong EdU incorporation signal (orange) is found in the oral and aboral

regions of the developing pharynx (oph and aph respectively), epithelial cells lining the

gastrovascular cavity (gvc), the developing tentacular median ridges (arrowheads) and lateral

ridges (lr) of the tentacle bulbs. (**R**) 25 min EdU pulse in 16-20 hpf KLF-Cas9 embryo. EdU

incorporation signal (orange) is diminished in regions that also express *MleKlf5a* and *MleKlf5b*

(refer to Fig. 1, Fig. S6) including the pharynx, gvc and tentacular median ridges. Note that lr

cell proliferation is not affected. (**s**) Comparison of phenotypic proportions in KLF-MO embryos

(51% no phenotype, 29% mild phenotype, 20% severe phenotype, $n = 45$) vs KLF-Cas9 embryos

(6% no phenotype, 19% mild phenotype, 75% severe phenotype, $n = 16$). Scale bars are 50 µm.

Fig 4. *MleKlf5a* and *MleKlf5b* double gene knockdown disrupts the development of

endodermally derived cell types and structures including lithocytes and the tentacular

median ridge. Live images of embryos at ~20 hpf. Schematic depiction of tentacular axis (TA)

and pharyngeal axis (PA) orientation are located in panel upper right. Aboral view in **A, C, E, H,**

J, L. Lateral view, oral up, in **B, D, F, I, K, M, O-S**. (**A, B**) Wildtype embryo with view of the

apical organ (ao) showing position of lithocytes (arrow) and dome cilia (arrowhead). (**C, D**)

Representative double gene knockdown KLF-MO embryo and (**E, F**) representative double gene

edited KLF-Cas9 embryo lacking lithocytes. Dome cilia (arrowheads) and balancer cilia are

present in both KLF-MO and KLF-Cas9 embryos. (**G**) Quantification of lithocyte production.

MleKlf5a and *MleKlf5b* double gene knockdown significantly reduces lithocyte production.

Centerlines show the medians; box limits indicate the 25th and 75th percentiles; whiskers extend

1.5 times the interquartile range from the 25th and 75th percentiles; * = two-tailed *t*-test, $t = 3.47$,

$p<0.005$; ** = two-tailed *t*-test, $t = 6.52$, $p<0.00001$. Individual counts are plotted as black dots

where n = 14, 33, and 15 samples respectively. (**H, I**) Wildtype tentacular median ridge

(arrowhead) and lateral ridge (lr). (**J, K**) Representative double gene knockdown KLF-MO

embryo and (**L, M**) representative double gene-edited KLF-Cas9 embryo with dramatically

reduced tentacular median ridge. The tentacle bulb lateral ridge remains present in both KLF-

MO and KLF-Cas9 embryos. (**N**) Quantification of median ridge width. *MleKlf5a* and *MleKlf5b*

double gene knockdown significantly reduces tentacular median ridge width. Centerlines show

the medians; box limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the

interquartile range from the 25th and 75th percentiles; * = two-tailed t-test, t = 4.01, p<0.0005;

** = two-tailed t-test, t = 8.32, p<0.00001. Individual measurements are plotted as black dots

where n = 28, 61, and 34 samples respectively. Each measurement represents a single tentacular

median ridge width, with a maximum of 2 (i.e., 2 tentacular median ridges) from each embryo. A

measurement of 0 indicates the absence of a tentacular median ridge and/or tentacle bulb. (**O-S**)

Representative images with a 100 μm^2 region of interest focused on the outer epidermal cell layer

of wildtype (**O**), KLF-MO mild (**P**) and severe (**Q**), and KLF-Cas9 mild (**R**), and severe (**S**)

embryos. (**T**) Quantification of epidermal nuclei cell counts. Centerlines show the medians; box

limits indicate the 25th and 75th percentiles; whiskers extend 1.5 times the interquartile range

from the 25th and 75th percentiles; crosses represent sample means; ***** = two-tailed *t*-test, *t*= -

5.35225, *p* < .005; ****** = two-tailed *t*-test, *t* = -4.99757, *p* < .005. Area nuclei counts are plotted as

black dots where n = 5, 5, 4, 5, and 5 area count samples respectively. KLF-MO severe (*M =*

100.40, *SD = 10.7*) and KLF-Cas9 severe (*M = 106.40*, *SD = 17.4*) both had a significantly

higher density of epidermal nuclei counts per 100 μm^2 area than control embryos (*M = 51.20*, *SD*

= 17.5). There was no significant difference in epidermal nuclei counts for KLF-MO mild (*M =*

54.60, *SD = 11.0*) or KLF-Cas9 mild (*M = 53.25*, *SD = 22.8*) relative to control embryos. Scale

bars are 50 µm.

Figure S1. Additional *MleKlf5b* and *MleKlfX* transcript expression domains. Schematic

depiction of tentacular axis (TA) and pharyngeal axis (PA) orientation are located in panel upper

right. Scale bars are 50 µm. (**A, B**) organogenesis stage 7-15 hpf embryos, lateral view, oral up.

Dashed lines bound *MleKlf5b* ectodermal expression (magenta) in rapidly dividing cells

(compare with Fig. S6c, g) that flank developing ctene row pairs (visible as aligned columns of

DAPI stained nuclei bounded by *MleKlf5b* expression). (**C, D**) Magnified aboral views of

MleKlfX expression (magenta) in epithelial floor cells of the apical organ. (**C**) Initially *MleKlfX*

expression is detected at 16 hpf in four small clusters of cells at the boundary of each developing

embryonic quadrant converging in the center of the apical organ. (**D**) Expression of *MleKlfX*

resolves into several cell clusters by 18 hpf in both the tentacular (TA) and pharyngeal (PA)

axes.

Figure S2. ICE analysis of KLF-Cas9 embryos. (**A**) Sham injection has no effect on

embryonic development. Bar graph comparison of percentage of normal development in

wildtype $n = 161$ (24 abnormal); devitellinized but uninjected $n = 217$ (42 abnormal), two-tailed

Fisher's exact test $P=0.2761$; control MO injected $n = 49$ (10 abnormal), two-tailed Fisher's

exact test $P=0.379$; Cas9 only injected $n = 7$ (1 abnormal), two-tailed Fisher's exact test $P=1$;

sgRNA only injected $n = 4$ (1 abnormal), two-tailed Fisher's exact test $P=0.4851$. (**B**)

Distribution of ICE scores (estimate of indel proportion in signal trace from genomic DNA of

individual KLF-Cas9 embryos) and KO scores (estimate of indel proportion that result in

frameshift mutation) for *MleKlf5a-sgRNA4* ($n = 14$) and *MleKlf5b-sgRNA3* ($n = 17$). Cas9 cut

sites were PCR amplified from genomic DNA prepared from individual KLF-Cas9 embryos and

subsequently Sanger sequenced. Each cut site data point represents Sanger trace analyses from

an individual embryo. Box plot center lines show the median, box limit at 25th and 75th

percentiles, whiskers extend 1.5 times interquartile range from the 25th and 75th percentiles, data

points are plotted as open circles.

Figure S3. Cas9 mediated genome editing of either *MleKlf5a* or *MleKlf5b* resulted in

reduced phenotype penetrance. Schematic depiction of tentacular axis (TA) orientation located

in panel upper right. Lateral view, oral up, scale bars are 50 µm. Single gene editing produced

primarily mild phenotypes for both *MleKlf5a* KLF-Cas9 embryos (**A**) and *MleKlf5b* KLF-Cas9

embryos (**B**). (**C**) Bar graph comparison of distribution of mild (gray) vs severe (black)

phenotypes in *MleKlf5a* KLF-Cas9 embryos ($n = 5$, 80% mild) and *MleKlf5b* KLF-Cas9

embryos ($n = 5$, 80% mild).

Figure S4. Aberrant pharyngeal patterning. Schematic depiction of tentacular axis (TA) and

pharyngeal axis (PA) orientation located in panel upper right. Lateral views, oral up, scale bars

are 50 µm. (**A, B**) 20 hpf double gene KLF-MO embryo with bifurcated (arrow) pharynx (ph).

(**C, D**) Same individual KLF-MO embryo 3 days post fertilization (dph) showing persistence of

pharyngeal bifurcation patterning defect just anterior of the pharyngeal-gastrovascular junction

resulting in a deletion of the pharyngeal folds.

Figure S5. Cydippid tentacle bulb schematic. Aboral view: top; Lateral view: bottom.

Tentacular median ridge colored light gray. Tentacular lateral ridges colored darker gray.

Tentacle (not shown) is rooted at the aboral medial base of the tentacle bulb. Location of

measurement taken for the tentacular median ridge width (refer to Fig. 4n) is denoted with red

dashed line.

Figure S6. Regions of rapid cell proliferation. A) Schematics highlighting major

morphological landmarks (e.g., ctene rows, pharynx, tentacle bulbs, apical organ) during *M.*

leidyi embryogenesis. The top row is an aboral view. The bottom row is a lateral view with oral

up and aboral down. Schematic depiction of tentacular axis (TA) and pharyngeal axis (PA)

orientation are located in panel upper right. (**B-I**) Edu incorporation after 25 min pulse during

embryogenesis. Orientation follows schematics from **A**. Aboral views in **B,D,F,H**. Lateral views

in **C,E,G,I**. EdU incorporation is localized to two cell clusters in the apical organ (**B,F**),

epithelial cells flanking the developing ctene rows bounded by dashed lines (**C,G**), epithelial

cells lining the gastrovascular cavity (gvc) (**E,I**) in the developing tentacular median ridges

(arrowheads), and the oral (oph) and aboral (aph) regions of the pharynx (**E**). (**H,I**) Later in

development, EdU incorporation is found in the tentacular median ridge (arrowheads) and lateral

ridges (lr) of the tentacle bulbs and in both oral (oph) and aboral (aph) regions of the pharynx.

(**L**) tentacle bulb after 25 min pulse:0 min chase. (**M,N**) tentacle bulb after 25 min pulse:24 hr

chase, EdU incorporation is primarily detected in emergent tentacle muscle cells. Scale bars: 50

µm. TA, tentacular axis; PA, pharyngeal axis.

Table 1. Primers and Oligonucleotides used in this study. All sequences are oriented 5'-3'.
Italicized nucleotide sequences correspond to T7 promoter. **Bold** nucleotide sequences
correspond to genomic *MleKlf* targets and include addition of two 5' G residues to aid T7
polymerase binding. <u>Underlined</u> nucleotide sequences denote region of complementary between
templated primers and Universal tracrRNA primer, which are annealed to the form the sgRNA
transcription template.

Name	Use	Forward	Reverse
MleKlf5a	ISH probe	ATGAGTGCTATGACATG	AAACGTGTTCAAATGCCTCTT
MleKlf5b	ISH probe	ATGGACGTTTCCACGC	AGACGAGCTAGGGGGAACG
MleKlfX	ISH probe	GGCAGTTTAGTTCGATCGG	TGCAGTGAGTGGTAGGTT
MleKlf5a sbMO#1	sbMO	TCTCGTGTCTGAAACAATTTTAAGT	n/a
MleKlf5a sbMO#2	sbMO	GTCTACCACCTGCAAGATTTTAAGT	n/a
MleKlf5b sbMO#1	sbMO	CAGTTGATTTCTCACCTGCCAAGAA	n/a
MleKlf5b sbMO#2	sbMO	CAAACAGACTTACCTTCAAATGTGA	n/a
Standard Control	sbMO	CCTCTTACCTCAGTTACAATTTATA	n/a
MleKlf5a sbMO RT-PCR	sbMO validation	CCCTTGTAAACTTGAGCA	TCTTCGTGTAAACCTTCG
MleKlf5b sbMO RT-PCR	sbMO validation	GACAAGTTCCAAAGACTAAC	TACAGTAGATGAGGAGGTTT
Universal tracrRNA	sgRNA synthesis	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTT<u>GCTATTTCTAGCTCTAAAAC</u>	n/a

sgRNA template	sgRNA synthesis	*GAAATTAATACGAC TCACTATA***GG[N$_{XX}$]** GTTTTAGAGCTAG AAATAGC	n/a
MleKlf5a sgRNA #1	sgRNA synthesis	AGCAACGGGTCCG TCCGT	n/a
MleKlf5a sgRNA #2	sgRNA synthesis	TTGAGGGACGCGG GAGCAA	n/a
MleKlf5a sgRNA #3	sgRNA synthesis	ACGGAGGGAATC GGCGAT	n/a
MleKlf5a sgRNA #4	sgRNA synthesis	TTAGACTCTGTCG CGGGG	n/a
MleKlf5b sgRNA #1	sgRNA synthesis	TGGTGATATACCA GGCG	n/a
MleKlf5b sgRNA #2	sgRNA synthesis	ATCTTTCACGCTT AGGGGC	n/a
MleKlf5b sgRNA #3	sgRNA synthesis	CGCTTGGAGGGAA CCTAA	n/a
MleKlf5b sgRNA #4	sgRNA synthesis	CTGAAACACCGGT CGCAG	n/a
MleKlf5a sgRNA #4 sequencing	sgRNA/Cas9 validation	AAGACGTCCGATA TTCTCTC	GGTGATCACTCCT ACTGAAA
MleKlf5b sgRNA #3 sequencing	sgRNA/Cas9 validation	GGTGTTCATACCT AGACGAT	TGTGTCTGTGTAT AGTCGAG

841 **Table 2. Off-target CRISPR/Cas9 loci with mismatches to either MleKlf5a or MleKl5b**
842 **target sequence and primers used for Sanger sequencing.** All sequences are oriented 5'-3'.

Locus ID	# Mismatches	Forward	Reverse
ML090223a	4 (*MleKlf5a-sgRNA4)*	AAATTGTTTGTGTTC ACACT	AGTTTTCTTTGTTTT CAGGG
ML021138a	5 (*MleKlf5a-sgRNA4)*	CAGCTTCATTGTAAA GAGTC	TTAGTTCTTATGTTT TCGCG
ML200217a	5 (*MleKlf5b-sgRNA3)*	GGATACTAGTTCCAT AGCAG	CTCCTTGTTGATAT TCTGGA
ML00363a	6 (*MleKlf5b-sgRNA3)*	TATGATTCTTGTTAC CAGGG	ACATACGTCATCCT ATTAGC
ML02979a	7 (*MleKlf5b-sgRNA3)*	CTGCATGATTACAA AGGTTT	ATGCTAAGAAGGAT GCAATA

A — early cleavage stages and gastrulation 0-6 hpf

ctene rows, tentacle bulb, median ridge, pharynx, TA, PA, gastrovascular cavity (gvc), lithocyte, apical organ epithelia, reserve lithocyte, statocyst dome cilia, balancer cilia

organogenesis 7 - 15 hpf

cydippid stage 16 - 24 hpf

B zygote — *MleKlf5a* / Dapi

C gastrula stage

D gastrula stage — *MleKlf5b* / Dapi

E gastrula stage

F TA / PA

G TA — oph / aph

H TA / PA

I TA — oph / aph

J

K oph / aph — gvc

L

M oph / aph — gvc

N

O oph / aph — gvc / lr

P

Q oph / aph — gvc / lr

wildtype | **control MO** | *MleKlf5a + MleKlf5b* sbMO **mild** / **severe**

A — TA / PA — Hoechst / Mitotracker / Lysotracker
C
E
G

B — ph — TA
D — ph
F — ph
H — ph

sgRNA | Cas9 | *MleKlf5a-sgRNA4 + MleKlf5b-sgRNA3* Cas9 **mild** / **severe**

I — TA / PA — Hoechst / Mitotracker / Lysotracker
K
M
O

J — ph — TA
L — ph
N — ph
P — ph

wildtype | Cas9 *MleKlf5a-sgRNA4 MleKlf5b-sgRNA3*

Q — TA — oph / aph / gvc / lr — EdU / Dapi
R — oph / aph / lr

S

%

5a+5b sbMO   5a+5b Cas9

■ no phenotype   □ mild phenotype   ■ severe phenotype

wildtype

sbMO
MleKlf5a
MleKlf5b

Cas9
MleKlf5a-sgRNA4
MleKlf5b-sgRNA3

A TA ⊕ PA

B TA ⊞

ao

C

D

ao

E

F

ao

G

Number of lithocytes

wildtype   sbMO
5a+5b   Cas9
5a+5b

*   **

wildtype

sbMO
MleKlf5a
MleKlf5b

Cas9
MleKlf5a-sgRNA4
MleKlf5b-sgRNA3

H TA ⊕ PA

lr

I TA ▧

lr

J

lr

K

lr

L

lr

M

lr

Hoechst
Lysotracker

N

Median ridge width (µm)

wildtype   sbMO
5a+5b   Cas9
5a+5b

*   **

wildtype

sbMO
MleKlf5a
MleKlf5b

Cas9
MleKlf5a-sgRNA4
MleKlf5b-sgRNA3

O

P *mild*

R *mild*

Q *severe*

S *severe*

T

Epidermal nuclei/100 µm²

wildtype   sbMO
5a+5b   Cas9
5a+5b   sbMO
5a+5b   Cas9
5a+5b

mild   *severe*

n.s.   n.s.   *   **