Relation between prescribing of different antibiotics and rates of bacteremia/septicemia and associated mortality in the US and England

Edward Goldstein

Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard TH Chan School of Public Health, Boston, MA 02115 USA. Email: egoldste@hsph.harvard.edu

Abstract

Background: Antibiotic use may contribute to the rates of bacteremia, sepsis and associated mortality through various mechanisms. Additionally, rates of prescribing of all antibiotics and rates of bacteremia, sepsis, and associated mortality are strongly correlated both within the US and England. At the same time, there is limited information on the effect of prescribing of some antibiotics vs. others on the rates of severe outcomes associated with bacterial infections.

Methods: For each of several antibiotic types/classes, we looked at correlations between the proportions (state-specific in the US, CCG-specific in England) of a given antibiotic type/class among all prescribed antibiotics in the outpatient setting, and rates of outcomes (mortality with septicemia, ICD-10 codes A40-41 present on the death certificate in different age groups of adults in the US, and *E. coli* or MSSA bacteremia in England) per unit of antibiotic prescribing (defined as the rate of outcome divided by the rate of prescribing of all antibiotics).

Results: In the US, prescribing of penicillins was positively associated with rates of mortality with septicemia for persons aged 75-84y and 85+y, while prescribing of fluoroquinolones was positively associated with rates of mortality with septicemia for persons aged 50-64y between 2014-2015. In England, prescribing of penicillins other than amoxicillin/co-amoxiclav was positively associated with rates of both MSSA and *E. coli* bacteremia for the period between financial years 2014/15 through 2017/18. Additionally, as time progressed, correlations between prescribing for both trimethoprim and co-amoxiclav and rates of bacteremia in England decreased, while correlations between amoxicillin prescribing and rates of bacteremia increased.

Conclusions: Our results suggest that prescribing of penicillins is associated with rates of *E. coli* and MSSA bacteremia in England, and rates of mortality with septicemia in older US adults, which agrees with our earlier findings. Further work is needed to better understand the relation between antibiotic prescribing, including antibiotic replacement and reduction in prescribing and rates of severe bacterial infections, including those for which a given antibiotic is prescribed infrequently.

Introduction

Rates of hospitalization with septicemia and sepsis in the diagnosis, associated mortality, as well as monetary costs of those hospitalizations have been rising rapidly during the past decades in the US [1-4]. A recent estimate from the US CDC suggests that about 270,000 Americans die annually as a result of sepsis [5]. Moreover, that estimate is expected to increase significantly if longer-term, e.g. 90-day mortality following sepsis diagnosis is accounted for [6]. In England, while rates of certain severe infections and related mortality, such as *Clostridium difficile* infections and MRSA bacteremia have declined during recent years [7,8], rates of *E. coli* and MSSA bacteremia and associated mortality were increasing [7-9].

Bacteremia outcomes in England are laboratory confirmed and there is less uncertainty about the interpretation of the recorded trends for those outcomes compared to trends for septicemia/sepsis hospitalization rates in the US. Part of the reason behind the rapid growth in the rates of hospitalization with septicemia/sepsis in the diagnosis in the US is changes in diagnostic practices, including the implementation of sepsis screening protocols [10,11]. However, changes in diagnostic practices in the US cannot fully explain the rise in the rates of hospitalization with septicemia/sepsis in the diagnosis, particularly prior to 2010 [12]. Indeed, trends in the rates of hospitalizations with any diagnosis of sepsis in the US between 2003-2009 closely resemble the trends in the rates of hospitalizations that involved infection and the use of mechanical ventilation (Figure 1 in [12]). Antibiotic use and resistance can contribute to the rates of bacteremia/sepsis hospitalization and mortality through several mechanisms, particularly lack of clearance of resistant infections/colonization following antibiotic treatment, with some of those infections subsequently devolving into bacteremia/sepsis, and lethal outcomes [13-19]. Some of the more direct evidence for the relation between antibiotic resistance and subsequent hospitalization with severe infections, including bacteremia/sepsis is described in [19,18]; evidence about the relation between antibiotic resistance for hospitalized patients with sepsis and mortality, particularly in the US is presented in

[16,17]. Antibiotic use is an important driver of the prevalence of antibiotic resistance [20-23,9], and as such, it further contributes to the rates of sepsis and associated mortality. Moreover, antibiotic use may also contribute to the prevalence of resistance not only to the drug class used, but to other drugs as well as resistance to different drug classes tends to cluster in bacterial populations, leading to the phenomenon of co-selection [24,25]. For example, fluoroquinolone use was found to be associated with methicillin-resistant *S. aureus* (MRSA) infections [26-28], while amoxicillin use was found to be associated with trimethoprim resistance in Enterobacteriaceae in England [20], with trimethoprim resistance in urinary tract infections (UTIs) being positively associated with bacteremia outcomes [21]. Furthermore, prevalence of co-amoxiclav resistance in *E. coli* bacteremia in England is very high [29], and use of co-amoxiclav [30] and possibly related penicillins may contribute to the incidence of co-amoxiclav-resistant *E. coli* infections/colonization and associated bacteremia outcomes.

Our earlier work [13,14] has shown associations between the use of different antibiotics and rates of septicemia hospitalization and mortality in the US, including the association between the use of penicillins and rates of septicemia hospitalization and mortality in older adults. We also note that prevalence of resistance to penicillins, particularly in older US adults, is high for a variety of infections with both Gram-negative and Gram-positive bacteria [31-33]. Additionally, rates of overall antibiotic prescribing are strongly correlated with rates of septicemia mortality in US adults, and rates of bacteremia in England (e.g. Tables 1 and 4 in this paper), with some of the geographic variability in antibiotic prescribing rates, at least in England, being associated with variability in the prevalence of underlying health conditions and certain demographic factors [34]. However, less is known about the effect of using some antibiotics vs. others in the treatment of various syndromes on the rates of bacteremia, septicemia, and associated mortality. In this paper, we examine how the proportions of overall antibiotic prescribing that are for different antibiotic types/classes are related to the rates of *E. coli* and MSSA bacteremia in England, and the rates mortality with septicemia in different age groups of US adults. Those analyses are based on state-level US CDC data on antibiotic prescribing and mortality between 2014-2015 in [35,36], and on the Clinical Commissioning Groups (CCG)-level English data (from Oxford U/PHE) on antibiotic prescribing and bacteremia [37,38]. We hope that such simple ecological analyses would lead to further work on the effect of antibiotic prescribing, including the effect of replacing some antibiotics by others, as well as of reduction in antibiotic prescribing on the rates of bacteremia, sepsis and associated mortality.

Methods

Data

1. US. Data on annual state-specific mortality with septicemia (ICD-10 codes A40-A41.xx representing either the underlying or a contributing cause of death) between 2014-2015 for different age groups of adults (18-49y, 50-64y, 65-74y, 75-84y, 85+y) were extracted from the US CDC Wonder database [36]. We note that for most of those deaths, septicemia is listed as a contributing rather than the underlying cause of death on the death certificate [39]. Data on the annual state-specific per capita rates of outpatient prescribing for four classes of antibiotics: fluoroquinolones, penicillins, macrolides, and cephalosporins, as well as overall antibiotic prescribing in different states in 2014 and 2015 were obtained from the US CDC Antibiotic Patient Safety Atlas database [35].

2. England. We've considered the following nine antibiotic types/classes:

1. Amoxicillin (British National Formulary (BNF) code 0501013B0)
2. Co-amoxiclav (Amoxicillin/Clavulanic acid) (BNF code 0501013K0)
3. Penicillins (BNF section 5.1.1) excluding amoxicillin/co-amoxiclav
4. Tetracyclines (BNF section 5.1.3)
5. Macrolides (BNF section 5.1.5)
6. Cephalosporins + other beta-lactams (BNF section 5.1.2)
7. Fluoroquinolones (BNF section 5.1.12)
8. Trimethoprim (BNF code 0501080W0)
9. Urinary Tract Infection antibiotics (BNF 5.1.13 -- nitrofurantoin/fosfomycin/methenamine)

For each antibiotic type/class above, we've extracted data for the different Clinical Commissioning Groups (CCGs) on the proportion of that antibiotic type/class among all antibiotic prescriptions (BNF classes 5.1.1 through 5.1.13) in the given CCG for each of the financial years (April through March) 2014/15 through 2017/18 [37]. We've also extracted CCG/year specific data on the prescribing of all antibiotics per 1,000 residents, as well as per 1,000 STAR-PUs [37,40]. In addition to prescribing data, we've extracted CCG/year-specific data on the (population-adjusted) rates of *E. coli* and MSSA bacteremia for each of the financial years 2014/15 through 2017/18 [38].

Correlation analysis

The contribution of prescribing of a given antibiotic type/class to the rates of severe outcomes associated with bacterial infections (e.g. bacteremia, or mortality with septicemia) is expected to be proportional to the rate of prescribing of that antibiotic type/class. One of the factors that modulates the relationship between the rates of antibiotic prescribing and rate of severe outcomes is the rate of infection that affects both prescribing and severe outcomes. Correspondingly, associations between rates of antibiotic prescribing and rates of severe outcomes are often positive (e.g. Tables 1 and 4 in this paper). The effect of the *proportion* of given antibiotic type/class among all antibiotics prescribed on the rates of severe outcomes associated with bacterial infections is expected to be proportional to the rates of severe outcomes per unit of antibiotic prescribing, defined as the rates of severe outcomes divided by the rates of prescribing of all antibiotics. If a unit of prescribing of a given antibiotic type/class affects the rate of a given severe outcome disproportionately (e.g. as a result of high prevalence of resistance to a given antibiotic) compared to the affect of prescribing of an average antibiotic dose, the association between the proportion of given antibiotic type/class among all antibiotics prescribed and the rate of a given severe outcome per unit of antibiotic prescribing is expected to be positive. We note that such disproportionate effects of prescribing of a unit of a given antibiotic can be the result not only of treatment of infections leading to a given outcome by a given antibiotic, but also of the contribution of the use of a given antibiotic to the rates of infection/colonization with different bacteria, and the prevalence of resistance to other antimicrobials (see Introduction). Additionally, correlations between proportions of a given antibiotic type/class among all antibiotics prescribed and the rates of severe outcomes per unit of antibiotic prescribing can also be affected by patterns of antibiotic prescribing in different locations, including changes in those patterns resulting from increases in resistance, introduction of new prescribing guidelines, etc. Correspondingly, we studied correlations between the proportions (state-specific in the US, CCG-specific in England) of a given antibiotic type/class among all prescribed antibiotics in the outpatient setting, and rates of outcome (mortality with septicemia in different age groups of adults in the US, and *E. coli* or MSSA bacteremia in England) per unit of antibiotic prescribing, with the caveats above regarding the causal relations underlying those correlations. The US analysis is done for the 2014-2015 period; given the ongoing changes in antibiotic prescribing patterns in England, correlations for the English data were computed for each financial year between 2014/15 through 2017/18.

Results

1. US

Table 1 shows, for each age group of adults, the mean (standard deviation) for the state-specific average annual rates of mortality with septicemia per 100,000 individuals in that age group between 2014-2015, as well as the linear correlation between those rates and state-specific rates of prescribing of all antibiotics. The latter correlations are high, ranging from 0.59(0.37,0.74) for ages 85+y to 0.77(0.62,0.68) for ages 65-74y. Additionally, annual rates of mortality with septicemia rise rapidly with age, from the mean of 8.31/100,000 for persons aged 18-49y to a mean of 750/100,000 for persons aged 85+y.

	Mean (standard deviation)	Linear correlation with rate of prescribing of all antibiotics
Septicemia mortality rate, ages 18-49y	8.31 (2.98)	**0.66(0.47,0.79)**
Septicemia mortality rate, ages 50-64y	55.8 (17.3)	**0.74(0.59,0.84)**
Septicemia mortality rate, ages 65-74y	143.4 (36.5)	**0.77(0.62,0.68)**
Septicemia mortality rate, ages 75-84y	330.8 (75.6)	**0.67(0.48,0.80)**
Septicemia mortality rate, ages 85+y	750 (161.9)	**0.59(0.37,0.74)**

Table 1: State-specific rates of mortality with septicemia (ICD-10 codes A40-41.xx present as either underlying or contributing causes on a death certificate) per 100,000 individuals in different age groups between 2014-2015 (mean + standard deviation), and the linear correlation between those rates and state-specific rates of prescribing of all antibiotics.

Table 2 shows correlations (both linear and Spearman) for each pair of antibiotic classes between state-specific percentages of all antibiotic prescriptions that were for each antibiotic class between 2014-2015, as well as the mean (standard deviation) for the state-specific percentages of all antibiotic prescriptions that were for each antibiotic class between 2014-2015. There is a strong negative correlation between the percentages of antibiotic prescribing that are for fluoroquinolones

and that are for penicillins; the Spearman correlation between percentages of antibiotic prescribing that are for penicillins and that are for cephalosporins was also negative. Those negative correlations suggest competition between certain antibiotics in prescriptions for various syndromes. We also note that on average, 66.2% of all antibiotic prescriptions in different states were for the four studied classes of antibiotics. Additionally, proportions of different antibiotic classes used in the US are notably different from the corresponding proportions in England (compare Table 2 with Table 5). Those differences, particularly for fluoroquinolones and cephalosporins may be related to differences in the rates of severe bacterial infections, particularly *Clostridium difficile* infections between the two countries [41].

	Mean (standard deviation)	Linear (Pearson) correlation with proportions of other antibiotic classes		
		Proportion penicillins	Proportion cephalosporins	Proportion macrolides
Proportion fluoroquinolones	11.81% (1.22%)	**-0.563 (-0.73,-0.34)**	-0.141 (-0.4,0.14)	0.123 (-0.16,0.39)
Proportion penicillins	22.57 (1.64%)		-0.262 (-0.5,0.01)	-0.235 (-0.48,0.04)
Proportion cephalosporins	13.56% (1.75%)			-0.164 (-0.42,0.12)
Proportion macrolides	18.22% (1.21%)			
	Mean (standard deviation)	Spearman correlation with proportions of other antibiotic classes		
		Proportion penicillins	Proportion cephalosporins	Proportion macrolides
Proportion fluoroquinolones	11.81% (1.22%)	**-0.436 (0.0015)**	-0.069 (0.63)	0.114 (0.42)
Proportion penicillins	22.57 (1.64%)		**-0.306 (0.03)**	-0.192 (0.18)
Proportion cephalosporins	13.56% (1.75%)			-0.197 (0.16)
Proportion macrolides	18.22% (1.21%)			

Table 2: Correlations (both linear (Pearson, with 95% confidence intervals) and Spearman (with p-values)) between state-specific proportions of prescriptions for each antibiotic class among all antibiotics prescribed in the state for different pairs of antibiotic classes, as well as the mean (standard deviation) for the state-specific percentages of all antibiotic prescriptions that were for each antibiotic class between 2014-2015.

Table 3 shows correlations (both Spearman and linear), for each antibiotic class and age group, between average annual state-specific proportions of a given antibiotic class among the overall outpatient antibiotic prescriptions and average annual state-specific rates of mortality with septicemia in a given age group per unit of prescribed antibiotics (Methods) between 2014-2015. The Spearman correlations are positive for penicillins for persons aged 75-84y and over 85y, and for fluoroquinolones for persons aged 50-64y; the Spearman correlations are negative for cephalosporins for persons aged 75-84y and over 85y, and for penicillins for persons aged 18-49y. Among those six significant Spearman correlations, all the corresponding linear correlations are also significant save for penicillins and mortality with septicemia in persons aged 75-84y.

		Proportion fluoroquinolones	Proportion penicillins	Proportion cephalosporins	Proportion macrolides
Age 18-49y	Spearman	0.219 (0.12)	**-0.36 (0.01)**	0.262 (0.064)	0.235 (0.097)
	Linear	0.142 (-0.14,0.4)	**-0.349 (-0.57,-0.08)**	0.168 (-0.11,0.42)	0.269 (-0.01,0.51)
Age 50-64y	Spearman	**0.419 (0.002)**	-0.233 (0.1)	-0.063 (0.66)	0.185 (0.19)
	Linear	**0.35 (0.08,0.57)**	-0.183 (-0.44,0.1)	-0.151 (-0.41,0.13)	0.205 (-0.07,0.46)
Age 65-74y	Spearman	0.274 (0.052)	0.003 (0.98)	-0.247 (0.08)	0.092 (0.52)
	Linear	0.258 (-0.02,0.5)	-0.011 (-0.29,0.27)	**-0.306 (-0.54,-0.03)**	0.109 (-0.17,0.37)
Age 75-84y	Spearman	0.081 (0.57)	**0.292 (0.038)**	**-0.418 (0.002)**	-0.004 (0.98)
	Linear	0.028 (-0.25,0.3)	0.229 (-0.05,0.47)	**-0.411 (-0.62,-0.15)**	0.028 (-0.25,0.3)
Age 85+y	Spearman	-0.023 (0.87)	**0.416 (0.003)**	**-0.422 (0.002)**	-0.048 (0.74)
	Linear	-0.04 (-0.31,0.24)	**0.354 (0.09,0.57)**	**-0.442 (-0.64,-0.19)**	-0.024 (-0.3,0.25)

Table 3: Correlations (both Spearman (p-value) and linear (Pearson, 95% CI)) between average annual state-specific percent of overall outpatient antibiotic prescribing that is for a given antibiotic

class and average annual state-specific rates of mortality with septicemia in a given age group per unit of prescribed antibiotics (Methods) between 2014-2015.

2. England

Table 4 shows annual rates of *E. coli* and MSSA bacteremia for the different CCGs (mean + standard error) for the 2014/15 through the 2017/18 financial years, as well as the correlation between those rates and the rates of antibiotic prescribing, both per 1,000 residents and per 1,000 STAR-PUs [40]. Table 4 suggests an ongoing increase in the rates of MSSA bacteremia [7], with the long-term growth in the rates of *E. coli* bacteremia [38,29,47,9] stalling in 2017/18. Table also 4 suggests that for each of the 4 years in the data, for both the *E. coli* and MSSA bacteremia rates, estimates of the correlation between those rates and antibiotic prescribing per 1,000 residents are higher than the estimates of the correlation between those rates and antibiotic prescribing per 1,000 STAR-PUs.

Outcome		2015/16	2016/17	2017/18	2017/18
E. coli bacteremia	Annual rate	67.72 (16.2)	71.76 (16.4)	75.81 (16.2)	76.15 (16.1)
	Correlation with antibiotic prescribing per 1,000 residents	**0.362 (0.23,0.48)**	**0.458 (0.34,0.56)**	**0.489 (0.37,0.59)**	**0.489 (0.37,0.59)**
	Correlation with antibiotic prescribing per 1,000 STAR-PUs	**0.311 (0.18,0.43)**	**0.417 (0.29,0.53)**	**0.443 (0.32,0.55)**	**0.435 (0.31,0.54)**
MSSA bacteremia	Annual rate	18.39 (5.3)	19.84 (6.0)	21.01 (5.8)	21.86 (5.9)
	Correlation with antibiotic prescribing per 1,000 residents	**0.246 (0.11,0.38)**	**0.459 (0.34,0.56)**	**0.449 (0.33,0.56)**	**0.389 (0.26,0.5)**
	Correlation with antibiotic prescribing per 1,000 STAR-PUs	**0.202 (0.06,0.34)**	**0.444 (0.32,0.55)**	**0.424 (0.3,0.53)**	**0.388 (0.26,0.5)**

Table 4: Annual rates of *E. coli* and MSSA bacteremia for the different CCGs (mean + standard error) for the 2014/15 through the 2017/18 financial years; correlations between those bacteremia rates and rates of overall antibiotic prescribing, both per 1,000 residents and per 1,000 STAR-PUs [40].

Table 5 shows annual CCG-specific rates of prescribing of different antibiotic types/classes per 1,000 residents (mean + standard error), as well as the annual CCG-specific proportions of those

antibiotic types/classes among all prescribed antibiotics (mean + standard error) for the 2014/15 through the 2017/18 financial years. Table 5 suggests significant temporal reduction in the rates/proportions of prescribing for trimethoprim, co-amoxiclav and cephalosporins/other beta lactams, as well as reduction in the rates/proportions of amoxicillin, fluroquinolone and macrolide prescribing. Prescribing of UTI antibiotics (BNF 5.1.13 -- nitrofurantoin/fosfomycin/methenamine) increased markedly (with a good amount of replacement of trimethoprim by nitrofurantoin in the treatment of UTIs taking place during the study period, [29]), with proportions of penicillins other than amoxicillin/co-amoxiclav and tetracyclines among all prescribed antibiotics also increasing.

	2014/15		2015/16		2016/17		2017/18	
	Percent	Rate	Percent	Rate	Percent	Rate	Percent	Rate
Amoxicillin	28.35% (3.8%)	187.6 (36.3)	26.8% (3.7%)	162.9 (32.5)	26.75% (3.5%)	161.1 (31.4)	26.04% (3.4%)	150 (29.7)
Co-amoxiclav	5.44% (2.2%)	35.4 (14.6)	4.8% (1.9%)	28.8 (11.4)	4.28% (1.5%)	25.4 (9.4)	4.16% (1.4%)	23.6 (8.4)
Penicillins except amoxicillin/co-amoxiclav	17.67% (1.5%)	116.7 (18.1)	18.41% (1.6%)	111.8 (17.9)	18.6% (1.7%)	111.8 (18.4)	19.18% (1.8%)	110.4 (18.9)
Tetracylines	11.59% (2.9%)	77.5 (23.7)	12.24% (2.9%)	75.4 (23)	12.74% (3%)	77.9 (24.1)	13.12% (3%)	77 (24)
Macrolides	12.66% (1.6%)	83.9 (16.6)	12.3% (1.5%)	75.1 (15.3)	12.19% (1.5%)	73.7 (15.4)	11.94% (1.4%)	69.1 (14.8)
Cephalosporins + other beta-lactams	3.3% (1.4%)	21.9 (10.3)	2.96% (1.2%)	18.1 (8.1)	2.7% (1.1%)	16.4 (7.5)	2.6% (1.1%)	15.2 (7.2)
Fluoroquinolones	1.96% (0.6%)	12.8 (3.8)	1.9% (0.5%)	11.4 (3.3)	1.87% (0.5%)	11.1 (3.1)	1.89% (0.5%)	10.8 (3)
Trimethoprim	10.33% (1.7%)	68.9 (16.6)	10.48% (1.9%)	64.5 (17)	9.8% (2.1%)	59.9 (17.7)	7.3% (1.8%)	42.8 (14.3)
UTI antibiotics	5.77% (1.4%)	38 (9.7)	7.01% (1.7%)	42.3 (10.3)	8.05% (2%)	48.1 (12.1)	10.67% (2.1%)	61.1 (13.9)

Table 5: Annual CCG-specific proportions of a given antibiotic type/class among all antibiotic prescriptions (mean + standard error), and annual CCG-specific rates of prescribing per 1,000 individuals (mean + standard error) for 9 antibiotic types/classes (Methods) during the 2014/15 through the 2017/18 financial years in England.

Tables 6 and 7 show correlations (both linear and Spearman) between CCG-specific proportions of different antibiotic types/classes among all prescribed antibiotics and rates of *E. coli* (Table 6) and MSSA (Table 7) bacteremia (per unit of antibiotic prescribing, Methods) for the 2014/15 through 2017/18 financial years. For penicillins other than amoxicillin/co-amoxiclav, correlations with rates of MSSA bacteremia were positive for all years, and correlations with rates of *E. coli* bacteremia were

positive for the 2014/15 through 2016/17 financial years. For macrolides and fluoroquinolones, the corresponding correlations were generally negative. Correlations with bacteremia rates increased with time for proportions of UTI antibiotics, cephalosporins, and, to a smaller extent, amoxicillin; the corresponding correlations declined for proportions of trimethoprim and co-amoxiclav prescribing, with all those relative changes presumably related more to changes in prescribing patterns rather than changes in the causal relation between the use of those antibiotics and bacteremia outcomes. In particular, Tables 6 and 7 suggest that relative reductions in trimethoprim and co-amoxiclav prescribing were greater in places with higher bacteremia rates compared to places with lower bacteremia rates. Finally, we note that positive correlations with bacteremia rates for proportions of prescribing for penicillins other than amoxicillin/co-amoxiclav, but not for proportions of co-amoxiclav or amoxicillin prescribing need not suggest that penicillins other than amoxicillin/co-amoxiclav have a stronger relative impact on bacteremia rates than co-amoxiclav or amoxicillin; those differences may also have to do with geographic/demographic variation in the choice of different antibiotics.

	2014/15		2015/16		2016/17		2017/18	
	Linear	Spearman	Linear	Spearman	Linear	Spearman	Linear	Spearman
Amoxicillin	-0.058 (-0.2,0.09)	-0.048 (0.5)	0.017 (-0.13,0.16)	0.036 (0.62)	0.01 (-0.13,0.16)	0.017 (0.82)	0.02 (-0.12,0.16)	-0.007 (0.93)
Co-amoxiclav	-0.056 (-0.2,0.09)	-0.056 (0.44)	-0.118 (-0.26,0.03)	**-0.146 (0.044)**	-0.14 (-0.28,0)	**-0.169 (0.02)**	-0.14 (-0.28,0)	**-0.175 (0.016)**
Penicillins except amoxicillin/ co-amoxiclav	**0.256 (0.12,0.38)**	**0.201 (0.006)**	**0.204 (0.06,0.34)**	**0.167 (0.02)**	**0.15 (0,0.28)**	0.117 (0.11)	0.09 (-0.05,0.23)	0.06 (0.41)
Tetracylines	**0.171 (0.03,0.31)**	**0.201 (0.006)**	0.095 (-0.05,0.23)	0.115 (0.12)	0.09 (-0.06,0.23)	0.104 (0.15)	-0.03 (-0.17,0.11)	0.02 (0.78)
Macrolides	**-0.208 (-0.34,-0.07)**	**-0.248 (0.0006)**	-0.14 (-0.28,0)	**-0.156 (0.03)**	-0.13 (-0.27,0.01)	**-0.153 (0.035)**	-0.07 (-0.21,0.08)	-0.091 (0.22)
Cephalosporins + other beta-lactams	-0.029 (-0.17,0.11)	0.023 (0.75)	0.03 (-0.11,0.17)	0.115 (0.11)	0.02 (-0.12,0.16)	0.039 (0.59)	0.12 (-0.02,0.26)	**0.15 (0.039)**
Fluoroquinolones	**-0.276 (-0.4,-0.14)**	**-0.306 (0.00002)**	**-0.207 (-0.34,-0.07)**	**-0.242 (0.0008)**	**-0.18 (-0.32,-0.04)**	**-0.203 (0.005)**	-0.12 (-0.26,0.02)	**-0.162 (0.026)**
Trimethoprim	-0.089 (-0.23,0.05)	-0.076 (0.30)	**-0.279 (-0.41,-0.14)**	**-0.26 (0.0003)**	**-0.28 (-0.41,-0.15)**	**-0.235 (0.001)**	**-0.26 (-0.39,-0.12)**	**-0.273 (0.0002)**
UTI antibiotics	0.038 (-0.1,0.18)	-0.065 (0.38)	**0.153 (0.01,0.29)**	0.044 (0.55)	**0.21 (0.07,0.34)**	0.135 (0.063)	**0.19 (0.05,0.33)**	**0.164 (0.024)**

Table 6: Correlations (both linear, with 95% CI, and Spearman, with p-value) between annual proportions of different antibiotic types/classes among all prescribed antibiotics and annual rates of *E. coli* bacteremia (per unit of antibiotic prescribing, Methods) for different CCGs in England, 2014/15 through 2017/18 financial years.

	2014/15		2015/16		2016/17		2017/18	
	Linear	Spearman	Linear	Spearman	Linear	Spearman	Linear	Spearman
Amoxicillin	-0.071 (-0.21,0.07)	-0.121 (0.10)	-0.047 (-0.19,0.1)	-0.076 (0.30)	-0.04 (-0.18,0.1)	-0.081 (0.27)	0.03 (-0.11,0.17)	-0.009 (0.90)
Co-amoxiclav	-0.114 (-0.25,0.03)	-0.112 (0.13)	**-0.177 (-0.31,-0.04)**	**-0.193 (0.008)**	**-0.26 (-0.39,-0.12)**	**-0.313 (0.00001)**	**-0.29 (-0.42,-0.16)**	**-0.282 (0.00009)**
Penicillins except amoxicillin/ co-amoxiclav	**0.259 (0.12,0.39)**	**0.261 (0.0003)**	**0.218 (0.08,0.35)**	**0.172 (0.018)**	**0.2 (0.06,0.33)**	**0.196 (0.007)**	**0.26 (0.12,0.39)**	**0.232 (0.001)**
Tetracylines	0.135 (-0.01,0.27)	**0.157 (0.032)**	**0.223 (0.08,0.35)**	**0.253 (0.0005)**	0.14 (0,0.28)	**0.234 (0.001)**	0 (-0.14,0.15)	0.057 (0.43)
Macrolides	**-0.155 (-0.29,-0.01)**	**-0.183 (0.012)**	**-0.183 (-0.32,-0.04)**	**-0.217 (0.003)**	**-0.17 (-0.31,-0.03)**	**-0.206 (0.005)**	-0.13 (-0.27,0.01)	**-0.151 (0.038)**
Cephalosporins + other beta-lactams	-0.044 (-0.19,0.1)	-0.01 (0.89)	-0.046 (-0.19,0.1)	-0.024 (0.74)	0 (-0.14,0.15)	0.028 (0.71)	0.04 (-0.1,0.18)	0.075 (0.31)
Fluoroquinolones	-0.1 (-0.24,0.04)	-0.084 (0.25)	**-0.206 (-0.34,-0.07)**	**-0.232 (0.001)**	-0.12 (-0.26,0.02)	**-0.18 (0.01)**	-0.12 (-0.26,0.03)	**-0.148 (0.04)**
Trimethoprim	0.005 (-0.14,0.15)	0.017 (0.82)	**-0.166 (-0.3,-0.02)**	-0.132 (0.07)	**-0.16 (-0.3,-0.02)**	-0.101 (0.17)	**-0.21 (-0.34,-0.07)**	**-0.214 (0.003)**
UTI antibiotics	0.001 (-0.14,0.14)	-0.029 (0.69)	0.09 (-0.05,0.23)	0.022 (0.77)	**0.16 (0.02,0.3)**	0.069 (0.35)	0.12 (-0.02,0.26)	0.111 (0.13)

Table 7: Correlations (both linear, with 95% CI, and Spearman, with p-value) between annual proportions of different antibiotic types/classes among all prescribed antibiotics and annual rates of MSSA bacteremia (per unit of antibiotic prescribing, Methods) for different CCGs in England, 2014/15 through 2017/18 financial years.

Discussion

Rates of mortality related to septicemia/sepsis in the US, as well as rates of *E. coli* bacteremia and associated mortality in England are high [1-3,5,7,8,29] and antimicrobial use may affect those rates through a variety of mechanisms (Introduction). At the same time, our understanding of the effect of the use of certain antibiotics vs. others for various indications on the rates of bacteremia, septicemia and associated mortality is still limited. Additionally, use of certain antibiotics may affect the rates of severe outcomes associated with syndromes for which a given antibiotic is rarely prescribed as use of antibiotics may affect prevalence of infection/colonization and resistance to different antibiotics in different bacterial pathogens that subsequently cause various syndromes (Introduction). In this paper, we relate the proportions of different antibiotic types/classes among the overall volume of outpatient antibiotic prescription in different US states and English CCGs [35,37] to rates of mortality with septicemia in different age groups of US adults [36] and rates of E. coli and MSSA

bacteremia in England [38]. Our results suggest, among other things, that prescribing of penicillins is associated with rates of *E. coli* and MSSA bacteremia in England, and rates of mortality with septicemia in older US adults, with the latter finding supporting our earlier results on the association between the use of penicillins and rates of septicemia hospitalization and mortality in older US adults [13,14]. Additionally, our analyses complement the evidence from [14] about the relation between fluoroquinolone use and rates of mortality with septicemia in non-elderly US adults. More granular analyses are needed to better understand the effect of prescribing of different antibiotics, including the effect of antibiotic, particularly penicillin replacement, as well as of reduction in antibiotic prescribing on the rates of bacteremia, septicemia and associated mortality, as explained further in the next two paragraphs.

Our findings about the positive associations between the use of penicillins and mortality with septicemia in older US adults is in agreement with the fact that prevalence of resistance to penicillins, particularly in older adults, is high for a variety of infections with both Gram-negative and Gram-positive bacteria [31-33]. Negative associations between the proportion of cephalosporins among all antibiotic prescriptions and rates of septicemia mortality in older US adults (Table 3) may be related to the competition in prescribing with other antibiotic classes, particularly penicillins and fluoroquinolones for which prevalence of resistance in the key syndromes leading to sepsis is higher than prevalence of resistance to cephalosporins, e.g. [31,42]. At the same time, prevalence of cephalosporin resistance and the frequency of extended-spectrum beta-lactamase (ESBL) production, including in Gram-negative bacteria is growing [43], and replacement of other antibiotics by cephalosporins might potentially lead to negative effects in the long run. Finally, we found no associations between the proportion of macrolides among all antibiotic prescriptions in the US and rates of mortality with septicemia in adults, which agrees with our earlier findings [13,14]. While macrolides are used relatively infrequently in the treatment of urinary tract and gastrointestinal infections, macrolides are commonly prescribed in the treatment of other sources of sepsis, particularly respiratory illness, both chronic [44] and acute, including pneumonia [45], with high prevalence of macrolide resistance in the corresponding infections [46].

In England, prevalence of resistance to trimethoprim in urinary tract infections (UTIs) is high [47], and trimethoprim use was found to be associated with UTI-related *E. coli* bacteremia [21]. Major reductions in trimethoprim prescribing in England took place in the recent years ([29]; Table 5 in this paper); moreover, prescribing of trimethoprim declined disproportionately in places in England with higher rates of *E. coli* and MSSA bacteremia (Results). All those changes might have played a

role in the fact that growth in the rates of *E. coli* bacteremia in England has stalled in 2017/2018 after many years of robust increases (Table 4; [38,29,47,9]). Prevalence of co-amoxiclav resistance in *E. coli* bacteremia in England is very high [29], and outpatient prescribing of co-amoxiclav was reduced significantly during the recent years ([29]; Table 5 in this paper), disproportionately in places with higher rates of bacteremia (Results). At the same time, prevalence of co-amoxiclav resistance in *E. coli* bacteremia and rates of the corresponding bacteremia outcomes are affected not only by GP prescribing of co-amoxiclav but also by other factors including clinic use of co-amoxiclav (inpatient/outpatient), and possibly the use of related penicillins. We note that amoxicillin use is also associated with trimethoprim resistance [20], which in turn affects the rates of *E. coli* bacteremia [21], while use of different penicillins may also affect prevalence of resistance to piperacillin/tazobactam in *E. coli* bacteremia [29]. Additionally, penicillins are widely prescribed in England (e.g. Table 5; [29]), and penicillin use/resistance to penicillins is therefor expected to affect prevalence of infection/colonization with different bacterial pathogens that subsequently lead to bacteremia outcomes. While positive correlations between rates of bacteremia and rates of prescribing for penicillins other than amoxicillin/co-amoxiclav, but not for amoxicillin or co-amoxiclav were found in this paper, it is uncertain which penicillins (including co-amoxiclav and amoxicillin) have a greater relative impact on rates of bacteremia, with the results of the correlation analyses in this paper potentially affected by patterns of prescribing of different antibiotics related to different geographic locations and demographic factors. Overall, our results support some replacement of penicillins in England by antibiotics for which prevalence of antimicrobial resistance is lower, as well as reduction in penicillin prescribing with the aim of reducing bacteremia rates. Finally, we note that additional factors besides antibiotic choice affect the rates of severe infections associated with bacterial pathogens. Those factors include prevalence of underlying health conditions and certain demographic factors [34]. Additionally, we have found that prescribing of metronidazole for skin infections (BNF 1310012K0) as a proportion of the overall antibiotic prescribing is correlated with the CCG-specific rates of both *E. coli* and MSSA bacteremia per unit of antibiotic prescribing for the four years in the data. This suggests the possibility that demographic/geographic differences result in differences in transmission of infections/colonization with bacterial pathogens (including transmission through skin infections), which in turn affect the rates of severe bacterial infections, including bacteremia. Further work is needed to better understand those differences in transmission, including the feasibility of mitigation efforts aimed at preventing infections.

Our paper has some limitations. Correlations between proportions of different antibiotic types/classes (particularly penicillins) and rates of severe outcomes associated with bacterial infections may be affected not only by the relative contributions of the use of a unit of different antibiotics to the rates of those severe outcomes but also by patterns of antibiotic prescribing in different locations. We note that penicillins are prescribed for a wide variety of indications, both in England and the US, affecting prevalence of infection/colonization with different bacterial pathogens, and that there is high prevalence of resistance to penicillins in both the Gram-negative and Gram-positive bacteria in the US (e.g. [31-33]), and high prevalence of co-amoxiclav resistance in *E. coli* bacteremia in England [29]. Coding practices for septicemia on death certificates may vary by US state [48]. Additionally, data on outpatient antibiotic prescribing in the whole population [35] were related to age-specific rates of mortality with septicemia in the US [36], while in England, no age-specific prescribing or bacteremia data were available for this study. We expect that those sources of noise/incompatibility should generally reduce precision and bias the correlations towards null rather than create spurious associations.

We believe that despite those limitations, our results suggest that prescribing of certain antibiotics, particularly penicillins is associated with rates of *E. coli* and MSSA bacteremia in England and rates of mortality with septicemia in older US adults, with the latter result supporting our earlier findings about the association between the rates of prescribing of penicillins and rates of hospitalization and mortality with septicemia in older US adults [13,14]. Further studies are needed to better understand the effect of replacement of certain antibiotics, particularly penicillins by others, well as of reduction in antibiotic prescribing in the treatment of certain syndromes on the rates of bacteremia, septicemia/sepsis and associated mortality.

Acknowledgment: We thank Koen Pouwels for helpful discussions.

References

[1] Elixhauser A, Friedman B, Stranges E. Septicemia in U.S. Hospitals, 2009: Statistical Brief #122. HealthCare Cost and Utilization Project (HCUP), Agency for Healthcare Research and Quality (AHRQ), 2011. Accessed on Jan. 12, 2019. Available from: https://www.hcup-us.ahrq.gov/reports/statbriefs/sb122.pdf

[2] McDermott KW, Elixhauser A, Sun R. Trends in Hospital Inpatient Stays in the United States, 2005–2014. Statistical Brief #225. HealthCare Cost and Utilization Project (HCUP), Agency for Healthcare Research and Quality (AHRQ), 2017. Accessed on Jan. 12, 2019. Available from: https://www.hcup-us.ahrq.gov/reports/statbriefs/sb225-Inpatient-US-Stays-Trends.pdf

[3] Iwashyna TJ, Cooke CR, Wunsch H, Kahn JM. Population burden of long-term survivorship after severe sepsis in older Americans. J Am Geriatr Soc. 2012;60(6):1070-7.

[4] Torio CM, Moore BJ. National Inpatient Hospital Costs: The Most Expensive Conditions by Payer, 2013: Statistical Brief #204. HealthCare Cost and Utilization Project (HCUP), Agency for Healthcare Research and Quality (AHRQ), 2016. Accessed on Jan. 12, 2019. Available from: https://www.hcup-us.ahrq.gov/reports/statbriefs/sb204-Most-Expensive-Hospital-Conditions.jsp

[5] US CDC. Sepsis. Data & Reports. (2018). Accessed on Jan. 12, 2019. Available from: https://www.cdc.gov/sepsis/datareports/index.html

[6] Iwashyna TJ, Cooke CR, Wunsch H, Kahn JM. Population burden of long-term survivorship after severe sepsis in older Americans. J Am Geriatr Soc. 2012;60(6):1070-7.

[7] Public Health England. Mandatory MRSA, MSSA and *E. coli* bacteraemia and *C. difficile* infection data 2015/16. Annual Epidemiological Commentary. 2016. Accessed on Jan. 12, 2019. Available from: https://www.researchgate.net/profile/Simon_Thelwall/publication/305392637_Annual_Epidemiological_Commentary_Mandatory_MRSA_MSSA_and_E_coli_bacteraemia_and_C_difficile_infection_data_201516/links/578ccb6a08ae254b1de8610a/Annual-Epidemiological-Commentary-Mandatory-MRSA-MSSA-and-E-coli-bacteraemia-and-C-difficile-infection-data-2015-16.pdf

[8] Thirty-day all-cause fatality subsequent to MRSA, MSSA and E. coli bacteraemia and C. difficile infection, 2017/18. Oct. 2018. Accessed on Jan. 12, 2019. Available from https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/765014/HCAI_fatality_report_201718_v2.pdf

[9] Livermore DM, Hope R, Reynolds R, Blackburn R, Johnson AP, Woodford N. Declining cephalosporin and fluoroquinolone non-susceptibility among bloodstream Enterobacteriaceae from the UK: links to prescribing change? J Antimicrob Chemother . 2013;68:2667-2674.

[10] Rhee C, Murphy MV, Li L, Platt R, Klompas M. Comparison of Trends in Sepsis Incidence and Coding Using Administrative Claims Versus Objective Clinical Data. Clin Infect Dis. 2015;60(1):88-95

[11] Umscheid CA, Betesh J, VanZandbergen C, Hanish A, Tait G, Mikkelsen ME, et al. Development, implementation, and impact of an automated early warning and response system for sepsis. J Hosp Med. 2015;10(1):26-31

[12] Walkey AJ, Lagu T, Lindenauer PK. Trends in sepsis and infection sources in the United States. A population-based study. Ann Am Thorac Soc. 2015;12(2):216-20

[13] Goldstein E, Olesen SW, Karaca Z, Steiner CA, Viboud C, Lipsitch M. Levels of outpatient prescribing for four major antibiotic classes and rates of septicemia hospitalization in different US states. BioRxiv 2018. Accessed on Jan. 12, 2019. Available from: https://www.biorxiv.org/content/early/2018/12/13/404046

[14] Goldstein E, Lipsitch M. Outpatient antibiotic prescribing and demographic factors associated with state-level septicemia mortality rates in US adults. BioRxiv 2018. Accessed on Jan. 12, 2019. Available from: https://www.biorxiv.org/content/early/2018/11/30/480137

[15] Goldstein E, MacFadden DR, Karaca Z, Steiner CA, Viboud C, Lipsitch M. Antimicrobial resistance prevalence and rates of hospitalization with septicemia in the diagnosis in adults in different US states. BioRxiv 2018. Accessed on Jan. 12, 2019. Available from: https://www.biorxiv.org/content/early/2018/08/29/404137

[16] Paul M, Shani V, Muchtar E, Kariv G, Robenshtok E, Leibovici L. Systematic review and meta-analysis of the efficacy of appropriate empiric antibiotic therapy for sepsis. Antimicrob Agents Chemother. 2010;54(11):4851-63

[17] Zilberberg MD, Shorr AF, Micek ST, Vazquez-Guillamet C, Kollef MH. Multi-drug resistance, inappropriate initial antibiotic therapy and mortality in Gram-negative severe sepsis and septic shock: a retrospective cohort study. Crit Care. 2014;18(6):596

[18] Lee YC, Hsiao CY, Hung MC, Hung SC, Wang HP, Huang YJ, et al. Bacteremic Urinary Tract Infection Caused by Multidrug-Resistant Enterobacteriaceae Are Associated With Severe Sepsis at Admission: Implication for Empirical Therapy. Medicine (Baltimore). 2016;95(20):e3694

[19] Hawkey PM, Warren RE, Livermore DM, McNulty CAM, Enoch DA, Otter JA, et al. Treatment of infections caused by multidrug-resistant Gram-negative bacteria: report of the British Society for Antimicrobial Chemotherapy/Healthcare Infection Society/British Infection Association Joint Working Party. J Antimicrob Chemother. 2018;73(suppl_3):iii2-iii78

[20] Pouwels KB, Freeman R, Muller-Pebody B, Rooney G, Henderson KL, Robotham JV, et al. Association between use of different antibiotics and trimethoprim resistance: going beyond the obvious crude association. J Antimicrob Chemother. 2018;73(6):1700-1707

[21] Lishman H, Costelloe C, Hopkins S, Johnson AP, Hope R, Guy R, et al. Exploring the relationship between primary care antibiotic prescribing for urinary tract infections, Escherichia coli bacteraemia incidence and antibiotic resistance: an ecological study. Int J Antimicrob Agents. 2018 Aug 23. doi: 10.1016/j.ijantimicag.2018.08.013

[22] Costelloe C , Metcalf C , Lovering A et al. Effects of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. BMJ 2010;340:c2096

[23] Bergman M, Huikko S, Huovinen P, Paakkari P, Seppälä H; Finnish Study Group for Antimicrobial Resistance (FiRe Network). Macrolide and azithromycin use are linked to increased macrolide resistance in Streptococcus pneumoniae. Antimicrob Agents Chemother. 2006;50(11):3646-50

[24] Chang HH, Cohen T, Grad YH, Hanage WP, O'Brien TF, Lipsitch M. Origin and proliferation of multiple-drug resistance in bacterial pathogens. Microbiol Mol Biol Rev 2015;79(1):101–16.

[25] Lehtinen S, Blanquart F, Lipsitch M, Fraser C, The Maela Pneumococcal Collaboration. On the evolutionary ecology of multidrug resistance in bacteria. BioRxiv 2019. Accessed on Jan. 19, 2019. Available from: https://www.biorxiv.org/content/early/2019/01/10/233957

[26] Tacconelli E, De Angelis G, Cataldo MA, Pozzi E, Cauda R. Does antibiotic exposure increase the risk of methicillin-resistant Staphylococcus aureus (MRSA) isolation? A systematic review and meta-analysis. J Antimicrob Chemother. 2008;61(1):26-38

[27] Weber SG, Gold HS, Hooper DC, Karchmer A, Carmeli Y. Fluoroquinolones and the risk for methicillin-resistant Staphylococcus aureus in hospitalized patients. Emerg Infect Dis. 2003;9:1415 - 22

[28] Knight GM, Budd EL, Whitney L, Thornley A, Al-Ghusein H, Planche T, et al. Shift in dominant hospital-associated methicillin-resistant Staphylococcus aureus (HA-MRSA) clones over time. J Antimicrob Chemother. 2012;67(10):2514-22.

[29] Public Health England. English Surveillance Programme for Antimicrobial Utilisation and Resistance (ESPAUR). Report 2018. Accessed on Jan. 12, 2019. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/759975/ESPAUR_2018_report.pdf

[30] Vihta KD, Stoesser N, Llewelyn M, Quan TP, Davies T, Fawcett NJ, et al. Trends over time in Escherichia coli bloodstream infections, urinary tract infections, and antibiotic susceptibilities in Oxfordshire, UK, 1998–2016: a study of electronic health records. Lancet Infect Dis. 2018;18(10):1138-1149

[31] Morrill HJ, Morton JB, Caffrey AR, Jiang L, Dosa D, Mermel LA, et al. Antimicrobial Resistance of Escherichia coli Urinary Isolates in the Veterans Affairs Health Care System. Antimicrob Agents Chemother. 2017;61(5) pii: e02236-16

[32] Barnato AE, Alexander SL, Linde-Zwirble WT, Angus DC. Racial variation in the incidence, care, and outcomes or severe sepsis: analysis of population, patient, and hospital characteristics. Am J Respir Crit Care Med. 2008;177:279–284

[33] Cheng MP, René P, Cheng AP, Lee TC. Back to the Future: Penicillin-Susceptible Staphylococcus aureus. Am J Med. 2016;129(12):1331-1333

[34] Hope EC, Crump RE, Hollingsworth TD, Smieszek T, Robotham JV, Pouwels KB. Identifying English Practices that Are High Antibiotic Prescribers Accounting for Comorbidities and Other Legitimate Medical Reasons for Variation. EClinicalMEdicine 2018. Online first, accessed on Jan. 19, 2019, available from: https://www.thelancet.com/pdfs/journals/eclinm/PIIS2589-5370(18)30053-1.pdf

[35] US CDC. Antibiotic Resistance Patient Safety Atlas. Outpatient Antibiotic Prescription Data. Accessed on Jan. 12, 2019. Available from: https://gis.cdc.gov/grasp/PSA/indexAU.html

[36] US CDC Wonder. Multiple Cause of Death, 1999-2016 Request. Accessed on Jan. 5, 2019. Available from: https://wonder.cdc.gov/mcd-icd10.html

[37] EBM DataLab, University of Oxford. OpenPrescribing data. Accessed on Jan. 12, 2019. Available from: https://openprescribing.net/analyse/

[38] Public Health England. AMR Local Indicators. Accessed on Jan. 12, 2019. Available from: https://fingertips.phe.org.uk/profile/amr-local-indicators/

[39] Epstein L, Dantes R, Magill S, Fiore A. Varying Estimates of Sepsis Mortality Using Death Certificates and Administrative Codes — United States, 1999–2014. MMWR Morb Mortal Wkly Rep 2016;65:342–345.

[40] National Health Service (NHS). STAR-PU weightings. Accessed on Jan. 12, 2019. Available from: https://www.nhs.uk/Scorecard/Pages/IndicatorFacts.aspx?MetricId=443

[41] King A, Mullish BH, Williams HRT, Aylin P. Comparative epidemiology of Clostridium difficile infection: England and the USA. Int J Qual Health Care. 2017;29(6):785-791

[42] Bidell MR, Palchak M, Mohr J, Lodise TP. Fluoroquinolone and Third-Generation-Cephalosporin Resistance among Hospitalized Patients with Urinary Tract Infections Due to Escherichia coli: Do

Rates Vary by Hospital Characteristics and Geographic Region? Antimicrob Agents Chemother. 2016;60(5):3170-3

[43] Park, SH. Third-generation cephalosporin resistance in gram-negative bacteria in the community: a growing public health concern. Korean J Intern Med. 2014;29(1):27-30

[44] Suresh Babu K, Kastelik J, Morjaria JB. Role of long term antibiotics in chronic respiratory diseases. Respir Med. 2013;107(6):800-15.

[45] Mandell LA, Wunderink RG, Anzueto A, Bartlett JG, Campbell GD, Dean NC, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. Clin Infect Dis. 2007;44 Suppl 2:S27-72

[46] Niederman MS. Macrolide-Resistant Pneumococcus in Community-acquired Pneumonia. Is There Still a Role for Macrolide Therapy? Am J Respir Crit Care Med. 2015;191(11):1216-7

[47] Public Health England. English Surveillance Programme for Antimicrobial Utilisation and Resistance (ESPAUR). Report 2017. Accessed on Jan. 12, 2019. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/656611/ESPAUR_report_2017.pdf

[48] Ong P, Gambatese M, Begier E, Zimmerman R, Soto A, Madsen A. Effect of cause-of-death training on agreement between hospital discharge diagnoses and cause of death reported, inpatient hospital deaths, New York City, 2008-2010. Prev Chronic Dis. 2015 Jan 15;12:E04