**Title: Parasite dispersal influences introgression rate: the case of dove wing**

**and body lice**

**Authors:** Jorge Doña[1]*, Andrew D. Sweet[1, 2], and Kevin P. Johnson[1]*

**Affiliations:**

[1]Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-
Champaign, 1816 S. Oak St., Champaign, Illinois 61820, USA.

[2]Department of Entomology, Purdue University, 901 W. State St., West
Lafayette, Indiana 47907, USA.

*Correspondence to: jorged@illinois.edu, kpjohnso@illinois.edu.

**Running title:** Dispersal influences introgression rate.

**Author contributions:** J.D., and K.P.J. conceived the study. J.D., A.D.S., and K.P.J. designed
the study. A.D.S. collected the data. J.D. and A.D.S. analysed the data. K.P.J. obtained financial
support for the project. J.D. wrote the manuscript and all authors contributed to editing the
manuscript.

**Acknowledgments:** This study was supported by the US National Science Foundation (DEB-
1239788 and DEB-1342604 to K.P.J). We declare that we have no conflict of interest.

**Data availability:** All data needed to evaluate the conclusions in the paper are present in the
paper and/or the Supplementary Materials. Additional data related to this paper may be requested
from the authors.

**Supplementary Materials:** Supplementary data are available at Figshare (DOI:
10.6084/m9.figshare.9176204).
22

**Abstract:** Organisms vary in their dispersal abilities, and these differences can have important

biological consequences, such as impacting the likelihood of hybridization events. However, the

factors shaping the frequency of hybridization are still poorly understood, and therefore how

dispersal ability affects the opportunities for hybridization is still unknown. Here, using the

ecological replicate system of dove wing and body lice (Insecta: Phthiraptera), we show that

species with higher dispersal abilities exhibited increased genomic signatures of introgression.

Specifically, we found a higher proportion of introgressed genomic reads and more reticulated

phylogenetic networks in wing lice, the louse group with higher dispersal abilities. Our results

illustrate how differences in dispersal ability can drive differences in the extent of introgression

through hybridization. The results from this study represent an important step for understanding

the factors driving hybridization. We expect our approach will stimulate future studies on the

ecological factors shaping hybridization to further understand this important process.

**Introduction**

Dispersal is the permanent movement of organisms away from their place of origin. It is a fundamental process in biology with significant implications at multiple scales of organization (Barton 1992; Clobert et al. 2001; Nathan 2001; Matthysen 2012), including the reproduction of individuals, the composition of populations and communities, and the geographical distribution of species (Clobert et al. 2001, 2012).

Organisms differ in their dispersal abilities, and these differences have an impact on their biology, such as on the distributional range of a species or gene flow between populations (Bohonak 1999). For example, organisms with lower dispersal abilities tend to have smaller distributional ranges and populations that are genetically more structured (Bohonak 1999; Dawson et al. 2014; DiBlasi et al. 2018).

Dispersal ability might also affect the opportunities for hybridization between species because the rates at which individuals encounter different species are likely to be higher in organisms with higher dispersal capabilities. Indeed, recent evidence supports this prediction by demonstrating that range expansion is associated with the extent of introgression (Currat et al. 2008; Nussberger et al. 2018). Similarly, dispersal differences explain more than 30% of the variation in the width of hybrid zones across animals (McEntee et al. 2018). However, overall the factors influencing hybridization events are poorly known (Randler 2006; Arnold 2015; Taylor and Larson 2019), and, in particular, the influence of dispersal ability on the rate of hybridization remains understudied.

Comparisons of the effect of dispersal on hybridization should ideally hold constant most factors other than dispersal. The ecological replicate system of wing and body lice (Insecta: Phthiraptera) of pigeons and doves (Aves: Columbidae) has proven to be an ideal system for

comparing the impact of dispersal differences on other aspects of biology, such as population

structure and codivergence (Clayton and Johnson 2003; Johnson and Clayton 2004; Clayton et

al. 2015; DiBlasi et al. 2018; Sweet and Johnson 2018). Specifically, this is an excellent system

in which to assess the effect of differences in dispersal capabilities on levels of introgression

because both of these two lineages of feather lice: 1) co-occur across the diversity of pigeons and

doves, 2) present highly comparable temporal patterns of diversification; specifically,

cophylogenetic analyses and bird time-calibrated trees indicate that both lineages originated on

the common ancestor of *Metropelia* doves (11.3-14.9 mya) and also share a cospeciation event

which occurred within the *Metriopelia* genus (5.2-7.4 mya; Sweet and Johnson 2015, 2018), 3)

have the same basic life history and diet (Clayton et al. 2015; Sweet and Johnson 2015, 2018),

but 4) they significantly differ in their dispersal ability (Harbison et al. 2008, 2009; Bartlow et al.

2016). Both wing and body lice disperse vertically between parents and offspring in the nest.

However, wing lice can also attach to and hitchhike on hippoboscid flies to disperse

"phoretically" between host individuals or host species (Harbison et al. 2008, 2009; Bartlow et

al. 2016). Indeed, this hitch-hiking dispersal mechanism profoundly influences their degree of

population structure and cophylogenetic history (Clayton and Johnson 2003; Sweet et al. 2017b;

DiBlasi et al. 2018; Sweet and Johnson 2018). In addition, wing lice have a higher rate of host-

switching (Clayton and Johnson 2003; Clayton et al. 2015; Sweet et al. 2017b) (i.e., successful

colonization of new host species) and of straggling (Whiteman et al. 2004) (i.e., dispersal to new

host species without reproduction on that new host).

To compare differences in the extent of introgression between wing and body lice, we

used whole-genome data from 71 louse individuals belonging to five taxa of wing lice

(*Columbicola*) and seven taxa of body lice (*Physconelloides*) occurring across the same host

species and have highly comparable patterns of diversification (Sweet and Johnson 2015, 2018). We predicted that wing lice, which have higher dispersal abilities and thus higher odds of encountering individuals of a different louse species on the same host, should show more extensive evidence of introgression (Fig. 1).

**Figure 1**. Diagram depicting the ecological replicate system and the hypothesis of this study. Wing lice (*Columbicola*) have higher dispersal abilities than body lice (*Physconelloides*), and thus higher odds of encountering individuals of a different louse sp.

**Materials and Methods**

<u>Data</u>

We studied whole genome data from 71 louse individuals belonging to five and seven taxa of *Columbicola* and *Physconelloides*, respectively (Supplementary Table S2). Data were available from previous studies (Boyd BM, et al. 2017; Sweet et al. 2017a; Sweet and Johnson 2018) and represent all described New World ground-dove wing and body louse species, most host species in this group, including samples across multiple biogeographic areas within species (Sweet and Johnson 2018; Supplementary Table S2). Illumina genome sequence data pre-processing

included several steps (Sweet and Johnson 2018). First, we discarded duplicate read pairs using

the fastqSplitDups script (https://github.com/McIntyre-Lab/mcscriptand

https://github.com/McIntyre-Lab/mclib). We then eliminated the Illumina sequencing adapters

with Fastx_clipper v0.014 from the FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit).

Also, we removed the first 5 nt from the 5' ends of reads using Fastx_trimmer v0.014 and

trimmed bases from the 3' ends of reads until reaching a base with a phred score ≥28 using

Fastq_quality_trimmer v0.014. Finally, we removed any reads less than 75 nt and analyzed the

cleaned libraries with Fastqc v0.11.5 to check for additional errors. We assembled nuclear loci in

aTRAM following previous studies (Allen et al. 2015; Boyd BM, et al. 2017; Sweet and Johnson

2018). In particular, we mapped modest coverage (25-60X), multiplexed genomic data to

reference loci from a closely related taxon. For our reference set of nuclear loci for wing lice, we

used 1,039 exons of *Columbicola drowni* generated in a previous study (Boyd BM, et al. 2017)

(raw data: SRR3161922). This data set was assembled de novo (Allen et al. 2015) using

orthologous protein-coding genes from the human body louse genome (*Pediculus humanus*

*humanus* (Kirkness et al. 2010)) as a set of target sequences. We mapped our newly generated

*Columbicola* reads and the reads obtained from GenBank to the *C. drowni* references using

Bowtie2 (Langmead and Salzberg 2012). For body lice, we obtained nuclear data using the same

pipeline and software parameters, except that we used 1,095 loci from *P. emersoni* as the

reference for mapping. To generate the input ultrametric gene trees for Phylonet v3.6.8 (Than et

al. 2008; Yu and Nakhleh 2015; Wen et al. 2018), we first aligned each nuclear locus in MAFFT

(Katoh 2002)(--auto) and removed columns with only ambiguous sequences ("N"). Then, we

estimated gene trees in RAxML v8.1.3 (Stamatakis 2006) with a GTR + Γ substitution model for

each gene alignment. Finally, we made trees ultrametric using the nnls method in the

force.ultrametric function within the "phytools" R package (Revell 2011).

Quantifying introgression

We used two different approaches to quantify differences in the extent of introgression between

the two louse genera. We employed methods suitable to both detect introgression between

species and individuals from the same species (i.e., we did not employ methods aimed to detect

differences at the population level, e.g., TreeMix; Pickrell and Pritchard 2012).  First, we used

sppIDer (Langdon et al. 2018) to quantify the genomic contributions of different louse species in

an individual louse genome. We built our reference for each genus using all the nuclear loci from

a single individual per species. For the reference, we selected those individuals for which we

assembled the highest number of genes for each genus. Finally, we estimated the extent of

introgression as the sum of the mean coverages of reads mapped from all the species excluding

the focal louse species, divided by the mean coverage of the focal louse species. Second, we

quantified introgression at the species level, while accounting for ILS (i.e., reticulations in this

method can be attributed to hybridization events), using a maximum pseudo-likelihood

framework with PhyloNet 3.6.1 (Than et al. 2008; Yu and Nakhleh 2015; Wen et al. 2018). We

trimmed the unrooted gene trees to the same individuals used as reference taxa in sppIDer, and

performed eleven independent analyses with a differing maximum number of reticulation nodes

(i.e., from zero to ten). We conducted ten runs per analysis. We then selected the optimal

network for each genus based on AIC values.

Analyses

We compared the sppIDer results using generalized linear models (GLMs). We used a Gaussian

distribution of errors and an identity link function. We performed one GLM for each simulation

iteration using the glm function of the "stats" R package (R Core Team 2013). The extent of

introgression for each louse genus was the dependent variable, the genus identity was the

independent variable, and we accounted for the introgression differences between louse species

including louse identity as a fixed factor. We confirmed assumptions underlying GLMs by

testing the normality of regression residuals for normality against a Q-Q plot. We also

considered the possibility that some of the reads mapping to other species were technical

contaminations, i.e., due to index-swapping (Carlsen et al. 2012; Esling et al. 2015; Schnell et al.

2015; Sinha et al. 2017). Previous studies have found that the misassignment of reads generally

ranges from 1 to 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al.

2017). Thus, to account for possible contaminants, we wrote a simulation in R that randomly

subtracted 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al. 2017)

from the mean coverage value of a particular sample (i.e., we subtracted a random proportion of

the mean coverage value for each sample until reaching 9 %). We ran 100 iterations of the

simulation and ran a GLM for each iteration (Table S1). Finally, we used the $\chi^2$ test to compare

the number of species in pairwise comparisons of each genus with the number of reticulations

found in each optimal phylogenetic network.

**Results**

We used two different approaches to quantify the overall (i.e., ancient plus recent) differences in

introgression between louse genera. First, in individual louse genomes, we quantified the

genomic contributions from different closely related louse species of the same genus (Langdon et

al. 2018). Second, we quantified introgression at the species level, while accounting for

incomplete lineage sorting (ILS), by inferring phylogenetic networks using a maximum pseudo-

likelihood framework (Than et al. 2008; Yu and Nakhleh 2015; Wen et al. 2018).

Both approaches revealed highly concordant results: higher levels of introgression among

species of wing lice compared to body lice. In particular, using a read-mapping based method,

the genomic signature of introgression was significantly higher in wing louse species than in

body louse species (GLM with the mean values of the simulations; F = 21.0705, df = 69, *P* =

2.367 x 10$^{-5}$; Fig. 2, Supplementary Table S1, Figs. S1-S12). Contrary to this effect, one body

louse individual (included in the GLMs) exhibited the highest level of introgression (Fig. 2,

Figs. S1-S12). However, the other individual from the same taxon, inhabiting the same host

species and collected in the same geographic region, did not show these elevated levels of

introgression (Supplementary Table S2, Figs. S1-S12).

**Figure 2**. Boxplot showing the differences in levels of
introgression between wing (green) and body (orange) lice. Level
of introgression represents the sum of the mean coverage of reads
mapped from all the species excluding the focal louse species,
divided by the mean coverage of the focal louse species (see
Methods). Black dots represent individual samples (horizontally
jittered).

Secondly, in a phylogenetic network framework, the optimal networks of wing lice were more

reticulated than those of body lice even though the number of taxa included in the networks was

lower (seven reticulations in *Columbicola* vs. four in *Physconelloides*, Fig. 3). Accordingly, the

number of reticulations given the number of potential combinations was significantly higher ($\chi^2$=

3.8132; df=1; *P*= 0.03). Also, the specific lineages involved in the reticulations were generally

congruent with signatures of introgression from the read-mapping based approach (Fig. S1-S12).

Figure 3. Optimal phylogenetic networks of feather lice genera. Orange branches depict reticulations. From left to right,

*Columbicola* (seven reticulations) and *Physconelloides* (four reticulations) networks (See Methods).

**Discussion**

Taken together, evidence from wing and body louse genomes suggests that differences in

dispersal ability drive differences in the extent of introgression in this system of ecological

replicate parasites. This work is among the first studies of introgression in a host-symbiont

system (Detwiler and Criscione 2010). Notably, recent studies have found that straggling and

host-switching are relatively common processes in host-symbiont systems (De Vienne et al.

2013; Bourguignon et al. 2018; Nylin et al. 2018; Doña et al. 2019). Our study suggests that in a

straggling/host-switching scenario, hybridization can provide further variation with important

eco-evolutionary consequences (Barton 2018). Indeed, we may have found a potential recent

hybridization event (i.e., the *Physconelloides* individual showing the highest level of

introgression), though this requires further study to rule out methodological issues (e.g., wet-lab

contamination). Overall, the results from this study represent a significant step towards

understanding the factors driving hybridization, because most previous studies focus on the

presence/absence of hybridization and the evolutionary consequences of hybridization events

(Arnold 2015; Folk et al. 2018; Taylor and Larson 2019). Further research is needed to

understand the factors shaping the frequency of hybridization and to ascertain their influence

across different scales (e.g., from ancient to recent hybridization events).

**References:**

Allen, J. M., D. I. Huang, Q. C. Cronk, and K. P. Johnson. 2015. aTRAM - automated target

restricted assembly method: a fast method for assembling loci across divergent taxa from

next-generation sequencing data. BMC Bioinformatics 16.

Arnold, M. L. 2015. Divergence with genetic exchange. OUP Oxford, Oxford, UK.

Bartlow, A. W., S. M. Villa, M. W. Thompson, and S. E. Bush. 2016. Walk or ride? Phoretic

behaviour of amblyceran and ischnoceran lice. Int J Parasitol 46:221–227.

Barton, N. H. 2018. The consequences of an introgression event. Mol Ecol 27:4973–4975.

Barton, N. H. 1992. The genetic consequences of dispersal. In: Animal Dispersal. Springer,

Netherlands.

Bohonak, A. J. 1999. Dispersal Gene Flow, and Population Structure. Q Rev Biol 74:21–45.

Bourguignon, T., N. Lo, C. Dietrich, J. Šobotník, S. Sidek, Y. Roisin, A. Brune, and T. A. Evans.

2018. Rampant host switching shaped the termite gut microbiome. Curr Biol 28:649–654.

Boyd BM, Allen JM, Nguyen N, Sweet AD, Warnow T, Shapiro MD, and et al. 2017.

Phylogenomics using Target-restricted Assembly Resolves Intra-generic Relationships of

Parasitic Lice (Phthiraptera: Columbicola). Syst Biol 66:896–911.

Carlsen, T., A. B. Aas, D. Lindner, T. Vrålstad, T. Schumacher, and H. Kauserud. 2012. Dont

make a mista(g)ke: is tag switching an overlooked source of error in amplicon

pyrosequencing studies? Fungal Ecol 5:747–749.

Clayton, D. H., S. E. Bush, and K. P. Johnson. 2015. Coevolution of life on hosts: integrating

ecology and history. University of Chicago Press, Chicago, IL.

Clayton, D. H., and K. P. Johnson. 2003. Linking coevolutionary history to ecological process:

doves and lice. Evolution 57:2335–2341.

Clobert, J., M. Baguette, T. G. Benton, and J. M. Bullock (eds). 2012. Dispersal Ecology and

Evolution. Oxford University Press.

Clobert, J., E. Danchin, A. A. Dhondt, and N. J. D. 2001. Dispersal. Oxford Univ. Press.

Currat, M., M. Ruedi, R. J. Petit, and L. Excoffier. 2008. The hidden side of invasions: massive

introgression by local genes. Evolution 62:1908–1920.

Dawson, M. N., C. G. Hays, R. K. Grosberg, and P. T. Raimondi. 2014. Dispersal potential and

population genetic structure in the marine intertidal of the eastern North Pacific. Ecol

Monogr 84:435–456.

De Vienne, D. M., G. Refrégier, M. López-Villavicencio, A. Tellier, M. E. Hood, and T. Giraud.

2013. Cospeciation vs host-shift speciation: methods for testing, evidence from natural

associations and relation to coevolution. New Phytol 198:347–385.

Detwiler, J. T., and C. D. Criscione. 2010. An infectious topic in reticulate evolution:

introgression and hybridization in animal parasites. Genes (Basel) 1:102–23.

DiBlasi, E., K. P. Johnson, S. A. Stringham, A. N. Hansen, A. B. Beach, and D. H. Clayton.

2018. Phoretic dispersal influences parasite population genetic structure. Mol Ecol

27:2770–2779.

Doña, J., D. Serrano, S. Mironov, A. Montesinos-Navarro, and R. Jovani. 2019. Unexpected

bird–feather mite associations revealed by DNA metabarcoding uncovers a dynamic

ecoevolutionary scenario. Mol Ecol 28:379–390.

Esling, P., F. Lejzerowicz, and J. Pawlowski. 2015. Accurate multiplexing and filtering for high-

throughput amplicon-sequencing. Nucleic Acids Res 43:2513–24.

Folk, R. A., P. S. Soltis, D. E. Soltis, and R. Guralnick. 2018. New prospects in the detection and

comparative analysis of hybridization in the tree of life. Am J Bot 105:364–375.

Harbison, C. W., S. E. Bush, J. R. Malenke, and D. H. Clayton. 2008. Comparative transmission

dynamics of competing parasite species. Ecology 89:3186–3194.

Harbison, C. W., M. V. Jacobsen, and D. H. Clayton. 2009. A hitchhiker's guide to parasite

transmission: The phoretic behaviour of feather lice. Int J Parasitol 39:569–575.

Johnson, K. P., and D. H. Clayton. 2004. Untangling coevolutionary history. Syst Biol 53:92–4.

Katoh, K. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast

Fourier transform. Nucleic Acids Res 30:3059–3066.

Kirkness, E. F., B. J. Haas, W. Sun, H. R. Braig, M. A. Perotti, and J. M. Clark. 2010. Genome

sequences of the human body louse and its primary endosymbiont provide insights into

the permanent parasitic lifestyle. Proc Natl Acad Sci U S A 107:12168–73.

Langdon, Q. K., D. Peris, B. Kyle, and C. T. Hittinger. 2018. sppIDer: A Species Identification

Tool to Investigate Hybrid Genomes with High-Throughput Sequencing. Mol Biol Evol

35:2835–2849.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat

Methods 9:357–9.

Matthysen, E. 2012. Multicausality of dispersal: a review. In: Dispersal Ecology and Evolution.

Oxford University Press.

McEntee, J. P., J. G. Burleigh, and S. Singhal. 2018. Dispersal predicts hybrid zone widths

across animal diversity: Implications for species borders under incomplete reproductive

isolation. bioRxiv, doi: 10.1101/472506.

Nathan, R. 2001. The challenges of studying dispersal. Trends Ecol Evol 16:481–483.

Nussberger, B., M. Currat, C. S. Quilodran, N. Ponta, and L. F. Keller. 2018. Range expansion as

an explanation for introgression in European wildcats. Biol Conserv 218:49–56.

Nylin, S., S. Agosta, S. Bensch, W. A. Boeger, M. P. Braga, D. R. Brooks, M. L. Forister, P. A.

Hambäck, E. P. Hoberg, and T. Nyman. 2018. Embracing colonizations: a new paradigm

for species association dynamics. Trends Ecol Evol 33:4–14.

Pickrell, J. K., and J. K. Pritchard. 2012. Inference of Population Splits and Mixtures from

Genome-Wide Allele Frequency Data. PLoS Genetics 8.

R Core Team. 2013. R: A language and environment for statistical computing.

Randler, C. 2006. Behavioural and ecological correlates of natural hybridization in birds. Ibis

148:459–467.

Revell, L. 2011. phytools: an R package for phylogenetic comparative biology (and other

things). Methods Ecol Evol 3:217–223.

Schnell, I. B., K. Bohmann, and M. T. Gilbert. 2015. Tag jumps illuminated–reducing sequence-

288           to-sample misidentifications in metabarcoding studies. Mol Ecol Resour 15:1289–303.

289    Sinha, R., G. Stanley, G. S. Gulati, C. Ezran, K. J. Travaglini, and E. Wei. 2017. Index

290           Switching Causes Spreading-Of-Signal Among. bioRxiv, doi: 10.1101/125724.

291    Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with

292           thousands of taxa and mixed models. Bioinformatics 22:2688–2690.

293    Sweet, A. D., B. M. Boyd, J. M. Allen, S. M. Villa, M. P. Valim, and J. L. Rivera-Parra. 2017a.

294           Integrating phylogenomic and population genomic patterns in avian lice provides a more

295           complete picture of parasite evolution. Evolution 72:95–112.

296    Sweet, A. D., R. T. Chesser, and K. P. Johnson. 2017b. Comparative cophylogenetics of

297           Australian phabine pigeons and doves (Aves: Columbidae) and their feather lice (Insecta:

298           Phthiraptera). Int J Parasitol 47:347–356.

299    Sweet, A. D., and K. P. Johnson. 2015. Patterns of diversification in small New World ground

300           doves are consistent with major geologic events. Auk 132:300–312.

301    Sweet, A. D., and K. P. Johnson. 2018. The role of parasite dispersal in shaping a host–parasite

302           system at multiple evolutionary scales. Mol Ecol 27:5104–5119.

303    Taylor, S. A., and E. L. Larson. 2019. Insights from genomes into the evolutionary importance

304           and prevalence of hybridization in nature. Nat Ecol Evol 3:170.

305    Than, C., D. Ruths, and L. Nakhleh. 2008. PhyloNet: a software package for analyzing and

306           reconstructing reticulate evolutionary relationships. BMC Bioinformatics 9:322.

307    Wen, D., Y. Yu, J. Zhu, and L. Nakhleh. 2018. Inferring Phylogenetic Networks Using

308           PhyloNet. Syst Biol 67:735–740.

309    Whiteman, N. K., D. Santiago-Alarcon, K. P. Johnson, and P. G. Parker. 2004. Differences in

310        straggling rates between two genera of dove lice (Insecta: Phthiraptera) reinforce

311        population genetic and cophylogenetic patterns. Int J Parasitol 34:1113–1119.

312    Yu, Y., and L. Nakhleh. 2015. A maximum pseudo-likelihood approach for phylogenetic

313        networks. BMC Genomics 16:Suppl 10: S10.