Title: Comparing rates of introgression in parasitic feather lice with differing

dispersal capabilities

Authors: Jorge Doña[1,2]*, Andrew D. Sweet[1,3], and Kevin P. Johnson[1]*

Affiliations:

[1]Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-
Champaign, 1816 S. Oak St., Champaign, Illinois 61820, USA.

[2]Departamento de Biología Animal, Universidad de Granada, Granada, 18001, Spain.

[3]Department of Entomology, Purdue University, 901 W. State St., West
Lafayette, Indiana 47907, USA.

*Correspondence to: jorged@illinois.edu, kpjohnso@illinois.edu.

Running title: Dispersal and introgression rate.

Author contributions: J.D., and K.P.J. conceived the study. J.D., A.D.S., and K.P.J. designed
the study. A.D.S. collected the data. J.D. and A.D.S. analysed the data. K.P.J. obtained financial
support for the project. J.D. wrote the manuscript and all authors contributed to editing the
manuscript.

Acknowledgments: This study was supported by the US National Science Foundation (DEB-
1239788, DEB-1342604, DEB-1926919, and DEB-1925487 to K.P.J). We declare that we have
no conflict of interest.

Data availability: All data needed to evaluate the conclusions in the paper are present in the
paper and/or the Supplementary Materials. Additional data related to this paper may be requested
from the authors.

Supplementary Materials: Supplementary data are available at GitHub (https://jorge-
dona.github.io/Comparing-rates-of-introgression-in-parasitic-feather-lice-with-differing-
dispersal-capabilities/supplementary.html) and at Figshare (DOI: 10.6084/m9.figshare.9176204).
Individual gene trees are available at Figshare (DOI: 10.6084/m9.figshare.9176204).
26

Abstract: Organisms vary in their dispersal abilities, and these differences can have important

biological consequences, such as impacting the likelihood of hybridization events. However,

there is still much to learn about the factors influencing hybridization, and specifically how

dispersal ability affects the opportunities for hybridization. Here, using the ecological replicate

system of dove wing and body lice (Insecta: Phthiraptera), we show that species with higher

dispersal abilities exhibited increased genomic signatures of introgression. Specifically, we

found a higher proportion of introgressed genomic reads and more reticulated phylogenetic

networks in wing lice, the louse group with higher dispersal abilities. Our results are consistent

with the hypothesis that differences in dispersal ability might drive the extent of introgression

through hybridization.

Introduction

Dispersal is the permanent movement of organisms away from their place of origin. It is a fundamental process in biology with significant implications at multiple scales of organization (Barton 1992; Clobert et al. 2001; Nathan 2001; Matthysen 2012), including the reproduction of individuals, the composition of populations and communities, and the geographical distribution of species (Clobert et al. 2001, 2012).

Organisms differ in their dispersal abilities, and these differences have an impact on their biology, such as on the distributional range of a species or gene flow between populations (Bohonak 1999). For example, organisms with lower dispersal abilities tend to have smaller distributional ranges and populations that are genetically more structured (Bohonak 1999; Dawson et al. 2014; DiBlasi et al. 2018).

Dispersal ability might also affect the opportunities for hybridization between species because the rates at which individuals encounter different species are likely to be higher in organisms with higher dispersal capabilities. Indeed, recent evidence supports this prediction by demonstrating that range expansion is associated with the extent of introgression (Currat et al. 2008; Nussberger et al. 2018). Similarly, dispersal differences explain more than 30% of the variation in the width of hybrid zones across animals (McEntee et al. 2018). However, overall there is still much to learn about the factors influencing hybridization (Randler 2006; Arnold 2015; Taylor and Larson 2019), and, in particular, the influence of dispersal ability on the rate of hybridization remains understudied.

Testing for the effect of dispersal on hybridization should ideally hold constant most factors other than dispersal. The ecological replicate system of wing and body lice (Insecta: Phthiraptera) of pigeons and doves (Aves: Columbidae) has proven to be an ideal system for

comparing the impact of dispersal differences on other aspects of biology, such as population structure and codivergence (Clayton and Johnson 2003; Johnson and Clayton 2004; Clayton et al. 2015; DiBlasi et al. 2018; Sweet and Johnson 2018). Specifically, this is an excellent system in which to assess the effect of differences in dispersal capabilities on levels of introgression because both of these lineages of feather lice: 1) significantly differ in their dispersal ability (Harbison et al. 2008, 2009; Bartlow et al. 2016), 2) co-occur across the diversity of pigeons and doves, and 3) have the same basic life history and diet (Clayton et al. 2015; Sweet and Johnson 2015, 2018). Both wing and body lice disperse vertically between parents and offspring in the nest. However, wing lice can also attach to and hitchhike on hippoboscid flies to disperse "phoretically" between host individuals or host species (Harbison et al. 2008, 2009; Bartlow et al. 2016). Indeed, this hitch-hiking dispersal mechanism profoundly influences their degree of population structure and cophylogenetic history (Clayton and Johnson 2003; Sweet et al. 2017b; DiBlasi et al. 2018; Sweet and Johnson 2018). In addition, wing lice have a higher rate of host-switching (Clayton and Johnson 2003; Clayton et al. 2015; Sweet et al. 2017b) (i.e., successful colonization of new host species) and of straggling (Whiteman et al. 2004) (i.e., dispersal to new host species without reproduction on that new host).

To compare differences in the extent of introgression between wing and body lice, we used whole-genome data from 71 louse individuals belonging to five species of wing lice (*Columbicola*) and seven species of body lice (*Physconelloides*) that occur across the same suite of host species and have highly comparable patterns of diversification (Sweet and Johnson 2015, 2018). Specifically, both lineages within these two groups of lice that are the focus of this study originated on the common ancestor of *Metriopelia* doves (11.3-14.9 mya) and have a correlated pattern of codiversification within the same group of hosts (including a shared cospeciation event

which occurred within the *Metriopelia* genus 5.2-7.4 mya; Sweet and Johnson 2015, 2018). We

predicted that wing lice, which have higher dispersal abilities and thus higher odds of

encountering individuals of a different louse species on the same host, should show more

extensive evidence of introgression (Fig. 1).

Figure 1. Diagram depicting the ecological replicate system and the hypothesis of this study. Wing lice (*Columbicola*) have

higher dispersal abilities than body lice (*Physconelloides*), and thus higher odds of encountering individuals of a different louse

sp.

Materials and Methods

<u>Data</u>

We analyzed Illumina whole genome sequence data (150 or 160bp paired-end reads) from 71

louse individuals belonging to five and seven taxa of *Columbicola* and *Physconelloides*,

respectively (Supplementary Table S2) hosted by the monophyletic clade of small New World

ground doves. This paper's taxonomic classification of lice is based on Sweet and Johnson

(2018) species delimitation analyses. In particular, they found most *Columbicola* OTUs matched

known species, and some *Physconelloides* OTUs were yet to be formally described as species

(and are named here following Sweet & Johnson, 2018). All raw sequence data used were

available from previous studies (Boyd et al. 2017; Sweet et al. 2017a; Sweet and Johnson 2018)

and represent all described New World ground-dove wing and body louse species (sampled from

most host species in this group) including lice samples across multiple biogeographic areas

within species of hosts (Sweet and Johnson 2018; Supplementary Table S2). Illumina genome

sequence data pre-processing included several steps (Sweet and Johnson 2018). First, we

discarded duplicate read pairs using the fastqSplitDups script (https://github.com/McIntyre-

Lab/mcscriptand https://github.com/McIntyre-Lab/mclib). We then eliminated the Illumina

sequencing adapters with Fastx_clipper v0.014 from the FASTX-Toolkit

(http://hannonlab.cshl.edu/fastx_toolkit). Also, we removed the first 5 nt from the 5' ends of

reads using Fastx_trimmer v0.014 and trimmed bases from the 3' ends of reads until reaching a

base with a phred score ≥28 (which is equivalent to a base call accuracy higher than 99.8 %)

using Fastq_quality_trimmer v0.014. Finally, we removed any reads less than 75 nt and analyzed

the cleaned libraries with Fastqc v0.11.5 to check for additional errors. We assembled nuclear

loci in aTRAM following previous studies (Allen et al. 2015; Boyd et al. 2017; Sweet and

Johnson 2018). In particular, we mapped modest coverage (25-60X), multiplexed genomic data

to reference loci from a closely related taxon. For our reference set of nuclear loci for wing lice,

we used 1,039 exons of *Columbicola drowni* (Boyd et al. 2017) (raw data: SRR3161922). This

data set was assembled de novo (Allen et al. 2015) using orthologous protein-coding genes from

the human body louse genome (*Pediculus humanus humanus* (Kirkness et al. 2010) as a set of

target sequences. We mapped *Columbicola* reads to the *C. drowni* references using Bowtie2

(Langmead and Salzberg 2012). For body lice, we obtained nuclear data using the same pipeline

and software parameters, except that we used 1,095 loci from *Physconelloides emersoni* as the

reference for mapping. To generate the input ultrametric gene trees for Phylonet v3.6.8 (Than et

al. 2008; Yu and Nakhleh 2015; Wen et al. 2018), we first aligned each nuclear locus in MAFFT

(Katoh 2002) (--auto) and removed columns with only ambiguous sequences ("N"). Then, we

estimated gene trees in RAxML v8.1.3 (Stamatakis 2006) with a GTR + Γ substitution model for

each gene alignment. Finally, we made trees ultrametric using the nnls method in the

force.ultrametric function within the "phytools" R package (Revell 2011).

<u>Quantifying introgression</u>

We used two different approaches to quantify differences in the extent of introgression (i.e.,

ancient plus recent) between the two louse genera. We employed methods suitable to detect

introgression between species and between individuals from the same species (i.e., we did not

employ methods aimed to detect differences at the population level, e.g., TreeMix; Pickrell and

Pritchard 2012). First, we used sppIDer (Langdon et al. 2018) to quantify the genomic

contributions of different louse species in an individual louse genome (Fig. 2). We built our

reference for each genus using all the nuclear loci from a single individual per species. For the

reference, we selected those individuals for which we assembled the highest number of genes for

each genus. We estimated the extent of introgression as the sum of the mean coverages of reads

mapped from all the species excluding the focal louse species, divided by the mean coverage of

the focal louse species (Fig. 2). Note that these mean coverage values are calculated using only

those reads that mapped with a mapping quality (MQ) > 3 (Li et al. 2009; Langdon et al. 2018;

Figs. S1-S12). Plots of coverage across the genomes suggested that reads mapping to other

species were not artificial mappings (e.g., high coverage mappings to short repetitive regions;

Figs. S1-S12). In addition, we used SPAdes v3.12.0 (default parameters) to perform a de-novo

assembly of the putatively introgressed reads detected by sppIDer).

Figure 2. Illustrative example of sppIDer results with an individual of *Columbicola passerinae* 1 (Cosp.Copas.11.7.2016.8).
Panel a) shows the average mean coverages of reads mapping to every species (i.e., the values that were used for the calculations
of introgression levels), and panel b) shows the mean coverages of reads mapping to each species across the whole set of loci.
These and additional visualizations of all the individuals can be found in the supplementary material (Figs. S1-S12).

Second, we quantified introgression at the species level, while accounting for incomplete lineage

sorting (ILS) using a maximum pseudo-likelihood framework with PhyloNet 3.6.1 (Than et al.

2008; Yu and Nakhleh 2015; Wen et al. 2018). Reticulations in this method can be attributed to

hybridization events. We trimmed the unrooted gene trees to the same individuals used as

reference taxa in sppIDer, and performed eleven independent analyses with a differing maximum

number of reticulation nodes (i.e., from zero to ten). We conducted ten runs per analysis. We

then selected the optimal network for each genus based on AIC values.

<u>Statistics and reproducibility</u>

We compared the sppIDer results using generalized linear models (GLMs). We used a Gaussian

distribution of errors and an identity link function. We performed one GLM for each simulation

iteration using the glm function of the "stats" R package (R Core Team 2013). The extent of

introgression for each louse genus was the dependent variable, the genus identity was the

independent variable, and we accounted for the introgression differences between louse species

including louse identity as a fixed factor. We confirmed assumptions underlying GLMs by

testing the normality of regression residuals for normality against a Q-Q plot. We also

considered the possibility that some of the reads mapping to other species were technical

contaminations, i.e., due to index-swapping (Carlsen et al. 2012; Esling et al. 2015; Schnell et al.

2015; Sinha et al. 2017). Previous studies have found that the misassignment of reads generally

ranges from 1 to 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al.

2017). Thus, to account for possible contaminants, we wrote a simulation in R that randomly

subtracted 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al. 2017)

from the mean coverage value of a particular sample (i.e., we subtracted a random proportion of

the mean coverage value for each sample until reaching 9%). We ran 100 iterations of the

simulation and ran a GLM for each iteration (Table S1). Finally, we used the χ^2 test to compare

the number of species in pairwise comparisons of each genus with the number of reticulations

found in each optimal phylogenetic network. Because we had an *a priori* prediction that

Physconelloides should exhibit less evidence of reticulation than *Columbicola*, we used a one-

tailed test; however, we also report the results of the two-tailed test equivalent.

Results

Both approaches revealed highly concordant results: higher levels of introgression among

species of wing lice compared to body lice. In particular, using a read-mapping based method,

the genomic signature of introgression was significantly higher in wing louse species than in

body louse species (GLM with the mean values of the simulations; $F = 21.0705$, df = 69, $P =$

2.367×10^{-5}, $R^2=0.58$; Fig. 3, Supplementary Table S1, Figs. S1-S12). The contigs assembled

from reads mapping to the non-focal species were in the size range of the loci used as reference

(mean max contig length= 1214 bp; mean contig length=292 bp; Table S3). Even though wing

lice showed more evidence for introgression, one body louse individual (included in the GLMs)

exhibited the highest level of introgression (Fig. 3, Figs. S1-S12). However, the other individual

from the same taxon, inhabiting the same host species and collected in the same geographic

region, did not show these elevated levels of introgression (Supplementary Table S2, Figs. S1-

S12).

Figure 3. Boxplot showing the differences in levels of introgression between wing (green) and body (orange) lice. Level of introgression represents the sum of the mean coverage of reads mapped from all the species excluding the focal louse species, divided by the mean coverage of the focal louse species (see Methods). Black dots show the levels of introgression (i.e., resulting from the equation) for each individual sample (horizontally jittered values).

Secondly, in a phylogenetic network framework, the optimal networks of wing lice were more

reticulated than those of body lice even though the number of taxa included in the networks was

lower (seven reticulations in *Columbicola* vs. four in *Physconelloides*, Fig. 4). Accordingly, the

number of reticulations given the number of potential combinations under a one-tailed test was

significantly higher in *Columbicola* (One-sided: χ^2= 3.8132; df=1; *P*= 0.03; CI=0.03–1), and the

p-value was still near 0.05 with a two-tailed test (Two-sided: χ^2= 3.8132; df=1; *P*= 0.05; CI=-

0.01–0.66). Also, the specific lineages involved in the reticulations were generally congruent

with signatures of introgression from the read-mapping based approach (Fig. S1-S12).

Figure 4. Optimal phylogenetic networks of feather lice genera. Orange branches depict reticulations: seven in *Columbicola* and four in *Physconelloides*.

Discussion

Estimates of introgression in two groups of ectoparasites that differ in their dispersal abilities, wing and body lice of doves, indicate that the lineage with higher dispersal ability (wing lice) shows more evidence of introgression. This evidence from wing and body louse genomes is

consistent with the hypothesis that dispersal differences might drive differences in the level of

introgression in this system of parasites. Admittedly, there may be some unknown factor, other

than dispersal, differing between these two groups of lice that causes the difference in the level

of introgression, but prior work on these groups of parasites points to dispersal as a crucial factor

underlying many of the ecological and evolutionary patterns in these parasites. Further research

on other taxa is needed to confirm the generality of these findings. This work is among the first

studies of introgression in a host-symbiont system (Detwiler and Criscione 2010). Notably,

recent studies have found that straggling and host-switching are relatively common processes in

host-symbiont systems (De Vienne et al. 2013; Bourguignon et al. 2018; Nylin et al. 2018; Doña

et al. 2019). Our study suggests that in a straggling/host-switching scenario, hybridization can

provide further genetic variation with important ecological and evolutionary consequences (e.g.,

facilitating adaptation to current hosts or facilitating the colonization of new ones) (Barton

2018). Indeed, we may have found a potential recent hybridization event (i.e., the

Physconelloides individual showing the highest level of introgression), though this requires

further study to rule out methodological issues (e.g., wet-lab contamination).

In this vein, a careful examination of the introgression history of these taxa (and

symbionts as a whole) is needed to better understand the patterns of introgression that we found.

Questions such as how much introgression can be expected or how the introgressed regions are

retained in parasite/symbiont genomes across time, among many others, require further attention.

For instance, in this study, the levels of introgression detected by the sppIDer analyses (i.e., the

magnitude but not the comparative pattern) may be unrealistic. It may be that some fraction of

the level of introgression detected by sppIDer may be due to ILS, and not introgression.

However, both louse genera are expected to have relatively similar rates of ILS (if any). It is also

possible that taxon age and interspecific divergence might affect introgression rates.

Nevertheless, Sweet and Johnson (2018) compared the degree of genetic divergence of two pairs

of species of both genera that inhabits the same host species and share a cospeciation event. In

this case of taxa of the same age, the pair of *Columbicola* species had lower interspecific genetic

distances than *Physconelloides*. This could be as a result of mutation rate differences between the

two genera. However, this could also be due to higher gene flow among host infrapopulations

due to higher dispersal capabilities of wing lice. The same pattern can be found overall across the

species studied here, i.e., on average, lower uncorrected interspecific genetic distances among

Columbicola than among *Physconelloides* species, though the range of the interspecific distances

does overlap (Table S4). Thus, if present, ILS could potentially be more prevalent in

Columbicola species (Pamilo & Nei, 1988; Maddison, 1997). The PhyloNet analysis, however,

does control for ILS, and showed highly congruent results. In addition, some individual gene

trees exhibit signatures suggestive of introgression with highly similar sequences shared by some

individuals of different species and much less likely to be a consequence of ILS (Figs. S13-S14).

Overall, the species of *Columbicola* and *Physconelloides* are from the same group of hosts and

thus are overall comparatively similar in levels of divergence, so it seems unlikely that these

small differences are driving the results.

Another caveat is that sppIDer can detect introgression from species that are not included

in the reference data. In those cases, the reads may map to the closest taxon available in the

reference set, and thus could artificially increase the level of introgression from a given species;

Langdon et al. 2018). Accordingly, the levels of introgression detected by sppIDer in certain

species could be an aggregate of introgression events from more than one species. Indeed, our

PhyloNet analysis supports this scenario, with several reticulations from ghost lineages and

species (Fig. 4). However, in this system we have nearly complete sampling of host taxa and are

missing few, if any, extant species making this concern less likely.

References:

Allen, J. M., D. I. Huang, Q. C. Cronk, and K. P. Johnson. 2015. aTRAM - automated target restricted assembly

method: a fast method for assembling loci across divergent taxa from next-generation sequencing data.

BMC Bioinformatics 16.

Arnold, M. L. 2015. Divergence with genetic exchange. OUP Oxford, Oxford, UK.

Bartlow, A. W., S. M. Villa, M. W. Thompson, and S. E. Bush. 2016. Walk or ride? Phoretic behaviour of

amblyceran and ischnoceran lice. Int J Parasitol 46:221–227.

Barton, N. H. 2018. The consequences of an introgression event. Mol Ecol 27:4973–4975.

Barton, N. H. 1992. The genetic consequences of dispersal. In: Animal Dispersal. Springer, Netherlands.

Bohonak, A. J. 1999. Dispersal Gene Flow, and Population Structure. Q Rev Biol 74:21–45.

Bourguignon, T., N. Lo, C. Dietrich, J. Šobotník, S. Sidek, Y. Roisin, A. Brune, and T. A. Evans. 2018. Rampant

host switching shaped the termite gut microbiome. Curr Biol 28:649–654.

Boyd BM, Allen JM, Nguyen N, Sweet AD, Warnow T, Shapiro MD, and et al. 2017. Phylogenomics using Target-

restricted Assembly Resolves Intra-generic Relationships of Parasitic Lice (Phthiraptera: Columbicola).

Syst Biol 66:896–911.

Carlsen, T., A. B. Aas, D. Lindner, T. Vrålstad, T. Schumacher, and H. Kauserud. 2012. Dont make a mista(g)ke: is

tag switching an overlooked source of error in amplicon pyrosequencing studies? Fungal Ecol 5:747–749.

Clayton, D. H., S. E. Bush, and K. P. Johnson. 2015. Coevolution of life on hosts: integrating ecology and history.

University of Chicago Press, Chicago, IL.

Clayton, D. H., and K. P. Johnson. 2003. Linking coevolutionary history to ecological process: doves and lice.

Evolution 57:2335–2341.

Clobert, J., M. Baguette, T. G. Benton, and J. M. Bullock (eds). 2012. Dispersal Ecology and Evolution. Oxford

University Press.

Clobert, J., E. Danchin, A. A. Dhondt, and N. J. D. 2001. Dispersal. Oxford Univ. Press.

Currat, M., M. Ruedi, R. J. Petit, and L. Excoffier. 2008. The hidden side of invasions: massive introgression by

local genes. Evolution 62:1908–1920.

Dawson, M. N., C. G. Hays, R. K. Grosberg, and P. T. Raimondi. 2014. Dispersal potential and population genetic

structure in the marine intertidal of the eastern North Pacific. Ecol Monogr 84:435–456.

De Vienne, D. M., G. Refrégier, M. López-Villavicencio, A. Tellier, M. E. Hood, and T. Giraud. 2013. Cospeciation

vs host-shift speciation: methods for testing, evidence from natural associations and relation to coevolution.

New Phytol 198:347–385.

Detwiler, J. T., and C. D. Criscione. 2010. An infectious topic in reticulate evolution: introgression and

hybridization in animal parasites. Genes (Basel) 1:102–23.

DiBlasi, E., K. P. Johnson, S. A. Stringham, A. N. Hansen, A. B. Beach, and D. H. Clayton. 2018. Phoretic dispersal

influences parasite population genetic structure. Mol Ecol 27:2770–2779.

Doña, J., D. Serrano, S. Mironov, A. Montesinos-Navarro, and R. Jovani. 2019. Unexpected bird–feather mite

associations revealed by DNA metabarcoding uncovers a dynamic ecoevolutionary scenario. Mol Ecol

28:379–390.

Esling, P., F. Lejzerowicz, and J. Pawlowski. 2015. Accurate multiplexing and filtering for high-throughput

amplicon-sequencing. Nucleic Acids Res 43:2513–24.

Folk, R. A., P. S. Soltis, D. E. Soltis, and R. Guralnick. 2018. New prospects in the detection and comparative

analysis of hybridization in the tree of life. Am J Bot 105:364–375.

Harbison, C. W., S. E. Bush, J. R. Malenke, and D. H. Clayton. 2008. Comparative transmission dynamics of

competing parasite species. Ecology 89:3186–3194.

Harbison, C. W., M. V. Jacobsen, and D. H. Clayton. 2009. A hitchhiker's guide to parasite transmission: The

phoretic behaviour of feather lice. Int J Parasitol 39:569–575.

Johnson, K. P., and D. H. Clayton. 2004. Untangling coevolutionary history. Syst Biol 53:92–4.

Katoh, K. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform.

Nucleic Acids Res 30:3059–3066.

Kirkness, E. F., B. J. Haas, W. Sun, H. R. Braig, M. A. Perotti, and J. M. Clark. 2010. Genome sequences of the

human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle.

Proc Natl Acad Sci U S A 107:12168–73.

Langdon, Q. K., D. Peris, B. Kyle, and C. T. Hittinger. 2018. sppIDer: A Species Identification Tool to Investigate

Hybrid Genomes with High-Throughput Sequencing. Mol Biol Evol 35:2835–2849.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357–9.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, and R. Durbin. 2009. The

sequence alignment/map format and SAMtools. Bioinformatics 25:2078–2079. Oxford University Press.

Maddison, W. P. 1997. Gene trees in species trees. Syst Biol, 46(3), 523-536.

Matthysen, E. 2012. Multicausality of dispersal: a review. In: Dispersal Ecology and Evolution. Oxford University

Press.

McEntee, J. P., J. G. Burleigh, and S. Singhal. 2018. Dispersal predicts hybrid zone widths across animal diversity:

Implications for species borders under incomplete reproductive isolation. bioRxiv, doi: 10.1101/472506.

Nathan, R. 2001. The challenges of studying dispersal. Trends Ecol Evol 16:481–483.

Nussberger, B., M. Currat, C. S. Quilodran, N. Ponta, and L. F. Keller. 2018. Range expansion as an explanation for

introgression in European wildcats. Biol Conserv 218:49–56.

Nylin, S., S. Agosta, S. Bensch, W. A. Boeger, M. P. Braga, D. R. Brooks, M. L. Forister, P. A. Hambäck, E. P.

Hoberg, and T. Nyman. 2018. Embracing colonizations: a new paradigm for species association dynamics.

Trends Ecol Evol 33:4–14.

Pamilo, P., & Nei, M. 1988. Relationships between gene trees and species trees. Mol Biol Evol, 5(5), 568-583.

Pickrell, J. K., and J. K. Pritchard. 2012. Inference of Population Splits and Mixtures from Genome-Wide Allele

Frequency Data. PLoS Genetics 8.

R Core Team. 2013. R: A language and environment for statistical computing.

Randler, C. 2006. Behavioural and ecological correlates of natural hybridization in birds. Ibis 148:459–467.

Revell, L. 2011. phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol

Evol 3:217–223.

Schnell, I. B., K. Bohmann, and M. T. Gilbert. 2015. Tag jumps illuminated–reducing sequence-to-sample

misidentifications in metabarcoding studies. Mol Ecol Resour 15:1289–303.

Sinha, R., G. Stanley, G. S. Gulati, C. Ezran, K. J. Travaglini, and E. Wei. 2017. Index Switching Causes Spreading-

Of-Signal Among. bioRxiv, doi: 10.1101/125724.

Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa

342   and mixed models. Bioinformatics 22:2688–2690.

344   Sweet, A. D., B. M. Boyd, J. M. Allen, S. M. Villa, M. P. Valim, and J. L. Rivera-Parra. 2017a. Integrating

345   phylogenomic and population genomic patterns in avian lice provides a more complete picture of parasite

346   evolution. Evolution 72:95–112.

347   Sweet, A. D., R. T. Chesser, and K. P. Johnson. 2017b. Comparative cophylogenetics of Australian phabine pigeons

348   and doves (Aves: Columbidae) and their feather lice (Insecta: Phthiraptera). Int J Parasitol 47:347–356.

349   Sweet, A. D., and K. P. Johnson. 2015. Patterns of diversification in small New World ground doves are consistent

350   with major geologic events. Auk 132:300–312.

351   Sweet, A. D., and K. P. Johnson. 2018. The role of parasite dispersal in shaping a host–parasite system at multiple

352   evolutionary scales. Mol Ecol 27:5104–5119.

353   Taylor, S. A., and E. L. Larson. 2019. Insights from genomes into the evolutionary importance and prevalence of

354   hybridization in nature. Nat Ecol Evol 3:170.

355   Than, C., D. Ruths, and L. Nakhleh. 2008. PhyloNet: a software package for analyzing and reconstructing reticulate

356   evolutionary relationships. BMC Bioinformatics 9:322.

357   Wen, D., Y. Yu, J. Zhu, and L. Nakhleh. 2018. Inferring Phylogenetic Networks Using PhyloNet. Syst Biol 67:735–

358   740.

359   Whiteman, N. K., D. Santiago-Alarcon, K. P. Johnson, and P. G. Parker. 2004. Differences in straggling rates

360   between two genera of dove lice (Insecta: Phthiraptera) reinforce population genetic and cophylogenetic

361   patterns. Int J Parasitol 34:1113–1119.

362   Yu, Y., and L. Nakhleh. 2015. A maximum pseudo-likelihood approach for phylogenetic networks. BMC Genomics

363   16:Suppl 10: S10.