1   **Title: Comparing rates of introgression in parasitic feather lice with differing**

2   **dispersal capabilities**

3   **Authors:** Jorge Doña[1]*, Andrew D. Sweet[1,2], and Kevin P. Johnson[1]*

Affiliations:

[1]Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-
Champaign, 1816 S. Oak St., Champaign, Illinois 61820, USA.

[2]Department of Entomology, Purdue University, 901 W. State St., West
Lafayette, Indiana 47907, USA.

*Correspondence to: jorged@illinois.edu, kpjohnso@illinois.edu.

Running title: Dispersal and introgression rate.

Author contributions: J.D., and K.P.J. conceived the study. J.D., A.D.S., and K.P.J designed
the study. A.D.S. collected the data. J.D. and A.D.S. analysed the data. K.P.J. obtained financial
support for the project. J.D. wrote the manuscript and all authors contributed to editing the
manuscript.

Acknowledgments: This study was supported by the US National Science Foundation (DEB-
1239788 and DEB-1342604 to K.P.J). We declare that we have no conflict of interest.

Data availability: All data needed to evaluate the conclusions in the paper are present in the
paper and/or the Supplementary Materials. Additional data related to this paper may be requested
from the authors.

Supplementary Materials: Supplementary data are available at Figshare (DOI:
10.6084/m9.figshare.9176204).
22

Abstract: Organisms vary in their dispersal abilities, and these differences can have important biological consequences, such as impacting the likelihood of hybridization events. However, there is still much to learn about the factors influencing hybridization, and specifically how dispersal ability affects the opportunities for hybridization. Here, using the ecological replicate system of dove wing and body lice (Insecta: Phthiraptera), we show that species with higher dispersal abilities exhibited increased genomic signatures of introgression. Specifically, we found a higher proportion of introgressed genomic reads and more reticulated phylogenetic networks in wing lice, the louse group with higher dispersal abilities. Our results are consistent with the hypothesis that differences in dispersal ability might drive the extent of introgression through hybridization. Overall, the results from this study represent an important step towards understanding the factors driving hybridization, and have major implications for coevolutionary biology.

Introduction

Dispersal is the permanent movement of organisms away from their place of origin. It is a fundamental process in biology with significant implications at multiple scales of organization (Barton 1992; Clobert et al. 2001; Nathan 2001; Matthysen 2012), including the reproduction of individuals, the composition of populations and communities, and the geographical distribution of species (Clobert et al. 2001, 2012).

Organisms differ in their dispersal abilities, and these differences have an impact on their biology, such as on the distributional range of a species or gene flow between populations (Bohonak 1999). For example, organisms with lower dispersal abilities tend to have smaller distributional ranges and populations that are genetically more structured (Bohonak 1999; Dawson et al. 2014; DiBlasi et al. 2018).

Dispersal ability might also affect the opportunities for hybridization between species because the rates at which individuals encounter different species are likely to be higher in organisms with higher dispersal capabilities. Indeed, recent evidence supports this prediction by demonstrating that range expansion is associated with the extent of introgression (Currat et al. 2008; Nussberger et al. 2018). Similarly, dispersal differences explain more than 30% of the variation in the width of hybrid zones across animals (McEntee et al. 2018). However, overall there is still much to learn about the factors influencing hybridization (Randler 2006; Arnold 2015; Taylor and Larson 2019), and, in particular, the influence of dispersal ability on the rate of hybridization remains understudied.

Comparisons of the effect of dispersal on hybridization should ideally hold constant most factors other than dispersal. The ecological replicate system of wing and body lice (Insecta: Phthiraptera) of pigeons and doves (Aves: Columbidae) has proven to be an ideal system for

comparing the impact of dispersal differences on other aspects of biology, such as population

structure and codivergence (Clayton and Johnson 2003; Johnson and Clayton 2004; Clayton et

al. 2015; DiBlasi et al. 2018; Sweet and Johnson 2018). Specifically, this is an excellent system

in which to assess the effect of differences in dispersal capabilities on levels of introgression

because both of these two lineages of feather lice: 1) significantly differ in their dispersal ability

(Harbison et al. 2008, 2009; Bartlow et al. 2016), 2) co-occur across the diversity of pigeons and

doves, 3) present highly comparable temporal patterns of diversification, specifically, both

lineages originated on the common ancestor of *Metriopelia* doves (11.3-14.9 mya) and have a

correlated pattern of codiversification within the same group of hosts (including a shared

cospeciation event which occurred within the *Metriopelia* genus 5.2-7.4 mya; Sweet and Johnson

2015, 2018), 3) and have the same basic life history and diet (Clayton et al. 2015; Sweet and

Johnson 2015, 2018). Both wing and body lice disperse vertically between parents and offspring

in the nest. However, wing lice can also attach to and hitchhike on hippoboscid flies to disperse

"phoretically" between host individuals or host species (Harbison et al. 2008, 2009; Bartlow et

al. 2016). Indeed, this hitch-hiking dispersal mechanism profoundly influences their degree of

population structure and cophylogenetic history (Clayton and Johnson 2003; Sweet et al. 2017b;

DiBlasi et al. 2018; Sweet and Johnson 2018). In addition, wing lice have a higher rate of host-

switching (Clayton and Johnson 2003; Clayton et al. 2015; Sweet et al. 2017b) (i.e., successful

colonization of new host species) and of straggling (Whiteman et al. 2004) (i.e., dispersal to new

host species without reproduction on that new host).

To compare differences in the extent of introgression between wing and body lice, we

used whole-genome data from 71 louse individuals belonging to five taxa of wing lice

(*Columbicola*) and seven taxa of body lice (*Physconelloides*) that occur across the same host

species and have highly comparable patterns of diversification (Sweet and Johnson 2015, 2018).

We predicted that wing lice, which have higher dispersal abilities and thus higher odds of

encountering individuals of a different louse species on the same host, should show more

extensive evidence of introgression (Fig. 1).

Figure 1. Diagram depicting the ecological replicate system and the hypothesis of this study. Wing lice (*Columbicola*) have

higher dispersal abilities than body lice (*Physconelloides*), and thus higher odds of encountering individuals of a different louse

sp.

Materials and Methods

Data

We studied Illumina whole genome sequence data (150 or 160bp paired-end reads) from 71

louse individuals belonging to five and seven taxa of *Columbicola* and *Physconelloides*,

respectively (Supplementary Table S2). All raw sequence data used were available from

previous studies (Boyd BM, et al. 2017; Sweet et al. 2017a; Sweet and Johnson 2018) and

represent all described New World ground-dove wing and body louse species–most host species

in this group–including lice samples across multiple biogeographic areas within species of hosts

(Sweet and Johnson 2018; Supplementary Table S2). Illumina genome sequence data pre-

processing included several steps (Sweet and Johnson 2018). First, we discarded duplicate read

pairs using the fastqSplitDups script (https://github.com/McIntyre-Lab/mcscriptand

https://github.com/McIntyre-Lab/mclib). We then eliminated the Illumina sequencing adapters

with Fastx_clipper v0.014 from the FASTX-Toolkit (http://hannonlab.cshl.edu/fastx_toolkit).

Also, we removed the first 5 nt from the 5' ends of reads using Fastx_trimmer v0.014 and

trimmed bases from the 3' ends of reads until reaching a base with a phred score $\geq$28 (which is

equivalent to a base call accuracy higher than 99.8 %) using Fastq_quality_trimmer v0.014.

Finally, we removed any reads less than 75 nt and analyzed the cleaned libraries with Fastqc

v0.11.5 to check for additional errors. We assembled nuclear loci in aTRAM following previous

studies (Allen et al. 2015; Boyd BM, et al. 2017; Sweet and Johnson 2018). In particular, we

mapped modest coverage (25-60X), multiplexed genomic data to reference loci from a closely

related taxon. For our reference set of nuclear loci for wing lice, we used 1,039 exons of

Columbicola drowni (Boyd BM, et al. 2017) (raw data: SRR3161922). This data set was

assembled de novo (Allen et al. 2015) using orthologous protein-coding genes from the human

body louse genome (*Pediculus humanus humanus* (Kirkness et al. 2010) as a set of target

sequences. We mapped *Columbicola* reads to the *C. drowni* references using Bowtie2

(Langmead and Salzberg 2012). For body lice, we obtained nuclear data using the same pipeline

and software parameters, except that we used 1,095 loci from *Physconelloides emersoni* as the

reference for mapping. To generate the input ultrametric gene trees for Phylonet v3.6.8 (Than et

al. 2008; Yu and Nakhleh 2015; Wen et al. 2018), we first aligned each nuclear locus in MAFFT

(Katoh 2002)(--auto) and removed columns with only ambiguous sequences ("N"). Then, we

estimated gene trees in RAxML v8.1.3 (Stamatakis 2006) with a GTR + Γ substitution model for

each gene alignment. Finally, we made trees ultrametric using the nnls method in the

force.ultrametric function within the "phytools" R package (Revell 2011).

<u>Quantifying introgression</u>

We used two different approaches to quantify differences in the extent of introgression (i.e.,

ancient plus recent) between the two louse genera. We employed methods suitable to detect

introgression between species and between individuals from the same species (i.e., we did not

employ methods aimed to detect differences at the population level, e.g., TreeMix; Pickrell and

Pritchard 2012). First, we used sppIDer (Langdon et al. 2018) to quantify the genomic

contributions of different louse species in an individual louse genome (Fig. 2). We built our

reference for each genus using all the nuclear loci from a single individual per species. For the

reference, we selected those individuals for which we assembled the highest number of genes for

each genus. We estimated the extent of introgression as the sum of the mean coverages of reads

mapped from all the species excluding the focal louse species, divided by the mean coverage of

the focal louse species (Fig. 2). Note that these mean coverage values are calculated using only

those reads that mapped with a mapping quality (MQ) > 3 (Li et al. 2009; Langdon et al. 2018;

Figs. S1-S12).

Figure 2. Illustrative example of sppIDer results with an individual of *Columbicola* (Cosp.Copas.11.7.2016.8). Panel a) shows the average mean coverages of reads mapping to every species (i.e., the values that were used for the calculations of introgression levels), and panel b) shows the mean coverages of reads mapping to each species across the whole set of loci. These and additional visualizations of all the individuals can be found in the supplementary material (Figs. S1-S12).

Second, we quantified introgression at the species level, while accounting for incomplete lineage

sorting (ILS) (i.e., reticulations in this method can be attributed to hybridization events), using a

maximum pseudo-likelihood framework with PhyloNet 3.6.1 (Than et al. 2008; Yu and Nakhleh

2015; Wen et al. 2018). We trimmed the unrooted gene trees to the same individuals used as

reference taxa in sppIDer, and performed eleven independent analyses with a differing maximum

number of reticulation nodes (i.e., from zero to ten). We conducted ten runs per analysis. We

then selected the optimal network for each genus based on AIC values.

<u>Statistics and reproducibility</u>

We compared the sppIDer results using generalized linear models (GLMs). We used a Gaussian

distribution of errors and an identity link function. We performed one GLM for each simulation

iteration using the glm function of the "stats" R package (R Core Team 2013). The extent of

introgression for each louse genus was the dependent variable, the genus identity was the

independent variable, and we accounted for the introgression differences between louse species

including louse identity as a fixed factor. We confirmed assumptions underlying GLMs by

testing the normality of regression residuals for normality against a Q-Q plot. We also

considered the possibility that some of the reads mapping to other species were technical

contaminations, i.e., due to index-swapping (Carlsen et al. 2012; Esling et al. 2015; Schnell et al.

2015; Sinha et al. 2017). Previous studies have found that the misassignment of reads generally

ranges from 1 to 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al.

2017). Thus, to account for possible contaminants, we wrote a simulation in R that randomly

subtracted 9% (Carlsen et al. 2012; Esling et al. 2015; Schnell et al. 2015; Sinha et al. 2017)

from the mean coverage value of a particular sample (i.e., we subtracted a random proportion of

the mean coverage value for each sample until reaching 9 %). We ran 100 iterations of the

simulation and ran a GLM for each iteration (Table S1). Finally, we used the χ^2 test to compare

the number of species in pairwise comparisons of each genus with the number of reticulations

found in each optimal phylogenetic network. Because we had an *a priori* prediction that

Physconelloides should exhibit less evidence of reticulation than *Columbicola,* we used a one-

tailed test; however, we also report the results of the two-tailed test equivalent.

Results

Both approaches revealed highly concordant results: higher levels of introgression among

species of wing lice compared to body lice. In particular, using a read-mapping based method,

the genomic signature of introgression was significantly higher in wing louse species than in

body louse species (GLM with the mean values of the simulations; F = 21.0705, df = 69, *P* =

2.367 x 10^{-5}, R^2=0.58; Fig. 3, Supplementary Table S1, Figs. S1-S12). Contrary to this effect,

one body louse individual (included in the GLMs) exhibited the highest level of introgression

(Fig. 3, Figs. S1-S12). However, the other individual from the same taxon, inhabiting the same

host species and collected in the same geographic region, did not show these elevated levels of

introgression (Supplementary Table S2, Figs. S1-S12).

Figure 3. Boxplot showing the differences in levels of introgression between wing (green) and body (orange) lice. Level of introgression represents the sum of the mean coverage of reads mapped from all the species excluding the focal louse species, divided by the mean coverage of the focal louse species (see Methods). Black dots show the levels of introgression (i.e., resulting from the equation) for each individual sample (horizontally jittered values).

Secondly, in a phylogenetic network framework, the optimal networks of wing lice were more

reticulated than those of body lice even though the number of taxa included in the networks was

lower (seven reticulations in *Columbicola* vs. four in *Physconelloides*, Fig. 4). Accordingly, the

number of reticulations given the number of potential combinations under a one-tailed test was

significantly higher in *Columbicola* (One-sided: χ^2= 3.8132; df=1; *P*= 0.03; CI=0.03–1), and the

p-value was still near 0.05 with a two-tailed test (Two-sided: χ^2= 3.8132; df=1; *P*= 0.05; CI=-

0.01–0.66). Also, the specific lineages involved in the reticulations were generally congruent

with signatures of introgression from the read-mapping based approach (Fig. S1-S12).

Figure 4. Optimal phylogenetic networks of feather lice genera. Orange branches depict reticulations: seven in *Columbicola* and

four in *Physconelloides*. Note that though visually overlapped, every reticulation in this figure involves at least two taxa (extant

or extinct). See the raw unedited networks (Figs. S13-S14) for more details on specific reticulations.

Discussion

Estimates of introgression in two groups of ectoparasites that differ in their dispersal abilities,

wing and body lice of doves, indicate that the lineage with higher dispersal ability (wing lice)

shows more evidence of introgression. This evidence from wing and body louse genomes is

consistent with the hypothesis that dispersal differences might drive differences in the level of

introgression in this system of parasites. Admittedly, there may be some unknown factor, other

than dispersal, differing between these two groups of lice that causes the difference in the level

of introgression, but prior work on these groups of parasites points to dispersal as a crucial factor

underlying many of the ecological and evolutionary patterns in these parasites. Further research

on other taxa is needed to confirm the generality of these findings. This work is among the first

studies of introgression in a host-symbiont system (Detwiler and Criscione 2010). Notably,

recent studies have found that straggling and host-switching are relatively common processes in

host-symbiont systems (De Vienne et al. 2013; Bourguignon et al. 2018; Nylin et al. 2018; Doña

et al. 2019). Our study suggests that in a straggling/host-switching scenario, hybridization can

provide further genetic variation with important ecological and evolutionary consequences (e.g.,

facilitating adaptation to current hosts or facilitating the colonization of new ones) (Barton

2018). Indeed, we may have found a potential recent hybridization event (i.e., the

Physconelloides individual showing the highest level of introgression), though this requires

further study to rule out methodological issues (e.g., wet-lab contamination).

In this vein, a careful examination of the introgression history of these taxa (and

symbionts as a whole) is needed to better understand the patterns of introgression that we found.

Questions such as how much introgression can be expected or how the introgressed regions are

retained in parasite/symbiont genomes across time, among many others, require further attention

and will certainly have a strong impact in coevolutionary biology theory. For instance, in this

study, the levels of introgression detected by the sppIDer analyses (i.e., the magnitude but not the

comparative pattern) may be unrealistic. It may be that some fraction of the level of introgression

detected by sppIDer may be due to ILS, and not introgression. However, both louse genera are

expected to have relatively similar rates of ILS (if any), and our PhyloNet analysis, which does

control for ILS, showed highly congruent results. On the other hand, sppIDer can detect

introgression from species that are not included in the reference data (and, in those cases, the

reads map to the closest taxon available in the reference, and thus could artificially increase the

level of introgression from a given species; Langdon et al. 2018). Accordingly, the levels of

introgression detected by sppIDer in certain species could be an aggregate of introgression

events from more than one species. Indeed, our PhyloNet analysis supports this scenario, with

several reticulations from ghost lineages and species (Figs. S13-S14), although in this system we

have nearly complete sampling of host taxa and are missing few, if any, extant species.

Overall, the results from this study represent an important step towards understanding the

factors driving hybridization, because most previous studies focus on the presence/absence of

hybridization and the evolutionary consequences of hybridization (Arnold 2015; Folk et al. 2018;

Taylor and Larson 2019). Further research is needed to understand the factors shaping the

frequency of hybridization and to ascertain their influence across different scales (e.g., from

ancient to recent hybridization events).

References:

Allen, J. M., D. I. Huang, Q. C. Cronk, and K. P. Johnson. 2015. aTRAM - automated target restricted assembly

method: a fast method for assembling loci across divergent taxa from next-generation sequencing data.

BMC Bioinformatics 16.

Arnold, M. L. 2015. Divergence with genetic exchange. OUP Oxford, Oxford, UK.

Bartlow, A. W., S. M. Villa, M. W. Thompson, and S. E. Bush. 2016. Walk or ride? Phoretic behaviour of

248   amblyceran and ischnoceran lice. Int J Parasitol 46:221–227.

249   Barton, N. H. 2018. The consequences of an introgression event. Mol Ecol 27:4973–4975.

250   Barton, N. H. 1992. The genetic consequences of dispersal. In: Animal Dispersal. Springer, Netherlands.

251   Bohonak, A. J. 1999. Dispersal Gene Flow, and Population Structure. Q Rev Biol 74:21–45.

252   Bourguignon, T., N. Lo, C. Dietrich, J. Šobotník, S. Sidek, Y. Roisin, A. Brune, and T. A. Evans. 2018. Rampant

253   host switching shaped the termite gut microbiome. Curr Biol 28:649–654.

254   Boyd BM, Allen JM, Nguyen N, Sweet AD, Warnow T, Shapiro MD, and et al. 2017. Phylogenomics using Target-

255   restricted Assembly Resolves Intra-generic Relationships of Parasitic Lice (Phthiraptera: Columbicola).

256   Syst Biol 66:896–911.

257   Carlsen, T., A. B. Aas, D. Lindner, T. Vrålstad, T. Schumacher, and H. Kauserud. 2012. Dont make a mista(g)ke: is

258   tag switching an overlooked source of error in amplicon pyrosequencing studies? Fungal Ecol 5:747–749.

259   Clayton, D. H., S. E. Bush, and K. P. Johnson. 2015. Coevolution of life on hosts: integrating ecology and history.

260   University of Chicago Press, Chicago, IL.

261   Clayton, D. H., and K. P. Johnson. 2003. Linking coevolutionary history to ecological process: doves and lice.

262   Evolution 57:2335–2341.

263   Clobert, J., M. Baguette, T. G. Benton, and J. M. Bullock (eds). 2012. Dispersal Ecology and Evolution. Oxford

264   University Press.

265   Clobert, J., E. Danchin, A. A. Dhondt, and N. J. D. 2001. Dispersal. Oxford Univ. Press.

266   Currat, M., M. Ruedi, R. J. Petit, and L. Excoffier. 2008. The hidden side of invasions: massive introgression by

267   local genes. Evolution 62:1908–1920.

268   Dawson, M. N., C. G. Hays, R. K. Grosberg, and P. T. Raimondi. 2014. Dispersal potential and population genetic

269   structure in the marine intertidal of the eastern North Pacific. Ecol Monogr 84:435–456.

270   De Vienne, D. M., G. Refrégier, M. López-Villavicencio, A. Tellier, M. E. Hood, and T. Giraud. 2013. Cospeciation

271   vs host-shift speciation: methods for testing, evidence from natural associations and relation to coevolution.

272   New Phytol 198:347–385.

273   Detwiler, J. T., and C. D. Criscione. 2010. An infectious topic in reticulate evolution: introgression and

274   hybridization in animal parasites. Genes (Basel) 1:102–23.

DiBlasi, E., K. P. Johnson, S. A. Stringham, A. N. Hansen, A. B. Beach, and D. H. Clayton. 2018. Phoretic dispersal

influences parasite population genetic structure. Mol Ecol 27:2770–2779.

Doña, J., D. Serrano, S. Mironov, A. Montesinos-Navarro, and R. Jovani. 2019. Unexpected bird–feather mite

associations revealed by DNA metabarcoding uncovers a dynamic ecoevolutionary scenario. Mol Ecol

28:379–390.

Esling, P., F. Lejzerowicz, and J. Pawlowski. 2015. Accurate multiplexing and filtering for high-throughput

amplicon-sequencing. Nucleic Acids Res 43:2513–24.

Folk, R. A., P. S. Soltis, D. E. Soltis, and R. Guralnick. 2018. New prospects in the detection and comparative

analysis of hybridization in the tree of life. Am J Bot 105:364–375.

Harbison, C. W., S. E. Bush, J. R. Malenke, and D. H. Clayton. 2008. Comparative transmission dynamics of

competing parasite species. Ecology 89:3186–3194.

Harbison, C. W., M. V. Jacobsen, and D. H. Clayton. 2009. A hitchhiker's guide to parasite transmission: The

phoretic behaviour of feather lice. Int J Parasitol 39:569–575.

Johnson, K. P., and D. H. Clayton. 2004. Untangling coevolutionary history. Syst Biol 53:92–4.

Katoh, K. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform.

Nucleic Acids Res 30:3059–3066.

Kirkness, E. F., B. J. Haas, W. Sun, H. R. Braig, M. A. Perotti, and J. M. Clark. 2010. Genome sequences of the

human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle.

Proc Natl Acad Sci U S A 107:12168–73.

Langdon, Q. K., D. Peris, B. Kyle, and C. T. Hittinger. 2018. sppIDer: A Species Identification Tool to Investigate

Hybrid Genomes with High-Throughput Sequencing. Mol Biol Evol 35:2835–2849.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357–9.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, and R. Durbin. 2009. The

sequence alignment/map format and SAMtools. Bioinformatics 25:2078–2079. Oxford University Press.

Matthysen, E. 2012. Multicausality of dispersal: a review. In: Dispersal Ecology and Evolution. Oxford University

Press.

McEntee, J. P., J. G. Burleigh, and S. Singhal. 2018. Dispersal predicts hybrid zone widths across animal diversity:

Implications for species borders under incomplete reproductive isolation. bioRxiv, doi: 10.1101/472506.

Nathan, R. 2001. The challenges of studying dispersal. Trends Ecol Evol 16:481–483.

Nussberger, B., M. Currat, C. S. Quilodran, N. Ponta, and L. F. Keller. 2018. Range expansion as an explanation for
305   introgression in European wildcats. Biol Conserv 218:49–56.

306   Nylin, S., S. Agosta, S. Bensch, W. A. Boeger, M. P. Braga, D. R. Brooks, M. L. Forister, P. A. Hambäck, E. P.
307   Hoberg, and T. Nyman. 2018. Embracing colonizations: a new paradigm for species association dynamics.
308   Trends Ecol Evol 33:4–14.

309   Pickrell, J. K., and J. K. Pritchard. 2012. Inference of Population Splits and Mixtures from Genome-Wide Allele
310   Frequency Data. PLoS Genetics 8.

311   R Core Team. 2013. R: A language and environment for statistical computing.

312   Randler, C. 2006. Behavioural and ecological correlates of natural hybridization in birds. Ibis 148:459–467.

313   Revell, L. 2011. phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol
314   Evol 3:217–223.

315   Schnell, I. B., K. Bohmann, and M. T. Gilbert. 2015. Tag jumps illuminated–reducing sequence-to-sample
316   misidentifications in metabarcoding studies. Mol Ecol Resour 15:1289–303.

317   Sinha, R., G. Stanley, G. S. Gulati, C. Ezran, K. J. Travaglini, and E. Wei. 2017. Index Switching Causes Spreading-
318   Of-Signal Among. bioRxiv, doi: 10.1101/125724.

319   Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa
320   and mixed models. Bioinformatics 22:2688–2690.

321   Sweet, A. D., B. M. Boyd, J. M. Allen, S. M. Villa, M. P. Valim, and J. L. Rivera-Parra. 2017a. Integrating
322   phylogenomic and population genomic patterns in avian lice provides a more complete picture of parasite
323   evolution. Evolution 72:95–112.

324   Sweet, A. D., R. T. Chesser, and K. P. Johnson. 2017b. Comparative cophylogenetics of Australian phabine pigeons
325   and doves (Aves: Columbidae) and their feather lice (Insecta: Phthiraptera). Int J Parasitol 47:347–356.

326   Sweet, A. D., and K. P. Johnson. 2015. Patterns of diversification in small New World ground doves are consistent
327   with major geologic events. Auk 132:300–312.

328   Sweet, A. D., and K. P. Johnson. 2018. The role of parasite dispersal in shaping a host–parasite system at multiple
329   evolutionary scales. Mol Ecol 27:5104–5119.

Taylor, S. A., and E. L. Larson. 2019. Insights from genomes into the evolutionary importance and prevalence of

hybridization in nature. Nat Ecol Evol 3:170.

Than, C., D. Ruths, and L. Nakhleh. 2008. PhyloNet: a software package for analyzing and reconstructing reticulate

evolutionary relationships. BMC Bioinformatics 9:322.

Wen, D., Y. Yu, J. Zhu, and L. Nakhleh. 2018. Inferring Phylogenetic Networks Using PhyloNet. Syst Biol 67:735–

740.

Whiteman, N. K., D. Santiago-Alarcon, K. P. Johnson, and P. G. Parker. 2004. Differences in straggling rates

between two genera of dove lice (Insecta: Phthiraptera) reinforce population genetic and cophylogenetic

patterns. Int J Parasitol 34:1113–1119.

Yu, Y., and L. Nakhleh. 2015. A maximum pseudo-likelihood approach for phylogenetic networks. BMC Genomics

16:Suppl 10: S10.