1
2 **Removal of Spo11 from meiotic DNA breaks *in vitro* but not *in vivo* by Tyrosyl DNA**
3 **Phosphodiesterase 2**
4

5 Dominic Johnson[1], Rachal M Allison[1], Elda Cannavo[2,3], Petr Cejka[2,3], and Matthew J
6 Neale[1,4]
7

8
9 **AFFILIATIONS**
10
1. Genome Damage and Stability Centre, School of Life Sciences, University of Sussex, Brighton UK
2. Institute for Research in Biomedicine, Università della Svizzera italiana, Bellinzona 6500,
Switzerland
3. Department of Biology, Institute of Biochemistry, Eidgenössische Technische Hochschule (ETH)
Zurich, Zurich 8093, Switzerland
4. To whom correspondence should be addressed: m.neale@sussex.ac.uk
17

ABSTRACT

Meiotic recombination events are initiated by DNA double-strand breaks (DSBs) created by

the topoisomerase-like protein, Spo11. Similar to type-II topoisomerases, Spo11 becomes

covalently linked to the 5′ ends generated on each side of the DSB. Whilst Spo11-oligos—the

product of nucleolytic removal by Mre11—have been detected in a number of biological

systems, the lifetime of the covalent Spo11-DSB precursor has not been systematically

determined and may be subject to alternative processing reactions. Here we explore the

activity of human Tyrosyl DNA Phosphodiesterase, TDP2, on Spo11-DSBs isolated from *S.*

cerevisiae cells. We demonstrate that TDP2 can remove Spo11 from natural ssDNA-oligos,

and dsDNA ends even when in the presence of excess competitor genomic DNA.

Interestingly, TDP2-processed Spo11-DSBs are refractory to resection by Exo1, suggesting

that ssDNA generated by Mre11 may be essential *in vivo* to facilitate resection-dependent HR

at Spo11-DSBs even if TDP2 were active. Moreover, although TDP2 can remove Spo11

peptides *in vitro*, TDP2 was unable to remove Spo11 *in vivo*—unlike during the repair of

topoisomerase-induced DNA lesions. These results suggest that Spo11-DNA, but not

topoisomerase-DNA cleavage complexes, are inaccessible to the TDP2 enzyme, perhaps due

to occlusion by higher order protein complexes resident at sites of meiotic recombination.

INTRODUCTION

In most organisms, meiotic recombination is essential to create genetically diverse, haploid

genomes suitable for sexual reproduction. In meiosis, homologous recombination (HR)

events are initiated by DNA double-strand breaks (DSBs) created by Spo11, a topoisomerase-

like enzyme most closely related to the catalytic 'A' subunit of the archael type-IIB

Topoisomerase VI (1–3). TopoVI functions as an A_2B_2 heterotetramer, and the recent

identification of the eukaryotic 'B' subunit, Top6BL, in mouse, plants and budding yeast

indicates that the active form of Spo11 is similarly heterotetrameric (4, 5). Like other type-II

and type-IIB topoisomerases, the Spo11 monomers become covalently attached via a

phosphotyrosine bond to the 5′ ends generated on each side of the DSB during cleavage (6–

8). In wildtype *S. cerevisiae* cells, such covalent Spo11-DSBs are transient, being rapidly

processed by the nuclease activity of the MRX/N complex (Mre11, Rad50 and Xrs2/Nbs1),

which is stimulated by Sae2, the yeast orthologue of human CtIP (9–14). This coordinated

reaction generates a covalent single-stranded Spo11-oligonucleotide complex, flanked by a

nick or short ssDNA gap on the 5′-ending strand that enables the DSB to be subsequently

channelled into homologous recombination repair via the onset of long-range ssDNA

resection catalysed by Exo1 (9, 15–17). Spo11-oligo complexes are formed in an

evolutionarily divergent yeast, *Schizosaccharomyces pombe* (18), and are also detectable in

mouse spermatocytes (10, 19), suggesting that nucleolytic release is the default mode of

Spo11 removal in both unicellular and multicellular eukaryotes.

However, whilst the product of this nucleolytic reaction—Spo11-oligos—have been detected

in numerous biological systems, the lifetime of the covalent DSB precursor has not been

systematically determined and may be subject to alternative processing reactions. For

example, in *S. pombe*—and unlike in *S. cerevisiae* (2)—covalent Spo11-DSBs are readily

detected in wildtype cells (20), indicating that DSB formation and the initiation of repair can

be temporally separated. Moreover, inactivation of the canonical Spo11 release reaction in *C.*

elegans (via mutation of COM-1, the CtIP/Sae2 orthologue) results in repair of DSBs by the

non-homologous endjoining (NHEJ) pathway and aberrant chromosome segregation (21, 22).

These observations suggest that delayed onset of meiotic DSB resection may be especially

problematic for meiotic cells, and further suggest there must be a COM-1-independent

mechanism to remove Spo11 in *C. elegans*, such that the DSBs can be ligated using the end-

joining apparatus.

In vegetative cells, the usually benign and transient protein-linked DNA breaks created by

Topoisomerase I (Top1) and II (Top2) as part of their reversible catalytic cycle may become

a stable and toxic covalent-DNA intermediates (23). This conversion is exploited in

chemotherapy by the use of drugs such as camptothecin and etoposide, which stabilise the

covalent cleavage-complex intermediate (Top1cc and Top2cc respectively), thereby

sensitising cells that have a relatively greater reliance on topoisomerase activity compared to

other tissue. Presumably because such topoisomerase failures are relatively common even in

unchallenged cells, two classes of enzymes exist that are capable of handling such lesions:

Tyrosine phosphodiesterase 1 (Tdp1), which primarily resolves the 3′ covalent

phosphotyrosine links created by Top1 (24), and tyrosine phosphodiesterase 2 (TDP2, also

named TTRAP), which primarily resolves 5′ phosphotyrosine links created by Top2 (25).

Some enzymatic overlap between the activities and substrates of Tdp1 and TDP2 exists,

suggesting some functional redundancy *in vivo* (25, 26). Nevertheless, the primary activity of

Tdp1 activity aids the conversion of Top1 lesions into ligatable single-strand breaks (SSBs),

whereas TDP2 converts Top2 lesions into DSB ends suitable for direct repair by the NHEJ

pathway. The reaction product created by TDP2 contrasts with the 3′ single-stranded ends

generated as a result of MRX/N activity, and which are suitable for HR (9, 15, 27).

Interestingly, TDP2 is expressed in mouse and human testes tissue (28), leading to the

possibility that TDP2 may be able to process a subset of Spo11-DSB intermediates.

Here we use ectopic expression and *in vitro* biochemistry to explore the activity of human

TDP2 on Spo11-DSBs generated in *S. cerevisiae* meiotic cells. Collectively, our results

indicate that while TDP2 is catalytically capable of removing Spo11 from both Spo11-oligos

and Spo11-DSB ends *in vitro*, TDP2 is unable to do so *in vivo*. This is despite TDP2

functioning efficiently to suppress the sensitivity of *mre11* or *sae2* mutants to topoisomerase

poisons. These results suggest that Spo11-DSBs, but not Topoisomerase-DSBs, are

inaccessible to the TDP2 active site, perhaps due to occlusion by higher order protein

complexes resident at sites of meiotic recombination. Moreover, we find that the 2 nt 5′

extensions revealed when Spo11 is removed from DSB ends by TDP2 *in vitro* are refractory

to nucleolytic resection by Exo1, suggesting that even were TDP2 to act during meiotic DSB

repair, it would nonetheless lead to DSBs being channelled away from the essential process

of homologous recombination. Overall, our findings provide insight into the mechanisms

involved in the resection and repair of protein-linked DSB ends created by topoisomerase-

like enzymes such as Spo11, and how they may differ to the repair of 'clean'-ended DNA

breaks created by site-specific nucleases.

RESULTS

TDP2 removes Spo11 protein and peptides from single-stranded Spo11-oligo complexes.

Meiotic recombination is initiated by DSBs created by the evolutionarily conserved Spo11

protein. As part of the catalytic reaction, Spo11 becomes covalently attached to the 5′ DSB

ends via a phosphotyrosyl bond. In *M. musculus*, *S. cerevisiae*, and *S. pombe*, Spo11 is

released from the DSB ends by the action of the evolutionarily conserved MRX/N nuclease

complex in conjunction with Sae2/CtIP/Ctp1, generating covalent Spo11-oligo complexes (9,

10, 18, 19, 29). However, whether alternative mechanisms of Spo11 end-processing are

possible has not been investigated.

To investigate whether Spo11-DSBs may also be a substrate for processing by TDP2—as are

Top2-induced DSBs (25)—we took advantage of the *S. cerevisiae* meiotic system in which

synchronised populations of cells undergo Spo11-DSB formation at hundreds of sites spread

across the genome (termed Spo11 hotspots), but in which TDP2 is not naturally expressed.

We first tested for the ability of recombinant TDP2 to remove Spo11 covalently attached to

ssDNA by enriching for the natural covalent Spo11-oligo complexes created by MRX and

Sae2-dependent processing of meiotic Spo11-DSBs in wildtype cells (Fig 1a). Cells in mid

meiotic prophase (4 hours after meiotic entry) were harvested, lysed, and Spo11-oligo

complexes isolated by immunoprecipitation using antibodies raised against a C-terminal

triple FLAG epitope (30).

Immunoprecipitates were washed and then the ssDNA component labelled at the terminal 3′

hydroxyl group using terminal deoxynucleotide transferase (TdT) and a radiolabelled

nucleotide (dCTP). Under the conditions employed, this reaction results in 1-4 radiolabelled

nucleotides being added to the 3′ OH end of the Spo11-oligo complex (D. Johnson and M.

Neale unpub. obs), which can be resolved via SDS-PAGE (Fig 1b). Recombinant human

TDP2 (25) was reacted with these "full-length" labelled Spo11-oligo complexes, or with

complexes pre-treated with either trypsin or proteinase K (Fig 1c-f). Based on substrate

specificity, trypsin is expected to leave twelve amino acid residues covalently attached to the

oligo complex via the phosphotyrosine bond, whereas proteinase K treatment will degrade all

but three amino acids in addition to the phosphotyrosine.

Following incubation with increasing concentrations of TDP2, reactions were halted by

adding formamide gel-loading buffer, then resolved on a 15% polyacrylamide gel under

DNA-denaturing conditions (Fig 1d-f). Under these conditions, untreated Spo11-oligo

complexes fail to enter the gel and are often lost from the wells during gel handling and

fixation (Fig 1d). By contrast, trypsin-treated molecules migrate as a heterogeneous smear,

presumably due to significant electrophoretic retardation caused by the remaining amino

acids bound to the DNA (Fig 1e). Proteinase K treated samples migrate more uniformly, as a

pair of broad peaks, consistent with the previously characterised dual size classes of Spo11-

oligo complex detected in *S. cerevisiae* (8-15 nt and 25-40 nt; Fig 1f; (9, 10)).

Consistent with experiments performed using synthetic phosphotyrosine ssDNA substrates

(25) or partially proteolysed Top2-DNA covalent complexes (31), we observed robust

activity of TDP2 against both the partially and fully proteolysed Spo11-oligo substrates (Fig

1e,f). In both cases, we observed the conversion of the radiolabelled signal to faster migrating

forms consistent with cleavage of the 5′ phosphotyrosine bond and complete removal of any

residual amino acids from the radiolabelled oligo.

Importantly, incubation of TDP2 with a radiolabelled oligononucleotide (i.e. without a

covalent protein attachment) resulted in no change in electrophoretic migration (Fig S1a),

confirming that the increased rate of migration observed above (Fig 1e,f) was not due to trace

nuclease activity within the TDP2 preparation, but instead was due to the expected cleavage

of the protein-DNA phosphotyrosyl bond.

Structural modelling of TDP2 indicates that the catalytic site is buried within the core of the

protein, which any phosphotyrosine substrate would be required to thread into to be cleaved

(32–34). Intriguingly, however, we also observed conversion of non-migrating to rapidly

migrating oligo complexes when TDP2 was reacted with full-length Spo11-oligos, albeit only

restricted to the reactions containing the higher enzyme concentrations (Fig 1d). These

observations support the view that TDP2 can remove both full-length and Spo11 protein

fragments from ssDNA ends, and that prior proteolytic degradation of the 5′-covalently

linked protein stimulates the enzymatic activity of TDP2, similar to what has been observed

for its activity on Top2-DNA intermediates (31).

TDP2 removes Spo11 peptides from DSB ends.

We next tested the ability of TDP2 to remove Spo11 from the naturally occurring double-

stranded DNA (dsDNA) ends generated during meiosis. As introduced above, in wildtype *S.*

cerevisiae cells, Spo11-DSBs are normally rapidly processed by MRX/Sae2 (liberating

Spo11-oligo complexes) and then subjected to Exo1-dependent resection to create long 3′

ending ssDNA tails. Thus, to enrich for a population of covalent Spo11-DSB molecules to

use as a substrate, and therefore to prevent such nucleolytic processing steps, we prepared

genomic DNA samples from *sae2Δ* cells in mid meiotic prophase (4 hours after meiotic

induction when DSBs are abundant). The resulting nucleic acid material is free from non-

covalently attached protein contaminants but, due to the *SAE2* gene deletion, retains covalent

Spo11-DSB intermediates (8, 35, 36). Samples were then incubated with and without TDP2

for up to 15 minutes, then subsequently reacted with lambda exonuclease—a 5′-3′

exonuclease with activity that depends on the presence of an unmodified 5′ phosphate group

on the substrate (37). Post-treatment with lambda exonuclease thus provides a sensitive test

for TDP2-dependent removal of the Spo11 moiety from the DSB end (Fig 2a).

To monitor the reactions, DNA was digested with *Pst*I, separated on an agarose gel,

transferred to nylon membrane and hybridised with a radiolabelled DNA probe that

recognises the *HIS4*::*LEU2* meiotic recombination hotspot (38). Preparation in this way

reveals three distinct bands, the upper most is the parental DNA molecule, and the lower

smaller bands are indicative of DSBs arising at the two strong hotspots present at this

genomic locus (Fig 2b). Treatment with lambda exonuclease alone resulted in no change in

the migration of the DSB bands—as expected due to the attached Spo11 protein preventing

5′-3′ resection (Fig 2b). By contrast, pretreatment with TDP2 led to complete disappearance

of the DSB bands—as expected if TDP2 was able to efficiently remove Spo11 from the DSB

end creating a clean 5′ phosphorylated substrate on which lambda exonuclease can act (Fig

2b). Notably, the parental DNA band was stable under these conditions indicating that the

exonuclease degradation was specific to the deprotected Spo11-DSB ends (Fig 2b). If we

assume there to be approximately 160 DSBs (39) within each replicated *S. cerevisiae* diploid

genome (~50 Mbp) there will be approximately one covalent Spo11-DSB end for every ~150

kb of genomic DNA. We conclude that recombinant TDP2 is able to efficiently remove

Spo11 from dsDNA ends even in the presence of a large excess of dsDNA.

Exo1 is unable to resect Spo11-DSBs that have been processed by TDP2.

As described above, the endogenous exonuclease involved in 5′-3′ resection of Spo11-DSBs

is Exo1 (9, 15, 17). We were thus interested to examine whether treatment of Spo11-DSBs

with TDP2 could enable resection by recombinant Exo1, perhaps indicating a potential

mechanism to bypass the requirement of MRX/N nuclease activity *in vivo*. To perform these

experiments, we isolated genomic DNA from *sae2Δ* cells as described above, incubated with

and without TDP2 for up to 15 minutes, then incubated with recombinant Exo1 rather than

lambda exonuclease. Unlike treatment with lambda exonuclease, Exo1 was unable to initiate

resection on Spo11-DSBs whether or not they had been processed by TDP2 (Fig 3a,b).

As a positive control to ensure that this inability to resect the Spo11-DSB end was not due to

a problem with our recombinant Exo1 preparation, we repeated the same set of assays with

genomic DNA isolated from an *exo1Δ* mutant (rather than *sae2Δ*). Under these conditions,

MRX and Sae2-mediated nucleolytic cleavage of Spo11 from the DSB *in vivo* results in a

short 3′ ssDNA tail (up to ~300 bp) (15). As expected, incubation of this substrate with

recombinant Exo1 resulted in specific degradation of the DSBs whether or not they were first

treated with TDP2 (Fig 3c,d).

As a final control, we confirmed that TDP2 was active in the Exo1 buffer by incubating the

Spo11-oligo substrate with TDP2 under the same conditions that the Exo1 resection assays

were performed (Fig S1b). Collectively these results demonstrate that whilst TDP2 may have

the necessary biochemical activity to remove covalently attached Spo11 from DSB ends, the

resulting clean DNA ends are a poor substrate for Exo1 to initiate resection.

Exo1 is unable to resect Spo11-DSBs that have been blunted by Klenow fragment.

Spo11 creates a 2 nt 5′ overhang at the DSB ends (7, 39, 40). Prior *in vitro* assays using

relatively short (~3.0 kb) plasmid fragments indicated that the preferred substrate for Exo1 is

a dsDNA end with a 3′ extension, similar in form to the resected DSBs created by

MRX/Sae2, and in agreement with our observations presented above (41). By contrast, Exo1

displayed reduced, but not abolished, activity on substrates with a 4 nt 5′ extension, and only

moderately reduced activity on a blunt ended substrate (41). These latter findings contrast

somewhat with our observations that even just a 2 nt 5′ overhang completely prevents Exo1-

catalysed resection of the TDP2-processed Spo11-DSBs (Fig 3a,b). A possible explanation

for the difference may be the ratio of dsDNA to DSB ends, which is ~100-fold greater in our

assay system—and a much closer mimic of the conditions present *in vivo* during meiosis.

To determine whether Exo1 was able to resect blunt DSB ends under these same 'in vivo-

like' conditions, we repeated the experiments utilising the TDP2-processed *sae2Δ* genomic

DNA, but this time including a preincubation step with Klenow fragment and dNTPs to blunt

the 2 nt 5′ overhang prior to incubation with Exo1 (Fig 3g,h). Unlike for the partially resected

substrates, no resection by Exo1 was detectable even when DSB ends were first blunted by

Klenow fragment (Fig 3g,h). Taken together our results suggest that Exo1 is acutely sensitive

to the structure of the DNA end, with partially resected ends—such as those created by

MRX/Sae2-dependent processing—being a significantly favoured substrate when excess

competitor dsDNA substrate is present. We suggest that our observations may represent a

closer match to the *in vivo* situation, where—even in meiosis—DSB ends are relatively

infrequent compared to the intact double-stranded genomic DNA.

Ectopic expression of TDP2 is unable to cleave Spo11 from DSB ends in *S. cerevisiae.*

An orthologue of the TDP2 protein is not present in *S. cerevisiae*. However, human TDP2

was identified via its ability to rescue the sensitivity of yeast cells to camptothecin (a Top1

poison), and to also suppress etoposide sensitivity (a Top2 poison) indicating that human

TDP2 is active when expressed in *S. cerevisiae* cells (25). Given our findings that TDP2 has

the appropriate biochemical activity to remove Spo11 from both ssDNA and dsDNA *in vitro*

(above), we were interested to investigate whether TDP2 expression during *S. cerevisiae*

meiosis would permit removal of Spo11 *in vivo*.

To test this idea, we placed His-tagged TDP2 under the control of the *ADH1* promoter in a

sae2Δ strain, and induced cells to enter meiosis (Fig 4a). Genomic DNA was harvested at

hourly intervals during meiotic prophase, and used to probe the DSB hotspot at the

HIS4::LEU2 locus as in earlier experiments (Fig 2 and Fig 3). We observed no difference in

the abundance or electrophoretic mobility of the DSB bands in either the presence or absence

of TDP2 expression, indicating that TDP2 was neither promoting resection nor repair of the

Spo11-DSBs (Fig 4a). Expression of TDP2 from an inducible *GAL1* promoter construct after

4 hours in meiosis in a *sae2Δ* strain also resulted in no detectable alteration in DSB signal

(data not shown).

A possible explanation for no visible resection or repair of Spo11-DSBs after TDP2

expression is that TDP2 has removed Spo11 from the DSB end, but these DSBs, with 2 nt 5′

overhangs, are resistant to nucleases such as Exo1 (as demonstrated *in vitro* in Fig 3), and

that NHEJ components may be downregulated in meiosis perhaps due to increasing CDK

activity as has been observed in cycling *S. pombe* cells (42). To test this possibility, we

isolated genomic DNA from a 10 h meiotic time point from cells expressing *TDP2,* or from

control cells, and incubated with lambda exonuclease (Fig 4b). At this time point, if TDP2

was capable of removing Spo11 from the 5′ end of the DSBs, lambda exonuclease would be

able to resect these DSBs (as described in Fig 2). However, we detected no visible resection

in the TDP2-expressing nor control cells (Fig 4b). These results suggest that despite the

ability of TDP2 to suppress topoisomerase-induced DNA damage in *S. cerevisiae* (25),

Spo11-DSBs are refractory to processing by the TDP2 enzyme.

Ectopic expression of TDP2 rescues sensitivity of *sae2Δ* and *mre11-H125N* mutant cells

to etoposide and camptothecin.

As a final investigation of TDP2 activity *in vivo* in *S. cerevisiae* cells, we expressed TDP2 in

drug-sensitive (*pdr1Δ)* haploid cells (43), and tested for the rescue of camptothecin (CPT)

and etoposide sensitivity, as well as to the DNA-alkylating agent methyl methanesulphonate

as a control (MMS). At the concentrations tested, wild type control cells (*pdr1Δ*) were not

sensitive to etoposide or camptothecin, and there was no effect of TDP2 expression (Fig 4c).

sae2Δ cells were sensitive to the higher concentration of etoposide and CPT, and were

rescued efficiently by ectopic TDP2 expression (Fig 4c). By contrast, whilst loss of Mre11

nuclease activity (*mre11-H125N*) resulted in drug sensitivities that were similar to the *sae2Δ*

mutant, TDP2 expression caused much weaker suppression than in the *sae2Δ* cells—

particularly following CPT treatment (Fig 4c). These observations suggest a separable role

for Sae2 and Mre11 in the repair of Top1- and Top2-induced DNA lesions, with TDP2 only

able to fully complement the functions of the Sae2 protein. Cells bearing a full deletion of

MRE11 were highly sensitive to even the lower drug concentrations, with no rescue of

growth observed upon expression of TDP2 (Fig 4c). This latter result is consistent with the

inability for TDP2 to complement some or all of the many functions of Mre11 (44). Notably,

TDP2 expression suppressed a very minor growth defect of *sae2Δ* and *mre11-H125N* cells in

the presence of MMS (compare colony sizes in Fig 4c with and without TDP2 expression),

suggesting either that MMS treatment leads to the accumulation of topoisomerase-lesions, or

that it generates directly or indirectly, other potentially toxic intermediates that are substrates

for the phosphodiesterase activity of TDP2.

DISCUSSION

TDP2 was identified in a screen for mammalian genes that rescued the sensitivity of *S.*

cerevisiae cells to the Top1 poison, camptothecin (25). Biochemical analyses subsequently

demonstrated the preferred activity of TDP2 against synthetic 5′ phosphotyrosine groups

(25), as well as larger peptides derived from partially proteolysed Top2 (31). Most recently,

TDP2 was shown to be proficient in catalysing the release of intact Top2 from dsDNA ends

in a reaction that is strongly dependent upon the ZATT protein (45). Here we have

characterised the activity of TDP2 against the 5′-linked Spo11 complexes that arise during

the induction of meiotic recombination. Recombinant human TDP2 displays robust removal

of partially proteolysed Spo11 protein from both ssDNA oligos and dsDNA molecules

derived from meiotic cell extracts. We find that non-proteolysed Spo11 is also released albeit

with reduced efficiency.

Under normal conditions in meiosis, Spo11-DSBs are processed by the nuclease activities of

Mre11 in conjunction with Sae2, followed by extensive 5′ to 3′ resection mediated by Exo1.

Surprisingly, despite efficient *in vitro* removal of Spo11 from DSB ends by TDP2, we have

found that the resulting protein-free DNA ends remain very poor substrates for Exo1, in

contrast to DSB substrates containing a terminal 3′ single-strand region of ~300 nt. These

observations agree with prior observations indicating a preference of Exo1 for partially

single-stranded substrates (12, 41). However, our experiments are the first that employ

physiological frequencies of DSB ends relative to genome content (approx. 1 DSB end for

every 150 kb), and lead us to conclude that excess dsDNA acts as a competitive inhibitor of

Exo1's ability to initiate resection at clean, relatively blunt, dsDNA ends—but not at dsDNA

ends that have already undergone limited resection.

Such competitive inhibition may also influence the activity of other proteins involved in DSB

end resection. For example, the Sae2-stimulated cleavage of protein-linked dsDNA by

Mre11, has thus far only been demonstrated on short, synthetic dsDNA substrates (11–13).

Attempts to reconstitute a similar Mre11 endonuclease reaction on natural Spo11-DSBs

isolated from meiotic cells has thus far proved unsuccessful in our laboratory (D. Johnson

and M. Neale, unpub. obs.)—perhaps in agreement with the reduced efficiency of the Mre11

endonuclease reaction on longer synthetic substrates (Ref 13 and E. Canavo and P. Cejka,

unpub. obs.). It is interesting to note that in *S. cerevisiae* Mre11 is loaded at meiotic

recombination hotspots prior to Spo11-DSB formation (46). It is tempting to speculate that

this interaction is important to ensure efficient nucleolytic processing of Spo11-DSB ends in

S. cerevisiae—thereby ensuring that meiotic DSB repair is directed down the HR repair

pathway.

In *C. elegans*, loss of COM-1 function (the Sae2/CtIP orthologue) leads to aberrant DSB

repair and meiotic chromosome segregation due to activation of the NHEJ repair pathway

(21, 22). Such observations suggest that mechanisms other than Mre11 are competent to

release Spo11 from meiotic DSB ends, generating ligatable DNA ends—at least under

conditions when the preferred Mre11-pathway is disrupted. An orthologue of human TDP2

exists in worms (33), but it is yet to be determined whether this activity is the one responsible

for enabling NHEJ repair of Spo11-DSBs in the *com-1* mutant.

Interestingly, whilst human TDP2 is capable of catalysing Spo11 removal from DSB ends *in*

vitro, we were unable to detect Spo11 removal *in vivo* following ectopic expression of TDP2

within meiotic *S. cerevisiae* cells. Whilst it is hard to rule out a technical reason for observing

this negative result, we *were* able to observe robust rescue of camptothecin and etoposide

sensitivity in *sae2Δ* and *mre11*-'nuclease-dead' mutant strains upon TDP2 expression. Thus,

our favoured interpretation is that Spo11-DSB ends in *S. cerevisiae*—unlike topoisomerase-

DNA lesions—are inaccessible to the TDP2 phosphodiesterase activity. Such inaccessibility

may be due to a higher order protein complex present at meiotic DSBs, which are known to

require directly or indirectly at least a dozen independent proteins (47). Such ideas are also

consistent with observations suggesting multimerisation of the Spo11 enzyme within

recombination hotspots (D. Johnson and M. Neale, manuscript in preparation). Alternatively,

or in addition, topoisomerase-induced lesions may be subject to greater rates of proteolysis

and/or denaturation than Spo11, thereby stimulating the end-cleaning activity of TDP2.

Recognition, and appropriate repair, of DNA lesions is critical for cell survival. During

meiotic prophase, the channelling of Spo11-DSBs down the HR pathway has further

importance to enable crossover formation, reductional chromosome segregation, and for the

generation of genetic diversity (48). Our observations here provide insight into the

mechanisms of repairing such protein-linked DSB ends created by topoisomerase-like

enzymes such as Spo11. Looking more broadly, our findings may also help to explain some

of the genetic differences observed for the resection and repair of 'clean'-ended site-specific

DSBs created by nucleases versus the complex DSB ends created by ionising radiation,

which may be a mixture not just of DNA ends with base damage, but also those with blunt or

partial 5′ extensions, or occluded by larger protein moieties.

ACKNOWLEDGEMENTS

We thank E. Hoffmann, N. Hunter, S. Keeney and J. Nitiss for yeast strains; L. Symington

for plasmids; K Caldecott for recombinant human TDP2 protein. D.J. is supported by an ERC

Consolidator Grant ('DNA-Repair-Chromatin') to M.J.N. M.J.N. was supported by this

award and a University Research Fellowship from the Royal Society, and a Career

Development Award from the Human Frontiers Science Program Organisation.

AUTHOR CONTRIBUTIONS

D.J. and M.J.N. designed and performed the experiments and wrote the paper. E.C. and P.C

provided essential reagents and technical supervision.

METHODS

Strains, Plasmids and Recombinant Proteins. Meiotic experiments utilised the *S.*

cerevisiae SK1 background (49), and harboured the *sae2Δ*::kanMX or *exo1Δ*::kanMX full

gene deletions (9). For Spo11-oligo enrichment, strains harboured the Spo11-

FLAG$_3$His$_6$::kanMX construct (30). The multicopy constitutive $P_{ADH1}TDP2$ plasmid (pDJ85)

was prepared after replacing the *URA3* gene in pVTU260TT (25) with a Hygromycin

resistance cassette. Cell viability spot tests utilized the S288c background harbouring the

dominant multi-drug sensitizing cassette *pdr1Δ*::*pdr1DBD-CYC8* (referred to as '*pdr1Δ*')

(43) and additionally harbouring the *sae2Δ*::kanMX, *mre11Δ*::*URA3* gene deletions or

mre11-H125N point mutation (50). Genetic modifications were introduced into yeast strains

using standard procedures (51). Recombinant human His-TDP2 was a gift from K. Caldecott,

and prepared as described previously (25). Recombinant *S. cerevisiae* Exo1-FLAG and RPA

was prepared as described previously (41).

Isolation and analysis of meiotic DNA breaks. SK1 cells were induced synchronously into

meiosis using standard procedures (52). At indicated timepoints, 10 ml of cell culture was

spheroplasted in 1M sorbitol / 50 mM EDTA / 50 mM NaHPO$_4$ buffer, and lysed by addition

of SDS to 0.5% in the presence of proteinase K at 60˚C. Nucleic acids were purified by

phenol extraction and ethanol precipitation, and resuspended in 1×TE. DSB signals at the

HIS4::*LEU2* recombination hotspot were detected using standard techniques (53). Briefly

DNA was digested with *Pst*I, electrophoresed on 0.7% agarose in 1×TAE for approximately

18 h at room temperature, transferred to nylon membrane under denaturing conditions then

hybridized with a probe that recognizes the *MXR2* locus. Prior to digestion, as appropriate,

samples were equilibrated in reaction buffer (25 mM HEPES, pH 8.0, 130 mM KCl, 1 mM

DTT, 10 mM MgCl2) and incubated with 300 nM recombinant TDP2 for 1 hour at 37˚C, 5

units of lambda exonuclease (NEB) for 1 hour at 37˚C, 15 units of Klenow (3′→5′ exo–)

polymerase (NEB) with 66 nM dNTPs for 30 minutes at 30˚C, 20 mM recombinant Exo1-

FLAG with 680 nM RPA (REF) for 30 minutes at 30˚C, or combinations thereof as indicated

in each figure. Reactions involving Exo1 were performed in a similar buffer (25 mM Tris

acetate, pH 7.5, 1 mM DTT, 5 mM magnesium acetate, 100 mM sodium acetate, 0.25 mg/mL

BSA) in which TDP2 maintains activity (Figure S1b). Reactions were stopped and DNA

extracted by the addition of 1 volume of water and 1 volume of phenol:chloroform:isoamyl

alcohol (25:24:1) prior to restriction digestion and electrophoresis. Radioactive signals were

collected on phosphor screens, scanned with a Fuji FLA5100 and quantified using

ImageGauge software.

Isolation and analysis of Spo11-oligo complexes. Spo11-oligonucleotide complexes were

detected by immunoprecipitation and end-labelling following established methods (54).

Briefly, cells were broken in 10% ice-cold TCA using zirconium beads and a BioSpec 24.

Precipitated material was dissolved in SDS buffer, diluted with Triton X100, and Spo11 was

immunoprecipitated from total soluble protein using anti-FLAG antibody (Sigma) and

protein-G-agarose (Roche). Oligonucleotide complexes were labelled with alpha-^{32}P dCTP

using terminal deoxynucleotidyl transferase (Fermentas) and fractionated on a 7.5% SDS-

PAGE gel. Alternatively, radiolabelled Spo11-oligo complexes were precipitated with 90%

ethanol at –80˚C overnight without or following digestion with trypsin or proteinase K

(Fisher) for 1 hour at 37˚C. Precipitated material was collected by centrifugation, solublised

in reaction buffer (25 mM HEPES, pH 8.0, 130 mM KCl, 1 mM DTT, 10 mM MgCl2), and

incubated with the stated concentration of recombinant TDP2 for 30 minutes at 30˚C prior to

denaturation in formamide loading dye and separation on 19% urea/PAGE/1×TBE.

Radioactive signals were collected on phosphor screens, scanned with a Fuji FLA5100 and

quantified using ImageGauge software.

Cell viability spot tests. YPD cultures were grown in the presence of 300 μg ml^{-1}

hygromycin (Invitrogen) to maintain selection of the plasmids. Overnight cultures were

diluted 40-fold in fresh YPD containing 300 μg ml^{-1} hygromycin and grown for 4 hours.

Cultures were diluted to 0.2 OD$_{600}$ and a 10-fold dilution series prepared before spotting out 3

μl onto the YPD plates containing 300 μg ml^{-1} hygromycin and the stated drug dose or

DMSO solvent control. Plates were incubated at 30 °C for 3 days prior to image acquisition.

FIGURE LEGENDS

Figure 1: TDP2 can cleave the covalent phosphotyrosyl bond between Spo11 and a

single-stranded oligonucleotide. a-b, Cartoon of the early stages of meiotic recombination.

Spo11 generates DSBs and is released covalently attached to short oligonucleotides that are

isolated by immunoprecipitation, and end-labelled at the 3′OH using alpha-^{32}P dCTP and

TdT, before resolving by SDS-PAGE (b). c-f, Schematic of the experiment (c). Labelled

Spo11-oligo complexes without (d) or after treatment with Trypsin (e) or Proteinase K (f),

were reacted with recombinant TDP2 at the indicated concentrations, then resolved on 18%

polyacrylamide gel under denaturing conditions. Solid arrowheads indicate the migration

position of the various Spo11-oligo species (prior to TDP2 treatment). Open triangles and

dashed lines indicate the migration position of Spo11-oligos after Spo11-peptide removal by

TDP2. Dotted connectors between gels indicate relative migration positions. Asterisks are a

non-specific band.

Figure 2: TDP2 can remove Spo11 peptides from the dsDNA end of natural Spo11-

DSBs enabling 5′-3′ resection by lambda exonuclease.

a, Schematic of experiment. b, Genomic DNA isolated from meiotic *sae2*Δ cells was

incubated with lambda exonuclease (λexo) without or with prior treatment with TDP2. DNA

was purified and digested with *Pst*I, separated by electrophoresis on a 0.7% agarose gel,

blotted to nylon membrane and hybridised with a probe (*MXR2* locus) close to the well-

characterised meiotic recombination hotspot *HIS4*::*LEU2*. The location of the two major

Spo11-DSBs at this locus are marked with solid arrowheads. Disappearance of these signals

following TDP2 and lambda exonuclease treatment is due to rapid exonucleolytic

degradation from the uncapped Spo11-DSB ends.

Figure 3: Recombinant Exo1 is unable to initiate 5′-3′ resection on 3′-recessed and

blunt-ended Spo11-DSBs even when uncapped by recombinant TDP2, but can

efficiently resect 5′-recessed Spo11-DSBs that have been processed by Mre11 *in vivo*.

a-b, Genomic DNA isolated from meiotic *sae2*Δ cells was incubated with recombinant Exo1

without (a) or with (b) prior treatment with TDP2. c-f, Genomic DNA isolated from meiotic

*exo1*Δ cells was incubated without (c), or with recombinant Exo1 (d), TDP2 (e), or TDP2

then Exo1 (f). c-d, As in (a, b), but TDP2-uncapped Spo11-DSB ends were blunted by

treating with Klenow polymerase + dNTPs prior to incubation with Exo1. DNA was purified

and digested with *Pst*I, separated by electrophoresis on a 0.7% agarose gel, blotted to nylon

membrane and hybridised with a probe (*MXR2* locus) close to the *HIS4*::*LEU2* meiotic

recombination hotspot. The location of the two major Spo11-DSBs at this locus are marked

with solid arrowheads. In *exo1*Δ cells, *in vivo* removal of Spo11 from DSB ends and partial

resection causes the DSB signals to migrate slightly faster than in *sae2*Δ samples. Further

smearing (open triangles) and subsequent disappearance of these signals following Exo1

treatment is due to rapid exonucleolytic degradation from the partially resected DSB ends.

Lower panels are schematics of each set of reactions.

Figure 4: Ectopic expression of *TDP2* cannot remove Spo11 from DSB ends *in vivo*, but

can suppress camptothecin and etoposide sensitivity.

a-b, Genomic DNA was isolated at the indicated timepoints from a synchronous meiotic

culture of *sae2*Δ cells harbouring either an empty vector, or one expressing *TDP2* from the

ADH1 promoter ($P_{ADH1}TDP2$). DNA was purified and digested with *Pst*I, separated by

electrophoresis on a 0.7% agarose gel, blotted to nylon membrane and hybridised with a

probe (*MXR2* locus) close to the *HIS4*::*LEU2* meiotic recombination hotspot. The location of

the two major Spo11-DSBs at this locus are marked with solid arrowheads. In *sae2*Δ cells,

Spo11 is not removed and resection cannot occur, causing DSB species to migrate as a tight

band. Expression of TDP2 does not alter this migration pattern (a). b, DNA from the 10 h

timepoints was incubated with lambda exonuclease for the indicated length of time. c, The

indicated strains harbouring either an empty vector, or one expressing *TDP2* from the *ADH1*

promoter ($P_{ADH1}TDP2$) were grown to log phase, serial diluted (10-fold), and spotted onto

plates containing 300 µg/ml hygromycin (HYG), to maintain plasmid selection, and the

indicated concentrations of etoposide, camptothecin (CPT) or methyl methanesulphonate

(MMS), and incubated for 3 days at 30˚C.

Figure 5: Model of DSB repair via HR or NHEJ depending on repair mechanism.

Covalently bound protein at DNA ends has to be removed in order for repair to occur. Mre11

and Sae2 remove the protein block (Spo11 or topoisomerases) nucleolytically, generating a 3′

ssDNA overhang substrate that is refractory to NHEJ, but the preferred substrate for long-

range resection by Exo1, thereby promoting homologous recombination (left and right

panels). Removal of covalently bound protein, for example, by hydrolytically cleaving the

phosphotyrosyl bond between the protein and the 5′ end of the DSB (e.g. by TDP2) generates

clean, short 5′ overhang ends that are refractory to resection by Exo1—thus potentially

inhibiting repair by HR—but the complementary nature of the two ends may promote repair

by NHEJ, for example at Top2-DSBs (central panel). The inability for ectopic expression of

TDP2 to suppress the requirement for Sae2 in S. *cerevisiae* meiosis suggests that Spo11-

DSBs, unlike Topoisomerase-DNA breaks, are inaccessible to TDP2, perhaps due to a large

complex at the Spo11-DSB site occluding access to TDP2 (dotted circle in left panel).

Figure S1: Recombinant TDP2 is free of nuclease activity, and is active in the preferred

Exo1 reaction buffer. a, To test for trace nuclease activity within the recombinant TDP2

preparation, a 30 nt oligonucleotide of random sequence was 3′-labelled using TdT and

radiolabelled dCTP then incubated with the stated concentrations of TDP2 at 37 °C for 1

hour then separated by denaturing urea-PAGE (19% acrylamide). The 30 nt oligo is indicated

with a filled arrowhead. **b,** Proteinase K-treated 3′-labelled Spo11-oligo complexes as in Fig

1f were incubated with the indicated concentration of TDP2 at 37˚C for 1 hour in Exo1

reaction buffer (25 mM Tris acetate, pH 7.5, 1 mM DTT, 5 mM magnesium acetate, 100 mM

sodium acetate, 0.25 mg/mL BSA) then separated by denaturing urea-PAGE (19%

acrylamide). Solid arrowheads indicate the migration position of Spo11-oligo species (prior

to TDP2 treatment). Open triangles indicate the migration position of Spo11-oligos after

Spo11-peptide removal by TDP2.

Table 1. *S. cerevisiae* strains used in this study.

Strain	Background	Genotype	Origin
VG296	SK1	*MATa/α, ho::LYS2/", lys2/", ura3/", arg4-nsp/arg4-bgl, leu2::hisG/", his4X::LEU2/" nuc1::LEU2/", SPO11-His6-FLAG3-loxP-KanMX-loxP/"*	This study
VG303	SK1	*MATa/α, ho::LYS2/", lys2/", ura3/", arg4/", leu2/", his4X::LEU2/" nuc1::LEU2/", SPO11-His6-FLAG3-loxP-KanMX-loxP/", sae2Δ::KanMX6/"*	This study
VG164	SK1	*MATa/α, ho::LYS2/", lys2", ura3/", arg4-nsp/", leu2::hisG/", his4X::LEU2/", nuc1::LEU2/", exo1Δ::KanMX4/"*	This study
DJ160	SK1	*MATa/α lys2/", ura3/", arg4/", leu2/", his4X::LEU2/" nuc1::LEU2/", SPO11-His6-FLAG3-loxP-KanMX-loxP/", sae2Δ::KanMX6/" + pDJ85 (P_{ADH1}His6-TDP2)*	This study
DJ199	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2 + pVG19 (empty HYG vector)*	This study
DJ209	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2 + pDJ85 (P_{ADH1}His6-TDP2)*	This study
DJ201	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, sae2Δ::KanMX6 + pVG19 (empty HYG vector)*	This study
DJ193	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, sae2Δ::KanMX6 + pDJ85 (P_{ADH1}His6-TDP2)*	This study
DJ203	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11Δ::KanMX4 + pVG19 (empty HYG vector)*	This study
DJ195	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11Δ::KanMX4 + pDJ85 (P_{ADH1}His6-TDP2)*	This study
DJ212	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11(H125N) + pVG19 (empty HYG vector)*	This study
DJ213	S288c	*MATa, ho, ura3Δ0, leu2Δ0, his3Δ1, met15Δ0, pdr1Δ::PDR1-DBD-CYC8::LEU2, mre11(H125N) pDJ85 (P_{ADH1}His6-TDP2)*	This study

537   **Reference List**
538

1.   Nichols,M.D., DeAngelis,K., Keck,J.L. and Berger,J.M. (1999) Structure and function
of an archaeal topoisomerase VI subunit with homology to the meiotic recombination
factor Spo11. *EMBO J* **18**, 6177-6188.
2.   Keeney,S., Giroux,C.N. and Kleckner,N. (1997) Meiosis-specific DNA double-strand
breaks are catalyzed by Spo11, a member of a widely conserved protein family. *Cell* **88**,
375-384.
3.   Bergerat,A., de Massy,B., Gadelle,D., Varoutas,P.C., Nicolas,A. and Forterre,P. (1997)
An atypical topoisomerase II from Archaea with implications for meiotic
recombination. *Nature* **386**, 414-417.
4.   Robert,T., Nore,A., Brun,C., Maffre,C., Crimi,B., Bourbon,H.M. and de Massy,B.
(2016) The TopoVIB-Like protein family is required for meiotic DNA double-strand
break formation. *Science* **351**, 943-949.
5.   Vrielynck,N., Chambon,A., Vezon,D., Pereira,L., Chelysheva,L., De Muyt,A.,
Mézard,C., Mayer,C. and Grelon,M. (2016) A DNA topoisomerase VI-like complex
initiates meiotic recombination. *Science* **351**, 939-943.
6.   de Massy,B., Rocco,V. and Nicolas,A. (1995) The nucleotide mapping of DNA double-
strand breaks at the CYS3 initiation site of meiotic recombination in Saccharomyces
cerevisiae. *EMBO J* **14**, 4589-4598.
7.   Liu,J., Wu,T.C. and Lichten,M. (1995) The location and structure of double-strand
DNA breaks induced during yeast meiosis: evidence for a covalently linked DNA-
protein intermediate. *EMBO J* **14**, 4599-4608.
8.   Keeney,S. and Kleckner,N. (1995) Covalent protein-DNA complexes at the 5' strand
termini of meiosis-specific double-strand breaks in yeast. *Proc Natl Acad Sci U S A* **92**,
11274-11278.
9.   Garcia,V., Phelps,S.E.L., Gray,S. and Neale,M.J. (2011) Bidirectional resection of
DNA double-strand breaks by Mre11 and Exo1. *Nature* **479**, 241-244.
10.   Neale,M.J., Pan,J. and Keeney,S. (2005) Endonucleolytic processing of covalent
protein-linked DNA double-strand breaks. *Nature* **436**, 1053-1057.
11.   Wang,W., Daley,J.M., Kwon,Y., Krasner,D.S. and Sung,P. (2017) Plasticity of the
Mre11-Rad50-Xrs2-Sae2 nuclease ensemble in the processing of DNA-bound
obstacles. *Genes Dev* **31**, 2331-2336.
12.   Reginato,G., Cannavo,E. and Cejka,P. (2017) Physiological protein blocks direct the
Mre11-Rad50-Xrs2 and Sae2 nuclease complex to initiate DNA end resection. *Genes*
Dev **31**, 2325-2330.
13.   Cannavo,E. and Cejka,P. (2014) Sae2 promotes dsDNA endonuclease activity within
Mre11-Rad50-Xrs2 to resect DNA breaks. *Nature* **514**, 122-125.
14.   Sartori,A.A., Lukas,C., Coates,J., Mistrik,M., Fu,S., Bartek,J., Baer,R., Lukas,J. and
Jackson,S.P. (2007) Human CtIP promotes DNA end resection. *Nature* **450**, 509-514.
15.   Zakharyevich,K., Ma,Y., Tang,S., Hwang,P.Y.-H., Boiteux,S. and Hunter,N. (2010)
Temporally and Biochemically Distinct Activities of Exo1 during Meiosis: Double-
Strand Break Resection and Resolution of Double Holliday Junctions. *Mol Cell* **40**,
1001-1015.
16.   Hodgson,A., Terentyev,Y., Johnson,R.A., Bishop-Bailey,A., Angevin,T., Croucher,A.
and Goldman,A.S.H. (2010) Mre11 and Exo1 contribute to the initiation and
processivity of resection at meiotic double-strand breaks made independently of Spo11.
DNA Repair (Amst)
17.   Tsubouchi,H. and Ogawa,H. (2000) Exo1 roles for repair of DNA double-strand breaks
and meiotic crossing over in Saccharomyces cerevisiae. *Mol Biol Cell* **11**, 2221-2233.

18.   Rothenberg,M., Kohli,J. and Ludin,K. (2009) Ctp1 and the MRN-complex are required
588   for endonucleolytic Rec12 removal with release of a single class of oligonucleotides in
589   fission yeast. *PLoS Genet* **5**, e1000722.
590   19.   Lange,J., Pan,J., Cole,F., Thelen,M.P., Jasin,M. and Keeney,S. (2011) ATM controls
591   meiotic double-strand-break formation. *Nature* **479**, 237-240.
592   20.   Hyppa,R.W., Cromie,G.A. and Smith,G.R. (2008) Indistinguishable landscapes of
593   meiotic DNA breaks in rad50+ and rad50S strains of fission yeast revealed by a novel
594   rad50+ recombination intermediate. *PLoS Genet* **4**, e1000267.
595   21.   Penkner,A., Portik-Dobos,Z., Tang,L., Schnabel,R., Novatchkova,M., Jantsch,V. and
596   Loidl,J. (2007) A conserved function for a Caenorhabditis elegans Com1/Sae2/CtIP
597   protein homolog in meiotic recombination. *EMBO J* **26**, 5071-5082.
598   22.   Lemmens,B.B.L.G., Johnson,N.M. and Tijsterman,M. (2013) COM-1 Promotes
599   Homologous Recombination during Caenorhabditis elegans Meiosis by Antagonizing
600   Ku-Mediated Non-Homologous End Joining. *PLoS Genet* **9**, e1003276.
601   23.   Pommier,Y., Sun,Y., Huang,S.N. and Nitiss,J.L. (2016) Roles of eukaryotic
602   topoisomerases in transcription, replication and genomic stability. *Nat Rev Mol Cell*
603   *Biol* **17**, 703-721.
604   24.   Yang,S.W., Burgin,A.B., Huizenga,B.N., Robertson,C.A., Yao,K.C. and Nash,H.A.
605   (1996) A eukaryotic enzyme that can disjoin dead-end covalent complexes between
606   DNA and type I topoisomerases. *Proc Natl Acad Sci U S A* **93**, 11534-11539.
607   25.   Cortes Ledesma,F., El Khamisy,S.F., Zuma,M.C., Osborn,K. and Caldecott,K.W.
608   (2009) A human 5'-tyrosyl DNA phosphodiesterase that repairs topoisomerase-
609   mediated DNA damage. *Nature* **461**, 674-678.
610   26.   Nitiss,K.C., Malik,M., He,X., White,S.W. and Nitiss,J.L. (2006) Tyrosyl-DNA
611   phosphodiesterase (Tdp1) participates in the repair of Top2-mediated DNA damage.
612   *Proc Natl Acad Sci U S A* **103**, 8953-8958.
613   27.   Mimitou,E.P. and Symington,L.S. (2008) Sae2, Exo1 and Sgs1 collaborate in DNA
614   double-strand break processing. *Nature* **455**, 770-774.
615   28.   Pype,S., Declercq,W., Ibrahimi,A., Michiels,C., Van Rietschoten,J.G., Dewulf,N., de
616   Boer,M., Vandenabeele,P., Huylebroeck,D. and Remacle,J.E. (2000) TTRAP, a novel
617   protein that associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF
618   receptor-associated factors (TRAFs), and that inhibits nuclear factor-kappa B
619   activation. *J Biol Chem* **275**, 18586-18593.
620   29.   Hartsuiker,E., Mizuno,K., Molnar,M., Kohli,J., Ohta,K. and Carr,A.M. (2009)
621   Ctp1CtIP and Rad32Mre11 nuclease activity are required for Rec12Spo11 removal, but
622   Rec12Spo11 removal is dispensable for other MRN-dependent meiotic functions. *Mol*
623   *Cell Biol* **29**, 1671-1681.
624   30.   Thacker,D., Mohibullah,N., Zhu,X. and Keeney,S. (2014) Homologue engagement
625   controls meiotic DNA break number and distribution. *Nature* **510**, 241-246.
626   31.   Gao,R., Schellenberg,M.J., Huang,S.Y., Abdelmalak,M., Marchand,C., Nitiss,K.C.,
627   Nitiss,J.L., Williams,R.S. and Pommier,Y. (2014) Proteolytic degradation of
628   topoisomerase II (Top2) enables the processing of Top2·DNA and Top2·RNA covalent
629   complexes by tyrosyl-DNA-phosphodiesterase 2 (TDP2). *J Biol Chem* **289**, 17960-
630   17969.
631   32.   Pommier,Y., Huang,S.Y., Gao,R., Das,B.B., Murai,J. and Marchand,C. (2014) Tyrosyl-
632   DNA-phosphodiesterases (TDP1 and TDP2). *DNA Repair (Amst)* **19**, 114-129.
633   33.   Shi,K., Kurahashi,K., Gao,R., Tsutakawa,S.E., Tainer,J.A., Pommier,Y. and Aihara,H.
634   (2012) Structural basis for recognition of 5'-phosphotyrosine adducts by Tdp2. *Nat*
635   *Struct Mol Biol* **19**, 1372-1377.

636   34.   Schellenberg,M.J., Appel,C.D., Adhikari,S., Robertson,P.D., Ramsden,D.A. and
637   Williams,R.S. (2012) Mechanism of repair of 5'-topoisomerase II-DNA adducts by
638   mammalian tyrosyl-DNA phosphodiesterase 2. *Nat Struct Mol Biol* **19**, 1363-1371.
639   35.   Prinz,S., Amon,A. and Klein,F. (1997) Isolation of COM1, a new gene required to
640   complete meiotic double-strand break-induced recombination in Saccharomyces
641   cerevisiae. *Genetics* **146**, 781-795.
642   36.   McKee,A.H. and Kleckner,N. (1997) A general method for identifying recessive
643   diploid-specific mutations in Saccharomyces cerevisiae, its application to the isolation
644   of mutants blocked at intermediate stages of meiotic prophase and characterization of a
645   new gene SAE2. *Genetics* **146**, 797-816.
646   37.   Sriprakash,K.S., Lundh,N., Huh,M.M.-O. and Radding,C.M. (1975) The specificity of
647   lambda exonuclease. Interactions with single-stranded DNA. *J Biol Chem* **250**, 5438-
648   5445.
649   38.   Cao,L., Alani,E. and Kleckner,N. (1990) A pathway for generation and processing of
650   double-strand breaks during meiotic recombination in S. cerevisiae. *Cell* **61**, 1089-1101.
651   39.   Pan,J., Sasaki,M., Kniewel,R., Murakami,H., Blitzblau,H.G., Tischfield,S.E., Zhu,X.,
652   Neale,M.J., Jasin,M., Socci,N.D., Hochwagen,A. and Keeney,S. (2011) A Hierarchical
653   Combination of Factors Shapes the Genome-wide Topography of Yeast Meiotic
654   Recombination Initiation. *Cell* **144**, 719-731.
655   40.   Murakami,H. and Nicolas,A. (2009) Locally, meiotic double-strand breaks targeted by
656   Gal4BD-Spo11 occur at discrete sites with a sequence preference. *Mol Cell Biol* **29**,
657   3500-3516.
658   41.   Cannavo,E., Cejka,P. and Kowalczykowski,S.C. (2013) Relationship of DNA
659   degradation by Saccharomyces cerevisiae Exonuclease 1 and its stimulation by RPA
660   and Mre11-Rad50-Xrs2 to DNA end resection. *Proc Natl Acad Sci U S A*
661   42.   Hentges,P., Waller,H., Reis,C.C., Ferreira,M.G. and Doherty,A.J. (2014) Cdk1 restrains
662   NHEJ through phosphorylation of XRCC4-like factor Xlf1. *Cell Rep* **9**, 2011-2017.
663   43.   Stepanov,A., Nitiss,K.C., Neale,G. and Nitiss,J.L. (2008) Enhancing drug accumulation
664   in Saccharomyces cerevisiae by repression of pleiotropic drug resistance genes with
665   chimeric transcription repressors. *Mol Pharmacol* **74**, 423-431.
666   44.   Stracker,T.H. and Petrini,J.H.J. (2011) The MRE11 complex: starting from the ends.
667   *Nat Rev Mol Cell Biol* **12**, 90-103.
668   45.   Schellenberg,M.J., Lieberman,J.A., Herrero-Ruiz,A., Butler,L.R., Williams,J.G.,
669   Muñoz-Cabello,A.M., Mueller,G.A., London,R.E., Cortés-Ledesma,F. and
670   Williams,R.S. (2017) ZATT (ZNF451)-mediated resolution of topoisomerase 2 DNA-
671   protein cross-links. *Science*
672   46.   Borde,V., Lin,W., Novikov,E., Petrini,J.H., Lichten,M. and Nicolas,A. (2004)
673   Association of Mre11p with double-strand break sites during yeast meiosis. *Mol Cell*
674   **13**, 389-401.
675   47.   Lam,I. and Keeney,S. (2015) Mechanism and regulation of meiotic recombination
676   initiation. *Cold Spring Harb Perspect Biol* **7**, a016634.
677   48.   Neale,M.J. and Keeney,S. (2006) Clarifying the mechanics of DNA strand exchange in
678   meiotic recombination. *Nature* **442**, 153-158.
679   49.   Kane,S.M. and Roth,R. (1974) Carbohydrate metabolism during ascospore
680   development in yeast. *J Bacteriol* **118**, 8-14.
681   50.   Krogh,B.O., Llorente,B., Lam,A. and Symington,L.S. (2005) Mutations in Mre11
682   phosphoesterase motif I that impair Saccharomyces cerevisiae Mre11-Rad50-Xrs2
683   complex stability in addition to nuclease activity. *Genetics* **171**, 1561-1570.
684   51.   Gietz,R.D. (2014) Yeast transformation by the LiAc/SS carrier DNA/PEG method.
685   *Methods Mol Biol* **1205**, 1-12.

52. Gray,S., Allison,R.M., Garcia,V., Goldman,A.S.H. and Neale,M.J. (2013) Positive
regulation of meiotic DNA double-strand break formation by activation of the DNA
damage checkpoint kinase Mec1(ATR). *Open Biol* **3**, 130019.
53. Garcia,V., Gray,S., Allison,R.M., Cooper,T.J. and Neale,M.J. (2015) Tel1(ATM)-
mediated interference suppresses clustered meiotic double-strand-break formation.
Nature **520**, 114-118.
54. Neale,M.J. and Keeney,S. (2009) End-labeling and analysis of Spo11-oligonucleotide
complexes in Saccharomyces cerevisiae. *Methods Mol Biol* **557**, 183-195.
694

Figure 1: TDP2 can cleave the covalent phosphotyrosyl bond between Spo11 and a single-stranded oligonucleotide. a-b, Cartoon of the early stages of meiotic recombination. Spo11 generates DSBs and is released covalently attached to short oligonucleotides that are isolated by immunoprecipitation, and end-labelled at the 3′OH using alpha-^{32}P dCTP and TdT, before resolving by SDS-PAGE (b). c-f, Schematic of the experiment (c). Labelled Spo11-oligo complexes without (d) or after treatment with Trypsin (e) or Proteinase K (f), were reacted with recombinant TDP2 at the indicated concentrations, then resolved on 18% polyacrylamide gel under denaturing conditions. Solid arrowheads indicate the migration position of the various Spo11-oligo species (prior to TDP2 treatment). Open triangles and dashed lines indicate the migration position of Spo11-oligos after Spo11-peptide removal by TDP2. Dotted connectors between gels indicate relative migration positions. Asterisks are a non-specific band.

Figure 2: TDP2 can remove Spo11 peptides from the dsDNA end of natural Spo11-DSBs enabling 5′-3′ resection by lambda exonuclease. a, Schematic of experiment. b, Genomic DNA isolated from meiotic *sae2*Δ cells was treated with proteinase K and incubated with lambda exonuclease (λexo) without or with prior treatment with TDP2. DNA was purified and digested with *Pst*I, separated by electrophoresis on a 0.7% agarose gel, blotted to nylon membrane and hybridised with a probe (*MXR2* locus) close to the well-characterised meiotic recombination hotspot *HIS4::LEU2*. The location of the two major Spo11-DSBs at this locus are marked with solid arrowheads. Disappearance of these signals following TDP2 and lambda exonuclease treatment is due to rapid exonucleolytic degradation from the uncapped Spo11-DSB ends.

Figure 3: Recombinant Exo1 is unable to initiate 5′-3′ resection on 3′-recessed and blunt-ended Spo11-DSBs even when uncapped by recombinant TDP2, but can efficiently resect 5′-recessed Spo11-DSBs that have been processed by Mre11 *in vivo*.

a-b, Genomic DNA isolated from meiotic *sae2*Δ cells was incubated with recombinant Exo1 without (a) or with (b) prior treatment with TDP2. c-f, Genomic DNA isolated from meiotic *exo1*Δ cells was incubated without (c), or with recombinant Exo1 (d), TDP2 (e), or TDP2 then Exo1 (f). c-d, As in (a, b), but TDP2-uncapped Spo11-DSB ends were blunted by treating with Klenow polymerase + dNTPs prior to incubation with Exo1. DNA was purified and digested with *Pst*I, separated by electrophoresis on a 0.7% agarose gel, blotted to nylon membrane and hybridised with a probe (*MXR2* locus) close to the *HIS4*::*LEU2* meiotic recombination hotspot. The location of the two major Spo11-DSBs at this locus are marked with solid arrowheads. In *exo1*Δ cells, *in vivo* removal of Spo11 from DSB ends and partial resection causes the DSB signals to migrate slightly faster than in *sae2*Δ samples. Further smearing (open triangles) and subsequent disappearance of these signals following Exo1 treatment is due to rapid exonucleolytic degradation from the partially resected DSB ends. Lower panels are schematics of each set of reactions.

Figure 4: Ectopic expression of *TDP2* cannot remove Spo11 from DSB ends *in vivo*, but can suppress camptothecin and etoposide sensitivity. a-b, Genomic DNA was isolated at the indicated timepoints from a synchronous meiotic culture of *sae2*Δ cells harbouring either an empty vector, or one expressing *TDP2* from the *ADH1* promoter (*P$_{ADH1}$TDP2*). DNA was purified and digested with *Pst*I, separated by electrophoresis on a 0.7% agarose gel, blotted to nylon membrane and hybridised with a probe (*MXR2* locus) close to the *HIS4::LEU2* meiotic recombination hotspot. The location of the two major Spo11-DSBs at this locus are marked with solid arrowheads. In *sae2*Δ cells, Spo11 is not removed and resection cannot occur, causing DSB species to migrate as a tight band. Expression of TDP2 does not alter this migration pattern (a). b, DNA from the 10 h timepoints was incubated with lambda exonuclease for the indicated length of time. c, The indicated strains harbouring either an empty vector, or one expressing *TDP2* from the *ADH1* promoter (*P$_{ADH1}$TDP2*) were grown to log phase, serial diluted (10-fold), and spotted onto plates containing 300 μg/ml hygromycin (HYG), to maintain plasmid selection, and the indicated concentrations of etoposide, camptothecin (CPT) or methyl methanesulphonate (MMS), and incubated for 3 days at 30˚C.

Figure 5: Model of DSB repair via HR or NHEJ depending on repair mechanism. Covalently bound protein at DNA ends has to be removed in order for repair to occur. Mre11 and Sae2 remove the protein block (Spo11 or topoisomerases) nucleolytically, generating a 3′ ssDNA overhang substrate that is refractory to NHEJ, but the preferred substrate for long-range resection by Exo1, thereby promoting homologous recombination (left and right panels). Removal of covalently bound protein, for example, by hydrolytically cleaving the phosphotyrosyl bond between the protein and the 5′ end of the DSB (e.g. by TDP2) generates clean, short 5′ overhang ends that are refractory to resection by Exo1—thus potentially inhibiting repair by HR—but the complementary nature of the two ends may promote repair by NHEJ, for example at Top2-DSBs (central panel). The inability for ectopic expression of TDP2 to suppress the requirement for Sae2 in S. *cerevisiae* meiosis suggests that Spo11-DSBs, unlike Topoisomerase-DNA breaks, are inaccessible to TDP2, perhaps due to a large complex at the Spo11-DSB site occluding access to TDP2 (dotted circle in left panel).

Figure S1: Recombinant TDP2 is free of nuclease activity, and is active in the preferred Exo1 reaction buffer. a, To test for trace nuclease activity within the recombinant TDP2 preparation, a 30 nt oligonucleotide of random sequence was 3′-labelled using TdT and radiolabelled dCTP then incubated with the stated concentrations of TDP2 at 37 °C for 1 hour then separated by denaturing urea-PAGE (19% acrylamide). The 30 nt oligo is indicated with a filled arrowhead. **b,** Proteinase K-treated 3′-labelled Spo11-oligo complexes as in Fig 1f were incubated with the indicated concentration of TDP2 at 37°C for 1 hour in Exo1 reaction buffer (25 mM Tris acetate, pH 7.5, 1 mM DTT, 5 mM magnesium acetate, 100 mM sodium acetate, 0.25 mg/mL BSA) then separated by denaturing urea-PAGE (19% acrylamide). Solid arrowheads indicate the migration position of Spo11-oligo species (prior to TDP2 treatment). Open triangles indicate the migration position of Spo11-oligos after Spo11-peptide removal by TDP2.