DNA methylation and histone H1 cooperatively repress transposable elements and aberrant intragenic transcripts

Jaemyung Choi[1], David B. Lyons[2], M. Yvonne Kim[2], Jonathan D. Moore[1] and Daniel Zilberman[1,*]

[1]Department of Cell & Developmental Biology, John Innes Centre, Norwich, NR4 7UH, UK

[2]Department of Plant & Microbial Biology, University of California, Berkeley, CA 94720, USA

*Correspondence: daniel.zilberman@jic.ac.uk

Summary

DNA methylation and histone H1 mediate transcriptional silencing of genes and transposable

elements, but how they interact is unclear. In plants and animals with mosaic genomic methylation,

functionally mysterious methylation is also common within constitutively active housekeeping

genes. Here we show that H1 is enriched in methylated sequences, including genes, of *Arabidopsis*

thaliana, yet this enrichment is independent of DNA methylation. Loss of H1 disperses

heterochromatin, globally alters nucleosome organization, and activates H1-bound genes, but only

weakly de-represses transposable elements. However, H1 loss strongly activates transposable

elements hypomethylated through mutation of DNA methyltransferase *MET1*. Loss of H1 also

activates antisense transcripts within demethylated genes. Our results demonstrate that H1 and

DNA methylation cooperatively maintain transcriptional homeostasis by silencing transposable

elements and aberrant intragenic transcripts. Such functionality plausibly explains why DNA

methylation, a well-known mutagen, has been maintained within coding sequences of crucial plant

and animal genes.

Keywords: DNA methylation, gene body methylation, histone H1, transcriptional silencing,

antisense transcription.

Highlights

• Histone H1 is enriched in methylated DNA independently of methylation

• Loss of H1 activates genes, alters nucleosome organization and disperses heterochromatin

• DNA methylation and H1 jointly silence transposons

• DNA methylation and H1 cooperatively suppress intragenic antisense transcripts

Introduction

Cytosine DNA methylation is an epigenetic modification that protects genome integrity by repressing transposable elements (TEs) in plants, vertebrates, fungi, and likely other eukaryotic groups (Du et al., 2015; Jones, 2012; Zemach and Zilberman, 2010). Methylation of regulatory sequences, such as promoters and enhancers, also causes gene silencing in plants and vertebrates (Hon et al., 2013; Jones, 2012; Schübeler, 2015; Zhang et al., 2018a). Proper patterns of DNA methylation are essential for plant and animal development (Iurlaro et al., 2017; Kawashima and Berger, 2014; Schmidt et al., 2015; Smith and Meissner, 2013), and their disruption is linked with cancer and other serious diseases (Jones et al., 2016; Rasmussen and Helin, 2016; Robertson, 2005).

Despite its association with transcriptional repression, DNA methylation is common within active genes (Bewick and Schmitz, 2017; Feng et al., 2010; Jones, 2012; Zemach et al., 2010). Vertebrate intragenic methylation occurs in the context of a globally methylated genome, in which methylation is the default state and some sequences, including active promoters and enhancers, are protected from methylation (Schübeler, 2015; Suzuki and Bird, 2008). Mammalian genes are co-transcriptionally hypermethylated by the *de novo* methyltransferase Dnmt3 (Baubec et al., 2015; Morselli et al., 2015; Neri et al., 2017), and – likely due to this activity – mammalian intragenic methylation is positively correlated with gene expression (Lister et al., 2009; Yang et al., 2014). Several functions have been ascribed to mammalian intragenic methylation, including the silencing of intragenic promoters and enhancers (Deaton et al., 2011; Hellman and Chess, 2007; Kulis et al., 2012; Maunakea et al., 2010; Yang et al., 2014), regulation of splicing (Lev Maor et al., 2015; Maunakea et al., 2013; Shukla et al., 2011), and suppression of aberrant transcripts (Neri et al., 2017). However, the latter function has been questioned, because strong DNA methylation

loss in mouse embryonic stem cells was not associated with activation of intragenic transcription

(Teissandier and Bourc'his, 2017).

In contrast to vertebrates, plant and invertebrate genomes usually have mosaic methylation

patterns, with specific sequences targeted for methylation (Schübeler, 2015; Suzuki and Bird,

2008). In flowering plants, TEs and gene bodies are preferentially methylated (Bewick and

Schmitz, 2017; Lister et al., 2008; Takuno and Gaut, 2013; Zemach and Zilberman, 2010). TEs

are affected by multiple pathways that mediate methylation in three sequence contexts (CG, CHG

and CHH, where $H \neq G$), whereas only CG sites are methylated in genes (Feng et al., 2010; Zemach

et al., 2010; Zhang et al., 2018a). Invertebrates preferentially or exclusively methylate gene bodies,

and essentially all methylation is in the CG context (Bewick et al., 2017; Dixon et al., 2016; Feng

et al., 2010; Suzuki et al., 2007; Zemach et al., 2010). Plant and invertebrate intragenic methylation

is not directly linked to transcription, but is instead a largely invariant genomic feature that

decorates the same sets of genes across developmental stages and tissues (Bartels et al., 2018;

Libbrecht et al., 2016; Suzuki et al., 2013; Zemach et al., 2018). This phenomenon, commonly

called gene body methylation (gbM), is concentrated in the exons of evolutionarily conserved,

stably and constitutively expressed genes, and is maintained by the methyltransferase Dnmt1

(called MET1 in plants) (Bewick and Schmitz, 2017; Bewick et al., 2018; Cokus et al., 2008;

Dixon et al., 2016; Lister et al., 2008; Sarda et al., 2012; Takuno and Gaut, 2012, 2013; Zemach

and Zilberman, 2010). GbM has been proposed to regulate splicing and transcriptional elongation,

and to suppress intragenic transcriptional initiation (Hunt et al., 2013; Regulski et al., 2013; Suzuki

et al., 2007; To et al., 2015; Tran et al., 2005; Zilberman et al., 2007). However, loss of gbM in

the flowering plant *Arabidopsis thaliana* and the insect *Oncopeltus fasciatus* is not associated with

transcriptional changes (Bewick et al., 2016, 2018; Kawakatsu et al., 2016), and gbM is commonly

thought to be a nonfunctional byproduct of TE methylation (Bewick and Schmitz, 2017; Bewick

et al., 2016; Teixeira and Colot, 2009). Overall, the functions of gbM – if any exist – remain

mysterious (Zhang et al., 2018a; Zilberman, 2017).

A potential explanation for the mystery and controversy surrounding gbM – and intragenic

methylation in general – is that its function is subtle and shared with other chromatin factors. This

hypothesis was tested in the flowering plant *Arabidopsis thaliana* by simultaneously eliminating

gbM and trimethylation of lysine 36 of histone H3, a histone modification that suppresses cryptic

intragenic transcripts and regulates splicing (Carrozza et al., 2005; Wagner and Carpenter, 2012),

but no transcriptional or RNA processing defects attributable to gbM were found (Bewick et al.,

2016). A plausible alternate candidate is histone H1, a protein that binds nucleosomes and the

intervening linker DNA (Fyodorov et al., 2018; Hergeth and Schneider, 2015; Over and Michaels,

2014). Like methylation, H1 is associated with transcriptional silencing but is nevertheless

abundant in active plant and animal genes (Krishnakumar et al., 2008; Rutowicz et al., 2015;

Torres et al., 2016). There is also evidence that H1 preferentially inhibits transcriptional initiation

from methylated DNA *in vitro* (Johnson et al., 1995; Levine et al., 1993). These data suggest that

H1 and DNA methylation may act cooperatively, including within gene bodies. However, whether

H1 affinity is regulated by DNA methylation is uncertain, with some studies finding preferential

association with methylated DNA (Levine et al., 1993; McArthur and Thomas, 1996), and others

reporting no preference (Campoy et al., 1995; Hashimshony et al., 2003; Nightingale and Wolffe,

1995). More generally, if and how H1 interacts with methylation *in vivo* to regulate transcription

is unknown.

Results

Histone H1 is enriched in methylated DNA independently of methylation

To investigate the functional relationship between DNA methylation and H1, we analyzed the

genomic distribution of the two major *Arabidopsis* H1 proteins, H1.1 and H1.2 (Over and

Michaels, 2014). Consistent with earlier work (Rutowicz et al., 2015), we did not find any salient

differences between H1.1 and H1.2 (Figure S1A), and therefore will refer simply to H1. As

expected, H1 preferentially associates with linker DNA (Figures 1A and S1B), is concentrated in

heavily methylated heterochromatic TEs (Figures 1B, 1C, S1C and Table S1), and is abundant

within genes, particularly those with low/no expression (Figures 1D and S1D). H1 is also enriched

in methylated genes compared to similarly expressed unmethylated genes (Figures 1E and S1E).

Thus, H1 is preferentially associated with methylated DNA in *Arabidopsis*.

To determine if DNA methylation affects H1 binding, we analyzed H1 distribution in *met1*

mutant plants. The *met1* mutation eliminates CG methylation, which includes the entirety of gbM

(Cokus et al., 2008; Lister et al., 2008). Many TEs also lose non-CG methylation and dimethylation

of lysine 9 of histone H3 (H3K9me2), a mark of heterochromatin (Feng and Michaels, 2015), and

become more accessible to DNase I (MET1-dependent TEs, Figure 1F) (Deleris et al., 2012; Zhang

et al., 2018b). We find that H1 distribution is well-correlated between wild type (*wt*) and *met1*

(Figures 1B, 1C, 1F, 1G and S1C, S1F). Genes methylated in *wt* retain higher levels of H1 in *met1*

plants (Figures 1E and S1E), as do TEs (Figures 1C and S1C). Therefore, H1 binding is largely

independent of DNA methylation. However, some TEs do lose H1 in *met1* mutants, particularly

MET1-dependent TEs that are transcriptionally activated (Figures 1F and S1G).

To determine how H1 distribution is regulated, we analyzed H1 association with various

genomic features using principal component analysis. H1 associates most strongly with GC

content (Figure 1H) independently of nucleosome density (Figure S1H). Methylated genes have

elevated GC content (Figure S1I), which likely explains their H1 enrichment (Figures 1E and S1E).

H1 also associates with heterochromatic features, including DNA methylation and H3K9me2,

along the first principal component (Figure 1H), and is well-separated from features of active

transcription (such as H3K4me3, Figure 1H). Based on these results, we used GC content and *wt*

H3K9me2 and H3K4me3 data to create a linear regression model of H1 abundance. This model

accurately describes *wt* H1 distribution across genes and TEs (Figures 1B, 1F, 1I, 1J, and S1J-L),

including H1 enrichment in methylated genes (Figures 1E and S1K). More importantly, our model

successfully predicts *met1* H1 distribution (Figures 1B, 1E, 1F, 1J and S1J-L), including the loss

of H1 from a subset of MET1-dependent (*i.e.* heterochromatin-deficient) TEs (Figures 1F and

S1G). Thus, H1 enrichment can largely be explained by sequence composition, transcription, and

heterochromatin, whereas DNA methylation has little, if any direct effect on H1 binding.

Histone H1 mediates global nucleosome organization

The distribution of *Arabidopsis* H1 (heterochromatin > silent genes > active genes, Figures 1B-D)

corresponds to an unexplained phenomenon in the human genome, in which genes expressed at

higher levels have shorter average nucleosome repeat length (NRL) than less expressed genes,

which have shorter average NRLs than heterochromatin (Valouev et al., 2011). Because loss of

H1 is known to reduce the average NRL in animal genomes (Baldi et al., 2018; Fan et al., 2003;

Woodcock et al., 2006), we tested the hypothesis that H1 causes the observed NRL differences by

comparing nucleosome spacing between plants with defective *H1.1* and *H1.2* genes (*h1* mutants)

(Zemach et al., 2013) and *wt* controls.

As in the human genome, *Arabidopsis* genes expressed at higher levels have shorter

average NRL in *wt* (Figure 2A). The NRL of less-expressed, H1-rich genes decreases substantially

in *h1* mutants (Figures 2A-C and S2A). In the most H1-rich genes, mean NRL is reduced from

185 bp to 169 bp (Figures 2B and 2C), which is consistent with the ability of H1 to protect about

20 bp of DNA *in vitro* (Simpson, 1978). Consistent with published results (Zhang et al., 2015), we

find that heterochromatic TEs have longer average NRLs than more euchromatic TEs (Figure 2D),

with the longest NRLs in the most H1-rich TEs (Figure 2E). Loss of H1 reduces mean NRL of

H1-rich TEs by 22 bp (from 189 bp to 167 bp), causing TEs in *h1* mutants to have similar NRLs

regardless of their *wt* H1 levels (Figures 2E, 2F and S2B). The average NRLs of TEs and silent

genes are also very similar in *h1* plants (Figure S2C). Thus, H1 is an important global regulator of

nucleosome placement that accounts for much of the average NRL difference across the

Arabidopsis genome.

Loss of H1 activates transcription and disperses heterochromatin

Histone H1 mediates gene silencing (Fyodorov et al., 2018; Krishnakumar et al., 2008; Torres et

al., 2016) and the 3D configuration of heterochromatin (Cao et al., 2013; Lu et al., 2009) in

animals, but the effects of H1 on transcription and higher order chromatin organization in plants

are unknown. To understand how H1 regulates transcription, we determined mRNA levels using

RNA-seq in *h1* mutants in comparison to *wt*. Most of the significantly mis-regulated genes in

leaves and seedlings (707/990, 71%) are upregulated in *h1* plants (Figures 3A, 3B and Table S2).

These genes are H1-enriched (Figures 3C, 3D and S3A) and are transcribed at low levels in *wt*

(Figure 3E), indicating that H1 can repress gene transcription in plants. However, given that H1 is

generally abundant in *Arabidopsis* genes (Figures 1C and 1D), its effects on transcription are fairly

selective, as is the case in animals (Fan et al., 2005). Several gene ontology functional categories

are overrepresented among the upregulated genes, most notably categories related to the meiotic

cell cycle and meiotic recombination (Figure S3B and Table S2). Upregulated genes include

ASYNAPTIC (ASY) 1 and 4, and other key meiotic genes (Figures 3B, 3F and Table S2)

(Chambon et al., 2018; Mercier et al., 2015). H1 is depleted in male and female *Arabidopsis*

meiocytes (She and Baroux, 2015; She et al., 2013), suggesting H1 removal may help to activate

the meiotic transcriptional program. Genes involved in epigenetic silencing are also preferentially

upregulated (Figure S3B and Table S2). Curiously, one of these is ARGONAUTE9 (Figure 3B),

which is involved in meiocyte specification (Olmedo-Monfil et al., 2010), further reinforcing the

link between H1 and meiocyte fate and function.

In *wt Arabidopsis*, heterochromatin is arranged into large DAPI-stained foci called

chromocenters (Figures 4A and S4A) (Ascenzi and Gantt, 1999; Fransz et al., 2002). We find that

in *h1* mutants nuclei typically have few chromocenters and many smaller DAPI-stained foci

(Figures 4A, 4B and S4A), indicating that the role of H1 in heterochromatin organization is

generally conserved. However, despite the dispersal of chromocenters, only 32 TEs are

significantly upregulated in *h1* plants (Table S2). An important consideration is that H1 is a major

regulator of *Arabidopsis* DNA methylation: depletion of H1 increases methylation of some TEs in

all sequence contexts, and reduces methylation of other TEs (Lyons and Zilberman, 2017;

Rutowicz et al., 2015; Wierzbicki and Jerzmanowski, 2005; Zemach et al., 2013). Most of the

upregulated TEs (29, 91%) are hypomethylated in *h1* mutants (Figures 4C-E and S4B), suggesting

that some of the observed activation is not directly caused by H1 depletion. Furthermore,

heterochromatin does not become substantially more accessible to DNase I in *h1* mutants (Figure

4F). Thus, although heterochromatin is altered (Figures 2D-F) and dispersed (Figures 4A, 4B and

S4A) by lack of H1, heterochromatic functionality remains largely intact.

DNA methylation and H1 cooperatively silence TEs

Limited TE activation in *h1* plants contrasts with the extensive TE upregulation in H1-deficient

Drosophila melanogaster (Iwasaki et al., 2016; Lu et al., 2013), but is similar to the effects of H1

depletion on mouse cells, which show few changes in TE expression (Fan et al., 2005). TEs are

silenced by DNA methylation in mouse and *Arabidopsis* (Law and Jacobsen, 2010), whereas

Drosophila lacks cytosine methylation (Raddatz et al., 2013; Zemach et al., 2010), suggesting that

the effects of H1 loss may be masked by methylation. To test this hypothesis, we investigated TE

expression in *h1met1* compound mutants and *met1* controls.

Because H1 can locally raise or lower DNA methylation (Rutowicz et al., 2015; Zemach

et al., 2013), we first identified active TEs with unchanged CHG and CHH methylation near the

transcriptional start site (TSS) between *met1* and *h1met1*, which turned out to be mostly TEs that

are fully demethylated in both genotypes (Figures 5A, 5B, S5A and Table S3). Unsupervised

clustering separated these TEs into two groups, both of which become much more accessible to

DNase I in *met1* and remain so in *h1met1* (Figures 5A and S5B). The larger group (nc; 272 TEs)

is expressed similarly in *met1* and *h1met1* (Figure 5A). However, the smaller group of TEs (up1;

134) is strongly upregulated in *h1met1* (Figures 5A and S5C). These TEs are significantly H1-

enriched in *met1* compared to TEs with unchanged expression, especially around the TSS (Figures

5C, 5D and S5D). These TEs are thus apparently transcriptionally repressed by H1 and DNA

methylation. Loss of methylation alone leads to mild activation and locus accessibility, but loss of

both methylation and H1 is required to create a highly active state (Figures 5A and S5B).

Analysis of TEs with altered TSS-proximal CHG or CHH methylation produced three

groups, all of which maintain robust non-CG methylation in *met1* and *h1met1* (Figure 5E and Table

S3). All three groups are more accessible to DNase I in *h1met1* compared to *met1* (Figures 5E and

S5B), indicating that loss of H1 increases the accessibility of partially demethylated sequences.

The first group (up2) is comprised of 63 TEs upregulated and hypomethylated in *h1met1* compared

to *met1* (Figures 5E and S5E). The second group (dn) consists of 147 TEs downregulated and

hypermethylated in *h1met1* (Figures 5E and S5F). Thus, both groups are likely differentially

expressed between *met1* and *h1met1* due to changes in DNA methylation. However, the third –

and largest – group (up3; 159 TEs) is upregulated in *h1met1* despite substantial hypermethylation

(Figures 5E, 5F and S5G). Compared to the dn group, these TEs are significantly H1-enriched

(Figures 5G, 5H and S5D). TEs in the up3 group, like those in group up1 described above, are

mildly activated in *met1* and require loss of H1 for strong expression (Figure 5E). These data

indicate that DNA methylation and H1 jointly create an inaccessible chromatin state that silences

TEs.

DNA methylation and H1 cooperatively suppress intragenic antisense transcripts

The ability of H1 and DNA methylation to cooperatively repress transcription, and enrichment of

H1 within methylated genes (Figure 1E), are consistent with the possibility that methylation and

H1 act together to reduce aberrant intragenic transcripts, a long-hypothesized function of gbM

(Tran et al., 2005; Zilberman et al., 2007). To test this hypothesis, we identified antisense

intragenic transcripts with significantly altered expression in *h1, met1* and *h1met1* mutants in

comparison to *wt*. We identified 17 such transcripts in *h1*, most of which (15, 88%) are upregulated

(Figure 6A). This result is consistent with the hypothesis that H1 suppresses intragenic

transcription, but the small number of identified transcripts limits confidence and prevents further

analysis. We found many more mis-expressed antisense transcripts in *met1* (468), and yet more in

h1met1 (1152; Figure 6A). In *met1,* only 32% of transcripts (149) are upregulated (Figure 6A). In

contrast, 91% of transcripts (1044) are upregulated in *h1met1* (Figure 6A), supporting the

hypothesis that intragenic transcription is de-repressed when DNA methylation and H1 are

simultaneously absent.

We next separated antisense transcripts for which the change in expression is positively

correlated with the sense transcript from those with an absent or negative correlation (some

antisense transcripts were ambiguously correlated and therefore were removed from this analysis).

A positive correlation suggests that antisense expression is altered due to increased or decreased

overall transcriptional activity at the locus, whereas uncorrelated or negatively correlated antisense

transcription must be regulated differently. In *met1* plants, sense and antisense expression is

correlated in most cases (65%, 262 out of 403 transcripts in Figure 6B and Table S4), indicating

that the antisense transcriptional changes we detected are mainly due to mis-regulation of sense

transcripts. However, negatively or un-correlated, upregulated antisense transcripts are much more

likely to initiate from methylated DNA than positively correlated, downregulated, or unchanged

transcripts (Figures 6B and 6C). This result suggests that antisense expression is activated by

methylation loss, but we identified just 47 (12%) such transcripts (Figures 6B and 6C). Thus, *met1*

data provide only modest support for the hypothesis that gbM suppresses antisense expression.

The situation is substantially different in *h1met1* plants, in which antisense expression is

most commonly upregulated and not positively correlated with sense expression (83%, 894 out of

1074 transcripts in Figure 6D and Table S4). As in *met1,* these transcripts are much more likely to

initiate from DNA methylated in *wt* than other antisense transcripts (Figures 6D-F and S6A, S6B),

indicating that gbM and H1 cooperatively silence antisense transcription.

To further analyze the link between gbM and antisense expression, we separated all the

antisense transcripts we annotated (regardless of their behavior in mutant genotypes) into three

groups based on their point of initiation: those from methylated DNA, from unmethylated DNA

within methylated genes, or from unmethylated genes (Figure 6G). As expected, transcripts that

initiate from methylated DNA have more H1 around the TSS (Figures 6H and S6C, S6D). In *met1*,

levels of transcripts arising from methylated genes are significantly elevated compared to

unmethylated genes (Figure 6I). Transcripts initiating from methylated sequences are also

significantly elevated compared to those initiating from unmethylated regions of methylated genes

(Figure 6I). In *h1met1*, levels of transcripts arising from methylated genes are also significantly

elevated, and transcripts initiating from methylated DNA are especially upregulated (Figure 6I),

specifically linking methylation loss with antisense transcript activation.

TEs that are highly active in the absence of H1 and DNA methylation tend to show weak

upregulation when methylation alone is lost (Figures 5A and 5E). We therefore asked if antisense

intragenic transcription is analogously affected. Indeed, antisense transcription tends to behave

similarly in *met1* and *h1met1* (Figure 6J). Of particular interest, transcripts that are significantly

upregulated only in *h1met1* are usually also activated to some extent in *met1* (Figures 6J and S6B).

Thus, as with TEs, loss of H1 hyperactivates antisense transcripts that are released from silencing

by elimination of intragenic methylation.

Discussion

DNA methylation and histone H1 are abundant and widely conserved constituents of eukaryotic chromatin associated with transcriptional inactivity (Fyodorov et al., 2018; Torres et al., 2016; Zhang et al., 2018a), and their interactions have been of interest for many years. There is now extensive evidence that H1 modulates DNA methylation pathways. Loss of H1 causes genome-wide hypermethylation in ascomycete fungi (Barra et al., Mol Cell Biol 2000; Seymour, G3 2016) and reduces methylation at specific loci in mouse (Fan et al., 2005; Geeven et al., 2015; MacLean et al., 2011; Yang et al., 2013). In *Arabidopsis*, loss of H1 reduces methylation of euchromatic TEs and increases methylation of heterochromatic elements in all sequence contexts (Rutowicz et al., 2015; Zemach et al., 2013). H1 also appears to impede DNA demethylation in *Arabidopsis* heterochromatin (He et al., 2018). H1 hinders heterochromatic methylation by restricting the access of DNA methyltransferases (Lyons and Zilberman, 2017), and likely interferes with demethylation in an analogous way. The ability of H1 to globally influence nucleosome positions in plants (Figure 2) and animals (Baldi et al., 2018; Fan et al., 2003; Woodcock et al., 2006) suggests an additional regulatory mechanism, because nucleosomes are substantial obstacles to DNA methylation (Baubec et al., 2015; Huff and Zilberman, 2014; Lyons and Zilberman, 2017).

The effects of DNA methylation on H1 are far less established. A number of studies evaluated whether H1 has preferential affinity for methylated DNA *in vitro* but came to opposing conclusions (Campoy et al., 1995; Hashimshony et al., 2003; Levine et al., 1993; McArthur and Thomas, 1996; Nightingale and Wolffe, 1995). Our results show that H1 is enriched at methylated loci in *Arabidopsis*, but this enrichment does not depend on DNA methylation *in vivo* (Figure 1). Our data indicate that the regulatory relationship between H1 and DNA methylation is

unidirectional: H1 modulates, but is not directly affected by, DNA methylation. However,

methylation indirectly influences H1 by, for example, enforcing TE silencing (Figure 1F).

The complex relationship between H1 and DNA methylation ultimately impinges on

chromatin accessibility and transcription. Our results indicate that H1 and DNA methylation

jointly maintain heterochromatic TEs in an inaccessible and silent state (Figure 5). Methylation is

the more powerful repressor, as few TEs are activated in *h1* mutants, and many of those that gain

activity also lose methylation (Figure 4). However, the role of H1 becomes obvious when

methylation is reduced, and loss of H1 can even overcome modest hypermethylation to activate

TEs (Figure 5E). These results may explain why H1 depletion has a far more drastic effect on TE

activity in *Drosophila* (Iwasaki et al., 2016; Lu et al., 2013), which lacks cytosine DNA

methylation (Raddatz et al., 2013; Zemach et al., 2010), than in mouse (Fan et al., 2005).

The cooperative silencing of transcription by methylation and H1 extends to gene bodies

(Figure 6). The function of gbM has been extensively debated for over a decade. One line of

argument is that gbM reprogramming regulates development (Herb et al., 2012), for example by

modulating alternative splicing (Lyko et al., 2010; Park et al., 2011). These ideas are supported by

developmental disruptions caused by erasure of methylation and/or downregulation of DNA

methyltransferases in species in which methylation is exclusive or near-exclusive to gene bodies

(Bewick et al., 2018; Biergans et al., 2017; Kucharski et al., 2008). However, in every species

examined so far, gbM is concentrated in constitutively expressed housekeeping genes (Coleman-

Derr and Zilberman, 2012; Dixon et al., 2016; Sarda et al., 2012; Zilberman, 2017). These genes

do not vary in expression during development and thus are not good candidates for differential

alternative splicing or other developmental regulation. There is also little evidence for substantial

shifts in gbM patterns during plant or animal development (Bartels et al., 2018; Libbrecht et al.,

2016; Suzuki et al., 2013; Zemach et al., 2018). Another argument is that gbM is a functionless

consequence of TE methylation (Bewick and Schmitz, 2017; Bewick et al., 2016; Teixeira and

Colot, 2009), motivated by the inability to detect transcriptional or RNA processing changes

caused by genetic removal of gbM. This argument is undermined by the well-established

observation that methylation is mutagenic in plants and animals (Alexandrov et al., 2013; Takuno

and Gaut, 2013). The maintenance of DNA methylation within the exons of some of the most

conserved and essential genes, especially in species that do not methylate other sequences, implies

a function that is important enough to outweigh the costs of increased mutation rates (Hunt et al.,

2013; Takuno and Gaut, 2013).

The last line of argument is that developmentally invariant methylation within

housekeeping genes should have one or more housekeeping functions (Zilberman, 2017). This is

supported by observations of small overall drops in gene expression associated with evolutionary

loss of gbM in plants (Muyle and Gaut, 2018; Takuno et al., 2017). One housekeeping function,

the inhibition of intragenic aberrant transcripts, was proposed when gbM was first discovered

(Tran et al., 2005). Mechanisms to silence such transcripts have been described in multiple species,

indicating that this is an important feature of transcriptional homeostasis (Carrozza et al., 2005;

Kaplan et al., 2003; Neri et al., 2017; Whitehouse et al., 2007). Here we provide the long-sought

experimental evidence for this gbM function. Our data show that intragenic antisense transcription

is upregulated in *h1met1* plants (Figure 6A). Most upregulated transcripts are anticorrelated with

sense expression (Figure 6D) and about half of these (55%) initiate from methylated DNA – far

more than is expected by random chance (Figure 6E). Conversely, antisense transcripts that initiate

from methylated DNA are preferentially upregulated in *h1met1* (Figure 6I). Antisense transcripts

activated in *h1met1* also tend to be upregulated, albeit more weakly, in *met1* (Figure 6J). Taken

together with the observed silencing of TEs by methylation and H1, the suppression of gene

transcription by H1, and the ability of H1 to associate with chromatin independently of DNA

methylation, our results strongly support the hypothesis that H1 and DNA methylation

cooperatively repress aberrant intragenic transcripts in *Arabidopsis*. The strong similarities

between gbM patterns of plants and animals (Feng et al., 2010; Zemach et al., 2010), and the

abundance of H1 in animal genes (Cao et al., 2013; Millán-Ariño et al., 2014), suggest that this

function is general, and potentially universal among eukaryotes with intragenic methylation.

Acknowledgements

We thank X. Feng, E. Hollwey and S. de-Bruijin for helpful comments on the manuscript. We also

thank S.L. McDevitt and M. Chung for sequencing and C. Wistrom for greenhouse assistance.

Funding: This work was supported by a fellowship from the Human Frontier Science Program

(LT000667/2013) and the National Research Foundation of Korea (2013022168) to J.C., by a

fellowship from the Helen Hay Whitney Foundation to D.B.L., by a European Research Council

grant (725746) and a Faculty Scholar grant from HHMI and the Simons Foundation (55108592)

to D.Z., and by the National Institutes of Health S10 program (1S10RR026866-01).

Author contributions

J.C. designed and performed experiments, analyzed data and wrote the manuscript. D.B.L.

performed cytology, image analysis, NRL calculation and MNase-seq analysis. M.Y.K. performed

H1 ChAP-seq experiments. J.D.M. defined the boundary of gbM regions. D.Z. conceived and

supervised the project and wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

357
358

References

Alexandrov, L.B., Nik-Zainal, S., Wedge, D.C., Aparicio, S.A.J.R., Behjati, S., Biankin, A. V.,
Bignell, G.R., Bolli, N., Borg, A., Børresen-Dale, A.L., et al. (2013). Signatures of mutational
processes in human cancer. Nature *500*, 415–421.

Ascenzi, R., and Gantt, J.S. (1999). Subnuclear distribution of the entire complement of linker
histone variants in Arabidopsis thaliana. Chromosoma *108*, 345–355.

Baldi, S., Krebs, S., Blum, H., and Becker, P.B. (2018). Genome-wide measurement of local
nucleosome array regularity and spacing by nanopore sequencing. Nat. Struct. Mol. Biol. *25*,
894–901.

Bartels, A., Han, Q., Nair, P., Stacey, L., Gaynier, H., Mosley, M., Huang, Q.Q., Pearson, J.K.,
Hsieh, T.F., An, Y.Q.C., et al. (2018). Dynamic DNA methylation in plant growth and
development. Int. J. Mol. Sci. *19*, 2144.

Baubec, T., Colombo, D.F., Wirbelauer, C., Schmidt, J., Burger, L., Krebs, A.R., Akalin, A., and
Schübeler, D. (2015). Genomic profiling of DNA methyltransferases reveals a role for DNMT3B
in genic methylation. Nature *520*, 243–247.

Bewick, A.J., and Schmitz, R.J. (2017). Gene body DNA methylation in plants. Curr. Opin. Plant
Biol. *36*, 103–110.

Bewick, A.J., Ji, L., Niederhuth, C.E., Willing, E.-M., Hofmeister, B.T., Shi, X., Wang, L., Lu,
Z., Rohr, N.A., Hartwig, B., et al. (2016). On the origin and evolutionary consequences of gene
body DNA methylation. Proc. Natl. Acad. Sci. USA *113*, 9111–9116.

Bewick, A.J., Vogel, K.J., Moore, A.J., and Schmitz, R.J. (2017). Evolution of DNA methylation
across insects. Mol. Biol. Evol. *34*, 654–665.

Bewick, A.J., Sanchez, Z., Mckinney, E.C., Moore, A.J., Moore, P.J., and Schmitz, R.J. (2018).
Gene-regulatory independent functions for insect DNA methylation. BioRxiv 355669.

Biergans, S.D., Claudianos, C., Reinhard, J., and Galizia, C.G. (2017). DNA methylation
mediates neural processing after odor learning in the honeybee. Sci. Rep. *7*, 43635.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-
seq quantification. Nat. Biotechnol. *34*, 525–527.

Burger, L., Gaidatzis, D., Schübeler, D., and Stadler, M.B. (2013). Identification of active
regulatory regions from DNA methylation data. Nucleic Acids Res. *41*, e155.

Campoy, F.J., Meehan, R.R., McKay, S., Nixon, J., and Bird, A. (1995). Binding of histone H1
to DNA is indifferent to methylation at CpG sequences. J. Biol. Chem. *270*, 26473–26481.

Cao, K., Lailler, N., Zhang, Y., Kumar, A., Uppal, K., Liu, Z., Lee, E.K., Wu, H., Medrzycki,
M., Pan, C., et al. (2013). High-Resolution Mapping of H1 Linker Histone Variants in

393 Embryonic Stem Cells. PLoS Genet. *9*, e1003417.

394 Carrozza, M.J., Li, B., Florens, L., Suganuma, T., Swanson, S.K., Lee, K.K., Shia, W.J.,
395 Anderson, S., Yates, J., Washburn, M.P., et al. (2005). Histone H3 methylation by Set2 directs
396 deacetylation of coding regions by Rpd3S to suppress spurious intragenic transcription. Cell *123*,
397 581–592.

398 Chambon, A., West, A., Vezon, D., Horlow, C., De Muyt, A., Chelysheva, L., Ronceret, A.,
399 Darbyshire, A.R., Osman, K., Heckmann, S., et al. (2018). Identification of ASYNAPTIC4, a
400 component of the meiotic chromosome axis. Plant Physiol. *178*, 233–246.

401 Chen, C., Li, C., Wang, Y., Renaud, J., Tian, G., Kambhampati, S., Saatian, B., Nguyen, V.,
402 Hannoufa, A., Marsolais, F., et al. (2017). Cytosolic acetyl-CoA promotes histone acetylation
403 predominantly at H3K27 in Arabidopsis. Nat. Plants *3*, 814–824.

404 Cheng, C.-Y., Krishnakumar, V., Chan, A.P., Thibaud-Nissen, F., Schobel, S., and Town, C.D.
405 (2017). Araport11: a complete reannotation of the *Arabidopsis thaliana* reference genome. Plant
406 J. *89*, 789–804.

407 Choi, K., Zhao, X., Tock, A.J., Lambing, C., Underwood, C.J., Hardcastle, T.J., Serra, H., Kim,
408 J., Cho, H.S., Kim, J., et al. (2018). Nucleosomes and DNA methylation shape meiotic DSB
409 frequency in Arabidopsis thaliana transposons and gene regulatory regions. Genome Res. *28*,
410 532–546.

411 Cokus, S.J., Feng, S., Zhang, X., Chen, Z., Merriman, B., Haudenschild, C.D., Pradhan, S.,
412 Nelson, S.F., Pellegrini, M., and Jacobsen, S.E. (2008). Shotgun bisulphite sequencing of the
413 Arabidopsis genome reveals DNA methylation patterning. Nature *452*, 215–219.

414 Coleman-Derr, D., and Zilberman, D. (2012). DNA methylation, H2A.Z, and the regulation of
415 constitutive expression. Cold Spring Harb. Symp. Quant. Biol. *77*, 147–154.

416 Davey, C., Pennings, S., and Allan, J. (1997). CpG methylation remodels chromatin structure in
417 vitro. J. Mol. Biol. *267*, 276–288.

418 Deaton, M., Webb, S., Kerr, A.R.W., Illingworth, R.S., Guy, J., Andrews, R., and Bird, A.
419 (2011). Cell type – specific DNA methylation at intragenic CpG islands in the immune system.
420 Genome Researh *21*, 1074–1086.

421 Deleris, A., Stroud, H., Bernatavichute, Y., Johnson, E., Klein, G., Schubert, D., and Jacobsen,
422 S.E. (2012). Loss of the DNA methyltransferase MET1 induces H3K9 hypermethylation at PcG
423 target genes and redistribution of H3K27 trimethylation to transposons in Arabidopsis thaliana.
424 PLoS Genet. *8*, e1003062.

425 Dixon, G.B., Bay, L.K., and Matz, M. V. (2016). Evolutionary consequences of DNA
426 methylation in a basal metazoan. Mol. Biol. Evol. *33*, 2285–2293.

427 Du, J., Johnson, L.M., Jacobsen, S.E., and Patel, D.J. (2015). DNA methylation pathways and
428 their crosstalk with histone methylation. Nat. Rev. Mol. Cell Biol. *16*, 519–532.

Fan, Y., Nikitina, T., Morin-Kensicki, E.M., Zhao, J., Magnuson, T.R., Woodcock, C.L., and
Skoultchi, A.I. (2003). H1 linker histones are essential for mouse development and affect
nucleosome spacing in vivo. Mol. Cell. Biol. *23*, 4559–4572.

Fan, Y., Nikitina, T., Zhao, J., Fleury, T.J., Bhattacharyya, R., Bouhassira, E.E., Stein, A.,
Woodcock, C.L., and Skoultchi, A.I. (2005). Histone H1 depletion in mammals alters global
chromatin structure but causes specific changes in gene regulation. Cell *123*, 1199–1212.

Feng, W., and Michaels, S.D. (2015). Accessing the inaccessible: The organization,
transcription, replication, and repair of heterochromatin in plants. Annu. Rev. Genet. *49*, 439–
459.

Feng, S., Cokus, S.J., Zhang, X., Chen, P.-Y., Bostick, M., Goll, M.G., Hetzel, J., Jain, J.,
Strauss, S.H., Halpern, M.E., et al. (2010). Conservation and divergence of methylation
patterning in plants and animals. Proc. Natl. Acad. Sci. USA *107*, 8689–8694.

Fransz, P., de Jong, J.H., Lysak, M., Castiglione, M.R., and Schubert, I. (2002). Interphase
chromosomes in Arabidopsis are organized as well defined chromocenters from which
euchromatin loops emanate. Proc. Natl. Acad. Sci. USA *99*, 14584–14589.

Fyodorov, D. V., Zhou, B.-R., Skoultchi, A.I., and Bai, Y. (2018). Emerging roles of linker
histones in regulating chromatin structure and function. Nat. Rev. Mol. Cell Biol. *19*, 192–206.

Geeven, G., Zhu, Y., Kim, B.J., Bartholdy, B.A., Yang, S.-M., Macfarlan, T.S., Gifford, W.D.,
Pfaff, S.L., Verstegen, M.J.A.M., Pinto, H., et al. (2015). Local compartment changes and
regulatory landscape alterations in histone H1-depleted cells. Genome Biol. *16*, 289.

Hashimshony, T., Zhang, J., Keshet, I., Bustin, M., and Cedar, H. (2003). The role of DNA
methylation in setting up chromatin structure during development. Nat. Genet. *34*, 187–192.

He, S., Vickers, M., Zhang, J., and Feng, X. (2018). Natural depletion of H1 in sex cells causes
DNA demethylation, heterochromatin decondensation and transposon activation. BioRxiv
451930.

Hellman, A., and Chess, A. (2007). Gene body–specific methylation on the active X
chromosome. Science *315*, 1141–1143.

Herb, B.R., Wolschin, F., Hansen, K.D., Aryee, M.J., Langmead, B., Irizarry, R., Amdam, G. V.,
and Feinberg, A.P. (2012). Reversible switching between epigenetic states in honeybee
behavioral subcastes. Nat. Neurosci. *15*, 1371–1373.

Hergeth, S.P., and Schneider, R. (2015). The H1 linker histones: multifunctional proteins beyond
the nucleosomal core particle. EMBO Rep. *16*, 1439–1453.

Hon, G.C., Rajagopal, N., Shen, Y., McCleary, D.F., Yue, F., Dang, M.D., and Ren, B. (2013).
Epigenetic memory at embryonic enhancers identified in DNA methylation maps from adult
mouse tissues. Nat. Genet. *45*, 1198–1206.

de Hoon, M.J.L., Imoto, S., Nolan, J., and Miyano, S. (2004). Open source clustering software.
Bioinformatics *20*, 1453–1454.

Huff, J.T., and Zilberman, D. (2014). Dnmt1-independent CG methylation contributes to
nucleosome positioning in diverse eukaryotes. Cell *156*, 1286–1297.

Hunt, B.G., Glastad, K.M., Yi, S. V., and Goodisman, M.A.D. (2013). Patterning and regulatory
associations of DNA methylation are mirrored by histone modifications in insects. Genome Biol.
Evol. *5*, 591–598.

Iurlaro, M., von Meyenn, F., and Reik, W. (2017). DNA methylation homeostasis in human and
mouse development. Curr. Opin. Genet. Dev. *43*, 101–109.

Iwasaki, Y.W., Murano, K., Ishizu, H., Shibuya, A., Iyoda, Y., Siomi, M.C., Siomi, H., and
Saito, K. (2016). Piwi modulates chromatin accessibility by regulating multiple factors including
histone H1 to repress transposons. Mol. Cell *63*, 408–419.

Johnson, C.A., Goddard, J.P., and Adams, R.L. (1995). The effect of histone H1 and DNA
methylation on transcription. Biochem. J. *305 (Pt 3)*, 791–798.

Jones, P.A. (2012). Functions of DNA methylation: islands, start sites, gene bodies and beyond.
Nat. Rev. Genet. *13*, 484–492.

Jones, P.A., Issa, J.P.J., and Baylin, S. (2016). Targeting the cancer epigenome for therapy. Nat.
Rev. Genet. *17*, 630–641.

Jühling, F., Kretzmer, H., Bernhart, S.H., Otto, C., Stadler, P.F., and Hoffmann, S. (2016).
metilene: fast and sensitive detection of differentially methylated regions from bisulfite
sequencing data. *26*, 256–262.

Kaplan, C.D., Laprade, L., and Winston, F. (2003). Transcription elongation factors repress
transcription inititaion from cryptic sites. Science *301*, 1096–1099.

Kawakatsu, T., Huang, S.C., Jupe, F., Sasaki, E., Schmitz, R.J., Urich, M.A., Castanon, R., Nery,
J.R., Barragan, C., He, Y., et al. (2016). Epigenomic diversity in a global collection of
Arabidopsis thaliana accessions. Cell *166*, 492–505.

Kawashima, T., and Berger, F. (2014). Epigenetic reprogramming in plant sexual reproduction.
Nat. Rev. Genet. *15*, 613–624.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low
memory requirements. Nat. Methods *12*, 357–360.

Krishnakumar, R., Gamble, M.J., Frizzell, K.M., Berrocal, J.G., Kininis, M., and Kraus, W.L.
(2008). Reciprocal binding of PARP-1 and histone H1 at promoters specifies transcriptional
outcomes. Science *319*, 819–821.

Kucharski, R., Maleszka, J., Foret, S., and Maleszka, R. (2008). Nutritional control of

reproductive status in honeybees via DNA methylation. Science *319*, 1827–1831.

Kulis, M., Heath, S., Bibikova, M., Queirós, A.C., Navarro, A., Clot, G., Martínez-Trillos, A.,
Castellano, G., Brun-Heath, I., Pinyol, M., et al. (2012). Epigenomic analysis detects widespread
gene-body DNA hypomethylation in chronic lymphocytic leukemia. Nat. Genet. *44*, 1236–1242.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S.L. (2009). Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. *10*, R25.

Law, J.A., and Jacobsen, S.E. (2010). Establishing, maintaining and modifying DNA
methylation patterns in plants and animals. Nat. Rev. Genet. *11*, 204–220.

Lev Maor, G., Yearim, A., and Ast, G. (2015). The alternative role of DNA methylation in
splicing regulation. Trends Genet. *31*, 274–280.

Levine, A., Yeivin, A., Ben-Asher, E., Aloni, Y., and Razin, A. (1993). Histone H1-mediated
inhibition of transcription initiation of methylated templates in vitro. J. Biol. Chem. *268*, 21754–
21759.

Libbrecht, R., Oxley, P.R., Keller, L., and Kronauer, D.J.C. (2016). Robust DNA methylation in
the clonal raider ant brain. Curr. Biol. *26*, 391–395.

Lister, R., O'Malley, R.C., Tonti-Filippini, J., Gregory, B.D., Berry, C.C., Millar, A.H., and
Ecker, J.R. (2008). Highly integrated single-base resolution maps of the epigenome in
Arabidopsis. Cell *133*, 523–536.

Lister, R., Pelizzola, M., Dowen, R.H., Hawkins, R.D., Hon, G., Tonti-Filippini, J., Nery, J.R.,
Lee, L., Ye, Z., Ngo, Q.-M., et al. (2009). Human DNA methylomes at base resolution show
widespread epigenomic differences. Nature *462*, 315–322.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

Lu, X., Wontakal, S.N., Emelyanov, A. V., Morcillo, P., Konev, A.Y., Fyodorov, D. V., and
Skoultchi, A.I. (2009). Linker histone H1 is essential for Drosophila development, the
establishment of pericentric heterochromatin, and a normal polytene chromosome structure.
Genes Dev. *23*, 452–465.

Lu, X., Wontakal, S.N., Kavi, H., Kim, B.J., Guzzardo, P.M., Emelyanov, A. V, Xu, N., Hannon,
G.J., Zavadil, J., Fyodorov, D. V, et al. (2013). Drosophila H1 regulates the genetic activity of
heterochromatin by recruitment of Su(var)3-9. Science *340*, 78–81.

Luo, C., Sidote, D.J., Zhang, Y., Kerstetter, R.A., Michael, T.P., and Lam, E. (2013). Integrative
analysis of chromatin states in Arabidopsis identified potential regulatory mechanisms for
natural antisense transcript production. Plant J. *73*, 77–90.

Lyko, F., Foret, S., Kucharski, R., Wolf, S., Falckenhayn, C., and Maleszka, R. (2010). The
honey bee epigenomes: Differential methylation of brain DNA in queens and workers. PLoS

Biol. *8*, e1000506.

Lyons, D.B., and Zilberman, D. (2017). DDM1 and Lsh remodelers allow methylation of DNA
wrapped in nucleosomes. Elife *6*, e30674.

MacLean, J.A., Bettegowda, A., Kim, B.J., Lou, C.-H., Yang, S.-M., Bhardwaj, A., Shanker, S.,
Hu, Z., Fan, Y., Eckardt, S., et al. (2011). The Rhox homeobox gene cluster is imprinted and
selectively targeted for regulation by histone H1 and DNA methylation. Mol. Cell. Biol. *31*,
1275–1287.

Maunakea, A.K., Nagarajan, R.P., Bilenky, M., Ballinger, T.J., D'Souza, C., Fouse, S.D.,
Johnson, B.E., Hong, C., Nielsen, C., Zhao, Y., et al. (2010). Conserved role of intragenic DNA
methylation in regulating alternative promoters. Nature *466*, 253–257.

Maunakea, A.K., Chepelev, I., Cui, K., and Zhao, K. (2013). Intragenic DNA methylation
modulates alternative splicing by recruiting MeCP2 to promote exon recognition. Cell Res. *23*,
1256–1269.

McArthur, M., and Thomas, J.O. (1996). A preference of histone H1 for methylated DNA.
EMBO J. *15*, 1705–1714.

Mercier, R., Mézard, C., Jenczewski, E., Macaisne, N., and Grelon, M. (2015). The Molecular
Biology of Meiosis in Plants. Annu. Rev. Plant Biol. *66*, 297–327.

Millán-Ariño, L., Islam, A.B.M.M.K., Izquierdo-Bouldstridge, A., Mayor, R., Terme, J.-M.,
Luque, N., Sancho, M., López-Bigas, N., and Jordan, A. (2014). Mapping of six somatic linker
histone H1 variants in human breast cancer cells uncovers specific features of H1.2. Nucleic
Acids Res. *42*, 4474–4493.

Morselli, M., Pastor, W.A., Montanini, B., Nee, K., Ferrari, R., Fu, K., Bonora, G., Rubbi, L.,
Clark, A.T., Ottonello, S., et al. (2015). In vivo targeting of de novo DNA methylation by
histone modifications in yeast and mouse. Elife *4*, e06205.

Muyle, A., and Gaut, B.S. (2018). Loss of gene body methylation in Eutrema salsugineum is
associated with reduced gene expression. Mol. Biol. Evol. doi: 10.1093/molbev/msy204.

Neri, F., Rapelli, S., Krepelova, A., Incarnato, D., Parlato, C., Basile, G., Maldotti, M., Anselmi,
F., and Oliviero, S. (2017). Intragenic DNA methylation prevents spurious transcription
initiation. Nature *543*, 72–77.

Nightingale, K., and Wolffe, A.P. (1995). Methylation at CpG sequences does not influence
histone H1 binding to a nucleosome including a Xenopus borealis 5 S rRNA gene. J. Biol. Chem.
270, 4197–4200.

Olmedo-Monfil, V., Durán-Figueroa, N., Arteaga-Vázquez, M., Demesa-Arévalo, E., Autran, D.,
Grimanelli, D., Slotkin, R.K., Martienssen, R.A., and Vielle-Calzada, J.P. (2010). Control of
female gamete formation by a small RNA pathway in Arabidopsis. Nature *464*, 628–632.

Over, R.S., and Michaels, S.D. (2014). Open and closed: The roles of linker histones in plants
and animals. Mol. Plant *7*, 481–491.

Park, J., Peng, Z., Zeng, J., Elango, N., Park, T., Wheeler, D., Werren, J.H., and Yi, S. V. (2011).
Comparative analyses of DNA methylation and sequence evolution using Nasonia genomes.
Mol. Biol. Evol. *28*, 3345–3354.

Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.-C., Mendell, J.T., and Salzberg, S.L.
(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat.
Biotechnol. *33*, 290–295.

Pertea, M., Kim, D., Pertea, G.M., Leek, J.T., and Salzberg, S.L. (2016). Transcript-level
expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown. Nat. Protoc.
11, 1650–1667.

Pimentel, H., Bray, N.L., Puente, S., Melsted, P., and Pachter, L. (2017). Differential analysis of
RNA-seq incorporating quantification uncertainty. Nat. Methods *14*, 687–690.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.

Raddatz, G., Guzzardo, P.M., Olova, N., Fantappie, M.R., Rampp, M., Schaefer, M., Reik, W.,
Hannon, G.J., and Lyko, F. (2013). Dnmt2-dependent methylomes lack defined DNA
methylation patterns. Proc. Natl. Acad. Sci. USA *110*, 8627–8631.

Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res. *44*, W160–W165.

Rasmussen, K.D., and Helin, K. (2016). Role of TET enzymes in DNA methylation,
development, and cancer. Genes Dev. *30*, 733–750.

Regulski, M., Lu, Z., Kendall, J., Donoghue, M.T.A., Reinders, J., Llaca, V., Deschamps, S.,
Smith, A., Levy, D., McCombie, W.R., et al. (2013). The maize methylome influences mRNA
splice sites and reveals widespread paramutation-like switches guided by small RNA. Genome
Res. *23*, 1651–1662.

Robertson, K.D. (2005). DNA methylation and human disease. Nat. Rev. Genet. *6*, 597–610.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

Rutowicz, K., Puzio, M., Halibart-Puzio, J., Lirski, M., Kotliński, M., Kroteń, M.A., Knizewski,
L., Lange, B., Muszewska, A., Śniegowska-Świerk, K., et al. (2015). A specialized histone H1
variant is required for adaptive responses to complex abiotic stress and related DNA methylation
in Arabidopsis. Plant Physiol. *169*, 2080–2101.

Sarda, S., Zeng, J., Hunt, B.G., and Yi, S. V. (2012). The evolution of invertebrate gene body

methylation. Mol. Biol. Evol. *29*, 1907–1916.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for
biological-image analysis. Nat. Methods *9*, 676–682.

Schmidt, A., Schmid, M.W., and Grossniklaus, U. (2015). Plant germline formation: common
concepts and developmental flexibility in sexual and asexual reproduction. Development *142*,
229–241.

Schmitz, R.J., Schultz, M.D., Lewsey, M.G., O'Malley, R.C., Urich, M.A., Libiger, O., Schork,
N.J., Ecker, J.R., and Epigenetic (2011). Transgenerational epigenetic instability is a source of
novel methylation variants. Science *334*, 369–373.

Schübeler, D. (2015). Function and information content of DNA methylation. Nature *517*, 321–
326.

She, W., and Baroux, C. (2015). Chromatin dynamics in pollen mother cells underpin a common
scenario at the somatic-to-reproductive fate transition of both the male and female lineages in
Arabidopsis. Front. Plant Sci. *6*, Article 294.

She, W., Grimanelli, D., Rutowicz, K., Whitehead, M.W.J., Puzio, M., Kotliński, M.,
Jerzmanowski, A., and Baroux, C. (2013). Chromatin reprograming during the somatic-to-
reproductive cell fate transition in plants. Development *140*, 4008–4019.

Shukla, S., Kavak, E., Gregory, M., Imashimizu, M., Shutinoski, B., Kashlev, M., Oberdoerffer,
P., Sandberg, R., and Oberdoerffer, S. (2011). CTCF-promoted RNA polymerase II pausing links
DNA methylation to splicing. Nature *479*, 74–79.

Simpson, R.T. (1978). Structure of the chromatosome, a chromatin particle containing 160 base
pairs of DNA and all the histones. Biochemistry *17*, 5524–5531.

Smith, Z.D., and Meissner, A. (2013). DNA methylation: Roles in mammalian development.
Nat. Rev. Genet. *14*, 204–220.

Stempor, P., and Ahringer, J. (2016). SeqPlots - Interactive software for exploratory data
analyses, pattern discovery and visualization in genomics. Wellcome Open Res. *1*, 14.

Suzuki, M.M., and Bird, A. (2008). DNA methylation landscapes: provocative insights from
epigenomics. Nat. Rev. Genet. *9*, 465–476.

Suzuki, M.M., Kerr, A.R.W., De Sousa, D., and Bird, A. (2007). CpG methylation is targeted to
transcription units in an invertebrate genome. Genome Res. *17*, 625–631.

Suzuki, S., Shaw, G., and Renfree, M.B. (2013). Postnatal epigenetic reprogramming in the
germline of a marsupial, the tammar wallaby. Epigenetics and Chromatin *6*, 14.

Takuno, S., and Gaut, B.S. (2012). Body-methylated genes in Arabidopsis thaliana are

functionally important and evolve slowly. Mol. Biol. Evol. *29*, 219–227.

Takuno, S., and Gaut, B.S. (2013). Gene body methylation is conserved between plant orthologs
and is of evolutionary consequence. Proc. Natl. Acad. Sci. USA *110*, 1797–1802.

Takuno, S., Seymour, D.K., and Gaut, B.S. (2017). The evolutionary dynamics of orthologs that
shift in gene body methylation between Arabidopsis species. Mol. Biol. Evol. *34*, 1479–1491.

Teissandier, A., and Bourc'his, D. (2017). Gene body DNA methylation conspires with
H3K36me3 to preclude aberrant transcription. EMBO J. *36*, 1471–1473.

Teixeira, F.K., and Colot, V. (2009). Gene body DNA methylation in plants: a means to an end
or an end to a means? EMBO J. *28*, 997–998.

Thorvaldsdottir, H., Robinson, J.T., and Mesirov, J.P. (2013). Integrative Genomics Viewer
(IGV): high-performance genomics data visualization and exploration. Brief. Bioinform. *14*,
178–192.

To, T.K., Saze, H., and Kakutani, T. (2015). DNA Methylation within transcribed regions. Plant
Physiol. *168*, 1219–1225.

Torres, C.M., Biran, A., Burney, M.J., Patel, H., Henser-Brownhill, T., Cohen, A.-H.S., Li, Y.,
Ben-Hamo, R., Nye, E., Spencer-Dene, B., et al. (2016). The linker histone H1.0 generates
epigenetic and functional intratumor heterogeneity. Science *353*, aaf1644.

Tran, R.K., Henikoff, J.G., Zilberman, D., Ditt, R.F., Jacobsen, S.E., and Henikoff, S. (2005).
DNA methylation profiling identifies CG methylation clusters in Arabidopsis genes. Curr. Biol.
15, 154–159.

Valouev, A., Johnson, S.M., Boyd, S.D., Smith, C.L., Fire, A.Z., and Sidow, A. (2011).
Determinants of nucleosome organization in primary human cells. Nature *474*, 516–520.

Wagner, E.J., and Carpenter, P.B. (2012). Understanding the language of Lys36 methylation at
histone H3. Nat. Rev. Mol. Cell Biol. *13*, 115–126.

Whitehouse, I., Rando, O.J., Delrow, J., and Tsukiyama, T. (2007). Chromatin remodelling at
promoters suppresses antisense transcription. Nature *450*, 1031–1035.

Wierzbicki, A.T., and Jerzmanowski, A. (2005). Suppression of histone H1 genes in Arabidopsis
results in heritable developmental defects and stochastic changes in DNA methylation. Genetics
169, 997–1008.

Woodcock, C.L., Skoultchi, A.I., and Fan, Y. (2006). Role of linker histone in chromatin
structure and function: H1 stoichiometry and nucleosome repeat length. Chromosom. Res. *14*,
17–25.

Xiao, W., Gehring, M., Choi, Y., Margossian, L., Pu, H., Harada, J.J., Goldberg, R.B., Pennell,
R.I., and Fischer, R.L. (2003). Imprinting of the MEA polycomb gene is controlled by

antagonism between MET1 methyltransferase and DME glycosylase. Dev. Cell *5*, 891–901.

Yang, S.-M., Kim, B.J., Norwood Toro, L., and Skoultchi, A.I. (2013). H1 linker histone
promotes epigenetic silencing by regulating both DNA methylation and histone H3 methylation.
Proc. Natl. Acad. Sci. USA *110*, 1708–1713.

Yang, X., Han, H., De Carvalho, D.D., Lay, F.D., Jones, P.A., and Liang, G. (2014). Gene body
methylation can alter gene expression and is a therapeutic target in cancer. Cancer Cell *26*, 577–
590.

Yelagandula, R., Stroud, H., Holec, S., Zhou, K., Feng, S., Zhong, X., Muthurajan, U.M., Nie,
X., Kawashima, T., Groth, M., et al. (2014). The histone variant H2A.W defines heterochromatin
and promotes chromatin condensation in Arabidopsis. Cell *158*, 98–109.

Zemach, A., and Zilberman, D. (2010). Evolution of eukaryotic DNA methylation and the
pursuit of safer sex. Curr. Biol. *20*, R780–R785.

Zemach, A., McDaniel, I.E., Silva, P., and Zilberman, D. (2010). Genome-wide evolutionary
analysis of eukaryotic DNA methylation. Science *328*, 916–919.

Zemach, A., Kim, M.Y., Hsieh, P.-H., Coleman-Derr, D., Eshed-Williams, L., Thao, K., Harmer,
S.L., and Zilberman, D. (2013). The Arabidopsis nucleosome remodeler DDM1 allows DNA
methyltransferases to access H1-containing heterochromatin. Cell *153*, 193–205.

Zemach, A., Lloyd, J.P.B., Harris, K.D., and Zilberman, D. (2018). Maximum DNA Methylation
Fidelity in the Germline Tolerates Global Non-Functional Gene Body Methylation Dynamics
During Development. SSRN Electron. J.

Zhang, H., Lang, Z., and Zhu, J.-K. (2018a). Dynamics and function of DNA methylation in
plants. Nat. Rev. Mol. Cell Biol. *19*, 489–506.

Zhang, T., Zhang, W., and Jiang, J. (2015). Genome-wide nucleosome occupancy and
positioning and their impact on gene expression and evolution in plants. Plant Physiol. *168*,
1406–1416.

Zhang, Y., Harris, C.J., Liu, Q., Liu, W., Ausin, I., Long, Y., Xiao, L., Feng, L., Chen, X., Xie,
Y., et al. (2018b). Large-scale comparative epigenomics reveals hierarchical regulation of non-
CG methylation in Arabidopsis. Proc. Natl. Acad. Sci. USA E1069–E1074.

Zilberman, D. (2017). An evolutionary case for functional gene body methylation in plants and
animals. Genome Biol. *18*, 87.

Zilberman, D., Gehring, M., Tran, R.K., Ballinger, T., and Henikoff, S. (2007). Genome-wide
analysis of Arabidopsis thaliana DNA methylation uncovers an interdependence between
methylation and transcription. Nat. Genet. *39*, 61–69.

Zilberman, D., Coleman-Derr, D., Ballinger, T., and Henikoff, S. (2008). Histone H2A.Z and
DNA methylation are mutually antagonistic chromatin marks. Nature *456*, 125–129.

 Figure Legends

Figure 1. Histone H1 is enriched in methylated DNA independently of methylation. (A)

Average H1.1 distribution around well-positioned nucleosomes. **(B)** Average H1.1, predicted H1,

and CG methylation (mCG) levels are plotted in 50 kb windows along *Arabidopsis* chromosome

4. **(C)** Box plots of H1.1 levels in heterochromatic TEs (hTEs), euchromatic TEs (eTEs), and genes.

The shaded area marks the middle 50% of genomic H1.1 levels. **(D)** H1.1 distribution around genes

with different expression levels. **(E)** Box plots of H1.1 abundance and predicted H1 abundance in

methylated (m) and unmethylated (u) genes in *wt* and *met1* plants. ** is $p < 0.01$, *** is $p < 0.001$,

**** is $p < 0.0001$, Student's *t*-test. **(F)** Heat maps of indicated chromatin features and expression

in MET1-independent and dependent hTEs. **(G)** Average H1.1 levels of 1 kb windows in *wt* and

met1. 1000 windows were randomly selected for plotting. "*r*" is Pearson's correlation coefficient.

(H) Principal component analysis of H1.1, H1.2, histone H3 lysine modifications, histone variant

H2A.W, DNA methylation (mCG, mCHG, mCHH), GC content (gc) and gene expression (exp).

(I) Linear model prediction of H1 distribution around the transcriptional start sites (TSS) of genes

in comparison to actual H1.1 distribution. **(J)** Example of actual H1.1 and predicted H1 distribution

at gene-body methylated genes in *wt* and *met1*. **(C and E)** Whiskers indicate 1.5X interquartile

range (IQR).

Figure 2. Loss of H1 alters nucleosome organization. (A) Nucleosome repeat length (NRL)

calculations for two biological replicates at genes grouped by their expression in *wt*. Phasograms

and linear regressions for the positions of nucleosome peaks at the genes with lowest expression

(shaded area) are in Figure S2A. **(B and E)** NRL calculations for two biological replicates at genes

(B) and TEs (E) grouped by H1 enrichment. **(C and F)** Phasograms and linear regressions for the

positions of nucleosome peaks at H1-rich genes (C) and TEs (F). **(D)** NRL calculations for two

biological replicates at TEs grouped by their H3K9me2 level in *wt*. Phasograms and linear

regressions for the positions of nucleosome peaks at H3K9me2-rich TEs (shaded area) are in

Figure S2B. **(A-B and D-E)** Error bars indicate standard error.

Figure 3. Loss of H1 activates H1-rich meiosis-related genes. (A) Number of up- and down-

regulated genes in *h1* plants vs. *wt*. **(B)** Heat maps of respective gene expression fold change vs.

wt in *h1* plants in leaves (L) and seedlings (S). Adjusted *p*-values for the expression change are

shown. **(C)** Box plots of *wt* H1 levels in genes upregulated (up), down-regulated (dn) and

unchanged (control; ct) in *h1* plants. "a" and "b" are significantly different ($p < 0.01$; ANOVA).

Whiskers indicate 1.5X IQR. **(D)** Distribution of H1 around control, up- and down-regulated genes.

Shaded areas represent 95% confidence intervals. **(E)** Violin plots of gene expression in *wt* plants

for control (ct), up- and down (dn)-regulated genes in (A). **(F)** An example meiosis-related gene

transcriptionally activated in *h1* plants. The dashed red line indicates average H1 level at genes.

Figure 4. Loss of H1 disperses heterochromatin and weakly activates TEs. (A) Nuclei of *wt*

and *h1* plants stained with DAPI. DAPI puncta volumes are highlighted in yellow. **(B)** Number

per nucleus and volume of DAPI puncta in *wt* and *h1* plants. Note that blurred chromocenter

boundaries in *h1* mutants cause some to be assigned very large volumes. The number of nuclei

(left) and DAPI puncta (right) used to generate the data were indicated. **(C)** Heat maps of

expression in leaves (L) and seedlings (S) with corresponding FDR (*h1* vs. *wt*) and TSS-proximal

DNA methylation change of TEs in *h1* plants vs. *wt*. *p*-value for methylation change was calculated

with Fisher's exact test. **(D)** Examples of upregulated TEs in *h1* plants. Note DNA methylation

loss near TSS (highlighted in yellow). **(E)** An example of upregulated TE in *h1* plants without

DNA methylation loss near TSS (highlighted in yellow). **(F)** Box plots of DNA accessibility of

hTEs, eTEs, and genes in *wt* and *h1*. Whiskers indicate 1.5X IQR.

Figure 5. DNA methylation and H1 cooperatively silence TE expression. (A and E) Expression,

TSS-proximal DNA methylation and DNA accessibility of hTEs in *wt*, *met1* and *h1met1* plants.

TEs with unchanged TSS non-CG methylation in *h1met1* compared to *met1* are shown in (A) and

TEs with altered non-CG methylation at the TSS are shown in (E). **(B and F)** Average CHG and

CHH methylation around TEs in the up1 (B) and up3 (F) clusters. Methylation from the TSS-

proximal regions shaded in yellow is shown in (A) and (E). **(C-D)** Box plots of *met1* H1 abundance

(C) and H1 distribution (D) in TEs in the nc and up1 clusters from (A). H1 from the TSS-proximal

region shaded in yellow in (D) is shown in (C). ******** is $p < 0.0001$, Student's *t*-test. **(G-H)** Box

plots of *met1* H1 abundance (G) and H1 distribution (H) in TEs in the dn and up3 clusters from

(E). H1 from the TSS-proximal region shaded in yellow in (H) is shown in (G). ******* is $p < 0.001$,

Student's *t*-test. **(C and G)** Whiskers indicate 1.5X IQR.

Figure 6. DNA methylation and H1 cooperatively suppress intragenic antisense transcripts.

(A) Number of differentially expressed antisense (AS) transcripts in *h1, met1,* and *h1met1* plants.

(B and D) Heat maps showing *wt* CG methylation at antisense TSS and expression of sense (S)

and antisense transcripts in *met1* (B) or *h1met1* (D) plants. **(C and E)** Box plots of TSS-proximal

wt CG methylation levels of upregulated (up) and downregulated (dn) antisense transcripts either

positively correlated (+) or negatively/un-correlated (0/-) with sense expression in *met1* (C) or

h1met1 (E) plants. Data are also shown for unchanged (control; ct) antisense transcripts. Magenta

asterisks indicate the percentage of antisense TSS that overlap gbM. "a", "b" and "c" are signi-

ficantly different; overlapping letters – i.e. "b" and "bc" – indicate that the difference is not

significant ($p < 0.01$; ANOVA). **(F)** *Wt* CG methylation around antisense TSS of indicated groups

from (E) with more than 100 transcripts. Shaded areas represent 95% confidence intervals. **(G-H)**

CG methylation (G) and H1.1 distribution (H) around the TSS of antisense transcripts initiated

from unmethylated genes, unmethylated regions within gbM genes, and methylated regions within

gbM genes. **(I)** Box plots of antisense expression in indicated groups from (G) in *met1* and *h1met1*

plants. "a", "b" and "c" are significantly different ($p < 0.01$; ANOVA). **(J)** Heat maps of *wt* CG

methylation at antisense TSS, expression of sense and antisense transcripts compared to *wt*, and

corresponding antisense expression *p*-values in *met1* and *h1met1* plants. **(C, E and I)** Whiskers

indicate 1.5X IQR.

Figure 1. Histone H1 is enriched in methylated DNA independently of methylation. (A) Average H1.1 distribution around well-positioned nucleosomes. **(B)** Average H1.1, predicted H1, and CG methylation (mCG) levels are plotted in 50 kb windows along *Arabidopsis* chromosome 4. **(C)** Box plots of H1.1 levels in heterochromatic TEs (hTEs), euchromatic TEs (eTEs), and genes. The shaded area marks the middle 50% of genomic H1.1 levels. **(D)** H1.1 distribution around genes with different expression levels. **(E)** Box plots of H1.1 abundance and predicted H1 abundance in methylated (m) and unmethylated (u) genes in *wt* and *met1* plants. ** is $p < 0.01$, *** is $p < 0.001$, **** is $p < 0.0001$, Student's *t*-test. **(F)** Heat maps of indicated chromatin features and expression in MET1-independent and dependent hTEs. **(G)** Average H1.1 levels of 1 kb windows in *wt* and *met1*. 1000 windows were randomly selected for plotting. "*r*" is Pearson's correlation coefficient. **(H)** Principal component analysis of H1.1, H1.2, histone H3 lysine modifications, histone variant H2A.W, DNA methylation (mCG, mCHG, mCHH), GC content (gc) and gene expression (exp). **(I)** Linear model prediction of H1 distribution around the transcriptional start sites (TSS) of genes in comparison to actual H1.1 distribution. **(J)** Example of actual H1.1 and predicted H1 distribution at gene-body methylated genes in *wt* and *met1*. **(C and E)** Whiskers indicate 1.5X interquartile range (IQR).

Figure 2. Loss of H1 alters nucleosome organization. (A) Nucleosome repeat length (NRL) calculations for two biological replicates at genes grouped by their expression in *wt*. Phasograms and linear regressions for the positions of nucleosome peaks at the genes with lowest expression (shaded area) are in Figure S2A. **(B and E)** NRL calculations for two biological replicates at genes (B) and TEs (E) grouped by H1 enrichment. **(C and F)** Phasograms and linear regressions for the positions of nucleosome peaks at H1-rich genes (C) and TEs (F). **(D)** NRL calculations for two biological replicates at TEs grouped by their H3K9me2 level in *wt*. Phasograms and linear regressions for the positions of nucleosome peaks at H3K9me2-rich TEs (shaded area) are in Figure S2B. **(A-B and D-E)** Error bars indicate standard error.

Figure 3. Loss of H1 activates H1-rich meiosis-related genes. (A) Number of up- and down-regulated genes in *h1* plants vs. *wt*. **(B)** Heat maps of respective gene expression fold change vs. *wt* in *h1* plants in leaves (L) and seedlings (S). Adjusted *p*-values for the expression change are shown. **(C)** Box plots of *wt* H1 levels in genes upregulated (up), down-regulated (dn) and unchanged (control; ct) in *h1* plants. "a" and "b" are significantly different ($p < 0.01$; ANOVA). Whiskers indicate 1.5X IQR. **(D)** Distribution of H1 around control, up- and down-regulated genes. Shaded areas represent 95% confidence intervals. **(E)** Violin plots of gene expression in *wt* plants for control (ct), up- and down (dn)-regulated genes in (A). **(F)** An example meiosis-related gene transcriptionally activated in *h1* plants. The dashed red line indicates average H1 level at genes.

Figure 4. Loss of H1 disperses heterochromatin and weakly activates TEs. (A) Nuclei of *wt* and *h1* plants stained with DAPI. DAPI puncta volumes are highlighted in yellow. **(B)** Number per nucleus and volume of DAPI puncta in *wt* and *h1* plants. Note that blurred chromocenter boundaries in *h1* mutants cause some to be assigned very large volumes. The number of nuclei (left) and DAPI puncta (right) used to generate the data were indicated. **(C)** Heat maps of expression in leaves (L) and seedlings (S) with corresponding FDR (*h1* vs. *wt*) and TSS-proximal DNA methylation change of TEs in *h1* plants vs. *wt*. *p*-value for methylation change was calculated with Fisher's exact test. **(D)** Examples of upregulated TEs in *h1* plants. Note DNA methylation loss near TSS (highlighted in yellow). **(E)** An example of upregulated TE in *h1* plants without DNA methylation loss near TSS (highlighted in yellow). **(F)** Box plots of DNA accessibility of hTEs, eTEs, and genes in *wt* and *h1*. Whiskers indicate 1.5X IQR.

Figure 5. DNA methylation and H1 cooperatively silence TE expression. (A and E) Expression, TSS-proximal DNA methylation and DNA accessibility of hTEs in *wt*, *met1* and *h1met1* plants. TEs with unchanged TSS non-CG methylation in *h1met1* compared to *met1* are shown in (A) and TEs with altered non-CG methylation at the TSS are shown in (E). **(B and F)** Average CHG and CHH methylation around TEs in the up1 (B) and up3 (F) clusters. Methylation from the TSS-proximal regions shaded in yellow is shown in (A) and (E). **(C-D)** Box plots of *met1* H1 abundance (C) and H1 distribution (D) in TEs in the nc and up1 clusters from (A). H1 from the TSS-proximal region shaded in yellow in (D) is shown in (C). **** is $p < 0.0001$, Student's *t*-test. **(G-H)** Box plots of *met1* H1 abundance (G) and H1 distribution (H) in TEs in the dn and up3 clusters from (E). H1 from the TSS-proximal region shaded in yellow in (H) is shown in (G). *** is $p < 0.001$, Student's *t*-test. **(C and G)** Whiskers indicate 1.5X IQR.

Figure 6. DNA methylation and H1 cooperatively suppress intragenic antisense transcripts.
(A) Number of differentially expressed antisense (AS) transcripts in *h1, met1,* and *h1met1* plants.
(B and D) Heat maps showing *wt* CG methylation at antisense TSS and expression of sense (S)
and antisense transcripts in *met1* (B) or *h1met1* (D) plants. **(C and E)** Box plots of TSS-proximal
wt CG methylation levels of upregulated (up) and downregulated (dn) antisense transcripts either
positively correlated (+) or negatively/un-correlated (0/-) with sense expression in *met1* (C) or
h1met1 (E) plants. Data are also shown for unchanged (control; ct) antisense transcripts. Magenta
asterisks indicate the percentage of antisense TSS that overlap gbM. "a", "b" and "c" are signi-
ficantly different; overlapping letters – i.e. "b" and "bc" – indicate that the difference is not
significant ($p < 0.01$; ANOVA). **(F)** *Wt* CG methylation around antisense TSS of indicated groups
from (E) with more than 100 transcripts. Shaded areas represent 95% confidence intervals. **(G-H)**
CG methylation (G) and H1.1 distribution (H) around the TSS of antisense transcripts initiated
from unmethylated genes, unmethylated regions within gbM genes, and methylated regions within
gbM genes. **(I)** Box plots of antisense expression in indicated groups from (G) in *met1* and *h1met1*
plants. "a", "b" and "c" are significantly different ($p < 0.01$; ANOVA). **(J)** Heat maps of *wt* CG
methylation at antisense TSS, expression of sense and antisense transcripts compared to *wt*, and
corresponding antisense expression *p*-values in *met1* and *h1met1* plants. **(C, E and I)** Whiskers
indicate 1.5X IQR.