Title: No consistent effects of humans on animal genetic diversity worldwide

Authors: Katie L. Millette[1,5,6,*], Vincent Fugère[1,2,5,6], Chloé Debyser[1,5,6], Ariel Greiner[3], Frédéric

J. J. Chain[4] and Andrew Gonzalez[1,5].

[1]McGill University, Department of Biology

[2]Université du Québec à Montréal, Département des Sciences Biologiques

[3]University of Toronto, Department of Ecology & Evolutionary Biology

[4]University of Massachusetts at Lowell, Department of Biological Sciences

[5]Quebec Centre for Biodiversity Science

[6]These authors contributed equally.

Author emails: katie.millette@mail.mcgill.ca; vincent.fugere@mail.mcgill.ca;

chloe.debyser@mail.mcgill.ca; ariel.greiner@mail.utoronto.ca; frederic_chain@uml.edu;

andrew.gonzalez@mcgill.ca

Running title: Human effects on genetic diversity

Keywords: intraspecific genetic diversity; land use; human density; spatial scale; cytochrome *c*

oxidase subunit I (COI); Anthropocene; nucleotide diversity

Type of article: Letters

Words in abstract (146); main text (4,988); number of references (73); number of figures (5);

number of tables (1); number of text boxes (0).

*Corresponding author: Katie Millette, 1205 Docteur Penfield, Department of Biology, McGill

University, Montreal QC, H3A 1B1, Canada; phone 1-514-398-1622; fax 1-514-398-5069.

Statement of authorship: All authors contributed to the idea and design of the project. KLM

and FJJC contributed to sequence data collection and analysis. VF led statistical analyses. CD led

spatial analyses. AGr collected taxonomic information. KLM drafted the manuscript. VF and CD

made the figures. AGo supervised the project. All authors contributed to data interpretation and

writing of the manuscript.

Data accessibility statement: All data used in this manuscript are available online in open

access, as described in the Methods section. Analysis code and formatted datasets used in

statistical analyses can be found at https://github.com/VFugere/GenDivProject.

Abstract

Human impacts on genetic diversity are poorly understood yet critical to biodiversity

conservation. We used 175,247 COI sequences collected between 1980 and 2016 to assess the

global effects of land use and human density on the intraspecific genetic diversity of 17,082

species of birds, fishes, insects, and mammals. Human impacts on mtDNA diversity were taxon

and scale-dependent, and were generally weak or non-significant. Spatial analyses identified

weak latitudinal diversity gradients as well as negative effects of human density on insect

diversity, and negative effects of intensive land use on fish diversity. The observed effects were

predominantly associated with species turnover. Time series analyses found nearly an equal

number of positive and negative temporal trends in diversity, resulting in no net monotonic trend

in diversity over this time period. Our analyses reveal critical data and theory gaps and call for

increased efforts to monitor global genetic diversity.

Introduction

Intraspecific genetic diversity, a measure of the genetic variation within populations, is an essential measure of biodiversity (Hughes *et al.* 2008). Genetic diversity reflects past and current evolutionary bottlenecks and indicates a population's potential for adaptation to future stressors (Hewitt 2000; Reed & Frankham 2003; Frankham 2005; Bijlsma & Loeschcke 2012). As such, understanding the drivers of genetic diversity change worldwide is of great interest to ecologists and conservation biologists (Hughes *et al.* 2008; Pereira *et al.* 2013; Mimura *et al.* 2017; Paz-Vinas *et al.* 2018). Human disturbances, acting as an evolutionary force by modifying rates of extinction and colonization (Palumbi 2001; Alberti 2015; Thomas 2015; Schlaepfer *et al.* 2018), may be altering the intraspecific genetic diversity of plants and animals around the world–yet no global assessment of temporal trends in genetic diversity has been conducted to date, nor have human impacts on such trends been quantified.

Theory predicts that human activities can affect intraspecific genetic diversity via demographic and evolutionary mechanisms (Kimura 1968, 1983; King & Jukes 1969). Depending on how human disturbances alter selection, drift, gene flow, and mutation rates, intraspecific genetic diversity may decrease, increase, or remain unchanged over time (DiBattista 2008). For example, habitat fragmentation and harvesting can reduce diversity directly due to high mortality and reduced population sizes, and indirectly due to population isolation and inbreeding (Banks *et al.* 2013). Alternatively, human disturbances can maintain or increase genetic diversity through time, for example by increasing connectivity between populations, magnifying temporal variation in selection, increasing mutation rates (e.g. mutagenic pollutants), or creating environments that favour hybridization and heterozygote advantage (Dubrova *et al.* 1996; Ellegren *et al.* 1997; Bickham *et al.* 2000; Crispo *et al.* 2011).

Trends in intraspecific genetic diversity are expected to be scale-dependent, as are trends in other dimensions of biodiversity like taxonomic, phylogenetic, and functional diversity (McGill *et al.* 2015; Jarzyna & Jetz 2018; Schlaepfer *et al.* 2018; Chase *et al*. 2019). Moreover, human disturbances occurring at different scales may have contrasting effects on genetic diversity. At local scales, high human population density and intensive land use can reduce genetic diversity by lowering habitat quality and population size (Vellend 2004; Coors *et al.* 2009; Helm *et al.* 2009). At regional scales, habitat fragmentation and increased habitat heterogeneity can contribute to genetic differentiation via reduced gene flow (e.g. Ripperger *et*

al. 2013) and diversifying selection (e.g. Merck *et al.* 2003), respectively. However, analyses at

spatial scales larger than the geographic extent of interbreeding populations will inflate diversity

values, particularly when individuals from distant and genetically divergent groups are

aggregated. Thus, if the effects of human activity on diversity are to be detected, they should be

apparent and strongest at small spatial scales.

Much remains to be learned regarding human impacts on intraspecific genetic diversity.

A recent global assessment of mammal and amphibian mtDNA diversity found evidence of

reduced diversity in human-impacted regions (Miraldo *et al.* 2016). Given the conservation-

relevance of this finding, it is paramount to confirm the robustness of this pattern across spatial

scales and taxa while controlling for important confounding variables such as the decay of

genetic similarity with distance (Gratton *et al.* 2017) and phylogenetic differences in mtDNA

mutation rates (Nabholz *et al.* 2009). Moreover, we are only beginning to understand how human

land use is affecting levels of genetic diversity. On a decadal timescale, landscape perturbations

would be expected to affect diversity mostly via their effect on population size (Whitlock 1992).

However, it remains unclear whether net overall trends in genetic diversity, and human impacts

on such trends, exist and can indeed be detected at the global scale.

Here, we evaluate the impacts of human population density and intensive land use on

animal intraspecific genetic diversity worldwide, at four spatial scales while accounting for

effects of geography, time, and spatial distance between sequences. We use location and time-

referenced mitochondrial cytochrome *c* oxidase subunit I (COI) sequences collected between

1980 and 2016 to estimate nucleotide diversity ($\hat{\pi}$), a measure of population genetic diversity, for

four animal classes–birds, inland and coastal bony fishes, insects, and mammals. We test two

main predictions: 1) genetic diversity is lower in animal populations exposed to intensive land

use and high human population density, and this relationship is strongest at small spatial scales;

2) genetic diversity has declined during the 1980-2016 time-period in highly human-impacted

regions.

Materials and methods

Sequence data and human impact estimates

Mitochondrial cytochrome *c* oxidase subunit I (mtDNA COI) sequences for birds

(Aves), fishes (Actinopterygii), insects (Insecta), and mammals (Mammalia) were downloaded

from the National Center for Biotechnology Information (NCBI) 'GenBank' (Benson *et al.*

2013) and from the 'Barcode of Life Data Systems' (BOLD; Ratnasingham & Hebert 2007) in

April 2017. Sequences from GenBank were retrieved with the Entrez Utilities Unix Command

Line, and from BOLD using the application platform interface. Only sequences with documented

geographic coordinates and sampling dates were downloaded. Sequences with ambiguous

taxonomic assignment (e.g. species name containing '.spp', 'var') were not considered, nor were

the few (< 0.1%) sequences collected before 1980. Species and year-specific sequence

alignments were performed using default parameters in MAFFT (Katoh & Standley 2013).

Global human population and land use estimates were obtained from the most recent

version of the 'History Database of the Global Environment' (HYDE 3.2; Klein Goldewijk *et al.*

2017) for years 1980-2016. HYDE 3.2 provides land use and human population density

estimates for all land masses divided into 5 arc-minute grid cells (~85 km^2 at the equator).

Estimates are available for every decade from 1980 to 2000, and for every year after 2000.

HYDE variables included in our analyses were: maximum land area (km^2/grid cell), human

population counts (inhabitants/grid cell), and four categories of intensive land use: cropland,

pasture, converted rangeland, and built-up area (km^2/grid cell). See Klein Goldewijk *et al.* (2017)

for a detailed description of these variables. HYDE datasets were converted to raster data

structures using the 'raster' package (Hijmans 2015) in R version 3.5.0 (R Core Team 2018).

Cropland, pasture, converted rangeland, and built-up areas were divided by grid cell land area to

estimate the proportion of each cell consisting of the respective category. We then computed two

grid cell-specific human impact variables: (1) 'human population density' (inhabitants/km^2), i.e.

human population counts divided by grid cell land area, and (2) 'land use intensity' (ranging

from 0 to 1), i.e. the summed proportions of cropland, pasture, converted rangeland, and built-up

area in a grid cell.

All COI sequences were attributed human population density and land use intensity

values based on the sequence's geographic coordinates and the corresponding HYDE grid cell.

Sequences sampled between 2000 and 2016 were assigned year-specific HYDE values, while

sequences from 1980-89 and 1990-99 were assigned values from the 1980 and 1990 HYDE

maps, respectively. Most sequences (92.3%) had year-specific HYDE values (mean +/- SD lag

time between year of sequence collection and HYDE 3.2 map = 0.4 +/- 1.7 years). Sequences

falling on the border of two grid cells, or in cells composed exclusively of water, were excluded.

138 We included data from aquatic animals in grid cells with some land, reasoning that land use can

139 affect inland and coastal waters and the species therein (Stoms *et al.* 2005; Dudgeon *et al.* 2006).

140 All data were processed using the 'tidyverse' collection of R packages (Wickham & RStudio

141 2017). We quantified the proportion of species, genera, families, and orders represented by our

142 final dataset (see 'Supplementary Methods') and visualized the distribution of COI sequences

143 according to geography, taxonomy, time, human population density, and land use intensity.

145 *Calculating genetic diversity and human impacts at the population scale*

146  To split species-specific sequences into distinct geographic populations, we grouped all

147 sequences from a given species across years and mapped their spatial distribution. A simple

148 agglomerative (bottom-up) hierarchical clustering algorithm was used to group sequences based

149 on spatial proximity following the 'single linkage' clustering rule (Sneath 1957). Sequences are

150 grouped together as long as the maximum geographical distance among all sequences in the

151 group does not exceed an *a priori* maximum distance threshold (x km). For example, with x = 10

152 km, two sequences, A and B, would be grouped in the same population if they are 5 km apart,

153 but a third sequence, C, would be placed in its own population if it is >10 km from both A and

154 B. This clustering method was implemented using the single 'hclust' function, and a distance

155 matrix of 'great circle' geographic distances among sequences with the 'spDists' R function ('sp'

156 package; Pebesma *et al.* 2018). To assess the scale-dependence of our results, we manipulated

157 'x' in the clustering algorithm to generate new populations with minimum spatial distances of

158 10, 100, 1,000, and 10,000 km (henceforth: 'scales'). At large scales, isolated populations may

159 be grouped into a single population, while at small scales, one (true) population may be split into

160 separate populations (see 'Supplementary Methods'). Without species-specific range and

161 dispersal information, altering the clustering distance criterion was the best approach to assess

162 the scale-dependence of results.

163  Each sequence was attributed a unique population ID at each spatial scale (e.g. common

164 redpoll, *Acanthis flammea*, from year 2001, in population '1' at scale = 10 km, with mean

165 geographical location 52.4°N and 4.54°W). Pairwise nucleotide differences were calculated

166 among all sequences in a population with > 50% sequence overlap as in Miraldo *et al.* (2016),

167 using adapted Julia script (https://github.com/mkborregaard/). The mean pairwise nucleotide

168 dissimilarity among sequences was calculated for all populations with > 1 sequence to estimate

nucleotide diversity ($\hat{\pi}$; henceforth: 'diversity'; Nei & Li 1979; Tajima 1993). Diversity was also

calculated using stricter inclusion criteria (> 5 sequences; see sensitivity analysis described

below). Diversity estimates with values 10 standard deviations greater than the mean of all

estimates were discarded. Taxonomic misidentification and/or sequencing error might explain

these outliers, which collectively accounted for < 0.1% of the full dataset. For populations with

multiple years of sequence data, separate $\hat{\pi}$ values were computed for each year to assess

temporal trends. It should be noted that spatial scale affects the number of sequences and

populations in the dataset by affecting both the number of populations into which species-

specific sequences are split, and the probability that a sequence is rejected from analysis for

being alone in its population.

Following data collection and filtration steps, our dataset consisted of 175,247 COI

sequences from 17,082 species of birds (Aves), inland and coastal bony fishes (Actinopterygii),

insects (Insecta), and mammals (Mammalia). These sequences were aggregated into 17,124–

27,588 'populations' depending on the spatial scale (Table 1). For each population and at each

scale, we calculated the geographical centroid of the population (latitude and longitude), the

mean great circle distance among sequences, and the mean land use intensity and human

population density of HYDE grid cells from which the sequences originated. Land use

heterogeneity within populations could not be incorporated in the analyses because it was

strongly collinear with the mean geographic distance among sequences, a well-known predictor

of genetic diversity (Gratton *et al.* 2017).

Spatial analyses of animal genetic diversity

Spatial analyses enabled us to test our first prediction that mtDNA diversity is lowest in

areas of intensive land use and high human population density, and that this relationship is

stronger at small spatial scales. We tested this hypothesis using generalized additive mixed

models (GAMMs), a flexible modelling approach, because we did not necessarily expect linear

relationships between diversity and human impact variables (e.g. Dornelas *et al.* 2014;

Daskalova *et al.* 2019). Models were constructed separately for each animal class and spatial

scale, and fitted with the 'bam' function in the 'mgcv' R package (Wood 2017). All GAMMs

used a Tweedie error structure (with a log link function) to account for the zero-inflated,

positive, and right-skewed distribution of diversity values. We included sequence number as

model weights because nucleotide diversity is likely better estimated with an increasing number

of pairwise sequence comparisons. Weights were calculated by dividing the (log-transformed)

number of sequences contributing to a diversity estimate by the mean (log-transformed) number

of sequences of all diversity estimates from a given taxonomic class and spatial scale. All smooth

terms in GAMMs were thin plate regression splines unless noted. The mgcv syntax of all models

and their ecological question are described in Supplementary Material (Table S1).

Our spatial GAMMs evaluated the effects of human activity on mtDNA diversity while

accounting for spatial and geographical factors known to affect diversity (i.e. spatial distance

among sequences, latitude, longitude). Models included the following smooth terms: mean

geographic (great circle) distance among sequences (log-transformed), latitude and longitude (a

Gaussian process smoothed to account for spatial autocorrelation), absolute latitude, mean

human population density (log-transformed), and mean land use intensity (log-transformed). All

predictors were rescaled from 0 to 1. Random effects included 'family' and 'order' to account for

phylogenetic influences on $\hat{\pi}$, and 'year of sequence collection', which could influence diversity

estimates if, for example, the error rate of sequencing technology has changed over time.

'Species' was not included as a random effect because most species had a single population in

the dataset; a single data point (population with the most sequences) was retained instead to

avoid non-independence.

Spatial GAMMs were used for two purposes. First, following Gratton *et al.* (2017), fitted

values from GAMMs were used to construct global maps of 'smoothed' COI nucleotide

diversity, after removing the confounding effect of mean spatial distance among sequences in a

population (setting this distance to 0 km). Diversity values in 2 x 2° grid cells (a resolution

facilitating visualization) were predicted for all combinations of latitudes and longitudes with

COI data available. Predicted values were obtained from GAMMs fitted at the 1,000 km scale,

which corresponded best to the grain size of the maps. Second, GAMMs were used to visualize

and test the significance of relationships between diversity and human variables (human density

and land use) after removing spatial effects. We used the 'summary.gam' function in mgcv to

obtain F and p values for smooth terms. Predicted relationships between diversity and smooth

terms in the model were estimated while setting values for all other predictors to their median

value in the taxon- and scale-specific dataset. A condensed summary of model predictions from

all 16 GAMMs (4 scales x 4 taxa) is given in the main text. Additional model information can be

found in Figs S2–S5. GAMM validation procedures are described in Supplementary Methods.

We conducted several sensitivity analyses to determine the robustness of our results.

First, we performed a rarefaction analysis, refitting all GAMMs after excluding diversity

estimates based on < 5 sequences. Second, we refitted models without weights and used the log-

transformed number of sequences as a fixed effect/smooth term. Finally, spatial and

anthropogenic effects on diversity might differ when considering broad biogeographical regions.

Diversity in the Northern and Southern Hemispheres may vary due to differences in geological

history (Hewitt 2000, 2004), while dissimilar land uses across biomes (tropical vs. temperate)

may bias human effects on diversity. Thus, we fitted additional GAMMs including 'hemisphere'

or 'biome' as factors. None of these analyses resulted in patterns that deviated strongly from the

described results (see Supplementary Methods, Results, Figs. S7-12).

Lastly, both species turnover and spatial variation in mtDNA diversity within species

may underlie patterns observed in spatial GAMMs. To determine the relative importance of these

two processes, we analyzed within-species changes in diversity in the subset of species with data

from at least five populations spanning a range of latitudes, human densities, and land use

intensities. This analysis focused on the 10 km scale because it provided the most populations

and because geographic influences on diversity are minimal at this scale (see Results). For each

taxon, we fitted a multi-level GAMM with latitude, human density, and land use intensity as

fixed effects, and species as a random effect/smooth, allowing the relationship between diversity

and all predictors to vary among species (Table S1). 'Year of sequence collection', 'family', and

'order' were also included as random effects. Predictor variables were centered within species by

subtracting, for each population and predictor, their species-specific mean values. This group-

mean-centering removes across-species effects when estimating within-species effects of

latitude, human density, and land use intensity on diversity.

Time series analyses

In order to test our second prediction that genetic diversity has declined between 1980

and 2016 in areas of high human impact, we investigated temporal trends in populations that

were sampled repeatedly over time ($\geq$ 4 years of data). This analysis was conducted exclusively

at the 1,000 km scale because smaller scales produced few time series (< 10 series for birds),

whilst at the largest scale (10,000 km), distant individuals could be sampled in different years,

conflating spatial and temporal effects. We chose 4 years as the minimum number of years and

retained all diversity estimates regardless of their number of sequences because stricter

thresholds led to too few time series for reliable analysis. These criteria led to 909 time series

from 873 species.

We estimated a Mann-Kendall trend coefficient (τ) for each time series; statistically-

significant coefficients indicate decreasing ($\tau < 0$) or increasing ($\tau > 0$) monotonic trends. We

also fitted one GAMM per taxon, using a Tweedie error structure and model weights calculated

as the mean-standardized number of sequences included in diversity estimates. Fixed effects in

these models included latitude, longitude, spatial distance among sequences, year of sequence

collection, human population density, land use intensity, and two tensor-product interactions

between year and human density, or year and land use intensity (Table S1). These tensor-product

interactions allow global temporal trends to vary along anthropogenic gradients, enabling us to

test the prediction that diversity is declining in impacted areas. Family and order were included

as random effects, as well as a factor-smooth interaction between year and population, to allow

population trends to deviate from the global average. We could not fit a random effect for

'species' since most species had a single time series, so one series per species (the longest) was

retained to avoid non-independence.

Results

Data distribution

We visualized the distribution of our sequence data across space, taxonomic groups, time,

and anthropogenic gradients. Biases in the spatial distribution of sequences were evident: 70.2%

of sequences originated from North America and Europe (Fig. 1a). The dataset was dominated

(74%) by insect sequences, while within each class, 1–3 speciose orders contributed a large

proportion of sequences (Table 1, Fig. 1b). In all taxonomic classes, COI sequences represented

a small proportion of the global number of species, but this proportion increased at higher

taxonomic levels (families, orders), suggesting a phylogenetically-broad pattern of sampling

(Table 1). Additional information on taxonomic biases can be found in Supplementary Results.

The number of sequences collected over time increased steadily in birds, fishes, and insects from

1980 to 2010 (Fig. 1c), while in mammals, the number reached a peak ~1990 and remained

relatively stable until 2005 (Fig. 1c). All groups demonstrate a recent (~5 year) decline in the

number of sequences, perhaps due to the lag between specimen collection and sequence

archiving (Fig. 1c). Sequences originated from grid cells generally representative of the global

distribution of intensive land use, but from areas with higher human population density than the

global average (Fig. 1d, e).

Spatial analyses of animal genetic diversity

To determine whether mtDNA diversity was lower in areas of high human impact,

particularly at small spatial scales, we constructed spatial GAMMs for all combinations of taxon

and spatial scale. GAMMs revealed a number of spatial influences on genetic diversity. First,

global maps of smoothed COI nucleotide diversity identified a negative latitudinal gradient in

birds, fishes, and mammals, but not insects (Fig. 2). Secondly, the mean geographic distance

among sequences had a strong positive effect on mtDNA diversity in all taxa (Fig. 3; Figs. S2-

S5). However, these spatial effects were scale-dependent: the negative effect of absolute latitude

on diversity varied with scale and differed among taxa, and geographic distance was only

significant at large spatial scales (Fig. 3). After accounting for spatial effects, we found that

human effects on genetic diversity were taxon-dependent and generally weak or non-significant

(Fig. 3; Figs. S2-S5). The two clearest effects were: 1) a negative effect of very high human

population density (> 5,000 inhabitants/km^2, or ~0.75 on the scaled axis) on insect mtDNA

diversity at all scales, and 2) the negative impact of heavy land use (grid cells with > 70%

intensive land use) on fish mtDNA diversity at all but the largest spatial scale (Fig. 3). A non-

monotonic effect of human density on fish genetic diversity was also found at the two largest

spatial scales (Fig. 3).

Rarefaction analysis confirmed the positive effect of geographic distance on genetic

diversity at large spatial scales, as well as the negative effects of latitude on mammal mtDNA

diversity, of human population density on insect diversity, and of land use on fish diversity (Fig.

S7). The latitudinal gradients in birds and fish, and effect of human density on fish genetic

diversity however were not robust to rarefaction (Fig. S7). Removing weights from models had

little influence on our results (Fig. S8), and incorporating biogeographical factors (biomes,

hemispheres) did not reveal clear patterns (Figs. S9-S11), in part due to limited data in the

Southern Hemisphere. Analysis of spatial trends in mtDNA diversity within species with

multiple population data points suggests that significant effects identified in Fig. 3 are driven

primarily by species turnover. After removing species differences in mean latitude, human

density, and land use intensity, these predictors had no effect on within-species variation in

mtDNA diversity (Fig. 4). Thus, species with relatively higher and lower genetic diversity occur

at different locations on latitudinal and anthropogenic gradients, but we find no evidence that

these gradients drive spatial variation in genetic diversity within species.

Time series analyses

To evaluate whether diversity has declined in areas of high human density and land use

intensity, we examined temporal changes in mtDNA diversity within populations sampled

repeatedly. Most time series showed no significant monotonic trend in diversity, regardless of

time series duration (Fig. 5a). In the 2.5% of time series with significant temporal trends, there

was nearly an equal number of time series with positive ($N = 10$) and negative ($N = 12$; Fig. 5a)

trends. When pooling all populations to estimate a global decadal trend, no significant change in

diversity was detected for birds, fishes, and mammals (Fig. 5b). Insects demonstrated a

decreasing trend in mtDNA diversity ($p < 0.01$), driven by a modest reduction in diversity from

2010 to 2016 (Fig. 5b; Fig. S12).

Several significant interactions between year and human density or land use intensity

were found (Figs. 5c, d). This indicates that temporal trends in mtDNA diversity often varied

along the two anthropogenic gradients–although effects were generally stronger for land use than

for human population density (Figs. 5c, d). Interactions were complex and none were entirely

consistent with our prediction of reduced diversity over time in heavily impacted areas. In birds,

diversity was highest in areas of intensive land use but declined over time (Fig. 5d). In fishes, an

increase in diversity was apparent at intermediate human densities (Fig. 5c), while areas with

more intensive land use demonstrated a slightly stronger decline in diversity than areas with less

intensive land use (Fig. 5d). Insect mtDNA diversity had a modest temporal increase in areas of

intensive land use, while mammal diversity increased over time in areas of high human

population densities (Fig. 5c, d).

Discussion

We found that human density and land use intensity had no clear effect on animal
mitochondrial genetic diversity worldwide. In all analyses, the direction and magnitude of human
effects were taxon-dependent and either non-significant or weak, even at small spatial scales. We
find minimal evidence of a net temporal changes in intraspecific genetic diversity. Time series
analyses revealed that the average temporal trend across populations was zero for all taxa but
insects, the latter showing a recent, weak, and uncertain decline in mtDNA diversity. Individual
populations do demonstrate temporal changes in diversity (2.5% of populations in our dataset),
however, with almost an equal proportion of increasing and decreasing trends. Thus, our initial
predictions regarding negative effects of humans on animal mtDNA diversity were mostly
unsupported, and in contrast with previous work (Miraldo *et al.* 2016).

The geographic distance among sequences was by far the strongest predictor of mtDNA
diversity in our global dataset of COI sequences from BOLD and GenBank. This confirms the
necessity to account for distance when measuring genetic diversity (Gratton *et al.* 2017),
especially when sequences are aggregated across large spatial scales. After accounting for the
distance-decay of similarity, latitude still had an influence on the distribution of mtDNA
diversity in birds, fishes, and mammals, although this effect was only robust to rarefaction in
mammals. This result is consistent with previous studies showing elevated genetic diversity in
mammals at low latitudes (Miraldo *et al.* 2016; Gratton *et al.* 2017).

Human impacts on the environment such as urbanization and land use intensification can
have both negative and positive effects on intraspecific variation (DiBattista 2008; Hendry *et al.*
2008; Alberti *et al.* 2017; Fugère & Hendry 2018) and other metrics of biodiversity such as
species richness (Pautasso 2007; Elahi *et al.* 2015; Daskalova *et al.* 2019). It is therefore not
surprising that we find complex and taxon-dependent human effects on mtDNA diversity–that is,
it is overly simplistic to expect an overall trend of lower genetic diversity in impacted areas. We
did find lower insect genetic diversity in areas with high human densities, which is consistent
with recent reports of large declines in insect abundances attributed in part to habitat loss,
agriculture, and urbanization (Hallmann *et al.* 2017; Lister & Garcia 2018). Moreover, we also
found intensive land use had a negative effect on fish genetic diversity, in both spatial and time
series analyses. However, the patterns we report are likely due to species turnover, such that it
remains unclear whether species in impacted areas 'lose' intraspecific diversity when colonizing
these habitats, or whether they have naturally lower diversity than species unable to persist in

human-altered environments. These findings stress the need for species-specific monitoring

programs across environmental and temporal gradients.

While our results do not support the general conclusion that intraspecific mtDNA

diversity is lower in human-dominated areas (Miraldo *et al.* 2016), human effects are currently

difficult to evaluate because of several data limitations. Areas with heterogeneous land use may

harbour higher genetic diversity than homogeneous landscapes; however, we could not

incorporate land use heterogeneity in statistical models due to its strong collinearity with the

mean pairwise distance among sequences. Land use estimates from the HYDE datasets are also

uncertain due to important model assumptions about mechanisms of land use change (Klein

Goldewijk *et al.* 2017). Another limitation is that a large fraction of populations in our dataset

originated from a single 'site' (geographic coordinates). Without spatial variation among

sequences, it is difficult to test the importance of spatial scale and land use on genetic diversity.

At large spatial scales (10,000 km), sequences do vary in geographic location, but to the extent

that isolation-by-distance becomes the main determinant of genetic diversity. Future assessments

of trends in global genetic diversity could incorporate species traits, dispersal potential,

phylogeny, and land use history, all of which could influence relationships between diversity and

anthropogenic drivers (Wiens 1989; Levin 1992; Chave 2013; Schiebelhut & Dawson 2018;

Tucker *et al.* 2018; Binks *et al.* 2019).

We acknowledge that the COI locus does not evolve under neutrality (Pentinsaari *et al.*

2016) and that sequence variation at this mitochondrial locus does not reflect genomic

intraspecific diversity, nor may it necessarily reflect anthropogenic pressures (Bazin *et al.* 2006;

Nabholz *et al.* 2008). More appropriate genetic metrics are known for measuring neutral

intraspecific diversity (e.g. microsatellites, allelic diversity) and are themselves more sensitive to

temporal dynamics in ecological processes; however, there is currently no global database for

these data (Leigh *et al.* 2019). COI is one of few genes with abundant sequences in common

databases (Porter & Hajibabaei 2018) with metadata readily available (e.g. spatial coordinates,

date of collection) for many species (likely because COI is used for species identification).

Despite the wealth of COI data with collection years in the databases, we noted taxonomic gaps

such as the small number of Coleoptera sequences, which as a group account for almost 40% of

insect species. As outlined in Supplementary Results, a few species contributed a

disproportionately large number of sequences. Moreover, the majority of populations in our

dataset were represented by < 10 sequences and mostly from a single year. These data limits

hinder the precision of diversity estimates, constrain time series analyses, and impede our ability

to assess global patterns of genetic diversity relative to human activity.

In conclusion, anthropogenic activity likely has complex, taxon-specific effects on

intraspecific genetic diversity; with the data at hand, one cannot conclude that genetic diversity is

generally lower in human-impacted areas. There is a clear need to establish a global and

systematic monitoring program to gather more data on intraspecific diversity (Mimura *et al.*

2017). More and longer time series of genetic diversity within populations are urgently needed,

as the lack of replication in time is a persistent problem in ecology that is constraining our ability

to assess global patterns of biodiversity change (Gonzalez *et al.* 2016; Estes *et al.* 2018; White

2019). Theory should also be developed to determine the conditions under which significant

human impacts on genetic diversity are to be expected, as the time scale and demographic

consequences of disturbances may not be sufficiently large to affect intraspecific mtDNA

diversity (Nabholz *et al.* 2009). Finally, we urge data collectors to upload metadata such as

collection year and spatial coordinates when depositing sequences in databases–a remarkably

large number of sequences in GenBank do not have a collection year (e.g. 95% of amphibian

sequences; Marques *et al.* 2013; Pope *et al.* 2015). Global monitoring of genetic diversity would

improve our ability to detect change and attribute the causes of worldwide patterns of spatial and

temporal variation in genetic diversity.

Acknowledgements

This work resulted from insightful discussions among past and present Gonzalez lab

members. We are grateful to Andreia Miraldo and Michael Borregaard for assistance with Julia

scripts. We thank Navin Ramankutty, Erle Ellis, and Kees Klein Goldewijk for help accessing

and interpreting the HYDE 3.2 dataset. Simon Joly, Arne Mooers, and Daniel Schoen

contributed insightful comments on earlier versions of the manuscript. Additional comments

from three anonymous reviewers greatly improved this manuscript. The authors acknowledge

support from the Natural Sciences and Engineering Research Council of Canada (NSERC) and

the Quebec Centre for Biodiversity Science (QCBS). AG is supported by the Liber Ero Chair in

Biodiversity Conservation and a Killam Fellowship.

Competing interests

The authors declare no competing interests.

References

Alberti, M. (2015). Eco-evolutionary dynamics in an urbanizing planet. *Trends Ecol. Evol.*, **30**, 114–126.

Alberti, M., Correa, C., Marzluff, J.M., Hendry, A.P., Palkovacs, E.P., Gotanda, K.M., *et al.* (2017). Global urban signatures of phenotypic change in animal and plant populations. *PNAS*, **114**, 8951–8956.

Banks, S.C., Cary, G.J., Smith, A.L., Davies, I.D., Driscoll, D.A., Gill, A.M., *et al.* (2013). How does ecological disturbance influence genetic diversity? *Trends Ecol. Evol.*, **28**, 670–679.

Bazin, E., Glémin, S. & Galtier, N. (2006). Population size does not influence mitochondrial genetic diversity in animals. *Science*, **312**, 570.

Binks, R.M., Gibson, N., Ottewell, K.M., Macdonald, B. & Byrne, M. (2019). Predicting contemporary range-wide genomic variation using climatic, phylogeographic and morphological knowledge in an ancient, unglaciated landscape. *J. Biogeogr.*, **46**, 503–514.

Benson, D.A., Cavanaugh, M., Clark, K., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., *et al.* (2013). GenBank. *Nucleic Acids Res.*, **41**, D36–D42.

Bickham, J.W., Sandhu, S., Hebert, P.D.N., Chikhi, L. & Athwal, R. (2000). Effects of chemical contaminants on genetic diversity in natural populations: implications for biomonitoring and ecotoxicology. *Mutat. Res. Rev. Mutat. Res.*, **463**, 33–51.

Bijlsma, R. & Loeschcke, V. (2012). Genetic erosion impedes adaptive responses to stressful environments. *Evol. Appl.*, **5**, 117–129.

Chase, J.M., McGill, B.J., Thompson, P.L., Antão, L.H., Bates, A.E., Blowes, S.A., *et al*. (2019). Species richness change across spatial scales. *Oikos*, **00**, 1–13.

Chave, J. (2013). The problem of pattern and scale in ecology: what have we learned in 20 years? *Ecol. Lett.*, **16**, 4–16.

Coors, A., Vanoverbeke, J., De Bie, T. & De Meester, L. (2009). Land use, genetic diversity and toxicant tolerance in natural populations of Daphnia magna. *Aquat. Toxicol.*, **95**, 71–79.

Crispo, E., Moore, J.-S., Lee-Yaw, J.A., Gray, S.M. & Haller, B.C. (2011). Broken barriers:
human-induced changes to gene flow and introgression in animals. *BioEssays*, **33**, 508–
518.

Daskalova, G.N., Myers-Smith, I.H., Bjorkman, A.D., Supp, S.R., Blowes, S.A., Magurran A., *et*
al. (2019). Forest loss as a catalyst of population and biodiversity change. *bioRxiv* doi
http://dx.doi.org/10.1101/473645.

DiBattista, J.D. (2008). Patterns of genetic variation in anthropogenically impacted populations.
Conserv. Genet., **9**, 141–156.

Dornelas, M., Gotelli, N.J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., *et al.* (2014).
Assemblage time series reveal biodiversity change but not systematic loss. *Science,* **344,**
296–299.

Dubrova, Y.E., Nesterov, V.N., Krouchinsky, N.G., Ostapenko, V.A., Neumann, R., Neil, D.L.,
et al. (1996). Human minisatellite mutation rate after the Chernobyl accident. *Nature*,
380, 683–686.

Dudgeon, D., Arthington, A.H., Gessner, M.O., Kawabata, Z.-I., Knowler, D.J., Lévêque, C., *et*
al. (2006). Freshwater biodiversity: importance, threats, status and conservation
challenges. *Biol. Rev.*, **81**, 163–182.

Elahi, R., O'Connor, M.I., Byrnes, J.E.K., Dunic, J., Eriksson, B.K., Hensel, M.J.S. & Kearns,
P.J. (2015). Recent trends in local-scale marine biodiversity reflect community structure
and human impacts. *Curr. Biol.*, **25**, 1938–1943.

Ellegren, H., Lindgren, G., Primmer, C.R. & Møller, A.P. (1997). Fitness loss and germline
mutations in barn swallows breeding in Chernobyl. *Nature*, **389**, 593.

Estes, L., Elsen, P.R., Treuer, T., Ahmed, L., Caylor, K., Chang, J., *et al.* (2018). The spatial and
temporal domains of modern ecology. *Nat. Ecol. Evol.*, **2**, 819–826.

Frankham, R. (2005). Genetics and extinction. *Biol. Conserv.*, **126**, 131–140.

Fugère, V. & Hendry, A.P. (2018). Human influences on the strength of phenotypic selection.
PNAS, **115**, 10070–10075.

Gonzalez, A., Cardinale, B.J., Allington, G.R.H., Byrnes, J., Endsley, K.A., Brown, D.G., *et al.*
(2016). Estimating local biodiversity change: a critique of papers claiming no net loss of
local diversity. *Ecology*, **97**, 1949–1960.

Gratton, P., Marta, S., Bocksberger, G., Winter, M., Keil, P., Trucchi, E., *et al.* (2017). Which
latitudinal gradients for genetic diversity? *Trends Ecol. Evol.*, **32**, 724–726.

Hallmann, C.A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., *et al.* (2017).
More than 75 percent decline over 27 years in total flying insect biomass in protected
areas. *PLoS One*, **12**, e0185809.

Helm, A., Oja, T., Saar, L., Takkis, K., Talve, T. & Pärtel, M. (2009). Human influence lowers
plant genetic diversity in communities with extinction debt. *J. Ecol.*, **97**, 1329–1336.

Hendry, A.P., Farrugia, T.J. & Kinnison, M.T. (2008). Human influences on rates of phenotypic
change in wild animal populations. *Mol. Ecol.*, **17**, 20–29.

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, **405**, 907–913.

Hewitt, G.M. (2004). Genetic consequences of climatic oscillations in the Quaternary. *Philos.*
Trans. Royal Soc. B, **359**, 183–195.

Hijmans, R.J. (2015). Raster: geographic data analysis and modeling. R package raster version
2.8-19. Available at: https://CRAN.R-project.org/package=raster.

Hughes, A.R., Inouye, B.D., Johnson, M.T.J., Underwood, N. & Vellend, M. (2008). Ecological
consequences of genetic diversity. *Ecol. Lett.*, **11**, 609–623.

Jarzyna, M.A. & Jetz, W. (2018). Taxonomic and functional diversity change is scale dependent.
Nat. Commun., **9**, 2565.

Katoh, K. & Standley, D.M. (2013). MAFFT Multiple Sequence Alignment Software version 7:
improvements in performance and usability. *Mol. Biol. Evol.*, **30**, 772–780.

Kimura, M. (1968). Evolutionary rate at the molecular level. *Nature*, **217**, 624–626.

Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. New York, Cambridge
University Press.

King, J. & Jukes, T. (1969). Non-Darwinian evolution. *Science*, **164**, 788–798.

Klein Goldewijk, K., Beusen, A., Doelman, J. & Stehfest, E. (2017). Anthropogenic land use
estimates for the Holocene – HYDE 3.2. *Earth Syst. Sci. Data*, **9**, 927–953.

Leigh, D.M., Hendry, A.P., Vázquez-Domínguez, E. & Friesen V.L. (2019). Estimated six per
cent loss of genetic variation in wild populations since the industrial revolution. *Evol.*
Appl., **10.1111/eva.12810**.

Levin, S.A. (1992). The problem of pattern and scale in ecology. *Ecology*, **73**, 1943–1967.

Lister, B.C. & Garcia, A. (2018). Climate-driven declines in arthropod abundance restructure a
rainforest food web. *PNAS*, **115**, E10397–E10406.

Marques, A.C., Maronna, M.M. & Collins, A.G. (2013). Putting GenBank data on the map.
Science, **341**, 1341–1341.

McGill, B.J., Dornelas, M., Gotelli, N.J. & Magurran, A.E. (2015). Fifteen forms of biodiversity
trend in the Anthropocene. *Trends Ecol. Evol.*, **30**, 104–113.

Merck, T., Van Dyck, H., Karlsson, B. & Leimar, O. (2003). The evolution of movements and
behaviour at boundaries in different landscapes: A common arena experiment with
butterflies. *Proc. Biol. Sci.*, **270**, 1815–1821.

Mimura, M., Yahara, T., Faith, D.P., Vázquez-Domínguez, E., Colautti, R.I., Araki, H., *et al.*
(2017). Understanding and monitoring the consequences of human impacts on
intraspecific variation. *Evol. Appl.*, **10**, 121–139.

Miraldo, A., Li, S., Borregaard, M.K., Flórez-Rodríguez, A., Gopalakrishnan, S., Rizvanovic,
M., *et al.* (2016). An Anthropocene map of genetic diversity. *Science*, **353**, 1532–1535.

Nabholz, B., Glémin, S. & Galtier, N. (2009). The erratic mitochondrial clock: variations of
mutation rate, not population size, affect mtDNA diversity across birds and mammals.
BMC Evol. Biol., **9**, 54.

Nabholz, B., Mauffrey, J.-F., Bazin, E., Galtier, N. & Glemin, S. (2008). Determination of
mitochondrial genetic diversity in mammals. *Genetics*, **178**, 351–361.

Nei, M. & Li, W.H. (1979). Mathematical model for studying genetic variation in terms of
restriction endonucleases. *Proc. Natl. Acad. Sci., USA*, **76**, 5269–5273.

Palumbi, S.R. (2001). Humans as the world's greatest evolutionary force. *Science*, **293**, 1786–
1790.

Pautasso, M. (2007). Scale dependence of the correlation between human population presence
and vertebrate and plant species richness. *Ecol. Lett.*, **10**, 16–24.

Paz-Vinas I., Loot, G., Hermoso, V., Veyssière, C., Poulet, N., Grenouillet, G., *et al.* (2018).
Systematic conservation planning for intraspecific genetic diversity. *Proc. R. Soc. B*, **285**,
20172746.

Pebesma, E., Bivand, R. S., Rowlingson, B., Gomez-Rubio, V., Hijmans, R., Sumner, M., *et al.*
(2018). sp: classes and methods for spatial data in R. *R News 5, ***2***.*

Pentinsaari, M., Salmela, H., Mutanen, M. & Roslin, T. (2016). Molecular evolution of a widely-
adopted taxonomic marker (COI) across the animal tree of life. *Sci. Rep.*, **6**, 35275.

Pereira, H.M., Ferrier, S., Walters, M., Geller, G.N., Jongman, R.H.G., Scholes, R.J., *et al.*
(2013). Essential biodiversity variables. *Science*, **339**, 277–278.

Pope, L.C., Liggins, L., Keyse, J., Carvalho, S.B. & Riginos, C. (2015). Not the time or the
place: the missing spatio-temporal link in publicly available genetic data. *Mol. Ecol.*, **24**,
3802–3809.

Porter, T.M. & Hajibabaei, M. (2018). Over 2.5 million COI sequences in GenBank and
growing. *PLoS One*, **13**, e0200177.

R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation
for Statistical Computing, Vienna, Austria. Available at: http://www.R-project.org.

Ratnasingham, S. & Hebert, P.D.N. (2007). BOLD: The Barcode of Life Data System
(http://www.barcodinglife.org). *Mol. Ecol. Notes*, **7**, 355–364.

Reed, D.H. & Frankham, R. (2003). Correlation between fitness and genetic diversity. *Conserv.*
Biol., **17**, 230–237.

Ripperger, S.P., Tschapka, M., Kalko, E.K.V., Rodriguez-Herrera, B. & Mayer, F. (2013). Life
in a mosaic landscape: Anthropogenic habitat fragmentation affects genetic population
structure in a frugivorous bat species. *Conserv. Genet.*, **14**, 925–934.

Schiebelhut, L.M. & Dawson, M.N. (2018). Correlates of population genetic differentiation in
marine and terrestrial environments. *J. Biogeogr.*, **45**, 2427–2441.

Schlaepfer, D.R., Braschler, B., Rusterholz, H.-P. & Baur, B. (2018). Genetic effects of
anthropogenic habitat fragmentation on remnant animal and plant populations: a meta-
analysis. *Ecosphere*, **9**, e02488.

Sneath, P.H. (1957). The application of computers to taxonomy. *J. Gen. Microbiol.*, **17**, 201–
226.

Stoms, D.M., Davis, F.W., Andelman, S.J., Carr, M.H., Gaines, S.D., Halpern, B.S., *et al.*
(2005). Integrated coastal reserve planning: making the land–sea connection. *Front. Ecol.*
Environ., **3**, 429–436.

Tajima, F. (1993). Statistical analysis of DNA polymorphism. *Jpn. J. Genet.*, **68**, 567–595.

Thomas, C.D. (2015). Rapid acceleration of plant speciation during the Anthropocene. *Trends*
Ecol. Evol., **30**, 448–455.

Tucker, M.A., Böhning-Gaese, K., Fagan, W.F., Fryxell, J.M., Moorter, B.V., Alberts, S.C., *et*

al. (2018). Moving in the Anthropocene: global reductions in terrestrial mammalian

movements. *Science*, **359**, 466–469.

Vellend, M. (2004). Parallel effects of land-use history on species diversity and genetic diversity

of forest herbs. *Ecology*, **85**, 3043–3055.

White, E.R. (2019). Minimum time required to detect population trends: the need for long-term

monitoring programs. *BioScience*, **69**, 40–46.

Whitlock, M.C. (1992). Temporal fluctuations in demographic parameters and the genetic

variance among populations. *Evolution*, **46**, 608–615.

Wickham, H. & RStudio. (2017). tidyverse: easily install and load the "tidyverse". R package

version 1.2.1. Available at: https://CRAN.R-project.org/package=tidyverse.

Wiens, J.A. (1989). Spatial scaling in ecology. *Funct. Ecol.*, **3**, 385–397.

Wood, S.N. (2017). *Generalized Additive Models: An Introduction with R*. 2nd edn. Chapman

and Hall/CRC.

Table 1. The number of COI sequences, populations, species, genera, families, and orders

included in the dataset of each animal class at each spatial scale.

scale	class	sequences	populations	species	genera	families	orders
10	Aves	6155	2026	1294 (12.50%)	677 (30.30%)	120 (52.86%)	31 (77.50%)
10	Actinopterygii	18057	4492	2297 (7.06%)	1022 (20.82%)	240 (49.18%)	43 (93.48%)
10	Insecta	105736	18904	9714 (1.05%)	4434 (5.53%)	430 (38.05%)	24 (85.71%)
10	Mammalia	15632	2166	531 (9.07%)	219 (17.03%)	42 (26.58%)	10 (34.48%)
100	Aves	6834	2174	1413 (13.64%)	711 (31.83%)	120 (52.86%)	31 (77.50%)
100	Actinopterygii	19194	3973	2366 (7.28%)	1044 (21.27%)	245 (50.20%)	44 (95.65%)
100	Insecta	114674	17677	10740 (1.16%)	4730 (5.90%)	442 (39.12%)	24 (85.71%)
100	Mammalia	16248	1517	547 (9.35%)	224 (17.42%)	43 (27.22%)	11 (37.93%)
1,000	Aves	7931	1977	1561 (15.07%)	746 (33.39%)	121 (53.30%)	31 (77.50%)
1,000	Actinopterygii	19919	2884	2401 (7.38%)	1054 (21.48%)	245 (50.20%)	44 (95.65%)
1,000	Insecta	125169	14591	12006 (1.29%)	5125 (6.40%)	452 (40.00%)	25 (89.29%)
1,000	Mammalia	16673	743	564 (9.64%)	228 (17.73%)	43 (27.22%)	11 (37.93%)
10,000	Aves	8597	1614	1607 (15.52%)	760 (34.02%)	123 (54.19%)	31 (77.50%)
10,000	Actinopterygii	20126	2427	2412 (7.42%)	1056 (21.52%)	246 (50.41%)	44 (95.65%)
10,000	Insecta	129705	12515	12496 (1.35%)	5261 (6.57%)	456 (40.35%)	25 (89.29%)
10,000	Mammalia	16819	568	567 (9.69%)	228 (17.73%)	43 (27.22%)	11 (37.93%)

Scale indicates maximum geographic distance (km) allowed between any two sequences when

defining populations. Percentages in parentheses for species, genera, families and orders are

estimates of the proportion of taxa present in the database relative to the total number of taxa

globally, as described in Supplementary Methods. Note that the percentage for Actinopterygii

orders is an overestimate caused by large discrepancies in order number in different taxonomic

databases.

Figure captions

Figure 1. Sequence distribution across geographical space (a), taxonomic classes (b), time (c), and anthropogenic parameters (d, e). COI sequences depicted are those available in NCBI or BOLD with geographic coordinates and year of collection documented in the databases after data filtering steps. (a) Global distribution of sequences. Bar height represents the total number of sequences from the four animal classes per HYDE 3.2 5' grid cell (min=1 sequence, max=3,258). (b) Distribution of sequences across the four taxonomic classes: birds (N=8,866 sequences), fish (N=20,388), insects (N=130,433), and mammals (N=16,890). Orders contributing ≥10% of a taxon's sequences are indicated. (c) Number of sequences in the dataset for each year and animal class. (d, e) Distribution of sequences (grey) according to human density (d) and land use intensity (e) relative to the frequency of these parameter values worldwide (blue).

Figure 2. Spatial variation in COI nucleotide diversity of birds, inland and coastal bony fishes, insects, and mammals. Diversity values are GAMM predictions when assuming that all sequences from a given population were collected at the same location, thus removing the confounding effect of population spatial extent on nucleotide diversity. Note that diversity scales differ among animal classes.

Figure 3. Relationship between COI nucleotide diversity and the mean spatial distance among sequences in a population, the absolute latitude of the population centroid, and the mean human population density, and land use intensity of HYDE 3.2 grid cells from which sequences originate. Panels on the same row belong to the same taxon. Lines represent fitted values from 16 scale and taxon-specific GAMMs (4 scales × 4 classes). Line colour indicates spatial scale; for each taxon, lines with the same colour belong to the same model. Line width indicates effect size (F value of predictor) standardized within taxa; thicker lines are stronger effects. Line type indicates statistical significance (solid line, $p < 0.01$; dashed line, $p > 0.01$). Predictor variables were scaled from 0 to 1. Predictions for one variable were made while setting the value of all other variables to their median value in the taxon and scale-specific dataset. Additional model details are provided in Figs. S2-S5.

Figure 4. GAMM results showing predicted variation in COI nucleotide diversity within species

along gradients of latitude, human population density, and land use intensity. Species included in

this analysis had data from ≥ 5 populations varying in their mean values of all three predictor

variables at the 10 km spatial scale. Predictor variables were centered using species-specific

mean values to 'align' all species trends along x axes, thus removing species differences in the

mean latitude, human population density, and land use intensity at which they occur. Panels on

the same row belong to the same taxon. Thick purple lines indicate overall trends (predicted

value ± 95% confidence intervals) and thin grey lines show fitted values for individual species. *F*

and *p* values for individual smooths and the overall model adjusted R^2 are shown.

Figure 5. Time series analysis of COI nucleotide diversity in 909 populations of birds, inland

and coastal bony fishes, insects, and mammals. Populations were defined at the 1,000 km scale

and included a minimum of 4 years of data. Panels on the same row belong to the same taxon. (a)

Mann-Kendall trend coefficients (τ) of time series vs. the number of years of data that they

contain. Purple symbols are time series with statistically-significant negative (τ < 0) or positive

(τ > 0) trends in diversity. A small offset was added on the y axis to aid visualization. (b-d)

GAMM results showing predicted values of diversity over time. Values are on the scale of the

linear predictor. (b) Effect of year on diversity. Thick purple lines indicate overall trends

(predicted value ± 95% confidence intervals) while thin grey lines show fitted values for

individual populations. *F* and *p* values for the 'time' smooth and the overall model adjusted R^2

are shown. (c, d) Predicted temporal trends in diversity as a function of mean human population

density (c) and land use intensity (d) in the HYDE 3.2 grid cells from which population-specific

sequences originate. A colour transition from yellow to purple indicates a reduction in nucleotide

diversity; for example, in insects (third row, column c), nucleotide diversity in areas of high

human population density transitions from high diversity (yellow) to low diversity (purple) over

time. The *F* and *p* values of both tensor-product interactions are provided; low *p* values indicate

that a given two-dimensional surface is not flat. Note that the scales for diversity and year differ

among animal classes.

log$_e$ COI diversity ($\hat{\pi}$)

mean spatial distance   absolute latitude   human density   land use intensity

10 km
100 km
1,000 km
10,000 km